World Journal of *Gastroenterology*

World J Gastroenterol 2024 January 28; 30(4): 286-423





Published by Baishideng Publishing Group Inc

WUG

World Journal of Gastroenterology

Contents

Weekly Volume 30 Number 4 January 28, 2024

EDITORIAL

286 Revolutionizing gastric cancer treatment: The potential of immunotherapy Christodoulidis G, Koumarelas KE, Kouliou MN

REVIEW

290 Portal hypertension in patients with nonalcoholic fatty liver disease: Current knowledge and challenges

Madir A, Grgurevic I, Tsochatzis EA, Pinzani M

ORIGINAL ARTICLE

Retrospective Study

308 Value of multiple models of diffusion-weighted imaging to predict hepatic lymph node metastases in colorectal liver metastases patients

Zhu HB, Zhao B, Li XT, Zhang XY, Yao Q, Sun YS

318 Hepatic arterial infusion chemotherapy with anti-angiogenesis agents and immune checkpoint inhibitors for unresectable hepatocellular carcinoma and meta-analysis

Cao YZ, Zheng GL, Zhang TQ, Shao HY, Pan JY, Huang ZL, Zuo MX

Observational Study

332 Circulating microRNA expression and nonalcoholic fatty liver disease in adolescents with severe obesity

Li YJ, Baumert BO, Stratakis N, Goodrich JA, Wu HT, He JX, Zhao YQ, Aung MT, Wang HX, Eckel SP, Walker DI, Valvi D, La Merrill MA, Ryder JR, Inge TH, Jenkins T, Sisley S, Kohli R, Xanthakos SA, Baccarelli AA, McConnell R, Conti DV, Chatzi L

Gastrointestinal manifestations of critical ill heatstroke patients and their associations with outcomes: A 346 multicentre, retrospective, observational study

Wang YC, Jin XY, Lei Z, Liu XJ, Liu Y, Zhang BG, Gong J, Wang LT, Shi LY, Wan DY, Fu X, Wang LP, Ma AJ, Cheng YS, Yang J, He M, Jin XD, Kang Y, Wang B, Zhang ZW, Wu Q

Basic Study

Amlodipine inhibits the proliferation and migration of esophageal carcinoma cells through the induction 367 of endoplasmic reticulum stress

Chen YM, Yang WQ, Gu CW, Fan YY, Liu YZ, Zhao BS

SYSTEMATIC REVIEWS

Current status of magnetic resonance imaging radiomics in hepatocellular carcinoma: A quantitative 381 review with Radiomics Quality Score

Brancato V, Cerrone M, Garbino N, Salvatore M, Cavaliere C



Contents

World Journal of Gastroenterology

Weekly Volume 30 Number 4 January 28, 2024

LETTER TO THE EDITOR

- Antiviral treatment standards for hepatitis B: An urgent need for expansion 418 Bao ZH, Dai ZK, Tang HX
- 421 Leukocyte immunoglobulin-like receptor B2: A promising biomarker for colorectal cancer Zhao WZ, Wang HG, Yang XZ



Contents

Weekly Volume 30 Number 4 January 28, 2024

ABOUT COVER

Editorial Board Member of World Journal of Gastroenterology, Giuseppe Sica, MD, PhD, Professor of Surgery, Department of Surgical Science, Tor Vergata University Hospital, Rome 00133, Italy. giuseppe.sica@ptvonline.it

AIMS AND SCOPE

The primary aim of World Journal of Gastroenterology (WJG, World J Gastroenterol) is to provide scholars and readers from various fields of gastroenterology and hepatology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online. WJG mainly publishes articles reporting research results and findings obtained in the field of gastroenterology and hepatology and covering a wide range of topics including gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, gastrointestinal oncology, and pediatric gastroenterology.

INDEXING/ABSTRACTING

The WJG is now abstracted and indexed in Science Citation Index Expanded (SCIE), MEDLINE, PubMed, PubMed Central, Scopus, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 edition of Journal Citation Reports[®] cites the 2022 impact factor (IF) for WJG as 4.3; Quartile category: Q2. The WJG's CiteScore for 2021 is 8.3.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Yu-Xi Chen; Production Department Director: Xiang Li; Editorial Office Director: Jia-Ru Fan.

NAME OF JOURNAL	INSTRUCTIONS TO AUTHORS
World Journal of Gastroenterology	https://www.wjgnet.com/bpg/gerinfo/204
ISSN	GUIDELINES FOR ETHICS DOCUMENTS
ISSN 1007-9327 (print) ISSN 2219-2840 (online)	https://www.wjgnet.com/bpg/GerInfo/287
LAUNCH DATE	GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH
October 1, 1995	https://www.wjgnet.com/bpg/gerinfo/240
FREQUENCY	PUBLICATION ETHICS
Weekly	https://www.wjgnet.com/bpg/GerInfo/288
EDITORS-IN-CHIEF	PUBLICATION MISCONDUCT
Andrzej S Tarnawski	https://www.wjgnet.com/bpg/gerinfo/208
EXECUTIVE ASSOCIATE EDITORS-IN-CHIEF	POLICY OF CO-AUTHORS
Xian-Jun Yu (Pancreatic Oncology), Jian-Gao Fan (Chronic Liver Disease), Hou- Bao Liu (Biliary Tract Disease)	https://www.wjgnet.com/bpg/GerInfo/310
EDITORIAL BOARD MEMBERS	ARTICLE PROCESSING CHARGE
http://www.wjgnet.com/1007-9327/editorialboard.htm	https://www.wjgnet.com/bpg/gerinfo/242
PUBLICATION DATE	STEPS FOR SUBMITTING MANUSCRIPTS
January 28, 2024	https://www.wjgnet.com/bpg/GerInfo/239
COPYRIGHT	ONLINE SUBMISSION
© 2024 Baishideng Publishing Group Inc	https://www.f6publishing.com
PUBLISHING PARTNER	PUBLISHING PARTNER'S OFFICIAL WEBSITE
Shanghai Pancreatic Cancer Institute and Pancreatic Cancer Institute, Fudan University Biliary Tract Disease Institute, Fudan University	https://www.shca.org.cn https://www.zs-hospital.sh.cn

© 2024 Baishideng Publishing Group Inc. All rights reserved. 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA E-mail: office@baishideng.com https://www.wjgnet.com



WJG

World Journal of Gastroenterology

Submit a Manuscript: https://www.f6publishing.com

World J Gastroenterol 2024 January 28; 30(4): 318-331

DOI: 10.3748/wjg.v30.i4.318

Retrospective Study

ISSN 1007-9327 (print) ISSN 2219-2840 (online)

ORIGINAL ARTICLE

Hepatic arterial infusion chemotherapy with anti-angiogenesis agents and immune checkpoint inhibitors for unresectable hepatocellular carcinoma and meta-analysis

Yu-Zhe Cao, Guang-Lei Zheng, Tian-Qi Zhang, Hong-Yan Shao, Jia-Yu Pan, Zi-Lin Huang, Meng-Xuan Zuo

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Unsolicited article; Externally peer reviewed

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): A Grade B (Very good): 0 Grade C (Good): 0 Grade D (Fair): 0 Grade E (Poor): 0

P-Reviewer: Saadi S, United States

Received: November 1, 2023 Peer-review started: November 1, 2023 First decision: December 4, 2023 Revised: December 11, 2023 Accepted: January 8, 2024 Article in press: January 8, 2024 Published online: January 28, 2024



Yu-Zhe Cao, Guang-Lei Zheng, Tian-Qi Zhang, Hong-Yan Shao, Jia-Yu Pan, Zi-Lin Huang, Meng-Xuan Zuo, Department of Minimally Invasive Interventional Therapy, Sun Yat-sen University Cancer Center, Guangzhou 510060, Guangdong Province, China

Yu-Zhe Cao, Guang-Lei Zheng, Tian-Qi Zhang, Hong-Yan Shao, Jia-Yu Pan, Zi-Lin Huang, Meng-Xuan Zuo, Department of Minimally Invasive Interventional Therapy, State Key Laboratory of Oncology in South China, Guangzhou 510060, Guangdong Province, China

Yu-Zhe Cao, Guang-Lei Zheng, Tian-Qi Zhang, Hong-Yan Shao, Jia-Yu Pan, Zi-Lin Huang, Meng-Xuan Zuo, Department of Minimally Invasive Interventional Therapy, Guangdong Provincial Clinical Research Center for Cancer, Guangzhou 510060, Guangdong Province, China

Corresponding author: Meng-Xuan Zuo, MD, Attending Doctor, Department of Minimally Invasive Interventional Therapy, Sun Yat-sen University Cancer Center, No. 651 Dongfeng Road East, Yuexiu District, Guangzhou 510060, Guangdong Province, China. zuomx@sysucc.org.cn

Abstract

BACKGROUND

Hepatic arterial infusion chemotherapy (HAIC) has been proven to be an ideal choice for treating unresectable hepatocellular carcinoma (uHCC). HAIC-based treatment showed great potential for treating uHCC. However, large-scale studies on HAIC-based treatments and meta-analyses of first-line treatments for uHCC are lacking.

AIM

To investigate better first-line treatment options for uHCC and to assess the safety and efficacy of HAIC combined with angiogenesis inhibitors, programmed cell death of protein 1 (PD-1) and its ligand (PD-L1) blockers (triple therapy) under real-world conditions.

METHODS

Several electronic databases were searched to identify eligible randomized controlled trials for this meta-analysis. Study-level pooled analyses of hazard ratios (HRs) and odds ratios (ORs) were performed. This was a retrospective single-center study involving 442 patients with uHCC who received triple therapy



or angiogenesis inhibitors plus PD-1/PD-L1 blockades (AIPB) at Sun Yat-sen University Cancer Center from January 2018 to April 2023. Propensity score matching (PSM) was performed to balance the bias between the groups. The Kaplan-Meier method and cox regression were used to analyse the survival data, and the log-rank test was used to compare the survival time between the groups.

RESULTS

A total of 13 randomized controlled trials were included. HAIC alone and in combination with sorafenib were found to be effective treatments (*P* values for ORs: HAIC, 0.95; for HRs: HAIC + sorafenib, 0.04). After PSM, 176 HCC patients were included in the analysis. The triple therapy group (n = 88) had a longer median overall survival than the AIPB group (n = 88) (31.6 months *vs* 14.6 months, P < 0.001) and a greater incidence of adverse events (94.3% *vs* 75.4%, P < 0.001).

CONCLUSION

This meta-analysis suggests that HAIC-based treatments are likely to be the best choice for uHCC. Our findings confirm that triple therapy is more effective for uHCC patients than AIPB.

Key Words: Unresectable hepatocellular carcinoma; Hepatic arterial infusion chemotherapy; Angiogenesis inhibitors; Programmed cell death protein 1; Programmed death ligand 1

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: The network meta-analysis showed the treatment based on hepatic arterial infusion chemotherapy (HAIC) had the best efficacy on unresectable hepatocellular carcinoma (uHCC). The retrospective, relatively large-scale study suggested HAIC combined with angiogenesis inhibitors and programmed cell death protein 1 (PD-1)/programmed death ligand 1 (PD-L1) blockers could improve the uHCC patients' prognosis. After propensity score matching, it demonstrated that triple therapy was able to prolong the uHCC patients' survival than angiogenesis inhibitors and PD-1/PD-L1 blockers.

Citation: Cao YZ, Zheng GL, Zhang TQ, Shao HY, Pan JY, Huang ZL, Zuo MX. Hepatic arterial infusion chemotherapy with antiangiogenesis agents and immune checkpoint inhibitors for unresectable hepatocellular carcinoma and meta-analysis. *World J Gastroenterol* 2024; 30(4): 318-331

URL: https://www.wjgnet.com/1007-9327/full/v30/i4/318.htm **DOI:** https://dx.doi.org/10.3748/wjg.v30.i4.318

INTRODUCTION

Primary liver cancer is a malignant tumor of the digestive system that is common worldwide. China has a particularly high incidence of liver cancer, with approximately 410000 new cases and 391000 deaths annually; liver cancer is the second largest cause of cancer-related death in the country[1]. Among primary liver cancers, hepatocellular carcinoma (HCC) is the main pathological type. In the subclinical phase, patients tend to be asymptomatic. Therefore, at the time of diagnosis, most patients have already reached advanced stages of the disease. Therefore, fewer than 30% of patients are candidates for surgical resection[2,3].

According to several clinical trials, sorafenib and lenvatinib have been recommended by multiple authoritative guidelines as first-line treatment options for advanced HCC for some time[4-6]. The results of the IMbrave150 trial ushered in a new era in HCC therapy, in which angiogenesis inhibitors were combined with programmed cell death protein 1/programmed cell death ligand 1 (PD-1/PD-L1) blockees, which is now becoming the new standard first-line therapy[7]. Researchers have conducted several clinical studies on various immune-related drugs, angiogenesis inhibitors, and various combination regimens for unresectable HCC (uHCC). The FOHAIC study also revealed that hepatic arterial perfusion chemotherapy (HAIC) using the FOLFOX regimen was more effective than sorafenib in patients with uHCC[8]. Several studies have suggested that triple therapy has the potential to further improve the prognosis in patients with uHCC[9-12]. As the results of multiple clinical studies have been published, several questions remain surrounding this type of therapy, such as which treatment approach has the best therapeutic effect on uHCC? How effective is triple therapy when used in large-scale real-world clinical applications?

In this study, we attempted to identify the optimal treatment for uHCC based on data from phase III randomized controlled trials (RCTs) through a network meta-analysis. We also investigated the safety and efficacy of triple therapy in patients with HCC from a Chinese population under real-world conditions. We then performed propensity score matching (PSM) to compare triple therapy to angiogenesis inhibitors plus PD1/PDL1 blockers (AIPB), which has been recommended as a first-line treatment for uHCC by some guidelines[13,14]. This study also confirmed the safety of triple therapy under real-world conditions.

Zaishidena® WJG | https://www.wjgnet.com

MATERIALS AND METHODS

Literature search, data extraction, and network meta-analysis

We performed an extensive literature search of the PubMed, Embase, and Cochrane Library databases for RCTs published from January 1, 2018, to January 1, 2023. The Supplementary material details the search strategy and inclusion criteria. Two authors independently screened the trials for eligibility and extracted information from each one. The included RCTs were then assessed for risk of bias using the Cochrane risk of bias 2 tool, which showed low risk levels for all the included studies (Supplementary Figure 1).

Triple therapy real-world study

Patients who were treated with triple therapy or AIPB as a first-line treatment for advanced HCC between January 2018 and April 2023 at the Department of Minimally Invasive Interventional Therapy, Sun Yat-sen University Cancer Center in Guangzhou, China, were screened for eligibility. HCC was diagnosed histologically or radiologically in accordance with the latest international guidelines[15]. The inclusion criteria were as follows: (1) Stage B (not applicable for surgery or progressed on locoregional therapy) or stage C HCC according to the Barcelona Clinic Liver Cancer (BCLC) staging system; (2) Child-Pugh score of A-C; (3) Eastern cooperative oncology group performance status (ECOG PS) of 0-2; (4) Age \geq 18 years; and (5) At least one available follow-up data point. The exclusion criteria were as follows: (1) History of receiving any systemic chemotherapy, angiogenesis inhibitors, or immunotherapy; (2) Lack of medical imaging data; and (3) History of a second primary malignant tumor. The details are shown in the Supplementary materials. This study was reviewed and approved by the Sun Yat-sen University Cancer Center Ethics Committee. Informed consent was waived due to the retrospective nature of the analysis.

Treatment regimens: Small-molecule tyrosine kinase inhibitors such as sorafenib, a type of angiogenesis inhibitor, were administered orally, and the doses were determined based on the manufacturers' instructions. Bevacizumab, another type of angiogenesis inhibitor, was administered intravenously at a dose of 15 mg/kg body weight every 3 wk. Atezolizumab, a type of programmed cell death ligand 1 blocker, was administered intravenously at a dose of 1200 mg every 3 wk. Programmed cell death protein 1 (PD-1) blockers, including pembrolizumab, camrelizumab, tislelizumab, and sintilimab, were administered intravenously at 200 mg every 3 wk. Toripalimab, another PD-1 blocker, was injected through an intravenous drip of 240 mg every 3 wk following the instructions. The HAIC regimen was based on FOLFOX and consisted of 85 mg/m² oxaliplatin, 200 mg/m² calcium folinate, and 2.5 g/m² 5-fluorouracil every 3 wk. HAIC was performed under the guidance of digital subtraction angiography by interventional radiologists. The celiac axis, superior mesenteric artery, inferior phrenic artery and right renal artery were selectively catheterized for angiography. If angiography revealed that the HCC blood supply originated from different vessels, the main feeding artery was reserved for super selective catheterization, and an indwelling microcatheter was inserted into the HAIC while the other feeding vessels were embolized. During the study period, dose modifications and treatment interruptions were sometimes initiated according to drug-related toxicity grades, as recommended relative to the physiological condition of each patient. HAIC was performed for 4-6 rounds in the absence of disease progression. Patients received angiogenesis inhibitors and PD-1/PD-L1 blockers during and after HAIC treatment to consolidate the therapeutic effects in the longterm.

Assessment of clinical outcomes: The patients involved in the study were followed up every 6-12 wk to assess treatment response. Radiological response was assessed according to the modified Response Evaluation Criteria in Solid Tumors (mRECIST) criteria based on liver dynamic computed tomography or magnetic resonance imaging data. The primary endpoint that was assessed was overall survival (OS), which was defined as the time from the start date of systemic chemotherapy or HAIC to death. progression-free survival (PFS) (the time from the start date of systemic chemotherapy or HAIC to the date of disease progression or death from any cause). The secondary endpoints that were determined included PFS and 6-, 12- and 24-mo OS rates; objective response rate (ORR); and adverse events (AEs). The ORR was defined as the proportion of patients who achieved a complete response (CR) or partial response (PR), and the disease control rate was defined as the proportion of patients who achieved CR, PR, or stable disease. AEs during treatment were identified using patient-reported symptom data, examination-based findings, and clinical laboratory test results. The National Cancer Institute Common Terminology Criteria for AEs, version 5.0, was used to classify AEs from any cause according to type and severity.

Statistical analysis

Unstratified hazard ratios (HRs) with 95% CI and odds ratios (ORs) with the number of responders and sample sizes that compared the different treatment regimens for treating uHCC were retrieved and synthesized to determine the overall treatment effects. Potential heterogeneity among the studies was assessed using I² statistics. Random effects models were used to calculate pooled ORR or HR in the presence of significant heterogeneity ($I^2 > 50\%$); otherwise, the fixed effects model was used.

To account for the different distributions of covariates between the two groups, we performed PSM. Then, 1:1 matching was performed using nearest-neighbor matching based on the performance status data. In this study, the caliper of the match was 0.03. OS, PFS and survival rates were calculated using the Kaplan-Meier method and were compared between the groups using the log-rank test. Cox regression was used to explore the potential risk factors associated with survival time. All real-world clinical data are expressed as the mean ± SD, median (range), or number (%), as appropriate. Continuous variables were compared using Student's t test (or the Mann-Whitney U test, if appropriate), and categorical variables were compared using the chi-square test (or Fisher's exact test, if appropriate).



WJG https://www.wjgnet.com

Table 1 P value of different interventions							
Intervention	OR	Intervention	HR				
HAIC	0.9520107	HAIC + Sorafenib	0.03757857				
HAIC + Sorafenib	0.9420071	HAIC	0.09333929				
SIRT	0.8131607	TACE + Lenvatinib	0.098075				
TACE + Lenvatinib	0.8014429	Sintilimab + Bevacizumabbiosimila	0.24973929				
Sintilimab + Bevacizumabbiosimila	0.6559643	Camrelizumab + Rivoceranib	0.30909286				
Lenvatinib + Pembrolizumab	0.5922857	Atezolizumab + Bevacizumab	0.35838929				
Camrelizumab + Rivoceranib	0.5853643	Lenvatinib + Pembrolizumab	0.51200714				
Durvalumab + Tremelimumab	0.523425	Durvalumab + Tremelimumab	0.52634286				
Durvalumab	0.3891893	Donafenib	0.62278571				
Atezolizumab + Bevacizumab	0.3369036	Tislelizumab	0.66164643				
Lenvatinib	0.3254286	Nivolumab	0.67260357				
Tislelizumab	0.2732107	Durvalumab	0.68975714				
Nivolumab	0.1859964	Lenvatinib	0.78121071				
Donafenib	0.1165357	Sorafenib	0.9135				
Sorafenib	0.007075	SIRT	0.97393214				

HAIC: Hepatic arterial infusion chemotherapy; TACE: Transcatheter arterial chemoembolization; SIRT: Selective internal radiation therapy; HR: Hazard ratio; OR: Odds ratio.

Statistical analyses were conducted using R v4.2.2 (R Core Team, Vienna, Austria). https://www.R-project.org/). Twosided *P* values < 0.05 were considered to indicate statistical significance.

RESULTS

Network meta-analysis on first-line treatment of uHCC

Literature search and screening results: Our initial literature search resulted 1735 articles. After deleting duplicate publications, 1079 articles remained. After screening the titles and abstracts, 68 articles were excluded. Our full-text review resulted in the removal of an additional seven articles. Ultimately, 13 studies involving 7817 patients were included in this network meta-analysis[6,8,16-25]. The literature selection process is described in Supplementary Figure 2, and the characteristics of the included patient population are shown in Supplementary Table 1.

Results of network-meta-analysis: ORRs per Response Evaluation Criteria in Solid Tumors 1.1 and HRs were reported in all 13 studies and included 15 different interventions. There was no significant heterogeneity between the studies (ORR: $I^2 = 5\%$; HR: $I^2 = 7\%$), so the fixed-effects model was adopted. The P scores for ORR showed that the best ORR outcomes were obtained with HAIC compared to sorafenib (OR: 35.66; 95%CI: 9.94–249.91; P: 0.952; Table 1, Supplementary Figure 3, Supplementary Table 2). The *P* values for HRs showed that the lowest HRs were obtained with HAIC plus sorafenib compared to those with sorafenib alone (HR: 0.36; 95%CI: 0.25–0.52; *P* = 0.891; Table 1, Supplementary Figure 4, Supplementary Table 3).

Retrospective study

Baseline characteristics of the patients: A total of 442 patients with uHCC were enrolled in the study; 324 patients underwent triple therapy, and 118 patients underwent AIPB. The median follow-up times were 14.6 months and 16.8 \pm 10.3 months in the triple therapy group and 8.25 months and 11.7 \pm 10.2 months in the AIPB group. The algorithm used for case enrollment is shown in Supplementary Figure 5. The average number of patients who received 5.08 \pm 1.61 rounds of HAIC in the triple therapy group. Based on our multivariable logistic regression model, baseline characteristics, including age, ECOG PS, Child–Pugh class, maximum tumor diameter, AFP level, tumor number, and vascular invasion and extrahepatic metastasis conditions, which were significantly different between the groups, were matched (Table 2). After PSM, 88 patients in the triple therapy group were matched to 88 patients in the AIPB group (Table 2). The median age in both groups was 55.0 years, and all the patients were evaluated as having an ECOG PS ranging from 0-1. Notably, some patients in the AIPB group were diagnosed at an earlier stage. In other words, the proportion of BCLC C patients in the AIPB cohort was lower (87.5% *vs* 92%).

Table 2 Baseline characteristics before or after propensity score matching, n (%)

	Prior to PSM			Following PSM		
Variable	Triple therapy (<i>n</i> = 324)	AIPB (<i>n</i> = 118)	<i>P</i> value	Triple therapy (<i>n</i> = 88)	AIPB (<i>n</i> = 88)	<i>P</i> value
Age						
Mean (SD)	50.2 (11.2)	55.4 (11.6)	< 0.001	54.2 (10.8)	53.9 (11.2)	0.859
Median [Min, Max]	51.0 [23.0, 80.0]	56.5 [23.0, 82.0]		55.0 [26.0, 78.0]	55.0 [23.0, 74.0]	
Sex						
Female	36 (11.1)	8 (6.8)	0.244	11 (12.5)	6 (6.8)	0.307
Male	288 (88.9)	110 (93.2)		77 (87.5)	82 (93.2)	
ECOG PS						
0	318 (98.1)	106 (89.8)	< 0.001	85 (96.6)	85 (96.6)	1
1	6 (1.9)	11 (9.3)		3 (3.4)	3 (3.4)	
2	0 (0)	1 (0.8)		0 (0)	0 (0)	
Hepatitis virus						
Negative	27 (8.3)	15 (12.7)	0.228	11 (12.5)	12 (13.6)	1
Positive	297 (91.7)	103 (87.3)		77 (87.5)	76 (86.4)	
ALT						
mean (SD)	62.4 (72.8)	53.6 (34.9)	0.0894	58.4 (60.5)	53.2 (36.1)	0.489
Median [Min, Max]	44.6 [8.90, 930]	45.4 [6.10, 196]		41.1 [8.90, 448]	44.6 [6.10, 196]	
AST						
mean (SD)	92.5 (83.1)	88.3 (78.5)	0.627	82.0 (65.3)	87.1 (84.9)	0.657
Median [Min, Max]	66.2 [13.5, 702]	66.8 [11.1, 470]		58.7 [13.5, 327]	63.7 [11.1, 470]	
Child-Pugh class						
А	296 (91.4)	94 (79.7)	0.00133	75 (85.2)	77 (87.5)	0.826
В	28 (8.6)	24 (20.3)		13 (14.8)	11 (12.5)	
AFP > 400 ng/mL						
No	142 (43.8)	60 (50.8)	0.229	45 (51.1)	42 (47.7)	0.763
Yes	182 (56.2)	58 (49.2)		43 (48.9)	46 (52.3)	
Maximum diameter of tumor/cm						
mean (SD)	10.5 (4.25)	9.38 (4.75)	0.0291	9.86 (4.07)	9.57 (4.83)	0.668
Median [Min, Max]	10.3 [1.90, 23.5]	9.25 [1.10, 21.0]		10.0 [2.10, 19.2]	9.10 [1.10, 20.5]	
Tumor number						
Single	122 (37.7)	19 (16.1)	< 0.001	21 (23.9)	19 (21.6)	0.857
Multiple	202 (62.3)	99 (83.9)		67 (76.1)	69 (78.4)	
Vascular invasion						
No	78 (24.1)	42 (35.6)	0.0221	25 (28.4)	30 (34.1)	0.515
Yes	246 (75.9)	76 (64.4)		63 (71.6)	58 (65.9)	
Extrahepatic metastasis						
No	171 (52.8)	45 (38.1)	0.00888	34 (38.6)	38 (43.2)	0.646
Yes	153 (47.2)	73 (61.9)		54 (61.4)	50 (56.8)	
BCLC						



Jaishideng® WJG | https://www.wjgnet.com

А	0 (0)	2 (1.7)	0.0632	0 (0)	2 (2.3)	0.309
В	30 (9.3)	11 (9.3)		7 (8.0)	9 (10.2)	
С	294 (90.7)	105 (89.0)		81 (92.0)	77 (87.5)	

AFP: Alpha-fetoprotein; ALT: Alanine aminotransferase; AST: Aspartate aminotransferase; AIPB: Angiogenesis inhibitors and programmed cell death protein 1/programmed cell death ligand 1 blockers; triple therapy: Hepatic arterial infusion chemotherapy plus angiogenesis inhibitors and programmed cell death protein 1/programmed cell death ligand 1 blockers; PSM: Propensity score matching; BCLC: Barcelona Clinic Liver Cancer; ECOG PS: Eastern Cooperative Oncology Group performance status.



DOI: 10.3748/wjg.v30.i4.318 Copyright ©The Author(s) 2024.

Figure 1 A patients with unresectable hepatocellular carcinoma and inferior vena cava tumor thrombus who has received triple therapy and reached complete response according to modified Response Evaluation Criteria in Solid Tumors and the tumor thrombus has shrank completely. A-C: Images taken before treatment; D-F: Images taken at 1 month after hepatic arterial infusion chemotherapy; G-I: Images taken at the latest follow-up. A, D and G is arterial phase in the axial view. B, E and H is venous phase in the axial view. C, F and I is venous phase in the coronal view. The arrow denotes the tumor thrombus. R: Right; RF: Right foot; L: Left; LH: Left head.

Efficacy of different treatments: The ORR was 62.9% (n = 204) in the triple group and 29.7% (n = 35) in the AIPB group in the primary database. After PSM, the ORR of the triple therapy group was still greater (55.7% vs 35.2%, P = 0.032) (Supplementary Table 4). CR was observed in 21 patients prior to PSM and in 6 patients following PSM in the triple therapy group (an example can be seen in Figure 1). According to the Kaplan-Meier analysis, the primary data showed longer median PFS (11.1 months vs 6.0 months, P < 0.001) and median OS (not reached vs 11.8 months, P < 0.001) with triple therapy. After PSM, the median PFSs were estimated to be 12.5 months and 7.8 months (P = 0.036), and the median OS were 31.6 months and 14.6 months (HR = 2.42, 95%CI = 1.49-3.92, P < 0.001) in the triple therapy group and AIPB group, respectively (Figure 2). The 6-months, 12-months, and 24-months survival rates of the patients receiving triple therapy were 96.5%, 82.2% and 57.0%, respectively, while they were 73.5%, 54.3% and 37.7%, respectively, in the AIPB group.

Univariate analysis revealed that four factors had effects on OS in the triple therapy group: Larger tumor diameter, multiple foci, extrahepatic metastasis, Child-Pugh grade B and number of rounds of HAIC (Figure 3A). Cox multivariate regression analysis revealed that Child-Pugh grade B (HR: 1.74, P < 0.001; Figure 3B) and multiple foci (HR: 2.17, P < 0.001; Figure 3B) were risk factors for poor long-term survival, and > 4 rounds of HAIC was a protective factor for survival (HR: 0.43, P < 0.001; Figure 3B). Survival analysis also revealed that patients who received > 4 rounds of HAIC (not reached vs 18.2 months; P < 0.001; Supplementary Figure 6A) or who were diagnosed with a single disease focus (not reached vs 24.6 months; P < 0.001; Supplementary Figure 6B) had longer OS. Subgroup analysis of OS using forest plots revealed that triple therapy was more effective in most patients, especially for males, Child-Pugh A patients, patients aged < 60 years, and patients diagnosed with multiple tumors or extrahepatic metastasis (Figure 3C).

Baishidena® WJG | https://www.wjgnet.com



Figure 2 Kaplan-Meier curves of progression-free survival and overall survival in the triple therapy group and angiogenesis inhibitors and programmed cell death protein 1/programmed cell death ligand 1 blockers group. A: Prior to propensity score matching (PSM), median progression-free survival (PFS) was 11.1 vs 6.0 mo, P < 0.001; B: Prior to PSM, median overall survival (OS) was not reached vs 11.8 mo, P < 0.001; C: Following PSM, median PFS was 12.5 vs 7.8 mo, P = 0.036; D: Following PSM, median OS was 31.6 vs 14.6 mo, P < 0.001. triple therapy: Hepatic arterial infusion chemotherapy plus angiogenesis inhibitors and programmed cell death protein 1/programmed cell death ligand 1 blockers; AIPB: Angiogenesis inhibitors and programmed cell death protein 1/programmed cell death ligand 1 blockers.

Safety of different treatments: After PSM, the incidence of AEs in the triple therapy group was greater than that in the AIPB group (94.3% *vs* 75.4%, P < 0.001). Although more Grade 3-4 AEs occurred in the triple therapy group, there was no significant difference in the incidence of Grade 3-4 AEs (56.8% *vs* 43.2%, P = 0.097), and there were no Grade 5 AEs (Table 3). The most common AE was abdominal pain in the triple therapy group, for which the incidence was 79.8% (259/ 324). When they started HAIC treatment, many patients had varying degrees of abdominal pain during the infusion of oxaliplatin. This was typically managed by slowing the speed of infusion or temporarily stopping it. In some cases of particularly severe and acute abdominal pain, anisodamine or lidocaine was administered through intravenous injection or an arterial catheter to relieve the pain. Two patients developed hepatic comas following HAIC but fully recovered during treatment. In addition, liver dysfunction, including increases in aminotransferases and/or bilirubin, was relatively common in both groups, not only because of drug-related side effects but also because of their own background of liver cirrhosis.

DISCUSSION

Although the first-line treatment recommended by authoritative clinical guidelines for uHCC is AIPB, such as atezolizumab plus bevacizumab, these treatments have a number of limitations in clinical applications. The default anti-inflammatory or immunotolerant immune status of the liver may interfere with the drugs that act on it[26], which may lead to a relatively low ORR. The main cause of death among patients with uHCC is liver tumor progression[27,28]. Although there are a number of different protocols for administering AIPB, the median patient survival time using this approach has remained less than 24 months[17,19,20,29]. In addition, the IMbrave150 studies suggested that the effects of AIPB treatment are likely to be severely diminished if patients are diagnosed with high-risk factors, such as tumor invasion of the main trunk of the portal vein (Vp4), bile duct invasion, or/or tumor occupancy of \geq 50% of the liver[17,30].

Table 3 Summary of adverse events, n (%)								
	Before PSM				After PSM			
	Triple therap 324)	y group (<i>n</i> =	AIPB group (<i>n</i> = 118)		Triple therapy group (<i>n</i> = 88)		AIPB group (<i>n</i> = 88)	
	Grade 1-2	Grade 3-4	Grade 1-2	Grade 3-4	Grade 1-2	Grade 3-4	Grade 1-2	Grade 3-4
Fever	67 (20.7)	23 (7.1)	4 (3.4)	2 (1.7)	20 (22.7)	8 (9.1)	4 (4.5)	0 (0)
Nausea	100 (30.9)	0 (0)	6 (5.1)	0 (0)	24 (27.3)	0 (0)	5 (5.7)	0 (0)
Vomit	32 (9.9)	15 (4.6)	7 (5.9)	6 (5.1)	7 (8.0)	3 (3.4)	6 (6.8)	4 (4.5)
Abdominal pain	97 (29.9)	162 (50.0)	18 (15.3)	18 (15.3)	24 (27.3)	44 (50.0)	15 (17.0)	13 (14.8)
ALT increased	79 (24.4)	125 (38.6)	32 (27.1)	39 (33.1)	27 (30.4)	33 (37.5)	22 (25.0)	29 (33.0)
AST increased	130 (40.1)	161 (49.7)	36 (30.5)	44 (37.3)	40 (45.5)	49 (55.7)	26 (29.5)	34 (38.6)
Hyperbilirubinemia	37 (11.4)	7 (2.2)	7 (5.9)	11 (9.3)	10 (11.4)	0 (0)	4 (4.5)	8 (9.1)
Anemia	37 (11.4)	8 (2.5)	7 (5.9)	4 (3.4)	11 (12.5)	1 (1.1)	5 (5.7)	3 (3.4)
Neutropenia	86 (26.5)	20 (6.2)	9 (7.6)	12 (10.2)	29 (33.0)	5 (5.7)	5 (5.7)	9 (10.2)
Thrombocytopenia	103 (31.8)	90 (27.8)	27 (22.9)	20 (16.9)	24 (27.3)	16 (18.2)	20 (22.7)	13 (14.8)
Bleeding	48 (14.8)	3 (0.9)	1 (0.8)	1 (0.8)	12 (13.6)	1 (1.1)	1 (1.1)	1 (1.1)
Diarrhea	62 (19.1)	60 (18.5)	22 (18.6)	3 (2.5)	17 (19.3)	5 (5.7)	17 (19.3)	3 (3.4)
Hoarseness	80 (24.7)	0 (0)	13 (11.0)	0 (0)	21 (23.9)	0 (0)	10 (11.4)	0 (0)
Rash	92 (28.4)	4 (1.2)	28 (23.7)	4 (3.4)	24 (27.3)	0 (0)	20 (22.7)	3 (3.4)
HFS	69 (21.3)	9 (2.8)	19 (16.1)	18 (15.3)	18 (20.5)	1 (1.1)	14 (15.9)	11 (12.5)
Hypertension	76 (23.5)	11 (3.4)	31 (26.3)	34 (28.8)	22 (25.0)	3 (3.4)	23 (26.1)	27 (30.7)
RCCEP	42 (13.0)	14 (4.3)	16 (13.6)	6 (5.1)	11 (12.5)	5 (5.7)	13 (14.8)	6 (6.8)
Hypothyroidism	64 (19.8)	8 (2.5)	2 (1.7)	0 (0)	18 (20.5)	2 (2.3)	0 (0)	0 (0)
Fatigue	32 (9.9)	31 (9.6)	4 (3.4)	4 (3.4)	6 (6.8)	8 (9.1)	3 (3.4)	2 (2.3)
Hepatitis	10 (3.1)	1 (0.3)	0 (0)	0 (0)	5 (5.7)	1 (1.1)	0 (0)	0 (0)
Coma	0 (0)	2 (0.6)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)

AFP: Alpha-fetoprotein; AIPB: Angiogenesis inhibitors and programmed cell death protein 1/programmed cell death ligand 1 blockers; triple therapy: Hepatic arterial infusion chemotherapy plus angiogenesis inhibitors and programmed cell death protein 1/programmed cell death ligand 1 blockers; ALT: Alanine aminotransferase; AST: Aspartate aminotransferase; HFS: Hand-foot syndrome; RCCEP: Reactive cutaneous capillary endothelial proliferation.

Many uHCC patients in some areas, especially in China, are diagnosed with vascular invasion or/or a high tumor burden. The most effective way to prolong survival is to control liver lesions. In terms of local hepatic treatment for uHCC, the most popular choice for most physicians is transcatheter arterial chemoembolization (TACE). Nonetheless, if the tumor burden is high, there is a very high probability of "TACE failure/refractoriness"[31-33]. If patients are diagnosed with reduced or absent portal vein blood supplies caused by portal vein tumor thrombi or severe cirrhosis, the use of TACE will be limited. Several studies have revealed that HAIC is more effective than TACE for large HCCs[34]. The FOHAIC study suggested that FOLFOX-HAIC had a significant effect on patients with uHCC and that HAIC could be used as an additional local hepatic treatment for uHCC[8]. According to our meta-analysis, HAIC plus sorafenib or HAIC alone was able to prolong the survival time of patients with uHCC more than AIPB regimens. Unfortunately, there is a scarcity of prospective or retrospective studies with large sample sizes on triple therapy.

Our retrospective data revealed that triple therapy was effective and safe. The ORR, PFS, and OS of patients receiving triple therapy (ORR: 33.2% per mRECIST; PFS: 6.9 months; OS: 19.2 months) outperformed those of patients receiving most AIPB regimens. For example, this was true for atezolizumab plus bevacizumab (ORR: 33.2% per mRECIST; PFS: 6.9 months; OS: 19.2 months) in the IMbrave150 trial; pembrolizumab plus lenvatinib (ORR: 40.8% per mRECIST; PFS: 8.2 months; OS: 21.2 months) in the LEAP002 trial; and camrelizumab plus rivoceranib (ORR: 33.1% per mRECIST; PFS: 5.6 months; OS: 22.1 months) in the CARES 310 study[7,17,19,29]. The survival benefit observed in this study may be due to the synergistic antitumor effects of these chemical agents. Transarterial chemotherapy can induce immunogenic cell death by releasing tumor-related antigens and supporting the evolution of tumor-specific CD8+ T cells, which may synergize with angiogenesis inhibitors to enhance the effect of PD-1/PD-L1 blockers[35-37]. Increased concentrations of drugs in the liver can cause liver lesions to shrink directly and slow the deterioration of liver function caused by disease progression. According to our survival analysis, the patients in the AIPB subgroup had shorter survival times than those

A

Factor			HR (95%CI)	P value
HAIC rounds>4				
No				
Yes			0.43(0.28 to 0.64)	<0.001
ECOG				
0				
1			1(0 to Inf)	1
Age	-		0.98(0.97 to 1.00)	0.073
Sex				
Female				
Male		•>	1.13(0.60 to 2.12)	0.7
Hepatitis virus				
Negative				
Positive		→	0.98(0.45 to 2.12)	0.96
Child-Pugh class				
A				
В		\longrightarrow	2.82(1.59 to 4.98)	<0.001
AST			1.00(1.00 to 1.00)	0.23
ALI			1.00(1.00 to 1.00)	0.45
AFP>400ng/mL				
NO				
Yes	_	-	1.21(0.81 to 1.80)	0.36
Maximum tumor diameter			1.05(1.00 to 1.10)	<0.05
namor number				
Single				
Multiple		\longrightarrow	2.14(1.37 to 3.34)	<0.001
Vascular invasion				
No				
Yes	-	→	1.54(0.94 to 2.52)	0.085
Extrahepatic metastasis				
No				
Tes			1.52(1.02 to 2.26)	<0.05
	0.5	1.5 2	2	

Prolong survival Shorten survival

В					
Factor				HR(95%CI)	P value
HAIC rounds>4					
No					
Yes				0.43(0.28 to 0.65)	<0.001
Child-Pugh class					
A					
В				→ 2.74(1.53 to 4.91)	<0.001
Maximum tumor diameter		-		1.04(0.99 to 1.09)	0.15
Tumor number					
Single					
Multiple				→ 2.17(1.39 to 3.40)	<0.001
Extrahepatic metastasis					
No					
Yes				1.20(0.80 to 1.80)	0.39
	0.5	1	1.5	2	

Prolong survival Shorten survival

С

Subgroup	Triple therapy (n = 88)	AIPB (<i>n</i> = 88)	1	HR (95%CI)	P value
Sex			1		
Female	11	6	-	0.25(0.04 to 1.56)	0.14
Male	77	82		0.44(0.27 to 0.73)	0.002
Age					
<60y	59	59		0.47(0.27 to 0.81)	0.007
≥60y	29	29		0.33(0.12 to 0.89)	0.028
ECOG					
0	85	85		0.43(0.26 to 0.70)	<0.001
1	3	3			1
Hepatitis virus					
Negative	11	12		→ 0.49(0.12 to 2.07)	0.33
Positive	77	76		0.40(0.24 to 0.68)	< 0.001
AFP					
<400ng/mL	45	42	_ -	0.43(0.21 to 0.90)	0.024
≥400ng/mL	43	46		0.41(0.21 to 0.78)	0.006
ChildPugh					
А	75	77		0.39(0.23 to 0.65)	<0.001
В	13	11		→ 0.65(0.16 to 2.64)	0.55
Maximum diamete	r of tumor				
<5cm	9	14			1
≥5cm	79	74		0.43(0.27 to 0.72)	<0.001
Tumor number					
Single	21	19		0.38(0.11 to 1.26)	0.11
Multiple	69	67		0.41(0.24 to 0.70)	0.001
Vacular invasion					
No	30	25		0.32(0.12 to 0.84)	0.021
Yes	58	63		0.42(0.24 to 0.74)	0.0027
Extrahepatic meta	stasis				
No	34	38		0.57(0.26 to 1.25)	0.16
Yes	54	50		0.34(0.18 to 0.62)	< 0.001
			0.5 1 1.5	2	
		←			

Favours triple thrapy Favours AIPB

DOI: 10.3748/wjg.v30.i4.318 Copyright ©The Author(s) 2024.

Figure 3 Forest plots of Cox regression analysis. A: The results of univariatable Cox regression in the triple therapy group prior to propensity score matching (PSM); B: The results of multivariable Cox regression in the triple therapy group prior to PSM; C: Subgroup analysis of overall survival after PSM. HR: Hazard ratio; ECOG: Eastern Cooperative Oncology Group; Inf: Infinite; EHM: Extrahepatic metastasis; AFP: Alpha-fetoprotein; ALT: Alanine aminotransferase; AST: Aspartate aminotransferase; triple therapy: Hepatic arterial infusion chemotherapy plus angiogenesis inhibitors and programmed cell death protein 1/programmed cell death protein 1/programmed cell death ligand 1 blockers; HAIC: Hepatic arterial infusion

Saishideng® WJG | https://www.wjgnet.com

chemotherapy

in the other trials on AIPB regimens, likely due to their poor baseline conditions prior to treatment. More than half of the patients (n = 58) in the AIBP group were diagnosed with major vein tumor thrombus, and the mean maximum tumor diameter was more than 9 cm, suggesting that the patients in this group had high tumor burdens. The median OS of similar patients in the IMbrave150 study was only 7.8 months, which is consistent with the results of our study[30].

We also attempted to identify which factors could influence the effect of triple therapy and found that > 4 rounds of HAIC were a protective factor. Four rounds of HAIC represent a regimen similar to the median number of HAIC rounds reported in several other studies[8,38-40]. The number of HAIC rounds performed was strongly affected by each patient's response to triple therapy because if tumors progress after the first few HAIC cycles, the HAIC cycles will be discontinued. Multiple liver lesions have also been recognized as risk factors because the presence of multiple lesions often implies the presence of multiple feeding vessels. Therefore, even if attempts are made to embolize other arteries until there is only a single blood supply, there is a high probability that some small arteries may be missed. To ensure that all lesions can be treated by HAIC, a microcatheter should be placed in a larger blood vessel branch, which implies that more normal liver tissue is likely to be damaged by the administered drugs, potentially harming liver function.

Overall, the incidence of adverse reactions to triple therapy was greater than that reported in the AIPB group. The combination of HAIC and systematic treatments was able to increase the incidence of AEs; another reasonable explanation is that most AIBP patients were treated and followed up as outpatients so that some AEs, especially some slight AEs, were ignored. Notably, abnormal liver function was the common AE in the triple therapy group. However, unlike many other local treatments, the effects of HAIC on liver function appear to be largely short-term, with few apparent adverse effects on long-term survival. However, we believe that the limitation of triple therapy in the Child-Pugh B population with poor hepatic functional reserve may result from irreversible liver injury secondary to chemotherapy. Another common AE, thrombocytopenia, is caused not only by myelosuppression due to chemotherapy but also by hypersplenism secondary to cirrhosis. A substantial proportion of the patients recovered from thrombocytopenia following splenic embolization.

A substantial amount of information was lost due to the limitations of this retrospective study. Our sample included only patients from China, so the study was inevitably affected by some degree of sampling bias. It remains unclear exactly which biomarkers can be used to judge patients' prognoses. Studies at the cellular or molecular level could not be carried out due to a lack of tumor biopsy samples. To prove the efficacy and safety of triple therapy, additional large-scale prospective RCTs on this topic are warranted.

CONCLUSION

This meta-analysis suggested that HAIC-based treatment regimens were able to effectively improve the prognosis in patients with uHCC. Our findings confirmed that even though the triple therapy protocol increased the incidence and severity of AEs, it yielded a higher ORR and longer PFS and OS than AIPB.

ARTICLE HIGHLIGHTS

Research background

Unresectable hepatocellular carcinoma had been difficult to be treated in the past, hepatic arterial chemotherapy infusion chemotherapy (HAIC) as well as angiogenesis inhibitors plus programmed cell death protein 1/programmed cell death ligand 1 (PD-1/PD-L1) blockers were proved to prolong the unresectable hepatocellular carcinoma (uHCC) patients' survival, respectively. Meanwhile, some phase II single arm suggested that the combination of HAIC and angiogenesis inhibitors and PD-1/PD-L1 blockers (AIPB) (triple therapy) was effective in treating uHCC. But which treatment is the best choice was still confused. The study was designed to answer the question.

Research motivation

The best first-line treatment for uHCC was unclear and there was lack of studies to compare the efficacy and safety between triple therapy and AIPB. There were so many choices that clinical staff may be confused when they need to treat uHCC patients. If we can find the relatively better regimen, it is helpful for the standardization of the uHCC treatment to improve the patients' prognosis.

Research objectives

The study aimed to identified the HAIC and HAIC-based treatments was the best choice for uHCC. Based on the result, we explored the efficacy and safety of one of HAIC-based treatment, triple therapy in the real-world condition compared to AIPB. The results of the study could be the evidence to guide clinical reasonable treatment and prospective clinical study.

WJG | https://www.wjgnet.com

Research methods

We have tried to perform a network meta-analysis to find the first choice to uHCC and identified the efficacy and safety of triple therapy compared to AIPB through a retrospective cohort study.

Research results

The network meta-analysis including 13 phase randomized controlled trials (RCTs) showed HAIC and HAIC-based treatments were likely to be the first choice to treat uHCC. HAIC plus camrelizumab plus AIPB (triple therapy) had better progression-free survival and overall survival than AIPB without HAIC for uHCC. Even though the incidence of adverse events in the triple therapy group was higher than the AIPB group, the safety of triple therapy was still acceptable.

Research conclusions

HAIC-based treatments were better than other regimens for treating uHCC. Triple therapy was more effective than AIPB in the Chinese uHCC population. All of the above results proved the significance of local treatments in the uHCC treating.

Research perspectives

There is absolutely a need for studies at the cellular or molecular level and additional large-scale prospective RCTs on this topic.

ACKNOWLEDGEMENTS

We are thankful for the support of the patients in this study. We are grateful to Dr. Fei Cao and Ms. Xiao-Ting Bei for their assistance with the clinical management of the patients.

FOOTNOTES

Co-first authors: Yu-Zhe Cao and Guang-Lei Zheng.

Co-corresponding authors: Zi-Lin Huang and Meng-Xuan Zuo.

Author contributions: Cao YZ, Zheng GL, Zuo MX and Huang ZL participated equally in conceptualization; Cao YZ, Zuo MX and Zheng GL made great contribution in the in the data curation, formal analysis, methodology and investigation; Zhang TQ, Shao HY and Pan JY has participated in the investigation and the validation; The project administration and supervision were accomplished by Huang ZL and Zuo MX; Cao YZ and Zheng GL had finished the original draft; Zuo MX and Huang ZL has been responsible for the review and editing; All authors have provided final approval of the manuscript version to be submitted for publication. Cao YZ and Zheng GL, contributed equally to this work and share first authorship for their contribution in conceptualization, data curation, formal analysis, methodology, investigation and draft writing. Huang ZL and Zuo MX, contributed equally to this work and are co-corresponding authors for their contribution in conceptualization, project administration, supervision, revision and approvement for final edition of the manuscript.

Supported by Natural Science Foundation of Guangdong Province, No. 2020A1515011539.

Institutional review board statement: This study was reviewed and approved by the Sun Yat-sen University Cancer Center Ethics Committee. Informed consent was waived due to the retrospective nature of the analysis.

Informed consent statement: Informed consent was waived due to the retrospective nature of the analysis, which has been approved by Ethic Committee of Sun Yat-sen University Cancer Center.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

Data sharing statement: The datasets of the manuscript is available from the corresponding authors at Sun Yat-sen University Cancer Center Research Data Deposit repository (https://www.researchdata.org.cn/). Consent was not obtained but the presented data are anonymized and risk of identification is low.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: https://creativecommons.org/Licenses/by-nc/4.0/

Country/Territory of origin: China

ORCID number: Tian-Qi Zhang 0000-0002-1528-5323; Zi-Lin Huang 0000-0002-1643-1994; Meng-Xuan Zuo 0000-0003-3589-7316.

S-Editor: Li L



REFERENCES

- 1 Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, Bray F. Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. CA Cancer J Clin 2021; 71: 209-249 [PMID: 33538338 DOI: 10.3322/caac.21660]
- Thomas MB, Jaffe D, Choti MM, Belghiti J, Curley S, Fong Y, Gores G, Kerlan R, Merle P, O'Neil B, Poon R, Schwartz L, Tepper J, Yao F, 2 Haller D, Mooney M, Venook A. Hepatocellular carcinoma: consensus recommendations of the National Cancer Institute Clinical Trials Planning Meeting. J Clin Oncol 2010; 28: 3994-4005 [PMID: 20679622 DOI: 10.1200/JCO.2010.28.7805]
- 3 Villanueva A. Hepatocellular Carcinoma. N Engl J Med 2019; 380: 1450-1462 [PMID: 30970190 DOI: 10.1056/NEJMra1713263]
- Kudo M, Arizumi T. Transarterial Chemoembolization in Combination with a Molecular Targeted Agent: Lessons Learned from Negative 4 Trials (Post-TACE, BRISK-TA, SPACE, ORIENTAL, and TACE-2). Oncology 2017; 93: 127-134 [PMID: 29258086 DOI: 10.1159/000481243
- Llovet JM, Ricci S, Mazzaferro V, Hilgard P, Gane E, Blanc JF, de Oliveira AC, Santoro A, Raoul JL, Forner A, Schwartz M, Porta C, 5 Zeuzem S, Bolondi L, Greten TF, Galle PR, Seitz JF, Borbath I, Häussinger D, Giannaris T, Shan M, Moscovici M, Voliotis D, Bruix J; SHARP Investigators Study Group. Sorafenib in advanced hepatocellular carcinoma. N Engl J Med 2008; 359: 378-390 [PMID: 18650514 DOI: 10.1056/NEJMoa07088571
- Kudo M, Finn RS, Qin S, Han KH, Ikeda K, Piscaglia F, Baron A, Park JW, Han G, Jassem J, Blanc JF, Vogel A, Komov D, Evans TRJ, 6 Lopez C, Dutcus C, Guo M, Saito K, Kraljevic S, Tamai T, Ren M, Cheng AL. Lenvatinib vs sorafenib in first-line treatment of patients with unresectable hepatocellular carcinoma: a randomised phase 3 non-inferiority trial. Lancet 2018; 391: 1163-1173 [PMID: 29433850 DOI: 10.1016/S0140-6736(18)30207-11
- Finn RS, Oin S, Ikeda M, Galle PR, Ducreux M, Kim TY, Kudo M, Breder V, Merle P, Kaseb AO, Li D, Verret W, Xu DZ, Hernandez S, Liu 7 J, Huang C, Mulla S, Wang Y, Lim HY, Zhu AX, Cheng AL; IMbrave150 Investigators. Atezolizumab plus Bevacizumab in Unresectable Hepatocellular Carcinoma. N Engl J Med 2020; 382: 1894-1905 [PMID: 32402160 DOI: 10.1056/NEJMoa1915745]
- 8 Lyu N, Wang X, Li JB, Lai JF, Chen QF, Li SL, Deng HJ, He M, Mu LW, Zhao M. Arterial Chemotherapy of Oxaliplatin Plus Fluorouracil Versus Sorafenib in Advanced Hepatocellular Carcinoma: A Biomolecular Exploratory, Randomized, Phase III Trial (FOHAIC-1). J Clin Oncol 2022; 40: 468-480 [PMID: 34905388 DOI: 10.1200/JCO.21.01963]
- Gu YK, Zhang TQ, Zuo MX, Geng ZJ, Li JB, Huang ZL, Wu PH. Hepatic artery infusion chemotherapy (HAIC) combined with apatinib and 9 camrelizumab for hepatocellular carcinoma (HCC) in BCLC stage c: A prospective, single-arm, phase II trial (TRIPLET study). Conference Abstract. Journal of Clinical Oncology40 [DOI: 10.1200/JCO.2022.40.16_suppl.4106]
- Lai Z, He M, Bu X, Xu Y, Huang Y, Wen D, Li Q, Xu L, Zhang Y, Wei W, Chen M, Kan A, Shi M. Lenvatinib, toripalimab plus hepatic 10 arterial infusion chemotherapy in patients with high-risk advanced hepatocellular carcinoma: A biomolecular exploratory, phase II trial. Eur J Cancer 2022; 174: 68-77 [PMID: 35981413 DOI: 10.1016/j.ejca.2022.07.005]
- Liu BJ, Gao S, Zhu X, Guo JH, Kou FX, Liu SX, Zhang X, Wang XD, Cao G, Chen H, Liu P, Xu HF, Gao QZ, Yang RJ. Real-world study of 11 hepatic artery infusion chemotherapy combined with anti-PD-1 immunotherapy and tyrosine kinase inhibitors for advanced hepatocellular carcinoma. Immunotherapy 2021; 13: 1395-1405 [PMID: 34607482 DOI: 10.2217/imt-2021-0192]
- Xu Y, Fu S, Mao Y, Huang S, Li D, Wu J. Efficacy and safety of hepatic arterial infusion chemotherapy combined with programmed cell death 12 protein-1 antibody and lenvatinib for advanced hepatocellular carcinoma. Front Med (Lausanne) 2022; 9: 919069 [PMID: 36117969 DOI: 10.3389/fmed.2022.919069]
- 13 Chen LT, Martinelli E, Cheng AL, Pentheroudakis G, Qin S, Bhattacharyya GS, Ikeda M, Lim HY, Ho GF, Choo SP, Ren Z, Malhotra H, Ueno M, Ryoo BY, Kiang TC, Tai D, Vogel A, Cervantes A, Lu SN, Yen CJ, Huang YH, Chen SC, Hsu C, Shen YC, Tabernero J, Yen Y, Hsu CH, Yoshino T, Douillard JY. Pan-Asian adapted ESMO Clinical Practice Guidelines for the management of patients with intermediate and advanced/relapsed hepatocellular carcinoma: a TOS-ESMO initiative endorsed by CSCO, ISMPO, JSMO, KSMO, MOS and SSO. Ann Oncol 2020; **31**: 334-351 [PMID: 32067677 DOI: 10.1016/j.annonc.2019.12.001]
- 14 Reig M, Forner A, Rimola J, Ferrer-Fàbrega J, Burrel M, Garcia-Criado Á, Kelley RK, Galle PR, Mazzaferro V, Salem R, Sangro B, Singal AG, Vogel A, Fuster J, Ayuso C, Bruix J. BCLC strategy for prognosis prediction and treatment recommendation: The 2022 update. J Hepatol 2022; 76: 681-693 [PMID: 34801630 DOI: 10.1016/j.jhep.2021.11.018]
- Rinella ME, Neuschwander-Tetri BA, Siddiqui MS, Abdelmalek MF, Caldwell S, Barb D, Kleiner DE, Loomba R. AASLD Practice Guidance 15 on the clinical assessment and management of nonalcoholic fatty liver disease. Hepatology 2023; 77: 1797-1835 [PMID: 36727674 DOI: 10.1097/HEP.00000000000323]
- He M, Li Q, Zou R, Shen J, Fang W, Tan G, Zhou Y, Wu X, Xu L, Wei W, Le Y, Zhou Z, Zhao M, Guo Y, Guo R, Chen M, Shi M. Sorafenib 16 Plus Hepatic Arterial Infusion of Oxaliplatin, Fluorouracil, and Leucovorin vs Sorafenib Alone for Hepatocellular Carcinoma With Portal Vein Invasion: A Randomized Clinical Trial. JAMA Oncol 2019; 5: 953-960 [PMID: 31070690 DOI: 10.1001/jamaoncol.2019.0250]
- 17 Cheng AL, Qin S, Ikeda M, Galle PR, Ducreux M, Kim TY, Lim HY, Kudo M, Breder V, Merle P, Kaseb AO, Li D, Verret W, Ma N, Nicholas A, Wang Y, Li L, Zhu AX, Finn RS. Updated efficacy and safety data from IMbrave150: Atezolizumab plus bevacizumab vs sorafenib for unresectable hepatocellular carcinoma. J Hepatol 2022; 76: 862-873 [PMID: 34902530 DOI: 10.1016/j.jhep.2021.11.030]
- Qin S, Chan LS, Gu S, Bai Y, Ren Z, Lin X, Chen Z, Jia W, Jin Y, Guo Y, Sultanbaev AV, Pazgan-Simon M, Pisetska M, Liang X, Chen C, 18 Nie Z, Wang L, Cheng AL, Kaseb A, Vogel A. LBA35 Camrelizumab (C) plus rivoceranib (R) vs sorafenib (S) as first-line therapy for unresectable hepatocellular carcinoma (uHCC): A randomized, phase III trial. Conference Abstract. Annals of Oncology 2022; 33: S1401-S1402 [DOI: 10.1016/j.annonc.2022.08.032]
- Finn RS, Kudo M, Merle P, Meyer T, Qin S, Ikeda M, Xu R, Edeline J, Ryoo BY, Ren Z, Cheng AL, Galle PR, Kaneko S, Kumada H, Wang 19 A, Mody K, Dubrovsky L, Siegel AB, Llovet J. LBA34 Primary results from the phase III LEAP-002 study: Lenvatinib plus pembrolizumab vs lenvatinib as first-line (1L) therapy for advanced hepatocellular carcinoma (aHCC). Conference Abstract. Annals of Oncology 2022; 33: S1401 [DOI: 10.1016/j.annonc.2022.08.031]



- Ren Z, Xu J, Bai Y, Xu A, Cang S, Du C, Li Q, Lu Y, Chen Y, Guo Y, Chen Z, Liu B, Jia W, Wu J, Wang J, Shao G, Zhang B, Shan Y, Meng 20 Z, Gu S, Yang W, Liu C, Shi X, Gao Z, Yin T, Cui J, Huang M, Xing B, Mao Y, Teng G, Qin Y, Xia F, Yin G, Yang Y, Chen M, Wang Y, Zhou H, Fan J; ORIENT-32 study group. Sintilimab plus a bevacizumab biosimilar (IBI305) vs sorafenib in unresectable hepatocellular carcinoma (ORIENT-32): a randomised, open-label, phase 2-3 study. Lancet Oncol 2021; 22: 977-990 [PMID: 34143971 DOI: 10.1016/S1470-2045(21)00252-7]
- Yau T, Park JW, Finn RS, Cheng AL, Mathurin P, Edeline J, Kudo M, Harding JJ, Merle P, Rosmorduc O, Wyrwicz L, Schott E, Choo SP, 21 Kelley RK, Sieghart W, Assenat E, Zaucha R, Furuse J, Abou-Alfa GK, El-Khoueiry AB, Melero I, Begic D, Chen G, Neely J, Wisniewski T, Tschaika M, Sangro B. Nivolumab vs sorafenib in advanced hepatocellular carcinoma (CheckMate 459): a randomised, multicentre, open-label, phase 3 trial. Lancet Oncol 2022; 23: 77-90 [PMID: 34914889 DOI: 10.1016/S1470-2045(21)00604-5]
- 22 Chow PKH, Gandhi M, Tan SB, Khin MW, Khasbazar A, Ong J, Choo SP, Cheow PC, Chotipanich C, Lim K, Lesmana LA, Manuaba TW, Yoong BK, Raj A, Law CS, Cua IHY, Lobo RR, Teh CSC, Kim YH, Jong YW, Han HS, Bae SH, Yoon HK, Lee RC, Hung CF, Peng CY, Liang PC, Bartlett A, Kok KYY, Thng CH, Low AS, Goh ASW, Tay KH, Lo RHG, Goh BKP, Ng DCE, Lekurwale G, Liew WM, Gebski V, Mak KSW, Soo KC; Asia-Pacific Hepatocellular Carcinoma Trials Group. SIRveNIB: Selective Internal Radiation Therapy Versus Sorafenib in Asia-Pacific Patients With Hepatocellular Carcinoma. J Clin Oncol 2018; 36: 1913-1921 [PMID: 29498924 DOI: 10.1200/JCO.2017.76.0892]
- Peng Z, Fan W, Zhu B, Wang G, Sun J, Xiao C, Huang F, Tang R, Cheng Y, Huang Z, Liang Y, Fan H, Qiao L, Li F, Zhuang W, Peng B, 23 Wang J, Li J, Kuang M. Lenvatinib Combined With Transarterial Chemoembolization as First-Line Treatment for Advanced Hepatocellular Carcinoma: A Phase III, Randomized Clinical Trial (LAUNCH). J Clin Oncol 2023; 41: 117-127 [PMID: 35921605 DOI: 10.1200/JCO.22.003921
- 24 Qin S, Bi F, Gu S, Bai Y, Chen Z, Wang Z, Ying J, Lu Y, Meng Z, Pan H, Yang P, Zhang H, Chen X, Xu A, Cui C, Zhu B, Wu J, Xin X, Wang J, Shan J, Chen J, Zheng Z, Xu L, Wen X, You Z, Ren Z, Liu X, Qiu M, Wu L, Chen F. Donafenib Versus Sorafenib in First-Line Treatment of Unresectable or Metastatic Hepatocellular Carcinoma: A Randomized, Open-Label, Parallel-Controlled Phase II-III Trial. J Clin Oncol 2021; 39: 3002-3011 [PMID: 34185551 DOI: 10.1200/JCO.21.00163]
- Qin S, Kudo M, Meyer T, Finn RS, Vogel A, Bai Y, Guo Y, Meng Z, Zhang T, Satoh T, Hiraoka A, Marino D, Assenat E, Wyrwicz L, 25 Campos MC, Hsing-Tao K, Boisserie F, Li S, Chen Y, Zhu AX. LBA36 Final analysis of RATIONALE-301: Randomized, phase III study of tislelizumab vs sorafenib as first-line treatment for unresectable hepatocellular carcinoma. Conference Abstract. Annals of Oncology 33: S1402-S1403 [DOI: 10.1016/j.annonc.2022.08.033]
- Kubes P, Jenne C. Immune Responses in the Liver. Annu Rev Immunol 2018; 36: 247-277 [PMID: 29328785 DOI: 26 10.1146/annurev-immunol-051116-052415
- 27 Zhu AX, Finn RS, Edeline J, Cattan S, Ogasawara S, Palmer D, Verslype C, Zagonel V, Fartoux L, Vogel A, Sarker D, Verset G, Chan SL, Knox J, Daniele B, Webber AL, Ebbinghaus SW, Ma J, Siegel AB, Cheng AL, Kudo M; KEYNOTE-224 investigators. Pembrolizumab in patients with advanced hepatocellular carcinoma previously treated with sorafenib (KEYNOTE-224): a non-randomised, open-label phase 2 trial. Lancet Oncol 2018; 19: 940-952 [PMID: 29875066 DOI: 10.1016/S1470-2045(18)30351-6]
- Gauci ML, Lanoy E, Champiat S, Caramella C, Ammari S, Aspeslagh S, Varga A, Baldini C, Bahleda R, Gazzah A, Michot JM, Postel-Vinay 28 S, Angevin E, Ribrag V, Hollebecque A, Soria JC, Robert C, Massard C, Marabelle A. Long-Term Survival in Patients Responding to Anti-PD-1/PD-L1 Therapy and Disease Outcome upon Treatment Discontinuation. Clin Cancer Res 2019; 25: 946-956 [PMID: 30297458 DOI: 10.1158/1078-0432.CCR-18-0793]
- 29 Qin S, Chan SL, Gu S, Bai Y, Ren Z, Lin X, Chen Z, Jia W, Jin Y, Guo Y, Hu X, Meng Z, Liang J, Cheng Y, Xiong J, Ren H, Yang F, Li W, Chen Y, Zeng Y, Sultanbaev A, Pazgan-Simon M, Pisetska M, Melisi D, Ponomarenko D, Osypchuk Y, Sinielnikov I, Yang TS, Liang X, Chen C, Wang L, Cheng AL, Kaseb A, Vogel A; CARES-310 Study Group. Camrelizumab plus rivoceranib vs sorafenib as first-line therapy for unresectable hepatocellular carcinoma (CARES-310): a randomised, open-label, international phase 3 study. Lancet 2023; 402: 1133-1146 [PMID: 37499670 DOI: 10.1016/S0140-6736(23)00961-3]
- Finn RS, Qin S, Ikeda M, Galle PR, Ducreux M, Kim TY, Kudo M, Lim HY, Breder V, Merle P, Kaseb A, Li D, Feng YH, Verret W, 30 Nicholas A, Li L, Ma N, Zhu AX, Cheng AL. IMbrave150: Updated efficacy and safety by risk status in patients (pts) receiving atezolizumab (atezo) + bevacizumab (bev) vs sorafenib (sor) as first-line treatment for unresectable hepatocellular carcinoma (HCC). Conference Abstract. Cancer Research 2021; 81 [DOI: 10.1158/1538-7445.AM2021-CT009]
- 31 Mazzaferro V, Llovet JM, Miceli R, Bhoori S, Schiavo M, Mariani L, Camerini T, Roayaie S, Schwartz ME, Grazi GL, Adam R, Neuhaus P, Salizzoni M, Bruix J, Forner A, De Carlis L, Cillo U, Burroughs AK, Troisi R, Rossi M, Gerunda GE, Lerut J, Belghiti J, Boin I, Gugenheim J, Rochling F, Van Hoek B, Majno P; Metroticket Investigator Study Group. Predicting survival after liver transplantation in patients with hepatocellular carcinoma beyond the Milan criteria: a retrospective, exploratory analysis. Lancet Oncol 2009; 10: 35-43 [PMID: 19058754 DOI: 10.1016/S1470-2045(08)70284-5]
- Yasui Y, Tsuchiya K, Kurosaki M, Takeguchi T, Takeguchi Y, Okada M, Wang W, Kubota Y, Goto T, Komiyama Y, Higuchi M, Takaura K, 32 Hayashi T, Takada H, Tamaki N, Nakanishi H, Itakura J, Takahashi Y, Asahina Y, Enomoto N, Himeno Y, Izumi N. Up-to-seven criteria as a useful predictor for tumor downstaging to within Milan criteria and Child-Pugh grade deterioration after initial conventional transarterial chemoembolization. Hepatol Res 2018; 48: 442-450 [PMID: 29278654 DOI: 10.1111/hepr.13048]
- Wang Q, Xia D, Bai W, Wang E, Sun J, Huang M, Mu W, Yin G, Li H, Zhao H, Li J, Zhang C, Zhu X, Wu J, Gong W, Li Z, Lin Z, Pan X, 33 Shi H, Shao G, Liu J, Yang S, Zheng Y, Xu J, Song J, Wang W, Wang Z, Zhang Y, Ding R, Zhang H, Yu H, Zheng L, Gu W, You N, Wang G, Zhang S, Feng L, Liu L, Zhang P, Li X, Chen J, Xu T, Zhou W, Zeng H, Huang W, Jiang W, Zhang W, Shao W, Li L, Niu J, Yuan J, Lv Y, Li K, Yin Z, Xia J, Fan D, Han G; China HCC-TACE Study Group. Development of a prognostic score for recommended TACE candidates with hepatocellular carcinoma: A multicentre observational study. J Hepatol 2019; 70: 893-903 [PMID: 30660709 DOI: 10.1016/j.jhep.2019.01.013]
- Li QJ, He MK, Chen HW, Fang WQ, Zhou YM, Xu L, Wei W, Zhang YJ, Guo Y, Guo RP, Chen MS, Shi M. Hepatic Arterial Infusion of 34 Oxaliplatin, Fluorouracil, and Leucovorin Versus Transarterial Chemoembolization for Large Hepatocellular Carcinoma: A Randomized Phase III Trial. J Clin Oncol 2022; 40: 150-160 [PMID: 34648352 DOI: 10.1200/JCO.21.00608]
- Yi M, Jiao D, Qin S, Chu Q, Wu K, Li A. Synergistic effect of immune checkpoint blockade and anti-angiogenesis in cancer treatment. Mol 35 Cancer 2019; 18: 60 [PMID: 30925919 DOI: 10.1186/s12943-019-0974-6]
- Zhu H, Shan Y, Ge K, Lu J, Kong W, Jia C. Oxaliplatin induces immunogenic cell death in hepatocellular carcinoma cells and synergizes with 36 immune checkpoint blockade therapy. Cell Oncol (Dordr) 2020; 43: 1203-1214 [PMID: 32797385 DOI: 10.1007/s13402-020-00552-2]
- Tesniere A, Schlemmer F, Boige V, Kepp O, Martins I, Ghiringhelli F, Aymeric L, Michaud M, Apetoh L, Barault L, Mendiboure J, Pignon 37



WJG | https://www.wjgnet.com

JP, Jooste V, van Endert P, Ducreux M, Zitvogel L, Piard F, Kroemer G. Immunogenic death of colon cancer cells treated with oxaliplatin. Oncogene 2010; 29: 482-491 [PMID: 19881547 DOI: 10.1038/onc.2009.356]

- 38 Lyu N, Kong Y, Mu L, Lin Y, Li J, Liu Y, Zhang Z, Zheng L, Deng H, Li S, Xie Q, Guo R, Shi M, Xu L, Cai X, Wu P, Zhao M. Hepatic arterial infusion of oxaliplatin plus fluorouracil/Leucovorin vs sorafenib for advanced hepatocellular carcinoma. J Hepatol 2018; 69: 60-69 [PMID: 29471013 DOI: 10.1016/j.jhep.2018.02.008]
- Liang RB, Zhao Y, He MK, Wen DS, Bu XY, Huang YX, Lai ZC, Xu YJ, Kan A, Wei W, Zhang YJ, Chen MS, Guo RP, Li QJ, Shi M. 39 Hepatic Arterial Infusion Chemotherapy of Oxaliplatin, Fluorouracil, and Leucovorin With or Without Sorafenib as Initial Treatment for Advanced Hepatocellular Carcinoma. Front Oncol 2021; 11: 619461 [PMID: 34055599 DOI: 10.3389/fonc.2021.619461]
- Zheng K, Zhu X, Fu S, Cao G, Li WQ, Xu L, Chen H, Wu D, Yang R, Wang K, Liu W, Wang H, Bao Q, Liu M, Hao C, Shen L, Xing B, 40 Wang X. Sorafenib Plus Hepatic Arterial Infusion Chemotherapy vs Sorafenib for Hepatocellular Carcinoma with Major Portal Vein Tumor Thrombosis: A Randomized Trial. Radiology 2022; 303: 455-464 [PMID: 35103539 DOI: 10.1148/radiol.211545]





Published by Baishideng Publishing Group Inc 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA Telephone: +1-925-3991568 E-mail: office@baishideng.com Help Desk: https://www.f6publishing.com/helpdesk https://www.wjgnet.com

