

World Journal of *Diabetes*

World J Diabetes 2020 September 15; 11(9): 370-399



EDITORIAL

- 370 Is there a role for glucagon-like peptide-1 receptor agonists in the management of diabetic nephropathy?
Veneti S, Tziomalos K

ORIGINAL ARTICLE**Basic Study**

- 374 Analysis of long noncoding RNA-associated competing endogenous RNA network in glucagon-like peptide-1 receptor agonist-mediated protection in β cells
Cui LJ, Bai T, Zhi LP, Liu ZH, Liu T, Xue H, Yang HH, Yang XH, Zhang M, Niu YR, Liu YF, Zhang Y

SYSTEMATIC REVIEWS

- 391 Lower extremity amputations and long-term outcomes in diabetic foot ulcers: A systematic review
Rathnayake A, Saboo A, Malabu UH, Falhammar H

ABOUT COVER

Editorial Board Member of *World Journal of Diabetes*, Dr. She-Yu Li, MD, is an Associate Professor and Specialist of Endocrinology and Diabetology at West China Hospital Sichuan University. He is the co-chief and co-founder of MAGIC China Center and board member of the Diabetes and Obesity Study Group of the Chinese Diabetes Society. He received his MD degree from the West China School of Medicine, Sichuan University in 2012, after which he worked as a visiting researcher at Ninewells Hospital, University of Dundee (Scotland; 2017 to 2019). His research involves electronic medical record-based real-world studies and evidence translation in the fields of diabetes, obesity, and related diseases. These efforts have led to the publication of over 100 papers in peer-reviewed journals. (L-Editor: Filipodia)

AIMS AND SCOPE

The primary aim of *World Journal of Diabetes* (*WJD*, *World J Diabetes*) is to provide scholars and readers from various fields of diabetes with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJD mainly publishes articles reporting research results and findings obtained in the field of diabetes and covering a wide range of topics including risk factors for diabetes, diabetes complications, experimental diabetes mellitus, type 1 diabetes mellitus, type 2 diabetes mellitus, gestational diabetes, diabetic angiopathies, diabetic cardiomyopathies, diabetic coma, diabetic ketoacidosis, diabetic nephropathies, diabetic neuropathies, Donohue syndrome, fetal macrosomia, and prediabetic state.

INDEXING/ABSTRACTING

The *WJD* is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports/Science Edition, PubMed, and PubMed Central. The 2020 Edition of Journal Citation Reports® cites the 2019 impact factor (IF) for *WJD* as 3.247; IF without journal self cites: 3.222; Ranking: 70 among 143 journals in endocrinology and metabolism; and Quartile category: Q2.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Yu-Jie Ma; Production Department Director: Xiang Li; Editorial Office Director: Jia-Ping Yan.

NAME OF JOURNAL

World Journal of Diabetes

ISSN

ISSN 1948-9358 (online)

LAUNCH DATE

June 15, 2010

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Timothy Koch

EDITORIAL BOARD MEMBERS

<https://www.wjnet.com/1948-9358/editorialboard.htm>

PUBLICATION DATE

September 15, 2020

COPYRIGHT

© 2020 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Is there a role for glucagon-like peptide-1 receptor agonists in the management of diabetic nephropathy?

Stavroula Veneti, Konstantinos Tziomalos

ORCID number: Stavroula Veneti 0000-0002-4970-6982; Konstantinos Tziomalos 0000-0002-3172-1594.

Author contributions: Veneti S drafted the editorial; and Tziomalos K critically revised the draft.

Conflict-of-interest statement: None of the authors has any conflict of interest to declare.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Received: July 5, 2020

Peer-review started: July 5, 2020

First decision: July 30, 2020

Revised: August 2, 2020

Accepted: September 3, 2020

Stavroula Veneti, Konstantinos Tziomalos, First Propedeutic Department of Internal Medicine, Medical School, Aristotle University of Thessaloniki, AHEPA Hospital, Thessaloniki 54636, Greece

Corresponding author: Konstantinos Tziomalos, MD, MSc, PhD, Associate Professor, First Propedeutic Department of Internal Medicine, Medical School, Aristotle University of Thessaloniki, AHEPA Hospital, 1 Stilponos Kyriakidi Street, Thessaloniki 54636, Greece. ktziomalos@yahoo.com

Abstract

Chronic kidney disease constitutes a major microvascular complication of diabetes mellitus. Accumulating data suggest that glucagon-like peptide-1 receptor agonists (GLP-1 RAs) might have a role in the management of diabetic kidney disease (DKD). GLP-1 RAs appear to reduce the incidence of persistent macro-albuminuria in patients with type 2 diabetes mellitus. This beneficial effect appears to be mediated not only by the glucose-lowering action of these agents but also on their blood pressure lowering, anti-inflammatory and antioxidant effects. On the other hand, GLP-1 RAs do not appear to affect the rate of decline of glomerular filtration rate. However, this might be due to the relatively short duration of the trials that evaluated their effects on DKD. Moreover, these trials were not designed nor powered to assess renal outcomes. Given that macroalbuminuria is a strong risk factor for the progression of DKD, it might be expected that GLP-1 RAs will prevent the deterioration in renal function in the long term. Nevertheless, this remains to be shown in appropriately designed randomized controlled trials in patients with DKD.

Key Words: Diabetic nephropathy; Type 2 diabetes mellitus; Glucagon-like peptide-1 receptor agonists; Liraglutide; Dulaglutide; Semaglutide

©The Author(s) 2020. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Glucagon-like peptide-1 receptor agonists prevent the development of persistent macroalbuminuria in patients with type 2 diabetes mellitus. However, it is unclear whether they delay the decline in glomerular filtration rate in this population. Long-term trials are needed to clarify the role of these agents in the management of diabetic nephropathy.

Article in press: September 3, 2020

Published online: September 15, 2020

P-Reviewer: Irons B

S-Editor: Huang P

L-Editor: A

P-Editor: Ma YJ



Citation: Veneti S, Tziomalos K. Is there a role for glucagon-like peptide-1 receptor agonists in the management of diabetic nephropathy? *World J Diabetes* 2020; 11(9): 370-373

URL: <https://www.wjgnet.com/1948-9358/full/v11/i9/370.htm>

DOI: <https://dx.doi.org/10.4239/wjcd.v11.i9.370>

INTRODUCTION

Chronic kidney disease (CKD) constitutes a major microvascular complication of diabetes mellitus (DM) and occurs both in type 1 and type 2 DM (T2DM)^[1]. The prevalence of diabetic nephropathy is 50% in patients with type 1 DM and 30%-50% in patients with T2DM^[2]. Diabetic kidney disease (DKD) is characterized by specific structural and functional changes in the kidneys of patients with DM. These changes result in a clinical presentation that includes hypertension, increased urinary albumin excretion and progressive deterioration in kidney function^[1]. It has been estimated that DKD is the leading cause of end-stage renal disease (ESRD) and 30%-40% of patients with DKD are expected to develop ESRD^[3]. More specifically, patients with higher levels of albuminuria, quick deterioration of glomerular filtration rate (GFR), uncontrolled hypertension, long duration of DM, presence of microvascular complications and positive family history of DKD are at higher risk of DKD progression to ESRD^[4]. Importantly, DKD is associated with increased cardiovascular morbidity and mortality^[4]. It has been shown that proteinuria and impaired GFR are independently associated with higher risk of adverse cardiovascular outcomes in patients with T2DM^[5]. The main goals of treatment of DKD are to delay the deterioration of kidney function and to prevent cardiovascular events. Lifestyle measures (*i.e.*, diet and exercise), strict glycemic control and blood pressure control using renin-angiotensin-aldosterone system inhibitors are the cornerstone of DKD treatment^[6].

Accumulating data suggest that glucagon-like peptide-1 receptor agonists (GLP-1 RAs) might have a role in the management of DKD. GLP-1 is secreted by the L-cells of small intestine after food intake and regulates glucose homeostasis^[7]. GLP-1 RAs are divided into short-acting (exenatide, liraglutide and lixisenatide) or long-acting (albiglutide, dulaglutide, exenatide long-acting release and semaglutide)^[8]. GLP-1 RAs induce substantial reductions in glucose levels without the risk of hypoglycemia and also reduce cardiovascular morbidity^[9]. Notably, several randomized, placebo-controlled trials in patients with T2DM and established cardiovascular disease, CKD or multiple cardiovascular risk factors reported a beneficial effect on DKD. In the Liraglutide Effect and Action in Diabetes: Evaluation of Cardiovascular Outcome Results (LEADER) trial ($n = 9340$), liraglutide reduced the incidence of the composite renal outcome (new-onset persistent macroalbuminuria, persistent doubling of the serum creatinine level and an estimated GFR ≤ 45 mL/min/1.73 m², the need for continuous renal-replacement therapy with no reversible cause of the renal disease, or death from renal disease) by 22% compared with placebo during a median follow-up of 3.8 years^[10]. This reduction was driven by the lower incidence of new-onset persistent macroalbuminuria whereas the other endpoints did not differ between patients treated with liraglutide and placebo^[10]. Liraglutide also reduced the incidence of new-onset microalbuminuria by 13%^[10]. Even though GFR declined and albuminuria increased during follow-up in both groups, these changes were smaller in patients treated with liraglutide^[10]. In the Trial to Evaluate Cardiovascular and Other Long-term Outcomes with Semaglutide in Subjects with Type 2 Diabetes (SUSTAIN-6, $n = 3297$), once-weekly semaglutide reduced the risk of new or worsening nephropathy (defined as a new onset of persistent macroalbuminuria, or persistent doubling of serum creatinine level and creatinine clearance) by 36% compared with placebo during a median follow-up of 2.1 years; this benefit was primarily due to the prevention of persistent macroalbuminuria^[11]. In the Researching Cardiovascular Events with a Weekly Incretin in Diabetes (REWIND) trial ($n = 9901$), once-weekly dulaglutide reduced the incidence of the renal component of the composite microvascular outcome (defined as first occurrence of new macroalbuminuria, a sustained decline in estimated GFR $\geq 30\%$ from baseline, or chronic renal replacement therapy) by 15% compared with placebo during a median follow-up of 5.4 years^[12]. Again, this benefit was due to a decreased risk of new macroalbuminuria in patients treated with dulaglutide whereas the incidence of sustained decline in GFR and chronic renal replacement therapy did not differ between the 2 groups^[12]. In a smaller randomized study in 577 patients with moderate-to-severe DKD, dulaglutide had

similar effects on albuminuria with insulin glargine but was associated with higher GFR at 52 wk^[13]. In a recent meta-analysis of 7 placebo-controlled, cardiovascular outcome trials in patients with T2DM ($n = 56004$), treatment with GLP-1 RAs reduced the risk of the composite renal outcome by 17%; again, this benefit was only due to a reduction in the incidence of macroalbuminuria by 24%^[14].

In addition to the glucose-lowering action of GLP-1 RAs, several other mechanisms appear to underpin the effects of these agents on renal function^[15]. GLP-1 RAs lower blood pressure both due to weight loss and due to direct effects on the kidney^[15]. Indeed, it has been reported that GLP-1 RAs promote natriuresis and diuresis due to the inhibition of the sodium-hydrogen exchanger 3, which is located in the renal proximal tubular cells^[16,17]. In addition, preclinical models suggest that GLP-1 RAs exert anti-inflammatory effects and decrease oxidative stress in the kidneys^[18,19].

CONCLUSION

In conclusion, GLP-1 RAs appear to reduce the incidence of persistent macroalbuminuria in patients with T2DM. On the other hand, these agents do not appear to affect the rate of decline of GFR. However, this might be due to the relatively short duration of the trials that evaluated these effects. Moreover, these trials were not designed nor powered to assess renal outcomes. Given that macroalbuminuria is a strong risk factor for the progression of DKD^[5,20], it might be expected that GLP-1 RAs will prevent the deterioration in renal function in the long term. However, this remains to be shown in appropriately designed randomized controlled trials in patients with DKD. The FLOW trial (NCT03819153) is currently evaluating the effects of semaglutide *vs* placebo on the progression of renal impairment in patients with DKD and is expected to be completed in 2024^[21].

REFERENCES

- 1 **Umanath K**, Lewis JB. Update on Diabetic Nephropathy: Core Curriculum 2018. *Am J Kidney Dis* 2018; **71**: 884-895 [PMID: 29398179 DOI: 10.1053/j.ajkd.2017.10.026]
- 2 **Gheith O**, Farouk N, Nampoory N, Halim MA, Al-Otaibi T. Diabetic kidney disease: world wide difference of prevalence and risk factors. *J Nephropharmacol* 2016; **5**: 49-56 [PMID: 28197499]
- 3 **Ninčević V**, Omanović Kolarić T, Roguljić H, Kizivat T, Smolić M, Bilić Čurčić I. Renal Benefits of SGLT 2 Inhibitors and GLP-1 Receptor Agonists: Evidence Supporting a Paradigm Shift in the Medical Management of Type 2 Diabetes. *Int J Mol Sci* 2019; **20** [PMID: 31757028 DOI: 10.3390/ijms20235831]
- 4 **Selby NM**, Taal MW. An updated overview of diabetic nephropathy: Diagnosis, prognosis, treatment goals and latest guidelines. *Diabetes Obes Metab* 2020; **22** Suppl 1: 3-15 [PMID: 32267079 DOI: 10.1111/dom.14007]
- 5 **Ninomiya T**, Perkovic V, de Galan BE, Zoungas S, Pillai A, Jardine M, Patel A, Cass A, Neal B, Poulter N, Mogensen CE, Cooper M, Marre M, Williams B, Hamet P, Mancia G, Woodward M, Macmahon S, Chalmers J; ADVANCE Collaborative Group. Albuminuria and kidney function independently predict cardiovascular and renal outcomes in diabetes. *J Am Soc Nephrol* 2009; **20**: 1813-1821 [PMID: 19443635 DOI: 10.1681/ASN.2008121270]
- 6 **Perkovic V**, Agarwal R, Fioretto P, Hemmelgarn BR, Levin A, Thomas MC, Wanner C, Kasiske BL, Wheeler DC, Groop PH; Conference Participants. Management of patients with diabetes and CKD: conclusions from a "Kidney Disease: Improving Global Outcomes" (KDIGO) Controversies Conference. *Kidney Int* 2016; **90**: 1175-1183 [PMID: 27884312 DOI: 10.1016/j.kint.2016.09.010]
- 7 **Meier JJ**. GLP-1 receptor agonists for individualized treatment of type 2 diabetes mellitus. *Nat Rev Endocrinol* 2012; **8**: 728-742 [PMID: 22945360 DOI: 10.1038/nrendo.2012.140]
- 8 **Cheang JY**, Moyle PM. Glucagon-Like Peptide-1 (GLP-1)-Based Therapeutics: Current Status and Future Opportunities beyond Type 2 Diabetes. *ChemMedChem* 2018; **13**: 662-671 [PMID: 29430842 DOI: 10.1002/cmdc.201700781]
- 9 **Aroda VR**. A review of GLP-1 receptor agonists: Evolution and advancement, through the lens of randomised controlled trials. *Diabetes Obes Metab* 2018; **20** Suppl 1: 22-33 [PMID: 29364586 DOI: 10.1111/dom.13162]
- 10 **Mann JFE**, Ørsted DD, Brown-Frandsen K, Marso SP, Poulter NR, Rasmussen S, Tornøe K, Zinman B, Buse JB; LEADER Steering Committee and Investigators. Liraglutide and Renal Outcomes in Type 2 Diabetes. *N Engl J Med* 2017; **377**: 839-848 [PMID: 28854085 DOI: 10.1056/NEJMoa1616011]
- 11 **Marso SP**, Bain SC, Consoli A, Eliaschewitz FG, Jódar E, Leiter LA, Lingvay I, Rosenstock J, Seufert J, Warren ML, Woo V, Hansen O, Holst AG, Pettersson J, Vilsbøll T; SUSTAIN-6 Investigators. Semaglutide and Cardiovascular Outcomes in Patients with Type 2 Diabetes. *N Engl J Med* 2016; **375**: 1834-1844 [PMID: 27633186 DOI: 10.1056/NEJMoa1607141]
- 12 **Gerstein HC**, Colhoun HM, Dagenais GR, Diaz R, Lakshmanan M, Pais P, Probstfield J, Botros FT, Riddle MC, Rydén L, Xavier D, Atisio CM, Dyal L, Hall S, Rao-Melacini P, Wong G, Avezum A, Basile J, Chung N, Conget I, Cushman WC, Franek E, Hancu N, Hanefeld M, Holt S, Jansky P, Keltai M, Lanis F, Leiter LA, Lopez-Jaramillo P, Cardona Munoz EG, Pirags V, Pogossova N, Raubenheimer PJ, Shaw JE, Sheu WH,

- Temelkova-Kurktschiev T; REWIND Investigators. Dulaglutide and renal outcomes in type 2 diabetes: an exploratory analysis of the REWIND randomised, placebo-controlled trial. *Lancet* 2019; **394**: 131-138 [PMID: 31189509 DOI: 10.1016/S0140-6736(19)31150-X]
- 13 **Tuttle KR**, Lakshmanan MC, Rayner B, Busch RS, Zimmermann AG, Woodward DB, Botros FT. Dulaglutide *versus* insulin glargine in patients with type 2 diabetes and moderate-to-severe chronic kidney disease (AWARD-7): a multicentre, open-label, randomised trial. *Lancet Diabetes Endocrinol* 2018; **6**: 605-617 [PMID: 29910024 DOI: 10.1016/S2213-8587(18)30104-9]
 - 14 **Giugliano D**, Maiorino MI, Bellastella G, Longo M, Chiodini P, Esposito K. GLP-1 receptor agonists for prevention of cardiorenal outcomes in type 2 diabetes: An updated meta-analysis including the REWIND and PIONEER 6 trials. *Diabetes Obes Metab* 2019; **21**: 2576-2580 [PMID: 31373167 DOI: 10.1111/dom.13847]
 - 15 **Greco EV**, Russo G, Giandalia A, Viazzi F, Pontremoli R, De Cosmo S. GLP-1 Receptor Agonists and Kidney Protection. *Medicina (Kaunas)* 2019; **55** [PMID: 31159279 DOI: 10.3390/medicina55060233]
 - 16 **Muskiet MHA**, Tonneijck L, Smits MM, van Baar MJB, Kramer MHH, Hoorn EJ, Joles JA, van Raalte DH. GLP-1 and the kidney: from physiology to pharmacology and outcomes in diabetes. *Nat Rev Nephrol* 2017; **13**: 605-628 [PMID: 28869249 DOI: 10.1038/nrneph.2017.123]
 - 17 **Sun F**, Wu S, Guo S, Yu K, Yang Z, Li L, Zhang Y, Quan X, Ji L, Zhan S. Impact of GLP-1 receptor agonists on blood pressure, heart rate and hypertension among patients with type 2 diabetes: A systematic review and network meta-analysis. *Diabetes Res Clin Pract* 2015; **110**: 26-37 [PMID: 26358202 DOI: 10.1016/j.diabres.2015.07.015]
 - 18 **Fujita H**, Morii T, Fujishima H, Sato T, Shimizu T, Hosoba M, Tsukiyama K, Narita T, Takahashi T, Drucker DJ, Seino Y, Yamada Y. The protective roles of GLP-1R signaling in diabetic nephropathy: possible mechanism and therapeutic potential. *Kidney Int* 2014; **85**: 579-589 [PMID: 24152968 DOI: 10.1038/ki.2013.427]
 - 19 **Kodera R**, Shikata K, Kataoka HU, Takatsuka T, Miyamoto S, Sasaki M, Kajitani N, Nishishita S, Sarai K, Hirota D, Sato C, Ogawa D, Makino H. Glucagon-like peptide-1 receptor agonist ameliorates renal injury through its anti-inflammatory action without lowering blood glucose level in a rat model of type 1 diabetes. *Diabetologia* 2011; **54**: 965-978 [PMID: 21253697 DOI: 10.1007/s00125-010-2028-x]
 - 20 **Fox CS**, Matsushita K, Woodward M, Bilo HJ, Chalmers J, Heerspink HJ, Lee BJ, Perkins RM, Rossing P, Sairenchi T, Tonelli M, Vassalotti JA, Yamagishi K, Coresh J, de Jong PE, Wen CP, Nelson RG; Chronic Kidney Disease Prognosis Consortium. Associations of kidney disease measures with mortality and end-stage renal disease in individuals with and without diabetes: a meta-analysis. *Lancet* 2012; **380**: 1662-1673 [PMID: 23013602 DOI: 10.1016/S0140-6736(12)61350-6]
 - 21 **Novo Nordisk A/S**. A Research Study to See How Semaglutide Works Compared to Placebo in People With Type 2 Diabetes and Chronic Kidney Disease (FLOW). In: ClinicalTrials.gov [Internet]. Bethesda (MD): National Library of Medicine (US). Available from: <https://clinicaltrials.gov/ct2/show/NCT03819153> NLM Identifier: NCT03819153



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

