

World Journal of *Gastrointestinal Surgery*

World J Gastrointest Surg 2023 October 27; 15(10): 2098-2381



Contents

Monthly Volume 15 Number 10 October 27, 2023

MINIREVIEWS

- 2098 Minimally invasive surgery for post cholecystectomy biliary stricture: current evidence and future perspectives
Kalayarasan R, Sai Krishna P
- 2108 From basic to clinical: Anatomy of Denonvilliers' fascia and its application in laparoscopic radical resection of rectal cancer
Chen Z, Zhang XJ, Chang HD, Chen XQ, Liu SS, Wang W, Chen ZH, Ma YB, Wang L

ORIGINAL ARTICLE

Basic Study

- 2115 Effects of thrombopoietin pre-treatment on peri-liver transplantation thrombocytopenia in a mouse model of cirrhosis with hypersplenism
Liu ZR, Zhang YM, Cui ZL, Tong W

Retrospective Cohort Study

- 2123 Effect of low anterior resection syndrome on quality of life in colorectal cancer patients: A retrospective observational study
Jin DA, Gu FP, Meng TL, Zhang XX
- 2133 Stent fracture after transjugular intrahepatic portosystemic shunt placement using the bare metal stent/stent-graft combination technique
Liu QJ, Cao XF, Pei Y, Li X, Dong GX, Wang CM

Retrospective Study

- 2142 Robotic natural orifice specimen extraction surgery I-type F method *vs* conventional robotic resection for lower rectal cancer
Tao F, Liu DN, He PH, Luo X, Xu CY, Li TY, Duan JY
- 2154 Gene polymorphisms associated with sudden decreases in heart rate during extensive peritoneal lavage with distilled water after gastrectomy
Yao S, Yuan Y, Zhang J, Yu Y, Luo GH
- 2171 Analgesic effect of ultrasound-guided bilateral transversus abdominis plane block in laparoscopic gastric cancer
Wang YY, Fu HJ
- 2179 Effects of an Omaha System-based follow-up regimen on self-care and quality of life in gastrointestinal surgery patients
Li YD, Qu N, Yang J, Lv CY, Tang Y, Li P

- 2191** Optimizing surgical outcomes for elderly gallstone patients with a high body mass index using enhanced recovery after surgery protocol
Gu YX, Wang XY, Chen Y, Shao JX, Ni SX, Zhang XM, Shao SY, Zhang Y, Hu WJ, Ma YY, Liu MY, Yu H
- 2201** Establishment and application of three predictive models of anastomotic leakage after rectal cancer sphincter-preserving surgery
Li HY, Zhou JT, Wang YN, Zhang N, Wu SF
- 2211** Identification of multiple risk factors for colorectal cancer relapse after laparoscopic radical resection
Luo J, He MW, Luo T, Lv GQ
- 2222** Examining the impact of early enteral nutritional support on postoperative recovery in patients undergoing surgical treatment for gastrointestinal neoplasms
Chen Z, Hong B, He JJ, Ye QQ, Hu QY
- 2234** Predicting lymph node metastasis in colorectal cancer: An analysis of influencing factors to develop a risk model
Lei YP, Song QZ, Liu S, Xie JY, Lv GQ
- 2247** Novel prognostic score based on the preoperative total bilirubin-albumin ratio and fibrinogen-albumin ratio in ampullary adenocarcinoma
Zhang XJ, Fei H, Sun CY, Li ZF, Li Z, Guo CG, Zhao DB
- 2259** Analysis of textbook outcomes for ampullary carcinoma patients following pancreaticoduodenectomy
Zhang XJ, Fei H, Guo CG, Sun CY, Li ZF, Li Z, Chen YT, Che X, Zhao DB
- 2272** Endoscopic retrograde cholangiopancreatography for diagnosing and treating pediatric biliary and pancreatic diseases
Qin XM, Yu FH, Lv CK, Liu ZM, Wu J

SYSTEMATIC REVIEWS

- 2280** Systematic review of diagnostic tools for peritoneal metastasis in gastric cancer-staging laparoscopy and its alternatives
Ho SYA, Tay KV
- 2294** Prediction of lymph node metastasis in early esophageal cancer
Li Y, Wang JX, Yibi RH
- 2305** Hepatobiliary tuberculosis in the developing world
Esguerra-Paculan MJA, Soldera J

META-ANALYSIS

- 2320** Timing of surgical operation for patients with intra-abdominal infection: A systematic review and meta-analysis
Song SR, Liu YY, Guan YT, Li RJ, Song L, Dong J, Wang PG

- 2331** Bariatric surgery reduces colorectal cancer incidence in obese individuals: Systematic review and meta-analysis

Liu YN, Gu JF, Zhang J, Xing DY, Wang GQ

CASE REPORT

- 2343** Postpolypectomy syndrome without abdominal pain led to sepsis/septic shock and gastrointestinal bleeding: A case report

Chen FZ, Ouyang L, Zhong XL, Li JX, Zhou YY

- 2351** Three-dimensional computed tomography reconstruction diagnosed digestive tract perforation and acute peritonitis caused by *Monopterus albus*: A case report

Yang JH, Lan JY, Lin AY, Huang WB, Liao JY

- 2357** Gastric adenosquamous carcinoma with an elevated serum level of alpha-fetoprotein: A case report

Sun L, Wei JJ, An R, Cai HY, Lv Y, Li T, Shen XF, Du JF, Chen G

- 2362** Mucocutaneous ulcer positive for Epstein-Barr virus, misdiagnosed as a small bowel adenocarcinoma: A case report

Song JH, Choi JE, Kim JS

- 2367** Hereditary hemorrhagic telangiectasia involving portal venous system: A case report and review of the literature

Wu JL, Zhao ZZ, Chen J, Zhang HW, Luan Z, Li CY, Zhao YM, Jing YJ, Wang SF, Sun G

- 2376** Giant dedifferentiated liposarcoma of the gastrocolic ligament: A case report

Kassi ABF, Yenon KS, Kassi FMH, Adjeme AJ, Diarra KM, Bombet-Kouame C, Kouassi M

ABOUT COVER

Editorial Board Member of *World Journal of Gastrointestinal Surgery*, Giuseppe Zimmiti, MD, PhD, Adjunct Professor, Attending Doctor, Postdoctoral Fellow, Surgical Oncologist, Department of General Surgery, Istituto Ospedaliero Fondazione Poliambulanza, Brescia 25124, Lombardia, Italy. giuseppe.zimmiti@poliambulanza.it

AIMS AND SCOPE

The primary aim of *World Journal of Gastrointestinal Surgery* (WJGS, *World J Gastrointest Surg*) is to provide scholars and readers from various fields of gastrointestinal surgery with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGS mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal surgery and covering a wide range of topics including biliary tract surgical procedures, biliopancreatic diversion, colectomy, esophagectomy, esophagostomy, pancreas transplantation, and pancreatectomy, etc.

INDEXING/ABSTRACTING

The WJGS is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports/Science Edition, PubMed, PubMed Central, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJGS as 2.0; IF without journal self cites: 1.9; 5-year IF: 2.2; Journal Citation Indicator: 0.52; Ranking: 113 among 212 journals in surgery; Quartile category: Q3; Ranking: 81 among 93 journals in gastroenterology and hepatology; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Zi-Hang Xu, Production Department Director: Xiang Li, Editorial Office Director: Jia-Ru Fan.

NAME OF JOURNAL

World Journal of Gastrointestinal Surgery

ISSN

ISSN 1948-9366 (online)

LAUNCH DATE

November 30, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Peter Schemmer

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/1948-9366/editorialboard.htm>

PUBLICATION DATE

October 27, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Retrospective Study

Establishment and application of three predictive models of anastomotic leakage after rectal cancer sphincter-preserving surgery

Hui-Yuan Li, Jiang-Tao Zhou, Ya-Nan Wang, Ning Zhang, Shao-Fen Wu

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Pazouki A, Iran;
Pedziwiatr M, Poland

Received: July 12, 2023

Peer-review started: July 12, 2023

First decision: August 2, 2023

Revised: August 9, 2023

Accepted: August 18, 2023

Article in press: August 18, 2023

Published online: October 27, 2023



Hui-Yuan Li, Jiang-Tao Zhou, Ya-Nan Wang, Ning Zhang, Department of General Surgery, Jincheng People's Hospital of Shanxi Province, Jincheng 048026, Shanxi Province, China

Shao-Fen Wu, Department of Gastroenterology, Jincheng People's Hospital of Shanxi Province, Jincheng 048026, Shanxi Province, China

Corresponding author: Shao-Fen Wu, BSc, Nurse, Department of Gastroenterology, Jincheng People's Hospital of Shanxi Province, No. 1168 Baishui East Street, Jincheng 048026, Shanxi Province, China. wushaofen3322@163.com

Abstract

BACKGROUND

Anastomotic leakage (AL) occurs frequently after sphincter-preserving surgery for rectal cancer and has a significant mortality rate. There are many factors that influence the incidence of AL, and each patient's unique circumstances add to this diversity. The early identification and prediction of AL after sphincter-preserving surgery are of great significance for the application of clinically targeted preventive measures. Developing an AL predictive model coincides with the aim of personalised healthcare, enhances clinical management techniques, and advances the medical industry along a more precise and intelligent path.

AIM

To develop nomogram, decision tree, and random forest prediction models for AL following sphincter-preserving surgery for rectal cancer and to evaluate the predictive efficacy of the three models.

METHODS

The clinical information of 497 patients with rectal cancer who underwent sphincter-preserving surgery at Jincheng People's Hospital of Shanxi Province between January 2017 and September 2022 was analyzed in this study. Patients were divided into two groups: AL and no AL. Using univariate and multivariate analyses, we identified factors influencing postoperative AL. These factors were used to establish nomogram, decision tree, and random forest models. The sensitivity, specificity, recall, accuracy, and area under the receiver operating characteristic curve (AUC) were compared between the three models.

RESULTS

AL occurred in 10.26% of the 497 patients with rectal cancer. The nomogram model had an AUC of 0.922, sensitivity of 0.745, specificity of 0.966, accuracy of 0.936, recall of 0.987, and accuracy of 0.946. The above indices in the decision tree model were 0.919, 0.833, 0.862, 0.951, 0.994, and 0.955, respectively and in the random forest model were 1.000, 1.000, 1.000, 0.951, 0.994, and 0.955, respectively. The DeLong test revealed that the AUC value of the decision-tree model was lower than that of the random forest model ($P < 0.05$).

CONCLUSION

The random forest model may be used to identify patients at high risk of AL after sphincter-preserving surgery for rectal cancer owing to its strong predictive effect and stability.

Key Words: Cancer of rectum; Anastomotic leakage; Nomogram; Decision tree; Random forest

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Anastomotic leakage (AL) is a very dangerous complication of rectal cancer surgery, which not only increases the recurrence rate of the tumor but also lowers the quality of life of affected patients. We examined the clinical data of 497 patients with rectal cancer to determine variables that influence AL. We established nomogram, decision tree, and random forest models to identify a prediction model tool for forecasting AL after rectal cancer surgery.

Citation: Li HY, Zhou JT, Wang YN, Zhang N, Wu SF. Establishment and application of three predictive models of anastomotic leakage after rectal cancer sphincter-preserving surgery. *World J Gastrointest Surg* 2023; 15(10): 2201-2210

URL: <https://www.wjgnet.com/1948-9366/full/v15/i10/2201.htm>

DOI: <https://dx.doi.org/10.4240/wjgs.v15.i10.2201>

INTRODUCTION

According to the most recent data from 2020, colorectal cancer has become the third most common cancer worldwide. In China, colorectal cancer has the third highest incidence and fatality rate among all malignancies. Cases of rectal cancer account for 39% of total colorectal cancer cases, making it a serious public health issue in China[1,2]. Radical surgery remains the first choice of treatment for rectal cancer, both for primary and secondary tumors[3]. With advances in medical technology, the prognosis of patients undergoing rectal cancer surgery has significantly improved, and the sphincter preservation rate has also continuously improved[4].

Nevertheless, anastomotic leakage (AL) remains the most common complication following sphincter-preserving surgery for rectal cancer. The perioperative mortality of patients with AL is as high as 18.6%, and these patients are more likely to experience other complications[5,6]. The early identification and prediction of AL after sphincter-preserving surgery are of great significance for the application of clinically targeted preventive measures. At present, a logistic regression analysis of the factors influencing AL in clinical practice is performed[7,8]; however, it cannot intuitively show the importance of each factor to the outcome. Therefore, this study developed nomogram, decision tree, and random forest models for predicting AL following rectal cancer sphincter-preserving surgery. The predictive power of the models were evaluated to identify a tool that would enable the identification of high-risk patients.

MATERIALS AND METHODS

Patient characteristics

For this retrospective analysis, we collected clinical data from 497 patients with rectal cancer who underwent sphincter-preserving surgery at Jincheng People's Hospital of Shanxi Province between January 2017 and September 2022. The inclusion criteria were: Rectal cancer diagnosed by colonoscopy or anal biopsy; tumor within 12 cm of the anal margin; age ≥ 18 years old; and complete clinical data. The exclusion criteria were: Extensive tumor metastasis or the presence of other malignant tumors; a history of rectal surgery, anal stenosis or anal fistula; and conversion to laparotomy. The patients were divided into two groups, those with AL and those without (no AL). The study was approved by the Jincheng People's Hospital of Shanxi Province (JCPH.No20230407001) and written informed consent was obtained from all study participants or their legal guardians.

Study variables

The following variables were analyzed: Sex, age, body mass index, diabetes mellitus, hypertension, smoking history, neoadjuvant treatment, hemoglobin level, albumin (Alb) level, tumor size, tumor-node-metastasis stage, American Society of Anesthesiologists score, tumor location, surgical approach, operative time, and blood loss.

Definition of AL

The diagnosis of AL was based on clinical manifestations (pain, persistent body temperature of $> 38^{\circ}\text{C}$, peritonitis, watery fecal matter, food residue in fecal matter, or pus in the drainage fluid), laboratory tests (elevated white blood cell count and neutrophil percentage), and imaging findings (computed tomography following an enema with a liquid, gas, or water-soluble contrast agent)[9].

Statistical analysis

Statistical analyses were performed using SPSS for Windows version 26.0 (IBM Corp., Armonk, NY, United States). All continuous variable data are presented as mean \pm SD, and student's *t*-tests were used to compare differences. Data from discrete variables are presented as numbers and percentages, and the χ^2 test was used to assess differences between groups. Variables associated with AL were identified using univariate and multivariate logistic regression analyses.

The prediction models were constructed using the R software, and the data were randomly divided between a training set and a verification set in a 7:3 ratio. The nomogram was created using the 'rms' package, the decision tree with the 'rpart' package, and the random forest using the 'random Forest' package. The model with the best predictive effect was selected by comparing the sensitivity, specificity, accuracy, recall rate, precision rate, and area under the receiver operating characteristic curve (AUC). AUCs were compared using the DeLong test. The statistical significance level was set at $P < 0.05$.

RESULTS

Patient characteristics

There were 271 men and 226 women among the 497 patients. The incidence of AL was 10.26% (51/497). Patients in the AL group had a mean age of 60.98 ± 10.83 years. The no AL group included 446 patients with a mean age of 59.27 ± 10.76 years.

Univariate analysis

Univariate analysis revealed statistically significant differences between the AL and no AL groups for the following variables: Sex, diabetes mellitus, smoking history, neoadjuvant treatment, Alb level, tumor size, and tumor location (Table 1).

Influencing factors of AL

Whether the patient had AL after surgery (not occurring = 0, occurring = 1) was used as the dependent variable, and the statistically significant factors identified by univariate analysis and shown in Table 1 (sex, diabetes mellitus, smoking history, neoadjuvant treatment, Alb level, tumor size, and tumor location) were used as independent variables. Table 2 lists the assignments of the indicators. A multivariate logistic regression analysis revealed sex [odds ratio (OR) = 3.656, 95% confidence interval (CI): 1.538-8.264, $P = 0.003$], diabetes mellitus (OR = 5.669, 95%CI: 2.455-13.092, $P < 0.001$), Alb level (OR = 0.898, 95%CI: 0.846-0.953, $P < 0.001$), tumor size (OR = 2.604, 95%CI: 1.840-3.684, $P < 0.001$), and tumor location (OR = 0.272, 95%CI: 0.180-0.413, $P < 0.001$) as factors that influence AL in patients with rectal cancer following sphincter-preserving surgery (Table 3).

Nomogram model

According to the results presented in Table 3, five variables (sex, diabetes mellitus, Alb level, tumor size, and tumor location) were used to construct a nomogram model for predicting AL after sphincter-preserving surgery for rectal cancer (Figure 1).

Decision tree model

A decision tree prediction model for AL after sphincter-preserving surgery for rectal cancer was also constructed, and four explanatory variables were screened: Tumor location, tumor size, Alb level, and sex. The results of the model showed that tumor location was the first-level factor influencing AL in patients with rectal cancer after sphincter-preserving surgery. The incidence of AL was 100% in patients with a distance of the tumor from the anal verge < 2.7 cm, and 78% in male patients with tumor location < 6.3 cm, Alb level < 41 g/L, and tumor size ≥ 5.1 cm (Figure 2).

Random forest model

According to the change in the overall prediction accuracy of the best model, the variables affecting AL in patients with rectal cancer after sphincter-preserving surgery were tumor location, tumor size, diabetes mellitus, sex, and Alb level (Figure 3).

Evaluation of the prediction efficacy of the three models

In the training dataset, the overall performance of the random forest model in predicting AL after sphincter-preserving surgery for rectal cancer was comparable to that of the decision tree model. The AUC of the random forest model was significantly higher than that of the decision tree model ($Z = -2.836$, $P = 0.004$) (Table 4 and Figure 4). In the validation dataset, the overall effectiveness of the three models was equivalent (Table 5 and Figure 5).

Table 1 Single factor analysis of anastomotic leakage, *n* (%)

Patient characteristics	AL (<i>n</i> = 51)	No AL (<i>n</i> = 446)	<i>t</i> / χ^2	<i>P</i> value
Sex			6.921	0.009
Male	33 (64.71)	244 (54.71)		
Female	18 (35.29)	202 (45.29)		
Age, year (mean \pm SD)	60.98 \pm 10.83	59.27 \pm 10.76	1.073	0.284
BMI, kg/m ² (mean \pm SD)	22.30 \pm 2.91	21.45 \pm 3.06	1.896	0.058
Diabetes mellitus			14.164	< 0.001
No	21 (41.18)	302 (67.71)		
Yes	30 (58.82)	144 (32.29)		
Hypertension			0.232	0.630
No	16 (31.37)	155 (34.75)		
Yes	35 (68.63)	291 (65.25)		
Smoking history			6.970	0.008
No	14 (27.45)	209 (46.86)		
Yes	37 (72.55)	237 (53.14)		
Neoadjuvant treatment			7.973	0.005
No	16 (31.37)	233 (52.24)		
Yes	35 (68.63)	213 (47.76)		
Hb, g/L (mean \pm SD)	135.60 \pm 10.46	136.75 \pm 10.41	0.746	0.456
Alb, g/L (mean \pm SD)	33.23 \pm 6.59	37.13 \pm 7.25	3.664	< 0.001
Tumor size, cm (mean \pm SD)	4.73 \pm 1.22	3.42 \pm 1.26	7.009	< 0.001
Tumor location, cm (mean \pm SD)	4.32 \pm 1.28	6.13 \pm 1.30	9.378	< 0.001
TNM stage			0.010	0.995
I	32 (62.75)	281 (63.00)		
II	12 (23.53)	106 (23.77)		
III	7 (13.72)	59 (13.23)		
ASA score			0.289	0.866
I	34 (66.67)	282 (63.23)		
II	12 (23.53)	111 (24.89)		
III	5 (9.80)	53 (11.88)		
Surgical approach			1.676	0.195
Open	13 (25.49)	154 (34.53)		
Laparoscopic	38 (74.51)	292 (65.47)		
Operation time, min (mean \pm SD)	182.19 \pm 6.25	181.87 \pm 5.79	0.378	0.705
Blood loss, mL (mean \pm SD)	230.45 \pm 17.62	232.74 \pm 20.58	0.761	0.447

AL: Anastomotic leakage; Alb: Albumin; ASA: American Society of Anesthesiologists; BMI: Body mass index; Hb: Hemoglobin; TNM: Tumor-node-metastasis.

DISCUSSION

Rectal carcinoma is a prevalent cancer of the digestive system. Currently, the treatment of this disease is mainly surgical [10]. AL is one of the most common and dangerous complications associated with rectal cancer surgery. According to studies, the incidence of AL after rectal cancer surgery ranges from 2.6% to 19.0% [11]. AL not only affects recovery from surgery, but also leads to a variety of complications, such as intra-abdominal abscesses, diffuse peritonitis, and sepsis,

Table 2 Assignment of each factor

Factor	Assignment
Sex	Female = 0, male = 1
Diabetes mellitus	No = 0, yes = 1
Smoking history	No = 0, yes = 1
Neoadjuvant treatment	No = 0, yes = 1
Albumin level	Enter the original value
Tumor size	Enter the original value
Tumor location	Enter the original value

Table 3 Analysis of influencing factors of anastomotic leakage

Factor	β	SE	Wald χ^2	P value	OR (95%CI)
Sex	1.271	0.429	8.778	0.003	3.656 (1.538-8.264)
Diabetes mellitus	1.735	0.427	16.504	< 0.001	5.669 (2.455-13.092)
Smoking history	1.758	0.967	3.309	0.069	5.801 (0.873-38.572)
Neoadjuvant treatment	-0.947	0.940	1.015	0.314	0.388 (0.062-2.448)
Albumin level	-0.108	0.030	12.627	< 0.001	0.898 (0.846-0.953)
Tumor size	0.957	0.177	29.204	< 0.001	2.604 (1.840-3.684)
Tumor location	-1.300	0.212	37.699	< 0.001	0.272 (0.180-0.413)

CI: Confidence interval; OR: Odds ratio.

Table 4 Prediction efficiency of the three model training sets

Model	Sensitivity	Specificity	Accuracy	Recall	Precision	AUC (95%CI)
Nomogram	0.745	0.966	0.936	0.987	0.946	0.922 (0.883-0.961)
Decision tree	0.833	0.862	0.951	0.994	0.955	0.919 (0.863-0.975)
Random forest	1.000	1.000	0.951	0.994	0.955	1.000 (1.000-1.000)

AUC: Area under the receiver operating characteristic curve; CI: Confidence interval.

Table 5 Prediction efficiency of the three model validation sets

Model	Sensitivity	Specificity	Accuracy	Recall	Precision	AUC (95%CI)
Nomogram	0.867	0.909	0.927	1.000	0.921	0.950 (0.908-0.992)
Decision tree	0.836	0.864	0.951	0.994	0.955	0.882 (0.797-0.968)
Random forest	0.984	0.727	0.893	1.000	0.889	0.934 (0.883-0.985)

AUC: Area under the receiver operating characteristic curve; CI: Confidence interval.

and can even cause tumor recurrence. In severe cases, secondary surgery is required, which worsens patient survival rates[12]. Therefore, it is critical to identify the factors that influence the development of AL in patients with rectal cancer after sphincter-preserving surgery and provide targeted interventions to reduce its occurrence.

The incidence of postoperative AL was analyzed in 497 patients with rectal cancer admitted to our hospital. A total of 51 patients developed postoperative AL, representing an incidence of 10.26%. This is similar to the incidence of AL reported by Degiuli *et al*[13] in 5398 patients with rectal cancer (10.2%), but lower than that reported by Peltrini *et al*[14] in 367 patients with rectal cancer (17.4%). These differences may be related to factors such as inclusion criteria and different




Figure 1 Nomogram for predicting anastomotic leakage following rectal cancer surgery.




Figure 2 Decision tree for predicting anastomotic leakage following rectal cancer surgery. AL: Anastomotic leakage.

populations. AL after sphincter-preserving surgery for rectal cancer results from multiple factors, including patient characteristics, tumor status, and operation-related factors. The results of this study showed that sex, diabetes mellitus, Alb level, tumor size, and tumor location influence AL.

According to this study, males were 3.656 times more likely to have postoperative AL than women, which is consistent with the results of most studies[15,16]. As the male pelvis is narrow, the visual field is not fully exposed during the operation, making surgery more challenging; this may cause the rectal stump to retract and cause secondary injury[17].

Penna *et al*[18] confirmed that diabetes mellitus is an independent risk factor for AL after transanal total mesorectal excision. Additionally, this study reported that individuals with diabetes had an AL risk 5.669 times greater than that of non-diabetic patients. Diabetes mellitus can affect the anastomotic blood supply as uncontrolled hyperglycemia leads to vascular damage, reduced blood flow, and cellular accumulation of toxic glucose-derived metabolites, resulting in a significant decrease in anastomotic healing and the ability to fight infection[19].




Figure 3 Importance of variables in the random forest model.




Figure 4 Receiver operating characteristic curves of the three models in the training set. A: Nomogram; B: Decision tree; C: Random forest. AUC: Area under the receiver operating characteristic curve; CI: Confidence interval.

According to the findings of this study, which are consistent with those of Shimura *et al*[20], there is a direct association between preoperative Alb levels and the risk of developing postoperative AL. This may be due to a low perioperative nutritional status leading to reduced immune function and a greater risk of infection and the spread of infection[21]. Yasui *et al*[22] found that patients with tumor sizes ≥ 4 cm were more likely to develop AL. This study also found that the probability of postoperative AL increases with tumor size. As more tissue has to be removed during surgery, more damage occurs and the chance of developing AL subsequently increases[23].

Both domestically and internationally, there is consensus on the impact of tumor location on AL following low anterior resection of rectal cancer; that is, the closer the tumor is to the anal margin, the higher the risk of AL[24-26]. The same conclusion was reached in the present study. This may be because the closer the tumor is to the anus, the larger the wound during resection. In addition, intraoperative electrocoagulation damage to the tissue and blood vessels causes exudation and bleeding, which reduces the blood supply to the anastomosis and increases the risk of postoperative AL [27].

Currently, there is no consensus regarding whether neoadjuvant therapy increases the incidence of AL. Arezzo *et al*[12] analyzed the effects of short- and long-term radiotherapy on AL and showed that the risk of postoperative AL significantly increased in patients receiving short-term radiotherapy. However, Chang *et al*[28] found no difference in the incidence of AL between patients with rectal cancer who underwent chemoradiotherapy before surgery, and those who did not. The present study also found that neoadjuvant therapy had no effect on AL occurrence.

With the development of computer software and artificial intelligence, machine learning has become a new direction for medical research. Studies have used machine learning models to predict the risk of anti-tuberculosis drug-induced liver injury in inpatients with tuberculosis[29], and machine learning can also predict the risk of essential hypertension [30]. In this study, five indicators with statistically significant differences in the multivariate analysis were used to




Figure 5 Receiver operating characteristic curves of the three models in the validation set. A: Nomogram; B: Decision tree; C: Random forest. AUC: Area under the receiver operating characteristic curve; CI: Confidence interval.

establish three prediction models of AL after sphincter-preserving surgery for rectal cancer using machine learning algorithms. In the training dataset, the sensitivity, specificity, accuracy, recall rate, and precision rate of the nomogram were lower than those of the random forest and decision tree models, and the prediction efficacy of the random forest model was better than that of the decision tree. But the sensitivity, specificity, and AUC of the random forest model reached 1, indicating that the random forest model may have overfitting or the generalization effect may be poor due to insufficient data in the training set. In the validation dataset, the three models exhibited similar prediction performances. Although the random forest model has better predictive performance, it also has some disadvantages. As a result, in practice, each of the three prediction models has benefits and limitations, and the most appropriate method should be chosen based on the situation.

As this was a single-center retrospective study, it has several limitations. The representativeness of single-center research is limited, and there may be some bias owing to time constraints. This study did not analyze additional risk factors for AL after sphincter-preserving surgery for rectal cancer, and the constructed model may have been overfitted. Future research will use a larger and more comprehensive sample set and multicenter studies to verify and build a more complete prediction model.

CONCLUSION

Overall, AL is a serious complication of rectal cancer surgery, with a high incidence rate. In this study, nomogram, random forest, and decision tree prediction models of AL after sphincter-preserving surgery for rectal cancer were established using machine learning algorithms. The random forest model was found to have excellent predictive effect and stability, and might serve as a reference for the clinical identification of high-risk groups for AL following sphincter-preserving surgery for rectal cancer.

ARTICLE HIGHLIGHTS

Research background

With advances in medical technology, the success rate of sphincter-preserving surgery in patients with rectal cancer is increasing. However, anastomotic leakage (AL) remains a devastating complication.

Research motivation

AL significantly lowers patients' quality of life. This study examines the elements that influence AL and establishes models to help doctors predict whether patients will develop AL, allowing the timely adoption of preventive measures.

Research objectives

This study aimed to identify the characteristics that influence AL and utilize these factors to build a prediction model for AL after sphincter-preserving surgery for rectal cancer.

Research methods

The clinical data of patients with rectal cancer who underwent sphincter-preserving surgery at our institution in the past five years were examined to analyze the factors influencing AL; nomogram, decision tree, and random forest prediction models were established; and the predictive efficacy of the three models was compared.

Research results

The factors influencing AL after sphincter-preserving surgery for rectal cancer were sex, diabetes mellitus, albumin level, tumor size, and tumor location. To predict the probability of postoperative AL, we constructed nomogram, decision tree, and random forest models.

Research conclusions

This study compared the predictive efficacy of the three prediction models. The random forest model performed the best and may be a useful alternative tool for predicting patients at a high risk of AL.

Research perspectives

Future research will include larger and more comprehensive cohorts across multiple centers, and build a more complete prediction model.

FOOTNOTES

Author contributions: Li HY designed the study and wrote the manuscript; Wu SF designed the study and reviewed the manuscript; Zhou JT, Wang YN, and Zhang N provided clinical advice.

Institutional review board statement: The study was reviewed and approved by the Jincheng People's Hospital of Shanxi Province (JCPH.No20230407001).

Informed consent statement: All study participants or their legal guardians provided written informed consent for personal and medical data collection before study enrolment.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

Data sharing statement: Dataset available from the corresponding author at wushaofen3322@163.com.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Hui-Yuan Li 0009-0004-6470-7088; Jiang-Tao Zhou 0009-0005-5873-7130; Ya-Nan Wang 0009-0004-4231-2165; Ning Zhang 0009-0008-3622-2437; Shao-Fen Wu 0009-0009-6849-723X.

S-Editor: Wang JJ

L-Editor: A

P-Editor: Wang JJ

REFERENCES

- 1 Manceau G, Brouquet A, Chaibi P, Passot G, Bouché O, Mathonnet M, Regimbeau JM, Lo Dico R, Lefèvre JH, Peschard F, Facy O, Volpin E, Chouillard E, Beyert-Berjot L, Verny M, Karoui M, Benoist S. Multicenter phase III randomized trial comparing laparoscopy and laparotomy for colon cancer surgery in patients older than 75 years: the CELL study, a Fédération de Recherche en Chirurgie (FRENCH) trial. *BMC Cancer* 2019; **19**: 1185 [PMID: 31801485 DOI: 10.1186/s12885-019-6376-8]
- 2 Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, Bray F. Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA Cancer J Clin* 2021; **71**: 209-249 [PMID: 33538338 DOI: 10.3322/caac.21660]
- 3 Ghaem H, Amiri Z, Kianpour F, Rezaianzadeh A, Hosseini SV, Khazraei H. Comparing Recurrence and Complications After Laparoscopy and Laparotomy Surgery among Patients Suffering from Colorectal Cancer, Shiraz, Iran. *Asian Pac J Cancer Prev* 2016; **17**: 3111-3116 [PMID: 27509938]
- 4 Jayne D, Pigazzi A, Marshall H, Croft J, Corrigan N, Copeland J, Quirke P, West N, Rautio T, Thomassen N, Tilney H, Gudgeon M, Bianchi PP, Edlin R, Hulme C, Brown J. Effect of Robotic-Assisted vs Conventional Laparoscopic Surgery on Risk of Conversion to Open Laparotomy Among Patients Undergoing Resection for Rectal Cancer: The ROLARR Randomized Clinical Trial. *JAMA* 2017; **318**: 1569-1580 [PMID: 29067426 DOI: 10.1001/jama.2017.7219]
- 5 Chiarello MM, Fransvea P, Cariati M, Adams NJ, Bianchi V, Brisinda G. Anastomotic leakage in colorectal cancer surgery. *Surg Oncol* 2022; **40**: 101708 [PMID: 35092916 DOI: 10.1016/j.suronc.2022.101708]
- 6 Stommel MWJ, Ten Broek RPG, Strik C, Slooter GD, Verhoef C, Grünhagen DJ, van Duijvendijk P, Bemelmans MHA, den Dulk M, Sietses C, van Heek TNT, van den Boezem PB, de Wilt JHW, van Goor H. Multicenter Observational Study of Adhesion Formation After Open-and Laparoscopic Surgery for Colorectal Cancer. *Ann Surg* 2018; **267**: 743-748 [PMID: 28207436 DOI: 10.1097/SLA.0000000000002175]
- 7 Kinugasa T, Nagasu S, Murotani K, Mizobe T, Ochi T, Isobe T, Fujita F, Akagi Y. Analysis of risk factors for anastomotic leakage after lower

- rectal Cancer resection, including drain type: a retrospective single-center study. *BMC Gastroenterol* 2020; **20**: 315 [PMID: [32977772](#) DOI: [10.1186/s12876-020-01462-1](#)]
- 8 **Yu XN**, Xu LM, Bin YW, Yuan Y, Tian SB, Cai B, Tao KX, Wang L, Wang GB, Wang Z. Risk Factors of Anastomotic Leakage After Anterior Resection for Rectal Cancer Patients. *Curr Med Sci* 2022; **42**: 1256-1266 [PMID: [36544033](#) DOI: [10.1007/s11596-022-2616-2](#)]
- 9 **Rahbari NN**, Weitz J, Hohenberger W, Heald RJ, Moran B, Ulrich A, Holm T, Wong WD, Tiet E, Moriya Y, Laurberg S, den Dulk M, van de Velde C, Büchler MW. Definition and grading of anastomotic leakage following anterior resection of the rectum: a proposal by the International Study Group of Rectal Cancer. *Surgery* 2010; **147**: 339-351 [PMID: [20004450](#) DOI: [10.1016/j.surg.2009.10.012](#)]
- 10 **Shiga M**, Maeda H, Oba K, Okamoto K, Namikawa T, Fujisawa K, Yokota K, Kobayashi M, Hanazaki K. Safety of laparoscopic surgery for colorectal cancer in patients over 80 years old: a propensity score matching study. *Surg Today* 2017; **47**: 951-958 [PMID: [28130643](#) DOI: [10.1007/s00595-017-1470-5](#)]
- 11 **Jannasch O**, Klinge T, Otto R, Chiapponi C, Udelnow A, Lippert H, Bruns CJ, Mroczkowski P. Risk factors, short and long term outcome of anastomotic leaks in rectal cancer. *Oncotarget* 2015; **6**: 36884-36893 [PMID: [26392333](#) DOI: [10.18632/oncotarget.5170](#)]
- 12 **Arezzo A**, Migliore M, Chiaro P, Arolo S, Filippini C, Di Cuozzo D, Cirocchi R, Morino M; REAL Score Collaborators. The REAL (REctal Anastomotic Leak) score for prediction of anastomotic leak after rectal cancer surgery. *Tech Coloproctol* 2019; **23**: 649-663 [PMID: [31240416](#) DOI: [10.1007/s10151-019-02028-4](#)]
- 13 **Deguli M**, Elmore U, De Luca R, De Nardi P, Tomatis M, Biondi A, Persiani R, Solaini L, Rizzo G, Soriero D, Cianflocca D, Milone M, Turri G, Rega D, Delrio P, Pedrazzani C, De Palma GD, Borghi F, Scabini S, Coco C, Cavaliere D, Simone M, Rosati R, Reddavid R; collaborators from the Italian Society of Surgical Oncology Colorectal Cancer Network Collaborative Group. Risk factors for anastomotic leakage after anterior resection for rectal cancer (RALAR study): A nationwide retrospective study of the Italian Society of Surgical Oncology Colorectal Cancer Network Collaborative Group. *Colorectal Dis* 2022; **24**: 264-276 [PMID: [34816571](#) DOI: [10.1111/codi.15997](#)]
- 14 **Peltrini R**, Carannante F, Costa G, Bianco G, Garbarino GM, Canali G, Mercantini P, Bracale U, Corcione F, Caricato M, Capolupo GT. Oncological outcomes of rectal cancer patients with anastomotic leakage: A multicenter case-control study. *Front Surg* 2022; **9**: 993650 [PMID: [36171821](#) DOI: [10.3389/fsurg.2022.993650](#)]
- 15 **Tsai KY**, Huang SH, You JF, Tang R, Chiang JM, Yeh CY, Hsieh PS, Tsai WS, Chiang SF, Lai CC. Smoking cessation for less than 10 years remains a risk factor of anastomotic leakage in mid-to-low rectal cancer patients undergoing sphincter-preserving surgery. *Langenbecks Arch Surg* 2022; **407**: 1131-1138 [PMID: [35079889](#) DOI: [10.1007/s00423-021-02381-9](#)]
- 16 **Zhang W**, Lou Z, Liu Q, Meng R, Gong H, Hao L, Liu P, Sun G, Ma J, Zhang W. Multicenter analysis of risk factors for anastomotic leakage after middle and low rectal cancer resection without diverting stoma: a retrospective study of 319 consecutive patients. *Int J Colorectal Dis* 2017; **32**: 1431-1437 [PMID: [28766076](#) DOI: [10.1007/s00384-017-2875-8](#)]
- 17 **Koyama M**, Murata A, Sakamoto Y, Morohashi H, Hasebe T, Saito T, Hakamada K. Risk Factors for Anastomotic Leakage After Intersphincteric Resection Without a Protective Defunctioning Stoma for Lower Rectal Cancer. *Ann Surg Oncol* 2016; **23** Suppl 2: S249-S256 [PMID: [25743332](#) DOI: [10.1245/s10434-015-4461-z](#)]
- 18 **Penna M**, Hompes R, Arnold S, Wynn G, Austin R, Warusavitarne J, Moran B, Hanna GB, Mortensen NJ, Tekkis PP; International TaTME Registry Collaborative. Incidence and Risk Factors for Anastomotic Failure in 1594 Patients Treated by Transanal Total Mesorectal Excision: Results From the International TaTME Registry. *Ann Surg* 2019; **269**: 700-711 [PMID: [29315090](#) DOI: [10.1097/SLA.0000000000002653](#)]
- 19 **Feinberg AE**, Chesney TR, Acuna SA, Sammour T, Quereshty FA. Oncologic Outcomes Following Laparoscopic versus Open Resection of pT4 Colon Cancer: A Systematic Review and Meta-analysis. *Dis Colon Rectum* 2017; **60**: 116-125 [PMID: [27926565](#) DOI: [10.1097/DCR.0000000000000641](#)]
- 20 **Shimura T**, Toiyama Y, Hiro J, Imaoka H, Fujikawa H, Kobayashi M, Ohi M, Inoue Y, Mohri Y, Kusunoki M. Monitoring perioperative serum albumin can identify anastomotic leakage in colorectal cancer patients with curative intent. *Asian J Surg* 2018; **41**: 30-38 [PMID: [27451010](#) DOI: [10.1016/j.asjsur.2016.07.009](#)]
- 21 **Liu YT**, Huang Y, Hao YG, Zhang PF, Yin X, Zhang JF, Hu XH, Li BK, Wang GY. [Current status of influencing factors for postoperative anastomotic leakage in low rectal cancer]. *Zhonghua Wei Chang Wai Ke Za Zhi* 2022; **25**: 1039-1044 [PMID: [36396382](#) DOI: [10.3760/cma.j.cn441530-20220517-00218](#)]
- 22 **Yasui M**, Takemasa I, Miyake Y, Hata T, Ikeda M, Hasegawa J, Ota H, Matsuda C, Mizushima T, Doki Y, Mori M; Clinical Study Group of Osaka University (CSGO), Colorectal Group. Tumor Size as an Independent Risk Factor for Postoperative Complications in Laparoscopic Low Anterior Resection for Advanced Rectal Cancer: A Multicenter Japanese Study. *Surg Laparosc Endosc Percutan Tech* 2017; **27**: 98-103 [PMID: [28141746](#) DOI: [10.1097/SLE.0000000000000377](#)]
- 23 **Yun JA**, Cho YB, Park YA, Huh JW, Yun SH, Kim HC, Lee WY, Chun HK. Clinical manifestations and risk factors of anastomotic leakage after low anterior resection for rectal cancer. *ANZ J Surg* 2017; **87**: 908-914 [PMID: [25925005](#) DOI: [10.1111/ans.13143](#)]
- 24 **Kim CW**, Baek SJ, Hur H, Min BS, Baik SH, Kim NK. Anastomotic Leakage After Low Anterior Resection for Rectal Cancer Is Different Between Minimally Invasive Surgery and Open Surgery. *Ann Surg* 2016; **263**: 130-137 [PMID: [25692355](#) DOI: [10.1097/SLA.0000000000001157](#)]
- 25 **Shi WK**, Qiu XY, Li YH, Lin GL. [Risk factor and early diagnosis of anastomotic leakage after rectal cancer surgery]. *Zhonghua Wei Chang Wai Ke Za Zhi* 2022; **25**: 981-986 [PMID: [36396373](#) DOI: [10.3760/cma.j.cn441530-20220827-00357](#)]
- 26 **Zhou S**, Pei W, Li Z, Zhou H, Liang J, Liu Q, Zhou Z, Wang X. Evaluating the predictive factors for anastomotic leakage after total laparoscopic resection with transrectal natural orifice specimen extraction for colorectal cancer. *Asia Pac J Clin Oncol* 2020; **16**: 326-332 [PMID: [32506809](#) DOI: [10.1111/ajco.13372](#)]
- 27 **Zheng H**, Wu Z, Wu Y, Mo S, Dai W, Liu F, Xu Y, Cai S. Laparoscopic surgery may decrease the risk of clinical anastomotic leakage and a nomogram to predict anastomotic leakage after anterior resection for rectal cancer. *Int J Colorectal Dis* 2019; **34**: 319-328 [PMID: [30470941](#) DOI: [10.1007/s00384-018-3199-z](#)]
- 28 **Chang JS**, Keum KC, Kim NK, Baik SH, Min BS, Huh H, Lee CG, Koom WS. Preoperative chemoradiotherapy effects on anastomotic leakage after rectal cancer resection: a propensity score matching analysis. *Ann Surg* 2014; **259**: 516-521 [PMID: [23598382](#) DOI: [10.1097/SLA.0b013e31829068c5](#)]
- 29 **Zhao P**, Chen J, Yang YQ, Peng Y. [Nomogram model for predicting risk of anti-tuberculosis drug-induced liver injury among inpatients with tuberculosis]. *Zhonghua Yi He He Hu Xi Za Zhi* 2022; **45**: 171-176 [PMID: [35135086](#) DOI: [10.3760/cma.j.cn112147-20210705-00467](#)]
- 30 **Pei Z**, Liu J, Liu M, Zhou W, Yan P, Wen S, Chen Y. Risk-Predicting Model for Incident of Essential Hypertension Based on Environmental and Genetic Factors with Support Vector Machine. *Interdiscip Sci* 2018; **10**: 126-130 [PMID: [29380342](#) DOI: [10.1007/s12539-017-0271-2](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

