

World Journal of *Gastrointestinal Surgery*

World J Gastrointest Surg 2023 December 27; 15(12): 2674-2961



Contents

Monthly Volume 15 Number 12 December 27, 2023

REVIEW

- 2674 Antimicrobial approach of abdominal post-surgical infections
Fiore M, Corrente A, Di Franco S, Alfieri A, Pace MC, Martora F, Petrou S, Mauriello C, Leone S

MINIREVIEWS

- 2693 Indocyanine green fluorescence in gastrointestinal surgery: Appraisal of current evidence
Kalayarasan R, Chandrasekar M, Sai Krishna P, Shanmugam D
- 2709 Post-cholecystectomy iatrogenic bile duct injuries: Emerging role for endoscopic management
Emara MH, Ahmed MH, Radwan MI, Emara EH, Basheer M, Ali A, Elfert AA

ORIGINAL ARTICLE

Case Control Study

- 2719 Multidisciplinary diagnosis and treatment nutritional support intervention for gastrointestinal tumor radiotherapy: Impact on nutrition and quality of life
Hui L, Zhang YY, Hu XD

Retrospective Cohort Study

- 2727 Association between the early high level of serum tacrolimus and recurrence of hepatocellular carcinoma in ABO-incompatible liver transplantation
Han JW, Choi JY, Jung ES, Kim JH, Cho HS, Yoo JS, Sung PS, Jang JW, Yoon SK, Choi HJ, You YK
- 2739 Determining the need for a thoracoscopic approach to treat a giant hiatal hernia when abdominal access is poor
Pérez Lara FJ, Zubizarreta Jimenez R, Prieto-Puga Arjona T, Gutierrez Delgado P, Hernández Carmona JM, Hernández Gonzalez JM, Pitarch Martinez M
- 2747 Predictive value of Hajibandeh index in determining peritoneal contamination in acute abdomen: A cohort study and meta-analysis
Hajibandeh S, Hajibandeh S, Evans L, Miller B, Waterman J, Ahmad SJ, Hale J, Higgi A, Johnson B, Pearce D, Helmy AH, Naguib N, Maw A

Retrospective Study

- 2757 Efficacy of pantoprazole plus perforation repair for peptic ulcer and its effect on the stress response
Leng ZY, Wang JH, Gao L, Shi K, Hua HB
- 2765 Application of electroacupuncture in the prevention of low anterior resection syndrome after rectal cancer surgery
Xu LL, Xiang NJ, Cheng TC, Li YX, Chen P, Jiang ZW, Liu XX

- 2774** Effects of remifentanyl combined with propofol on hemodynamics and oxidative stress in patients undergoing resection of rectal carcinoma
Huang J, Tian WJ
- 2783** Percutaneous microwave ablation and transcatheter arterial chemoembolization for serum tumor markers and prognostics of middle-late primary hepatic carcinoma
Lin ZP, Huang DB, Zou XG, Chen Y, Li XQ, Zhang J
- 2792** Novel invagination procedure for pancreaticojejunostomy using double purse string sutures: A technical note
Li J, Niu HY, Meng XK
- 2799** Laparoscopic resection and endoscopic submucosal dissection for treating gastric ectopic pancreas
Zheng HD, Huang QY, Hu YH, Ye K, Xu JH
- 2809** Prediction of the lymphatic, microvascular, and perineural invasion of pancreatic neuroendocrine tumors using preoperative magnetic resonance imaging
Liu YL, Zhu HB, Chen ML, Sun W, Li XT, Sun YS
- 2820** Impact of hepatectomy and postoperative adjuvant transarterial chemoembolization on serum tumor markers and prognosis in intermediate-stage hepatocellular carcinoma
Hu YD, Zhang H, Tan W, Li ZK

Observational Study

- 2831** Analysis of nutritional risk, skeletal muscle depletion, and lipid metabolism phenotype in acute radiation enteritis
Ma CY, Zhao J, Qian KY, Xu Z, Xu XT, Zhou JY

Randomized Controlled Trial

- 2844** Holistic conditions after colon cancer: A randomized controlled trial of systematic holistic care *vs* primary care
Wang J, Qiao JH

Basic Study

- 2855** Mutational separation and clinical outcomes of *TP53* and *CDH1* in gastric cancer
Liu HL, Peng H, Huang CH, Zhou HY, Ge J
- 2866** Hepatic vagotomy blunts liver regeneration after hepatectomy by downregulating the expression of interleukin-22
Zhou H, Xu JL, Huang SX, He Y, He XW, Lu S, Yao B

META-ANALYSIS

- 2879** Recent evidence for subcutaneous drains to prevent surgical site infections after abdominal surgery: A systematic review and meta-analysis
Ishinuki T, Shinkawa H, Kouzu K, Shinji S, Goda E, Ohyanagi T, Kobayashi M, Kobayashi M, Suzuki K, Kitagawa Y, Yamashita C, Mohri Y, Shimizu J, Uchino M, Haji S, Yoshida M, Ohge H, Mayumi T, Mizuguchi T

- 2890** Prognostic role of serum carcinoembryonic antigen in patients receiving liver resection for colorectal cancer liver metastasis: A meta-analysis

Tang F, Huang CW, Tang ZH, Lu SL, Bai T, Huang Q, Li XZ, Zhang B, Wu FX

- 2907** Significance of carcinoembryonic antigen detection in the early diagnosis of colorectal cancer: A systematic review and meta-analysis

Wang R, Wang Q, Li P

CASE REPORT

- 2919** Primary repair of esophageal atresia gross type C *via* thoracoscopic magnetic compression anastomosis: A case report

Zhang HK, Li XQ, Song HX, Liu SQ, Wang FH, Wen J, Xiao M, Yang AP, Duan XF, Gao ZZ, Hu KL, Zhang W, Lv Y, Zhou XH, Cao ZJ

- 2926** Portal vein embolization for closure of marked arteriportal shunt of hepatocellular carcinoma to enable radioembolization: A case report

Wang XD, Ge NJ, Yang YF

- 2932** Removal of a large rectal polyp with endoscopic submucosal dissection-trans-anal rectoscopic assisted minimally invasive surgery hybrid technique: A case report

Polese L

- 2938** Successful treatment of invasive liver abscess syndrome caused by *Klebsiella variicola* with intracranial infection and septic shock: A case report

Zhang PJ, Lu ZH, Cao LJ, Chen H, Sun Y

- 2945** Duodenojejunostomy treatment of groove pancreatitis-induced stenosis and obstruction of the horizontal duodenum: A case report

Zhang Y, Cheng HH, Fan WJ

- 2954** Awake robotic liver surgery: A case report

Delvecchio A, Pavone G, Conticchio M, Piacente C, Varvara M, Ferraro V, Stasi M, Casella A, Filippo R, Tedeschi M, Pullano C, Inchingolo R, Delmonte V, Memeo R

ABOUT COVER

Editorial Board Member of *World Journal of Gastrointestinal Surgery*, Lapo Bencini, PhD, General Surgery Unit, AOUC, Department of Oncology and Robotics, Careggi University Hospital, Florence 350134, Italy. lapbenc@tin.it

AIMS AND SCOPE

The primary aim of *World Journal of Gastrointestinal Surgery* (WJGS, *World J Gastrointest Surg*) is to provide scholars and readers from various fields of gastrointestinal surgery with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGS mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal surgery and covering a wide range of topics including biliary tract surgical procedures, biliopancreatic diversion, colectomy, esophagectomy, esophagostomy, pancreas transplantation, and pancreatectomy, etc.

INDEXING/ABSTRACTING

The WJGS is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports/Science Edition, PubMed, PubMed Central, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJGS as 2.0; IF without journal self cites: 1.9; 5-year IF: 2.2; Journal Citation Indicator: 0.52; Ranking: 113 among 212 journals in surgery; Quartile category: Q3; Ranking: 81 among 93 journals in gastroenterology and hepatology; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Rui-Rui Wu, Production Department Director: Xu Guo, Editorial Office Director: Jia-Ru Fan.

NAME OF JOURNAL

World Journal of Gastrointestinal Surgery

ISSN

ISSN 1948-9366 (online)

LAUNCH DATE

November 30, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Peter Schemmer

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/1948-9366/editorialboard.htm>

PUBLICATION DATE

December 27, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Recent evidence for subcutaneous drains to prevent surgical site infections after abdominal surgery: A systematic review and meta-analysis

Tomohiro Ishinuki, Hiroji Shinkawa, Keita Kouzu, Seiichi Shinji, Erika Goda, Toshio Ohyanagi, Masahiro Kobayashi, Motomu Kobayashi, Katsunori Suzuki, Yuichi Kitagawa, Chizuru Yamashita, Yasuhiko Mohri, Junzo Shimizu, Motoi Uchino, Seiji Haji, Masahiro Yoshida, Hiroki Ohge, Toshihiko Mayumi, Toru Mizuguchi

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B, B, B
Grade C (Good): C, C
Grade D (Fair): D
Grade E (Poor): E

P-Reviewer: Jiang L, China; Manesis EK, Greece; Meng Y, China; Valek V, Czech Republic; Osman Nuri Dilek, Turkey

Received: July 26, 2023

Peer-review started: July 26, 2023

First decision: October 9, 2023

Revised: October 23, 2023

Accepted: November 27, 2023

Article in press: November 27, 2023

Published online: December 27, 2023



Tomohiro Ishinuki, Toru Mizuguchi, Department of Nursing, Surgical Sciences, Sapporo Medical University, Sapporo 060-8556, Hokkaido, Japan

Hiroji Shinkawa, Department of Hepatobiliary-Pancreatic Surgery, Osaka Metropolitan University Graduate School of Medicine, Abeno-Ku 545-0051, Japan

Keita Kouzu, Department of Surgery, National Defense Medical College, Tokorozawa 359-8513, Japan

Seiichi Shinji, Department of Gastrointestinal and Hepato-Biliary-Pancreatic Surgery, Nippon Medical School, Tokyo 113-8602, Japan

Erika Goda, Department of Nursing, Japan Health Care University, Sapporo 062-0053, Japan

Toshio Ohyanagi, Department of Liberal Arts and Sciences, Center for Medical Education, Sapporo Medical University, Sapporo 060-8556, Japan

Masahiro Kobayashi, Department of Clinical Pharmacokinetics, Research and Education Center for Clinical Pharmacy, Kitasato University, Tokyo 108-8641, Japan

Motomu Kobayashi, Department of Anesthesiology, Hokushinkai Megumino Hospital, Eniwa 061-1395, Japan

Katsunori Suzuki, Department of Infectious Disease Medicine, School of Medicine, University of Occupational and Environmental Health, Kitakyushu 807-8555, Japan

Yuichi Kitagawa, Department of Gastrointestinal Surgery, National Center for Geriatrics and Gerontology, Obu 474-8511, Japan

Chizuru Yamashita, Department of Anesthesiology and Critical Care Medicine, Fujita Health University, Toyoake 470-1192, Japan

Yasuhiko Mohri, Department of Gastrointestinal Surgery, Mie Prefectural General Medical Center, Yokkaichi 510-8561, Japan

Junzo Shimizu, Department of Surgery, Toyonaka Municipal Hospital, Toyonaka 560-8565, Japan

Motoi Uchino, Department of Gastroenterological Surgery, Hyogo Medical University, Nishinomiya 663-8501, Japan

Seiji Haji, Department of Surgery, Soseikai General Hospital, Kyoto 612-8473, Japan

Masahiro Yoshida, Department of Hepato-Biliary-Pancreatic & Gastrointestinal Surgery, International University of Health and Welfare, Ichikawa 272-0827, Japan

Hiroki Ohge, Department of Infectious Disease, Hiroshima University Hospital, Hiroshima 734-8551, Japan

Toshihiko Mayumi, Department of Intensive Care Unit, Chukyo Hospital, Japan Community Health Care Organization, Nagoya 457-8510, Japan

Corresponding author: Toru Mizuguchi, MD, PhD, Doctor, Professor, Surgical Oncologist, Department of Nursing, Surgical Sciences, Sapporo Medical University, S1, W17, Chuo-Ku, Sapporo 060-8556, Hokkaido, Japan. tmizu@sapmed.ac.jp

Abstract

BACKGROUND

Surgical site infections (SSIs) increase mortality, hospital stays, additional medical treatment, and medical costs. Subcutaneous drains prevent SSIs in gynecological and breast surgeries; however, their clinical impact in abdominal surgery remains unclear.

AIM

To investigate whether subcutaneous drains were beneficial in abdominal surgery using a systematic review and meta-analysis.

METHODS

The database search used PubMed, MEDLINE, and the Cochrane Library. The following inclusion criteria were set for the systematic review: (1) Randomized controlled trial studies comparing SSIs after abdominal surgery with or without subcutaneous drains; and (2) Studies that described clinical outcomes, such as SSIs, seroma formation, the length of hospital stays, and mortality.

RESULTS

Eight studies were included in this meta-analysis. The rate of total SSIs was significantly lower in the drained group (54/771, 7.0%) than in the control group (89/759, 11.7%), particularly in gastrointestinal surgery. Furthermore, the rate of superficial SSIs was slightly lower in the drained group (31/517, 6.0%) than in the control group (49/521, 9.4%). No significant differences were observed in seroma formation between the groups. Hospital stays were shorter in the drained group than in the control group.

CONCLUSION

Subcutaneous drains after abdominal surgery prevented SSIs and reduced hospital stays but did not significantly affect seroma formation. The timing of drain removal needs to be reconsidered in future studies.

Key Words: Abdominal surgery; Mortality; Seroma formation; Subcutaneous drain; Surgical site infections

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: This review supports the beneficial effects of subcutaneous drains after abdominal surgery. Subcutaneous drains may not prevent seroma formation. Most studies removed drains within 3 d regardless of the amount of fluid discharge. The timing of drain removal may affect the clinical outcome. Nevertheless, subcutaneous drains can prevent surgical site infections and shorten the length of hospital stays after abdominal surgery.

Citation: Ishinuki T, Shinkawa H, Kouzu K, Shinji S, Goda E, Ohyanagi T, Kobayashi M, Kobayashi M, Suzuki K, Kitagawa Y, Yamashita C, Mohri Y, Shimizu J, Uchino M, Haji S, Yoshida M, Ohge H, Mayumi T, Mizuguchi T. Recent evidence for subcutaneous drains to prevent surgical site infections after abdominal surgery: A systematic review and meta-analysis. *World J Gastrointest Surg* 2023; 15(12): 2879-2889

URL: <https://www.wjgnet.com/1948-9366/full/v15/i12/2879.htm>

DOI: <https://dx.doi.org/10.4240/wjgs.v15.i12.2879>

INTRODUCTION

Surgical site infections (SSIs) are a common complication after gastrointestinal surgery and increase patient mortality[1, 2]. They were previously reported to occur in approximately 20% of patients after colorectal and hepatobiliary surgeries [3-5]. SSIs prolong hospital stays and increase additional medical management with high medical costs[6]. Therefore, their prevention during the perioperative period after surgery is essential.

Fluid collection in subcutaneous tissue is a risk factor for the development of SSIs through its provision of an environment for bacterial growth[7]. The Centers for Disease Control and Prevention guidelines recommend the use of preoperative antimicrobial agents, the removal of serous fluid with drains, including negative pressure wound therapy, and antibacterial-coated absorbent sutures for the prevention of SSIs[8]. The World Health Organization global guidelines also recommend the use of negative pressure wound therapy and antibacterial, absorbent sutures in any surgical procedure[9]. Subcutaneous drains are commonly used after gynecological and breast surgeries[10]; however, conflicting findings have been reported on their clinical impact on SSIs after abdominal surgery[11-15].

The majority of previous studies that examined the clinical benefits of subcutaneous drains were retrospective analyses without cause responsiveness. Therefore, further evidence is needed from randomized controlled trials (RCTs). This systematic review and meta-analysis were designed to compare clinical outcomes among subcutaneous drained and control groups after abdominal surgery.

MATERIALS AND METHODS

Search strategy

The review followed the statement of Preferred Reporting Items for Systematic Reviews and Meta-analyses[16]. The protocol was registered in INPLASY (INPLASY No. 202350115). Ethical approval was not required for this review because of the observational meta-analysis. The protocol was augmented and evaluated with the peer review of electronic search strategies guidelines before performing the search[17]. A comprehensive literature search was conducted using PubMed, MEDLINE, and the Cochrane Library for articles published between January 1, 1999 and April 1, 2023, and the search strategy is described in [Supplementary Table 1](#).

Screening and eligibility

Five independent authors (Ishinuki T, Kouzu K, Kobayashi Mo, Suzuki K, and Yamashita C) reviewed the title and abstract of each article based on the following criteria: (1) RCT studies published in 1999 or thereafter; (2) Patients underwent abdominal surgery, defined as gastrointestinal, hepatobiliary, and hernia repair surgery; (3) Comparisons between subcutaneous drains and no subcutaneous drains; and (4) Incidence of SSIs was presented. Exclusion criteria were duplicate studies and studies in the same database or population. In the case of a conflict of agreement among authors, a group consensus was used to reach acceptance or rejection. Six authors (Ishinuki T, Goda E, Kitagawa Y, Mohri Y, Uchino M, and Haji S) reviewed the full texts of studies that passed the first screening. In addition, references were checked to ensure that all studies were included in eligible studies.

Data selection and quality assessment

Data selection was performed by four independent authors (Ishinuki T, Shinkawa H, Suzuki K, and Yoshida M). A data entry form was prepared in advance. The following information was selected: study details (title, first author, journal name, and year of publication), study design (purpose, study period, eligibility and exclusion criteria, age, and total number of participants), interventions delivered (procedure, surgical site, type of drain, timing of drain removal, length of hospital stays, and type of antimicrobial agents), and study results [number of SSIs, type of SSI (superficial or deep/organ), number of seroma formed, sample size, and *P* value]. The authors independently assessed the risk of bias for studies using the criteria outlined in the Cochrane Handbook for Systematic Reviews of Interventions and Cochrane's effective practice and organization of care guidance, such as random sequence generation, allocation concealment, blinding of participants, blinding of personnel, blinding of outcome assessments, incomplete outcome data, selective reporting, and other bias[18]. Conflicts of opinion among authors were discussed with all authors, and a consensus was reached.

Statistical analysis

The primary outcome was the rate of SSIs, including individual superficial or deep/organ SSIs. Secondary outcomes were the rate of seroma formation, the length of hospital stays, and mortality. Dichotomous data were analyzed for risk ratios using a random effects model and the Mantel-Haenszel method. Continuous data, such as the length of hospital stays, were analyzed using non-parametric tests[19-21]. Data were presented as means with standard deviations using a random effects model. Median and interquartile range data were calculated as means and standard deviations[22]. In addition, a subgroup analysis of SSIs was performed according to each type of drain or surgical site. Data synthesis and statistical analyses were performed with Review Manager (version 5.4; The Nordic Cochrane Centre, The Cochrane Collaboration, Copenhagen, Denmark).

RESULTS

Findings

The Preferred Reporting Items for Systematic Reviews and Meta-analyses flow diagram for this review is shown in [Figure 1](#). Database search results showed that 9462 studies were eligible, with two additional studies being added to the references. We excluded 834 studies because of duplication, and 8621 were excluded after the screening of titles and abstracts. As a result, nine studies were reviewed as full texts[11,19-21,23-27]. Eight studies were ultimately included in the meta-analysis[11,19-21,23-26] after the exclusion of one study without SSI data[27]. The bias summary is shown in [Supplementary Figure 1](#).

Summary and characteristics of studies

A summary of data in each study is shown in [Table 1](#). Five studies disclosed the outcomes of superficial or deep/organ SSIs[19,23,24,26]. Three studies reported the number of seromas formed[21,24,25], and three described the length of hospital stays[19-21]. Although three studies discussed mortality[19,25,26], only one mortality occurred in the drained group; therefore, a further analysis of mortality was impossible[25,26].

The characteristics of each study are shown in [Supplementary Table 2](#). There were three studies on lower gastrointestinal surgery[21,24,26], one on hepatobiliary surgery[19], three on mixed surgery[11,20,23], and one on hernia repair[25]. The types of drains were the Redon drain[23], Hemovac drain[11], Blake drain[19,26], Penrose drain[24], and Tubular drain[25]. Two studies did not specify the type of drain used[20,21]. The timing of drain removal ranged between 2-5 postoperative days or when the drainage volume was less than 40 mL/day. Cefuroxime and metronidazole were used in one study[23], cefazoline in three[11,19,25], cefmetazole in one[24], ampicillin/sulbactam in one[21], and flomoxef in one[26]. One study did not describe the type of antibiotic prophylaxis[20]. An intent-to-treat analysis was performed in three out of seven studies[19,23,25].

Results on the primary outcome

The rate of total SSIs was significantly lower in the drained group (54/771, 7.0%) than in the control group (89/759, 11.7%) without heterogeneity ($I^2 = 0\%$) [risk ratio (RR): 0.63, 95% confidence interval (95%CI): 0.45-0.87, $P = 0.005$] ([Figure 2A](#)). Furthermore, the rate of superficial SSIs was slightly lower in the drained group (31/517, 6.0%) than in the control group (49/521, 9.4%) (RR: 0.64, 95%CI: 0.37-1.11, $P = 0.11$) ([Figure 2B](#)). In addition, no significant differences were observed in deep/organ SSIs between the groups ([Figure 2C](#)). Publication bias of the SSIs was assessed by the funnel plot ([Supplementary Figure 2](#)).

Results on secondary outcomes

No significant differences were observed in seroma formation between the drained and control groups (RR: 1.21, 95%CI: 0.65-2.26, $P = 0.54$) ([Figure 3](#)). However, the length of hospital stays significantly differed (mean difference: -1.34, 95%CI: -2.33 to -0.34, $P = 0.008$) ([Figure 4](#)). Publication bias of the seroma formation and the hospital stays were assessed by the funnel plot ([Supplementary Figures 3 and 4](#), respectively).

Results of subgroup analyses

The results of subgroup analyses of the types of drains are shown in [Figure 5](#). SSIs with the Blake drain (RR: 0.56, 95%CI: 0.24-1.29, $P = 0.18$) and other closed suction methods (RR: 0.71, 95%CI: 0.47-1.07, $P = 0.52$) did not significantly differ from those with the control, whereas SSIs using passive drainage significantly decreased more than with the control (RR: 0.33, 95%CI: 0.11-0.99, $P = 0.05$) ([Figure 5](#)). The results of subgroup analyses of surgical procedures are shown in [Figure 6](#). Subcutaneous drains in gastrointestinal surgery decreased SSIs more than the control (RR: 0.48, 95%CI: 0.26-0.88, $P = 0.02$), whereas those in the hepatobiliary surgery (RR: 0.82, 95%CI: 0.37-1.83, $P = 0.63$), mixed surgery (RR: 0.64, 95%CI: 0.33-1.24, $P = 0.22$), and hernia repair surgery (RR: 0.80, 95%CI: 0.25-2.57, $P = 0.71$) did not ([Figure 6](#)).

DISCUSSION


We identified eight RCTs that compared SSIs after abdominal surgery with or without subcutaneous drains. Although the rate of total SSIs was significantly lower in the drained group than in the control group, no significant differences were observed in superficial or deep/organ SSIs because of statistical power. According to the surgical site, drains effectively prevented SSIs in gastrointestinal surgery. No significant differences were observed in SSIs among the drain types examined. We also showed that seroma formation did not significantly differ between the drained and control groups. On the other hand, subcutaneous drains effectively reduced hospital stays.

Subcutaneous drains remove interstitial fluid from a wound thereby reducing the risk of SSIs[28]. Watanabe *et al*[26] showed the benefits of subcutaneous drains in preventing SSIs after colorectal surgery, and putative reasons for reducing SSIs were proposed. The first reason is the specific shape of the drain itself. The classic shape of a subcutaneous drain is a tube with multiple small holes[11,21,23,25]. This drain type may damage the hypodermis because of high negative pressure from the holes[29]. The Blake drain is a closed suction drain made of a silicone elastomer with a solid core in the center and four slits along the sides. Since the Blake drain was developed to drain without causing tissue damage, it may be more beneficial than other drains[26]. Therefore, this review investigated whether the Blake drain significantly reduced SSIs; however, no significant differences were observed between the drain types examined. The small holes of a

Table 1 Summary of clinical outcomes in each study for the subcutaneous drain group vs the control group after abdominal surgery

Ref.	Year	Drained, <i>n</i>	Control, <i>n</i>	Total SSI, <i>n</i>	Superficial SSI, <i>n</i>	Deep/organ SSI, <i>n</i>	Seroma, <i>n</i>	Hospital stay, mean ± SD	Mortality, <i>n</i>
Baier <i>et al</i> [23]	2010	100	100	10:9	9:6	1:3	ND	ND	ND
Kaya <i>et al</i> [11]	2010	210	192	12:19	ND	ND	ND	ND	ND
Nakayama <i>et al</i> [19]	2014	131	129	10:12	10:12	0:0	ND	26.3 ± 39.7 : 30.3 ± 49.5	1:0
Numata <i>et al</i> [24]	2014	124	122	4:12	4:12	0:0	0:0	ND	ND
Westphalen <i>et al</i> [25]	2015	21	21	4:5	ND	ND	11:9	ND	0:0
Arer <i>et al</i> [20]	2016	23	25	2:8	ND	ND	ND	5.9 ± 2.4 : 7.9 ± 3.8	ND
Lauscher <i>et al</i> [21]	2016	50	53	7:9	4:7	3:2	1:1	8.3 ± 2.3 : 9.3 ± 3.8	ND
Watanabe <i>et al</i> [26]	2017	112	117	5:15	4:12	1:3	ND	ND	0:0

ND: Not described; SD: Standard deviation; SSI: Surgical site infection.



DOI: 10.4240/wjgs.v15.i12.2879 Copyright ©The Author(s) 2023.

Figure 1 Flow diagram of systematic reviews and meta-analyses. SSI: Surgical site infection.

drain may become blocked with fat tissue during the flow of fluid, which eventually prevents drainage[30]. Therefore, drainage methods that are less likely to become blocked need to be developed.

The second proposed reason is the timing of drain removal. Long-term drain placement may increase the risk of contamination[26]. A previous study published in 2004 reported a high infection rate associated with drain placement [31]. On the other hand, studies published from the 2010s demonstrated that a sufficiently long drain time, such as longer than 72 h, prevented wound infection[12,30]. The majority of studies analyzed in the present study removed drains within 72 h, which may have been too short to prevent superficial infection[11,19,21,23,24]. Nakayama *et al*[19] also reported that the timing of drain removal was too early to detect clinical effects and showed that 41% of SSIs occurred within 72 h of drain removal. Watanabe *et al*[26] removed drains 120 h after surgery and found no incidence of SSIs. Therefore, the timing of drain removal needs to be investigated in further studies.




Figure 2 Surgical site infections in the subcutaneous drain group vs the control group after abdominal surgery. A: Total surgical site infections; B: Superficial surgical site infections; C: Deep surgical site infections. CI: Confidence interval.




Figure 3 Seroma formation in the subcutaneous drain group vs the control group after abdominal surgery. CI: Confidence interval.

Although subcutaneous drains significantly decreased the rate of total SSIs, no significant differences were observed in the rate of superficial or deep/organ SSIs. Since two previous studies did not describe data on superficial or deep/organ SSIs, the statistical power in the subgroup analysis may have been lower than that in the total analysis[11,25]. The present study also examined clinical effects, which were slightly better in the drained group than in the control group. This result may be attributed to improvements in drain materials and management.

This review showed that the effectiveness of subcutaneous drains was dependent on the surgical site. The usefulness of subcutaneous drains for the prevention of SSIs has been demonstrated in cholecystectomy[32] and gastrointestinal surgery[11,23]. Baier *et al*[23] reported no benefit of subcutaneous drains in the abdominal field. On the other hand, Kaya *et al*[11] showed a significant reduction in SSIs after surgery for colorectal malignancies and lower abdominal incisions in their subgroup analysis. Previous studies also found that drains prevented SSIs in contaminated surgery, such as lower




Figure 4 Hospital stays in the subcutaneous drain group vs the control group after abdominal surgery. CI: Confidence interval.





Figure 5 Subgroup analysis of surgical site infections depending on drain types. CI: Confidence interval.

gastrointestinal surgery[33,34]. The present results provide support for the use of drains in lower gastrointestinal surgery.

The meta-analysis showed that subcutaneous drains did not significantly prevent seroma formation. Westphalen *et al* [25] noted that seroma formation mostly occurred after drain removal. The peak incidence of seroma formation was previously reported to be approximately 2 wk after surgery, at which time the majority of drains had already been removed[35-37]. Shima *et al*[12] demonstrated that the timing of drain removal in breast cancer surgery was important to avoid wound complications. They found that seroma formation may be prevented when subcutaneous drains were removed after 7 d or later without an increase in SSIs. Therefore, the timing of drain removal needs to be individually selected for specific surgical procedures.

The use of subcutaneous drains affects the length of hospital stays. The length of hospital stays is a socioeconomic factor that is increased by any complication[21]. Hospital stays were prolonged by SSIs, large seromas, hematomas, and other complications[21]. Therefore, the prevention of SSIs by appropriately placing a subcutaneous drain may shorten hospital stays.

To ensure the quality of evidence, this review performed a meta-analysis of RCTs only. However, the interpretation of the present results was limited. The small number of studies included and insufficient published data affected the statistical results obtained. Therefore, a subgroup analysis of factors, such as the material of the drain and the timing of drain removal, was not possible. In addition, surgical techniques may have improved over the study period. Furthermore, putative adverse effects, such as pain, cosmetic evaluations, and management costs, were not analyzed due to the lack of data. It was not possible to investigate a publication bias due to the lack of sufficiently large publications to estimate. A limitation of the present study was that some factors may not have been included in the analysis, such as the age of patients, preoperative conditions, the length of surgery, the surgical technique, and the surgeon's experience.



DOI: 10.4240/wjgs.v15.i12.2879 Copyright ©The Author(s) 2023.

Figure 6 Subgroup analysis of surgical site infections depending on surgical procedures. CI: Confidence interval.

CONCLUSION

In conclusion, the present study supports the use of subcutaneous drains to prevent SSIs and shorten the length of hospital stays after abdominal surgery. On the other hand, subcutaneous drains may not prevent seroma formation. Several specific clinical outcomes, such as the timing of drain removal, pain control, cosmetic evaluations, medical cost, and the quality of life of patients need to be clarified in future studies.

ARTICLE HIGHLIGHTS

Research background

Surgical site infections (SSIs) are a common complication after gastrointestinal surgery and result in a worse clinical prognosis for the patients. Subcutaneous drains are commonly used after abdominal surgery to prevent SSIs. Nevertheless, further evidence is needed about subcutaneous drains. Therefore, this systematic review and meta-analysis were designed to compare clinical outcomes among subcutaneous drains and control groups after abdominal surgery.

Research motivation

There is a shortage of evidence as to whether subcutaneous draining is beneficial in preventing the development of SSIs, and the results are different in each of the previous studies.

Research objectives

The objective for this review was to determine whether subcutaneous drainage is beneficial in preventing the development of SSIs.

Research methods

Independent authors reviewed the previous studies and data selection. The primary outcome was the rate of SSIs, including individual superficial or deep/organ SSIs. Secondary outcomes were the rate of seroma formation, the length of hospital stays, and mortality. Data were presented as means with standard deviations using a random effects model.

Research results

We identified eight RCTs that compared SSIs after abdominal surgery with or without subcutaneous drains. Although the rate of total SSIs was significantly lower in the drained group than in the control group, no significant differences were observed in superficial or deep/organ SSIs because of statistical power. According to the surgical site, drains effectively prevented SSIs in gastrointestinal surgery. No significant differences were observed in SSIs among the drain types examined. We also showed that seroma formation did not significantly differ between the drained and control groups. On the other hand, subcutaneous drains effectively reduced hospital stays.

Research conclusions

This review supports the use of subcutaneous drains to prevent SSIs and shorten the length of hospital stays after abdominal surgery. On the other hand, subcutaneous drains may not prevent seroma formation.

Research perspectives

Impact of subcutaneous drains for pain control, cosmetic evaluations, medical cost, and the quality of life of patients need to be clarified.

FOOTNOTES

Author contributions: Ishinuki T conducted the acquisition of data, abstract evaluation, full-text evaluation, interpretation of data, result discussion, and drafting the article; Shinkawa H conducted the acquisition of data, interpretation of data, and drafting the article; Kouzu K conducted the acquisition of data, abstract evaluation, and result discussion; Shinji S conducted the acquisition of data bias evaluation; Goda E conducted the full-text evaluation and graphic design; Ohyanagi T conducted statistical analyses and graphic design; Kobayashi Ma conducted the acquisition of data and bias evaluation; Kobayashi Mo conducted the abstract evaluation and bias evaluation; Suzuki K conducted the abstract evaluation and interpretation of data; Kitagawa Y conducted the full-text evaluation and bias evaluation; Yamashita C conducted the abstract evaluation and bias evaluation; Mohri Y conducted the full-text evaluation and bias evaluation; Shimizu J conducted the abstract evaluation and bias evaluation; Uchino M conducted the full-text evaluation and graphic design; Haji S conducted the full-text evaluation; Yoshida M conducted the interpretation of data and statistical analyses; Ohge H conducted result discussion; Mayumi T conducted a design of the review and result discussion; Mizuguchi T conducted conception of the review and result discussion.

Supported by Grants-in-Aid from JSPS KAKENHI, No. JP 21K10715 and No. JP 20K10404; Northern Advancement Center for Science & Technology, No. T-2-2; the Yasuda Medical Foundation, No. 31010316; the Okawa Foundation for Information and Telecommunications, No. 41111042; Taiju Life Social Welfare Foundation, No. 50811490; Japan Keirin Autorace Foundation, No. 2023M-378; Project Mirai Cancer Research Grants, No. 31010269; Takahashi Industrial and Economic Research Foundation, No. 50411278; Sapporo Doto Hospital, No. 50311211; Noguchi Hospital, No. 40310551; Doki-kai Tomakomai Hospital, No. 40710739; and Tsuchida Hospital, No. 50811478.

Conflict-of-interest statement: All authors have nothing to disclose.

PRISMA 2009 Checklist statement: The authors have read the PRISMA 2009 Checklist, and the manuscript was prepared and revised according to the PRISMA 2009 Checklist.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Japan

ORCID number: Tomohiro Ishinuki 0000-0003-3225-9781; Hiroji Shinkawa 0000-0002-0164-2414; Keita Kouzu 0000-0002-6433-3184; Seiichi Shinji 0000-0002-9204-6393; Erika Goda 0009-0008-5184-690X; Toshio Ohyanagi 0000-0001-8335-3087; Masahiro Kobayashi 0000-0002-4714-2183; Motomu Kobayashi 0009-0008-6565-3500; Katsunori Suzuki 0000-0002-1906-7577; Yuichi Kitagawa 0009-0001-6463-1018; Chizuru Yamashita 0000-0002-0903-0180; Yasuhiko Mohri 0000-0001-8818-944X; Junzo Shimizu 0000-0002-6613-5341; Motoi Uchino 0000-0002-2934-1680; Seiji Haji 0000-0001-8320-3257; Masahiro Yoshida 0000-0003-3035-3539; Hiroki Ohge 0000-0001-7267-9823; Toshihiko Mayumi 0000-0002-4462-9375; Toru Mizuguchi 0000-0002-8225-7461.

S-Editor: Lin C

L-Editor: Filipodia

P-Editor: Yuan YY

REFERENCES

- 1 **Mangram AJ**, Horan TC, Pearson ML, Silver LC, Jarvis WR. Guideline for Prevention of Surgical Site Infection, 1999. Centers for Disease Control and Prevention (CDC) Hospital Infection Control Practices Advisory Committee. *Am J Infect Control* 1999; **27**: 97-132; quiz 133 [PMID: 10196487]

- 2 **Kirkland KB**, Briggs JP, Trivette SL, Wilkinson WE, Sexton DJ. The impact of surgical-site infections in the 1990s: attributable mortality, excess length of hospitalization, and extra costs. *Infect Control Hosp Epidemiol* 1999; **20**: 725-730 [PMID: [10580621](#) DOI: [10.1086/501572](#)]
- 3 **Romy S**, Eisenring MC, Bettschart V, Petignat C, Francioli P, Troillet N. Laparoscope use and surgical site infections in digestive surgery. *Ann Surg* 2008; **247**: 627-632 [PMID: [18362625](#) DOI: [10.1097/SLA.0b013e3181638609](#)]
- 4 **Nagino M**, Ebata T, Yokoyama Y, Igami T, Sugawara G, Takahashi Y, Nimura Y. Evolution of surgical treatment for perihilar cholangiocarcinoma: a single-center 34-year review of 574 consecutive resections. *Ann Surg* 2013; **258**: 129-140 [PMID: [23059502](#) DOI: [10.1097/SLA.0b013e3182708b57](#)]
- 5 **Uchiyama K**, Ueno M, Ozawa S, Kiriya S, Kawai M, Hirono S, Tani M, Yamaue H. Risk factors for postoperative infectious complications after hepatectomy. *J Hepatobiliary Pancreat Sci* 2011; **18**: 67-73 [PMID: [20676699](#) DOI: [10.1007/s00534-010-0313-1](#)]
- 6 **Haruki K**, Shiba H, Fujiwara Y, Furukawa K, Wakiyama S, Ogawa M, Ishida Y, Misawa T, Yanaga K. Negative impact of surgical site infection on long-term outcomes after hepatic resection for colorectal liver metastases. *Anticancer Res* 2013; **33**: 1697-1703 [PMID: [23564818](#)]
- 7 **Panici PB**, Zullo MA, Casalino B, Angioli R, Muzii L. Subcutaneous drainage vs no drainage after minilaparotomy in gynecologic benign conditions: a randomized study. *Am J Obstet Gynecol* 2003; **188**: 71-75 [PMID: [12548198](#) DOI: [10.1067/mob.2003.103](#)]
- 8 **Berrios-Torres SI**, Umscheid CA, Bratzler DW, Leas B, Stone EC, Kelz RR, Reinke CE, Morgan S, Solomkin JS, Mazuski JE, Dellinger EP, Itani KMF, Berbari EF, Segreti J, Parvizi J, Blanchard J, Allen G, Kluytmans JAJW, Donlan R, Schechter WP; Healthcare Infection Control Practices Advisory Committee. Centers for Disease Control and Prevention Guideline for the Prevention of Surgical Site Infection, 2017. *JAMA Surg* 2017; **152**: 784-791 [PMID: [28467526](#) DOI: [10.1001/jamasurg.2017.0904](#)]
- 9 World Health Organization. Global Guidelines for the Prevention of Surgical Site Infection, second edition. Dec 1, 2018. [cited 15 June 2023]. Available from: <https://www.who.int/publications/i/item/9789241550475>
- 10 **Xue DQ**, Qian C, Yang L, Wang XF. Risk factors for surgical site infections after breast surgery: a systematic review and meta-analysis. *Eur J Surg Oncol* 2012; **38**: 375-381 [PMID: [22421530](#) DOI: [10.1016/j.ejso.2012.02.179](#)]
- 11 **Kaya E**, Paksoy E, Ozturk E, Sigirli D, Bilgel H. Subcutaneous closed-suction drainage does not affect surgical site infection rate following elective abdominal operations: a prospective randomized clinical trial. *Acta Chir Belg* 2010; **110**: 457-462 [PMID: [20919669](#) DOI: [10.1080/00015458.2010.11680655](#)]
- 12 **Shima H**, Kutomi G, Sato K, Kuga Y, Wada A, Satomi F, Uno S, Nisikawa N, Kameshima H, Ohmura T, Mizuguchi T, Takemasa I. An Optimal Timing for Removing a Drain After Breast Surgery: A Systematic Review and Meta-Analysis. *J Surg Res* 2021; **267**: 267-273 [PMID: [34171562](#) DOI: [10.1016/j.jss.2021.05.031](#)]
- 13 **Al-Inany H**, Youssef G, Abd ElMaguid A, Abdel Hamid M, Naguib A. Value of subcutaneous drainage system in obese females undergoing cesarean section using pfannenstiel incision. *Gynecol Obstet Invest* 2002; **53**: 75-78 [PMID: [11961377](#) DOI: [10.1159/000052996](#)]
- 14 **Magann EF**, Chauhan SP, Rodts-Palenik S, Bufkin L, Martin JN Jr, Morrison JC. Subcutaneous stitch closure vs subcutaneous drain to prevent wound disruption after cesarean delivery: a randomized clinical trial. *Am J Obstet Gynecol* 2002; **186**: 1119-1123 [PMID: [12066083](#) DOI: [10.1067/mob.2002.123823](#)]
- 15 **Ramsey PS**, White AM, Guinn DA, Lu GC, Ramin SM, Davies JK, Neely CL, Newby C, Fonseca L, Case AS, Kaslow RA, Kirby RS, Rouse DJ, Hauth JC. Subcutaneous tissue reapproximation, alone or in combination with drain, in obese women undergoing cesarean delivery. *Obstet Gynecol* 2005; **105**: 967-973 [PMID: [15863532](#) DOI: [10.1097/01.AOG.0000158866.68311.d1](#)]
- 16 **Moher D**, Liberati A, Tetzlaff J, Altman DG; PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009; **6**: e1000097 [PMID: [19621072](#) DOI: [10.1371/journal.pmed.1000097](#)]
- 17 **McGowan J**, Sampson M, Salzweid DM, Cogo E, Foerster V, Lefebvre C. PRESS Peer Review of Electronic Search Strategies: 2015 Guideline Statement. *J Clin Epidemiol* 2016; **75**: 40-46 [PMID: [27005575](#) DOI: [10.1016/j.jclinepi.2016.01.021](#)]
- 18 **Higgins JP**, Savović J, Page MJ, Elbers RG, Sterne JA. Assessing risk of bias in a randomized trial. In: Cochrane Handbook for Systematic Reviews of Interventions. 8th ed. Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA, editor. Hoboken: Wiley, 2019 [DOI: [10.1002/9781119536604.ch8](#)]
- 19 **Nakayama H**, Takayama T, Okubo T, Higaki T, Midorikawa Y, Moriguchi M, Aramaki O, Yamazaki S. Subcutaneous drainage to prevent wound infection in liver resection: a randomized controlled trial. *J Hepatobiliary Pancreat Sci* 2014; **21**: 509-517 [PMID: [24519844](#) DOI: [10.1002/jbhp.93](#)]
- 20 **Arer IM**, Yabanoglu H, Aytac HO, Ezer A. The effect of subcutaneous suction drains on surgical site infection in open abdominal surgery A prospective randomized study. *Ann Ital Chir* 2016; **87**: 49-55 [PMID: [27025777](#)]
- 21 **Lauscher JC**, Schneider V, Lee LD, Stroux A, Buhr HJ, Kreis ME, Ritz JP. Necessity of subcutaneous suction drains in ileostomy reversal (DRASTAR)-a randomized, controlled bi-centered trial. *Langenbecks Arch Surg* 2016; **401**: 409-418 [PMID: [27138020](#) DOI: [10.1007/s00423-016-1436-x](#)]
- 22 **Wan X**, Wang W, Liu J, Tong T. Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range. *BMC Med Res Methodol* 2014; **14**: 135 [PMID: [25524443](#) DOI: [10.1186/1471-2288-14-135](#)]
- 23 **Baier PK**, Glück NC, Baumgartner U, Adam U, Fischer A, Hopt UT. Subcutaneous Redon drains do not reduce the incidence of surgical site infections after laparotomy. A randomized controlled trial on 200 patients. *Int J Colorectal Dis* 2010; **25**: 639-643 [PMID: [20140620](#) DOI: [10.1007/s00384-010-0884-y](#)]
- 24 **Numata M**, Godai T, Shirai J, Watanabe K, Inagaki D, Hasegawa S, Sato T, Oshima T, Fujii S, Kunisaki C, Yukawa N, Rino Y, Taguri M, Morita S, Masuda M. A prospective randomized controlled trial of subcutaneous passive drainage for the prevention of superficial surgical site infections in open and laparoscopic colorectal surgery. *Int J Colorectal Dis* 2014; **29**: 353-358 [PMID: [24385026](#) DOI: [10.1007/s00384-013-1810-x](#)]
- 25 **Westphalen AP**, Araújo AC, Zacharias P, Rodrigues ES, Fracaro GB, Lopes Filho Gde J. Repair of large incisional hernias. To drain or not to drain. Randomized clinical trial. *Acta Cir Bras* 2015; **30**: 844-851 [PMID: [26735057](#) DOI: [10.1590/S0102-865020150120000009](#)]
- 26 **Watanabe J**, Ota M, Kawamoto M, Akikazu Y, Suwa Y, Suwa H, Momiyama M, Ishibe A, Watanabe K, Masui H, Nagahori K. A randomized controlled trial of subcutaneous closed-suction Blake drains for the prevention of incisional surgical site infection after colorectal surgery. *Int J Colorectal Dis* 2017; **32**: 391-398 [PMID: [27783162](#) DOI: [10.1007/s00384-016-2687-2](#)]
- 27 **Peiper C**, Conze J, Ponschek N, Schumpelick V. Value of subcutaneous drainage in repair of primary inguinal hernia. A prospective randomized study of 100 cases. *Chirurg* 1997; **68**: 63-67 [PMID: [9132351](#) DOI: [10.1007/s001040050151](#)]
- 28 **Colli A**, Camara ML. First experience with a new negative pressure incision management system on surgical incisions after cardiac surgery in high risk patients. *J Cardiothorac Surg* 2011; **6**: 160 [PMID: [22145641](#) DOI: [10.1186/1749-8090-6-160](#)]
- 29 **Braakenburg A**, Obdeijn MC, Feitz R, van Rooij IA, van Griethuysen AJ, Klinkenbijn JH. The clinical efficacy and cost effectiveness of the

- vacuum-assisted closure technique in the management of acute and chronic wounds: a randomized controlled trial. *Plast Reconstr Surg* 2006; **118**: 390-7; discussion 398 [PMID: [16874208](#) DOI: [10.1097/01.prs.0000227675.63744.af](#)]
- 30 **Tsujita E**, Yamashita Y, Takeishi K, Matsuyama A, Tsutsui S, Matsuda H, Taketomi A, Shirabe K, Ishida T, Maehara Y. Subcuticular absorbable suture with subcutaneous drainage system prevents incisional SSI after hepatectomy for hepatocellular carcinoma. *World J Surg* 2012; **36**: 1651-1656 [PMID: [22411085](#) DOI: [10.1007/s00268-012-1524-1](#)]
- 31 **Liu CL**, Fan ST, Lo CM, Wong Y, Ng IO, Lam CM, Poon RT, Wong J. Abdominal drainage after hepatic resection is contraindicated in patients with chronic liver diseases. *Ann Surg* 2004; **239**: 194-201 [PMID: [14745327](#) DOI: [10.1097/01.sla.0000109153.71725.8c](#)]
- 32 **Chowdri NA**, Qadri SA, Parray FQ, Gagloo MA. Role of subcutaneous drains in obese patients undergoing elective cholecystectomy: a cohort study. *Int J Surg* 2007; **5**: 404-407 [PMID: [17720642](#) DOI: [10.1016/j.ijssu.2007.05.011](#)]
- 33 **Kumar S**, Chatterjee S, Gupta S, Satpathy A, Ray U. Role of subcutaneous closed vacuum drain in preventing surgical site infection in emergency surgery for perforative peritonitis: A randomized control study. *Bangladesh Journal of Medical Science* 2017; **16**: 85-90 [DOI: [10.3329/bjms.v16i1.31138](#)]
- 34 **Pan HD**, Wang L, Peng YF, Li M, Yao YF, Zhao J, Zhan TC, Du CZ, Gu J. Subcutaneous vacuum drains reduce surgical site infection after primary closure of defunctioning ileostomy. *Int J Colorectal Dis* 2015; **30**: 977-982 [PMID: [25700809](#) DOI: [10.1007/s00384-015-2168-z](#)]
- 35 **Baroudi R**, Ferreira CA. Seroma: how to avoid it and how to treat it. *Aesthet Surg J* 1998; **18**: 439-441 [PMID: [19328174](#) DOI: [10.1016/s1090-820x\(98\)70073-1](#)]
- 36 **Nahas FX**, Ferreira LM, Ghelfond C. Does quilting suture prevent seroma in abdominoplasty? *Plast Reconstr Surg* 2007; **119**: 1060-4; discussion 1065 [PMID: [17312514](#) DOI: [10.1097/01.prs.0000242493.11655.68](#)]
- 37 **Bercial ME**, Sabino Neto M, Calil JA, Rossetto LA, Ferreira LM. Suction drains, quilting sutures, and fibrin sealant in the prevention of seroma formation in abdominoplasty: which is the best strategy? *Aesthetic Plast Surg* 2012; **36**: 370-373 [PMID: [21858593](#) DOI: [10.1007/s00266-011-9807-8](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

