

World Journal of *Gastrointestinal Surgery*

World J Gastrointest Surg 2023 June 27; 15(6): 1007-1261



OPINION REVIEW

- 1007 Diverticulitis is a population health problem: Lessons and gaps in strategies to implement and improve contemporary care
Stovall SL, Kaplan JA, Law JK, Flum DR, Simianu VV

REVIEW

- 1020 Distal pancreatectomy with or without radical approach, vascular resections and splenectomy: Easier does not always mean easy
Bencini L, Minuzzo A

MINIREVIEWS

- 1033 Endoscopic ultrasound-guided portal pressure gradient measurement in managing portal hypertension
Lesmana CRA
- 1040 Robotic surgery in elderly patients with colorectal cancer: Review of the current literature
Teo NZ, Ngu JCY
- 1048 Median arcuate ligament syndrome often poses a diagnostic challenge: A literature review with a scope of our own experience
Giakoustidis A, Moschonas S, Christodoulidis G, Chourmouzi D, Diamantidou A, Masoura S, Louri E, Papadopoulos VN, Giakoustidis D
- 1056 Surgical complications of oncological treatments: A narrative review
Fico V, Altieri G, Di Grezia M, Bianchi V, Chiarello MM, Pepe G, Tropeano G, Brisinda G

ORIGINAL ARTICLE**Basic Study**

- 1068 Impact of interstitial cells of Cajal on slow wave and gallbladder contractility in a guinea pig model of acute cholecystitis
Ding F, Guo R, Chen F, Liu LP, Cui ZY, Wang YX, Zhao G, Hu H

Retrospective Cohort Study

- 1080 Fascia- vs vessel-oriented lateral lymph node dissection for rectal cancer: Short-term outcomes and prognosis in a single-center experience
Zhao W, Wang ZJ, Mei SW, Chen JN, Zhou SC, Zhao FQ, Xiao TX, Huang F, Liu Q
- 1093 Prognostic value of 11-factor modified frailty index in postoperative adverse outcomes of elderly gastric cancer patients in China
Xu ZY, Hao XY, Wu D, Song QY, Wang XX

Retrospective Study

- 1104** Long-term outcomes and failure patterns after laparoscopic intersphincteric resection in ultralow rectal cancers
Qiu WL, Wang XL, Liu JG, Hu G, Mei SW, Tang JQ
- 1116** Predictors for success of non-operative management of adhesive small bowel obstruction
Ng ZQ, Hsu V, Tee WWH, Tan JH, Wijesuriya R
- 1125** Preoperative albumin-bilirubin score is a prognostic factor for gastric cancer patients after curative gastrectomy
Szor DJ, Pereira MA, Ramos MFKP, Tustumi F, Dias AR, Zilberstein B, Ribeiro Jr U
- 1138** Ability of lactulose breath test results to accurately identify colorectal polyps through the measurement of small intestine bacterial overgrowth
Li L, Zhang XY, Yu JS, Zhou HM, Qin Y, Xie WR, Ding WJ, He XX
- 1149** Treatment outcome analysis of bevacizumab combined with cyclophosphamide and oxaliplatin in advanced pseudomyxoma peritonei
Zhang Y, Zhao X, Gao C, Lin LY, Li Y
- 1159** Surgical management of duodenal Crohn's disease
Yang LC, Wu GT, Wu Q, Peng LX, Zhang YW, Yao BJ, Liu GL, Yuan LW
- 1169** Influences of dexmedetomidine on stress responses and postoperative cognitive and coagulation functions in patients undergoing radical gastrectomy under general anesthesia
Ma XF, Lv SJ, Wei SQ, Mao BR, Zhao XX, Jiang XQ, Zeng F, Du XK
- 1178** Dissimilar survival and clinicopathological characteristics of mucinous adenocarcinoma located in pancreatic head and body/tail
Li Z, Zhang XJ, Sun CY, Li ZF, Fei H, Zhao DB

SYSTEMATIC REVIEWS

- 1191** Gallbladder perforation with fistulous communication
Quiroga-Garza A, Alvarez-Villalobos NA, Muñoz-Leija MA, Garcia-Campa M, Angeles-Mar HJ, Jacobo-Baca G, Elizondo-Omana RE, Guzman-Lopez S

META-ANALYSIS

- 1202** Efficacy of transanal drainage tube in preventing anastomotic leakage after surgery for rectal cancer: A meta-analysis
Fujino S, Yasui M, Ohue M, Miyoshi N

CASE REPORT

- 1211** Percutaneous transhepatic cholangial drainage-guided methylene blue for fistulotomy using dual-knife for bile duct intubation: A case report
Tang BX, Li XL, Wei N, Tao T

- 1216** Optimal resection of gastric bronchogenic cysts based on anatomical continuity with adherent gastric muscular layer: A case report
Terayama M, Kumagai K, Kawachi H, Makuuchi R, Hayami M, Ida S, Ohashi M, Sano T, Nunobe S
- 1224** Intrahepatic cholangiocarcinoma in patients with primary sclerosing cholangitis and ulcerative colitis: Two case reports
Miyazu T, Ishida N, Asai Y, Tamura S, Tani S, Yamade M, Iwaizumi M, Hamaya Y, Osawa S, Baba S, Sugimoto K
- 1232** Massive bleeding from a gastric artery pseudoaneurysm in hepatocellular carcinoma treated with atezolizumab plus bevacizumab: A case report
Pang FW, Chen B, Peng DT, He J, Zhao WC, Chen TT, Xie ZG, Deng HH
- 1240** Bedside ultrasound-guided water injection assists endoscopically treatment in esophageal perforation caused by foreign bodies: A case report
Wei HX, Lv SY, Xia B, Zhang K, Pan CK
- 1247** Modified stomach-partitioning gastrojejunostomy for initially unresectable advanced gastric cancer with outlet obstruction: A case report
Shao XX, Xu Q, Wang BZ, Tian YT
- 1256** Small bowel diverticulum with enterolith causing intestinal obstruction: A case report
Wang LW, Chen P, Liu J, Jiang ZW, Liu XX

ABOUT COVER

Editorial Board Member of *World Journal of Gastrointestinal Surgery*, Danko Mikulic, FEBS, MD, PhD, Assistant Professor, Surgeon, Department of Surgery, University Hospital Merkur, Zagreb 10000, Croatia.
danko.mikulic@zg.t-com.hr

AIMS AND SCOPE

The primary aim of *World Journal of Gastrointestinal Surgery* (*WJGS, World J Gastrointest Surg*) is to provide scholars and readers from various fields of gastrointestinal surgery with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGS mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal surgery and covering a wide range of topics including biliary tract surgical procedures, biliopancreatic diversion, colectomy, esophagectomy, esophagostomy, pancreas transplantation, and pancreatectomy, etc.

INDEXING/ABSTRACTING

The *WJGS* is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports/Science Edition, PubMed, PubMed Central, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for *WJGS* as 2.505; IF without journal self cites: 2.473; 5-year IF: 3.099; Journal Citation Indicator: 0.49; Ranking: 104 among 211 journals in surgery; Quartile category: Q2; Ranking: 81 among 93 journals in gastroenterology and hepatology; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Rui-Rui Wu, Production Department Director: Xiang Li, Editorial Office Director: Jia-Ru Fan.

NAME OF JOURNAL

World Journal of Gastrointestinal Surgery

ISSN

ISSN 1948-9366 (online)

LAUNCH DATE

November 30, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Peter Schemmer

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/1948-9366/editorialboard.htm>

PUBLICATION DATE

June 27, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Retrospective Study

Ability of lactulose breath test results to accurately identify colorectal polyps through the measurement of small intestine bacterial overgrowth

Lan Li, Xue-Yuan Zhang, Jin-Sheng Yu, Hui-Min Zhou, Yan Qin, Wen-Rui Xie, Wen-Jing Ding, Xing-Xiang He

Specialty type: Medicine, research and experimental

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Bordonaro M, United States; Pavlidis TE, Greece

Received: February 19, 2023

Peer-review started: February 19, 2023

First decision: March 24, 2023

Revised: April 5, 2023

Accepted: April 18, 2023

Article in press: April 18, 2023

Published online: June 27, 2023



Lan Li, Xue-Yuan Zhang, Hui-Min Zhou, Yan Qin, Wen-Rui Xie, Xing-Xiang He, Department of Gastroenterology, The First Affiliated Hospital of Guangdong Pharmaceutical University, Guangzhou 510080, Guangdong Province, China

Lan Li, Hui-Min Zhou, Yan Qin, Wen-Rui Xie, Xing-Xiang He, Washing Microbiota Transplantation, Engineering Techniques of Microbiota-Targeted Therapies of Guangdong Province, Guangzhou 510080, Guangdong Province, China

Xue-Yuan Zhang, Department of Gastroenterology, People's Hospital of Nanxiong County, Nanxiong 512400, Guangdong Province, China

Jin-Sheng Yu, Department of Genetics, Washington University School of Medicine, Saint Louis, MO 63110, United States

Wen-Jing Ding, North America Medical Education Foundation, California, CA 91710, United States

Corresponding author: Xing-Xiang He, MD, PhD, Chief Doctor, Professor, Department of Gastroenterology, The First Affiliated Hospital of Guangdong Pharmaceutical University, No. 19 Nonglinxia Road, Guangzhou 510080, Guangdong Province, China.

hexingxiang@gdpu.edu.cn

Abstract

BACKGROUND

While colorectal polyps are not cancerous, some types of polyps, known as adenomas, can develop into colorectal cancer over time. Polyps can often be found and removed by colonoscopy; however, this is an invasive and expensive test. Thus, there is a need for new methods of screening patients at high risk of developing polyps.

AIM

To identify a potential association between colorectal polyps and small intestine bacteria overgrowth (SIBO) or other relevant factors in a patient cohort with lactulose breath test (LBT) results.

METHODS

A total of 382 patients who had received an LBT were classified into polyp and non-polyp groups that were confirmed by colonoscopy and pathology. SIBO was diagnosed by measuring LBT-derived hydrogen (H) and methane (M) levels according to 2017 North American Consensus recommendations. Logistic regression was used to assess the ability of LBT to predict colorectal polyps. Intestinal barrier function damage (IBFD) was determined by blood assays.

RESULTS

H and M levels revealed that the prevalence of SIBO was significantly higher in the polyp group than in the non-polyp group (41% *vs* 23%, $P < 0.01$; 71% *vs* 59%, $P < 0.05$, respectively). Within 90 min of lactulose ingestion, the peak H values in the adenomatous and inflammatory/hyperplastic polyp patients were significantly higher than those in the non-polyp group ($P < 0.01$, and $P = 0.03$, respectively). In 227 patients with SIBO defined by combining H and M values, the rate of IBFD determined by blood lipopolysaccharide levels was significantly higher among patients with polyps than those without (15% *vs* 5%, $P < 0.05$). In regression analysis with age and gender adjustment, colorectal polyps were most accurately predicted with models using M peak values or combined H and M values limited by North American Consensus recommendations for SIBO. These models had a sensitivity of ≥ 0.67 , a specificity of ≥ 0.64 , and an accuracy of ≥ 0.66 .

CONCLUSION

The current study made key associations among colorectal polyps, SIBO, and IBFD and demonstrated that LBT has moderate potential as an alternative noninvasive screening tool for colorectal polyps.

Key Words: Lactulose breath test; Colorectal polyp; Small intestine bacteria overgrowth; Intestinal barrier function; Retrospective study

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: As the lactulose breath test (LBT) is an indirect method of measuring bacteria in the digestive tract, it is primarily used to support small intestine bacteria overgrowth (SIBO) diagnosis but is implemented as a new method for screening colorectal polyps in this study. A total of 382 patients with LBT results were classified into polyp and non-polyp groups that were confirmed by colonoscopy and pathology. First, it applied the LBT for assessment of its utility as a noninvasive screening tool for colorectal polyps as well as for diagnosis of SIBO. Second, the results revealed certain key associations among colorectal polyps, SIBO and Intestinal barrier function damage (IBFD), such as SIBO was more prevalent in patients with colorectal polyp than those without polyp and IBFD was more susceptible in patients with colorectal polyp than those without polyp only when SIBO was evident. Third, in regression analysis with age and gender adjustment, colorectal polyp was best predicted by models using plain methane peak values or combined hydrogen and methane values limited by the North American Consensus for SIBO. One of the most important result was moderate potential of LBT as an alternative noninvasive screening tool for colorectal polyps.

Citation: Li L, Zhang XY, Yu JS, Zhou HM, Qin Y, Xie WR, Ding WJ, He XX. Ability of lactulose breath test results to accurately identify colorectal polyps through the measurement of small intestine bacterial overgrowth. *World J Gastrointest Surg* 2023; 15(6): 1138-1148

URL: <https://www.wjgnet.com/1948-9366/full/v15/i6/1138.htm>

DOI: <https://dx.doi.org/10.4240/wjgs.v15.i6.1138>

INTRODUCTION

Colorectal polyps are caused by colorectal mucosal proliferation that creates pedunculated or sessile outgrowths. They become more common as people age and are prevalent in individuals > 40 years of age[1,2]. While most polyps are benign, some can become cancerous and may even metastasize to other parts of the body[3]. Adenomatous polyps are known precursors of colon cancer but can be difficult to diagnose in their early stages. Moreover, most colorectal cancers develop from focal changes in benign polyps through a multistep process involving genetic, histological, morphological, and intestinal microbiome changes that accumulate over more than 10 years[4,5]. A long precancerous state provides an opportunity to screen for polyps and successfully prevent or treat any cancerous lesions that develop. Thus, new methods that can identify precancerous colorectal lesions can play an important role

in early-stage colorectal cancer treatment and prevention.

Several methods are used to screen for colorectal cancer, including fecal occult blood testing (FOBT), flexible sigmoidoscopy, and colonoscopy, each with its own merits and disadvantages. A pooled meta-analysis of randomized trials found that FOBT and flexible sigmoidoscopy screening reduced colorectal cancer mortality by 16% and 30%, respectively[6]. While colonoscopy is the best method for visualizing focal lesions and taking biopsies for diagnosis[7], it is invasive, costly, and can be uncomfortable, especially for asymptomatic participants with low compliance. Thus, colonoscopy may not be suitable for primary screening of colorectal polyps and cancers. Despite the benefits of these modalities, there are overwhelming limitations, which highlight a need for more accurate, noninvasive screening tools for colorectal cancer and precancerous polyps.

The lactulose breath test (LBT) is an indirect method of measuring bacteria in the digestive tract. It uses equipment to determine the concentration in parts per million (ppm) of hydrogen (H) and methane (M) gas in the breath[8]. The LBT can indicate the approximate population size and location of the microbiome, as well as some information about the types of bacteria present. While jejunal aspiration culture remains the gold standard for diagnosing small intestine bacterial overgrowth (SIBO), LBT is widely used as a noninvasive method of diagnosing SIBO due to its safety, accessibility, and affordability. However, there is limited data on the association between SIBO and colorectal polyps.

SIBO is a condition in which the small bowel is colonized by excessive aerobic and anaerobic microbes that are normally present in the colon[9]. SIBO and intestinal microbiota are associated with several conditions, including Crohn's disease[10], irritable bowel syndrome (IBS)[11], functional gastrointestinal disorders (FGID)[12], nonalcoholic fatty liver disease[13], diabetes[14], and hepatic encephalopathy[15]. Recent studies have found a link between the gut microbiome and the pathogenesis of adenomatous polyps and colorectal cancer[16,17], offering a promising avenue for personalized prevention[18]. For example, higher numbers of some bacterial species are found in patients with adenomatous polyps than in those without[17]. The current study analyzed a patient cohort with LBT testing data to characterize potential associations among colorectal polyps, SIBO, and other relevant factors and assessed the use of LBT as a potential screening tool for colorectal polyps.

MATERIALS AND METHODS

Study subjects

Medical records from patients in registry database of The First Affiliated Hospital of Guangdong Pharmaceutical University who received an LBT for gastrointestinal symptoms from July 2017 to February 2019 were reviewed. A total of 382 patients (213 males and 169 females) were included in the study. The subjects ranged in age from 22 to 92 years (mean \pm SD, 57 \pm 14 years). All patients signed an informed consent prior to inclusion. The study was approved by the Ethics Committee of the First Affiliated Hospital of Guangdong Pharmaceutical University.

The patients were classified into a polyp group ($n = 169$) and a non-polyp group ($n = 213$). All colorectal polyps were diagnosed by colonoscopy and pathology. Patients with no polyps or other intestinal lesions identified by colonoscopy were included in the non-polyp group. Individuals with: (1) Acute intestinal infection; (2) antibiotic use within 4 wk before the test; (3) severe heart, lung, brain, and other diseases who are unable to tolerate colonoscopy; (4) susceptibility to hypoglycemia; and (5) age < 18 years were excluded from the study.

Blood assays to evaluate intestinal barrier function damage

Intestinal barrier function damage (IBFD) was assessed using the instruments and assay kits from Beijing Zhongsheng Jinyu Diagnostic Technology Co., Ltd. Blood samples were taken after 8 h of fasting and stored at 4 °C. Within 4 h, the blood samples were tested for diamine oxidase (DAO), D-lactate (D-lac), and lipopolysaccharide (LPS) concentration according to the manufacturer's instructions. Patients whose blood samples had values above the reference for DAO (10 U/L), D-lac (15 mg/L), and LPS (20 U/L) were defined as having IBFD.

Lactulose breath test

The LBT was completed using the Quintron Breath Tracker (SC model) to determine the concentration of H and M. Procedures were performed with common standards[9,19]. In brief, all patients fasted for 12 h and brushed their teeth prior to the test. Lactulose (10 g) in warm water was provided and breath samples were collected every 30 min for 150 min. No drink, food, or exercise was permitted during the test, but subjects were allowed to sleep.

Diagnosis of SIBO and prediction of colorectal polyps by LBT

Diagnosis of SIBO by LBT was made qualitatively according to the following definitions of a positive result recommended by the 2017 North American Consensus[19]: (1) A rise of > 20 ppm H within 90 min of substrate ingestion; and (2) ≥ 10 ppm methane. A patient was determined as having SIBO if either or

both standards were met.

LBT quantitative measurements were also used to predict the presence of colorectal polyps. The performance of prediction models was assessed with logistic regression supported by the R program, pROC. Each model was tested by 100-time repeated re-sampling to ensure its accuracy.

Statistical analyses

All data were tested for statistical differences using IBM SPSS software (v22.0). An ANOVA test was used to assess differences in measurements between groups, and a one-side Fisher Exact test was used to measure differences in frequency between one group and another. $P < 0.05$ was considered statistically significant.

RESULTS

Characteristics of patients and colorectal polyps

As shown in [Table 1](#), patients in the polyp group were 9 years older than those in the non-polyp group (mean 62 *vs* 53 years, $P < 0.001$), and were more often male (64% *vs* 49%, $P < 0.01$). Colorectal polyps were least prevalent among patients 19-45 years of age (7%) and most common among those 61-92 years of age (55%). The polyp group also had a higher proportion of patients with constipation than the non-polyp group (22% *vs* 14%, $P < 0.05$), and more often had metabolic disorders, including diabetes (19% *vs* 10%, $P < 0.01$), hyperlipidemia (20% *vs* 13%, $P < 0.05$), and fatty liver/cirrhosis (41% *vs* 27%, $P < 0.01$), in addition to hypertension (38% *vs* 21%, $P < 0.001$). However, patients in the polyp group were less likely to have non-organic disorders, such as FGID (5% *vs* 13%, $P < 0.01$), IBS (8% *vs* 16%, $P < 0.05$) and gastroesophageal reflux disease (11% *vs* 16%, $P = 0.096$), than those in the non-polyp group.

Colonoscopy showed that the colorectal mucosa from 213 patients had a normal appearance, while colorectal polyps were found in 169 patients, including 81 with inflammatory/hyperplastic polyps, and 88 with adenomatous polyps. Polyp size was < 1.0 cm in 136 patients, 1.0-2.0 cm in 25 patients, and > 2.0 cm in eight patients. While 71 patients had single polyps, 98 patients had multiple polyps. In 114 patients, the polyps were found on the left side of the colon, including on the descending colon, sigmoid colon, and rectum, and in 55 patients, polyps were located on other parts of the colon.

Ability of LBT to detect SIBO and predict colorectal polyps

According to H, M measurements, alone or in combination, the prevalence of SIBO by LBT was all significantly higher in the polyp group than in the non-polyp group [H: 41% (70/169) *vs* 23% (49/213), $P < 0.001$; M: 71% (120/169) *vs* 59% (125/213), $P < 0.05$; combined: 80% (136/169) *vs* 67% (143/213), $P < 0.01$] ([Table 2](#)). Within 90 min of substrate ingestion, the peak values of hydrogen were significantly higher in patients with adenomatous or inflammatory/hyperplastic polyps than those in the non-polyp group ($P < 0.01$, and $P = 0.03$, respectively; [Table 3](#)). The peak values of methane were similar in all three groups ($P = 0.168$), and there was no significant difference in the number of patients with SIBO by polyp type ($P > 0.05$).

Associations between IBFD, SIBO, and colorectal polyps

A total of 311 of the 382 patients were evaluated for IBFD by blood assays, including measurements of DAO, D-lac, and LPS. Of these, 174 (56%) of the patients, including 82 in the polyp group (58%) and 92 in the non-polyp group (54%), were characterized as having potential IBFD using a combination of the three assays ($P > 0.05$) or each assay alone (all $P > 0.05$). Of the 311 patients, 227 were positive for SIBO using combined H and M measurements. Among patients with SIBO, the rate of IBFD using all three blood assays was marginally higher in the polyp group than in the non-polyp group (57% *vs* 48%, $P = 0.13$), but differed significantly when IBFD was defined using LPS alone (polyp = 15% *vs* non-polyp = 5%, $P < 0.05$; [Figure 1](#)). Among the remaining 84 patients without SIBO, there was no significant difference in the rate of IBFD between patients in the polyp and non-polyp groups using all three assays together or individually (all $P > 0.05$).

Prediction performance of LBT results for colorectal polyps

LBT was also assessed quantitatively for its prediction performance as a screening tool for colorectal polyps. Using different H and M cutoff values, 17 models were built using different subsets of the patient population ([Table 4](#)). Peak values in H and M were obtained during the tests and rise values were got from baseline subtracted peak values. To account for the effects of age and gender on the model performance, 7 of the 17 models with differences in the mean LBT value between the polyp and non-polyp groups ($P < 0.01$) were selected for further assessment (model # 1, 2, 4, 5, 11, 12 and 17; [Table 4](#) and [Figure 2A-G](#)). Differences in the area under the receiver operating characteristic curve between age and gender-adjusted and unadjusted models were statistically significant (all $P < 0.01$). These models performed similarly well when age and gender were used as covariates, with almost all of them showing an accuracy of $> 65\%$ ([Table 5](#)). Models with a methane peak value with or without a ≥ 5

Table 1 Demographics and comorbidity of the study subjects, *n* (%)

	Overall (<i>N</i> = 382)	Polyps (<i>n</i> = 169)	Non-polyps (<i>n</i> = 213)	<i>P</i> value
Age, yr	57.0 ± 14.0	62.1 ± 11.7	53.0 ± 14.4	0
19-45	70 (18)	11 (7)	59 (28)	0
46-60	150 (39)	65 (38)	85 (40)	0.428
61-92	162 (43)	93 (55)	69 (32)	0
Male	213 (56)	108 (64)	105 (49)	0.003
Bilestone	34 (9)	18 (11)	16 (8)	0.187
Constipation	67 (18)	38 (22)	29 (14)	0.017
Diabetes	53 (14)	32 (19)	21 (10)	0.008
Fatty liver/cirrhosis	127 (33)	70 (41)	57 (27)	0.002
FGID	37 (10)	9 (5)	28 (13)	0.007
GERD	54 (14)	19 (11)	35 (16)	0.096
Hyperlipidemia	60 (16)	33 (20)	27 (13)	0.046
Hypertension	108 (28)	64 (38)	44 (21)	0
Hyperuricemia	42 (11)	20 (12)	22 (10)	0.38
IBS	49 (13)	14 (8)	35 (16)	0.013
PU	31 (8)	24 (14)	7 (3)	0

Values presented as mean ± SD, or *n* (%) of observations. *P* values were from one-side Fisher exact statistics, with bold font for those less than 0.05. FGID: Functional gastrointestinal disorders; GERD: Gastroesophageal reflux disease; IBS: Irritable bowel syndrome; PU: Peptic ulcer.

Table 2 Small intestine bacteria overgrowth distribution between polyp & non-polyp groups

SIBO (+)	Overall (<i>N</i> = 382)	Polyps (<i>n</i> = 169)	Non-polyps (<i>n</i> = 213)	<i>P</i> value
By methane	245 (64)	120 (71)	125 (59)	0.014 ^a
By hydrogen 90 min	119 (31)	70 (41)	49 (23)	0.000 ^a
By combined M and H	279 (73)	136 (80)	143 (67)	0.004 ^a

^a*P* < 0.05, polyps vs non-polyps.

SIBO: Small intestine bacteria overgrowth.

Table 3 The peak values of methane and hydrogen in inflammatory/hyperplastic polyp, adenomatous polyp and non-polyp groups

Peak values	Polyps (<i>n</i> = 169)		Non-polyps (<i>n</i> = 213)		<i>P</i> value
	Inflammatory/hyperplastic polyp	Adenomatous polyp	Inflammatory/hyperplastic polyp	Adenomatous polyp	
Methane	208.2	197.86	182.52	182.52	0.168
Hydrogen within 90 min	209.53 ^b	220.87 ^b	172.51	172.51	0.001 ^a

^a*P* < 0.05 was considered to indicate a statistically significant difference between inflammatory/hyperplastic polyp, adenomatous polyp and non-polyp groups.

^b*P* < 0.05 was considered to indicate a statistically significant difference between inflammatory/hyperplastic polyp and non-polyp groups or between adenomatous polyp and non-polyp groups. Univariate analysis was performed using the nonparametric tests (Kruskal-Wallis independent samples).

ppm cutoff (Figure 2A and D) and the model using the SIBO subpopulation (Figure 2G) performed best.

Table 4 Performance of prediction models for colorectal polyps

Model No.	Value cutoff (ppm) for subset	N	n (polyp/non-polyp)	Mean ppm (polyp)	Mean ppm (non-polyp)	Mean ppm (P value)	Sensitivity	Specificity	Accuracy
1	Methane peak value (not applied)	382	169/213	12.82	12.06	0.074^a	0.427	0.578	0.511
2	Methane peak value (≥ 5)	359	160/199	13.34	12.66	0.084^a	0.417	0.571	0.502
3	Methane peak value (≥ 10)	245	120/125	15.28	15.55	0.905	0.443	0.390	0.416
4	Hydrogen peak value (not applied)	380	169/211	52.34	45.48	0.091^a	0.408	0.604	0.517
5	Hydrogen peak value (≥ 10)	310	140/170	62.05	55.24	0.090^a	0.383	0.590	0.496
6	Hydrogen peak value (≥ 20)	259	125/134	67.85	66.02	0.776	0.400	0.439	0.421
7	Hydrogen rise value (not applied)	372	165/207	41.94	37.40	0.121	0.391	0.582	0.498
8	Hydrogen rise value (≥ 10)	280	130/150	52.12	49.93	0.406	0.380	0.477	0.432
9	Hydrogen rise value (≥ 20)	217	108/109	60.06	63.39	0.479	0.492	0.375	0.433
10	Hydrogen rise value (≥ 20 by 90 min)	119	70/49	47.74	50.10	0.929	0.460	0.323	0.403
11	Combined M peak & H peak (M ≥ 5 &/or H ≥ 10)	373	166/207	66.15	58.48	0.068^a	0.412	0.615	0.525
12	Combined M peak & H peak (M ≥ 5 & H ≥ 10)	294	134/160	76.49	68.28	0.055^a	0.391	0.607	0.508
13	Combined M peak & H risen (M ≥ 5 &/or H ≥ 10)	370	166/204	54.58	50.11	0.166	0.403	0.578	0.499
14	Combined M peak & H rise (M ≥ 5 & H ≥ 10)	267	124/143	66.54	62.50	0.239	0.379	0.530	0.460
15	Combined M peak & H peak (M ≥ 10 &/or H ≥ 20)	177	96/81	86.28	85.49	0.674	0.390	0.395	0.392
16	Combined M peak & H rise (M ≥ 10 & H ≥ 20)	149	84/65	77.32	83.17	0.723	0.522	0.346	0.445
17	Combined M peak & H rise (M ≥ 10 &/or H ≥ 20 by 90 min)	279	136/143	42.29	35.71	0.008^a	0.382	0.702	0.546

^a $P < 0.1$ for difference in mean value of lactulose breath test between polyp and non-polyp groups for further assessment.

Rise values are baseline-subtracted peak values during the tests. Bold P values indicate the 7 best models in further assessment. ppm: Parts per million.

DISCUSSION

Recent studies have shown that the gut microbiome is associated with certain gastrointestinal symptoms [12], colon polyps, and colorectal cancer [18,20]. However, little is known about the relationship between SIBO and colorectal polyps. The current study analyzed a patient cohort that had recently received LBT for uncertain gastrointestinal symptoms. The findings revealed certain key associations among colorectal polyps, SIBO, and IBFD while demonstrating that LBT had moderate potential as an alternative noninvasive screening tool for colorectal polyps. SIBO was more prevalent in patients with colorectal polyps than those without and IBFD was worse in patients with colorectal polyps than those without only when SIBO was evident.

SIBO is caused by gut microbiota dysregulation and is characterized by the excessive density and/or abnormal composition of bacteria in the small intestine. The current study was the first to demonstrate that patients with colorectal polyps had a higher prevalence of SIBO than those without, defined by methane and hydrogen test results alone or in combination. These findings suggest that SIBO may be a risk factor for colorectal polyps. While this study showed no difference in SIBO by polyp type, further investigation is needed to confirm this finding. The results also showed a higher rate of IBFD among patients with colorectal polyps than those without, however this was only true for patients with SIBO. This suggests that patients with polyps are more susceptible to IBFD when SIBO are present.

Table 5 Model performance with key parameters (area under the receiver operating characteristic curve, accuracy, sensitivity, and specificity)

Panel ID	AUC, % (95%CI)	Accuracy	Sensitivity	Specificity
a	71.6 (66.5-76.7)	0.659	0.666	0.653
b	71.4 (66.3-76.5)	0.642	0.663	0.625
c	72.0 (66.9-77.1)	0.650	0.669	0.634
d	71.7 (66.4-77.0)	0.663	0.679	0.651
e	72.9 (67.3-78.4)	0.651	0.677	0.629
f	72.6 (66.9-78.4)	0.650	0.683	0.622
g	71.7 (65.7-77.7)	0.658	0.673	0.643

AUC: Area under the curve.





Figure 1 Rate of intestinal barrier function damage in 227 small intestine bacteria overgrowth patients. ^aThe rate of intestinal barrier function damage (IBFD) by 3 blood assays altogether was marginally higher in polyp group than that in non-polyp group, but no significance, $P = 0.13$. ^bIt was significantly different between polyp group and non-polyp group when IBFD defined by lipopolysaccharide alone, $P < 0.05$. SIBO: Small intestine bacteria overgrowth; DAO: Diamine oxidase; LPS: Lipopolysaccharide.

The culture of small bowel aspirates is the gold standard for SIBO diagnosis, but this is an invasive method and it can be a challenge to culture gut flora[21,22]. As a result, noninvasive testing using LBT results is often used. While the diagnostic criteria for SIBO by LBT are not yet standardized, the 2017 North American Consensus guidelines used in this study can make the results comparable across studies with similar data and analysis. LBT is primarily used to support SIBO diagnosis but is implemented here as a new method for screening colorectal polyps. To our knowledge, this is the first study to use quantitative LBT measurements for prediction analysis. LBT had moderate potential as a screening tool to identify patients with polyps in the large intestine. The best fit models were greatly improved after adjusting for age and gender. It is worth noting that models that only included the peak methane values without filtering by cutoff values performed as well as the model with combined H and M values limited by North American Consensus guidelines (Figure 2H). This suggests that methane peak values were as useful as combined hydrogen and methane values in patients with SIBO when using LBT as a screening tool for colorectal polyps.

In this retrospective study, patients with colorectal polyps were about 9 years older and more often male than those without polyps. These findings are consistent with the characteristics of polyp development and support the results of prior studies. Most studies report that men have almost twice the prevalence of polyps as women[23,24], and this tends to increase with age[3]. In addition, comorbidity analysis showed that patients with colorectal polyps were more often complicated with metabolic disorders and less likely with non-organic abnormalities. This is consistent with recent studies indicating that metabolic syndrome is a high-risk factor for colorectal adenomatous polyps and cancer and should be included in colorectal cancer screening programs[25,26]. These findings suggest that metabolic disorders can be an inherent characteristic among certain patients with colorectal polyps.

The incidence of constipation was significantly higher in the polyp than non-polyp groups. Patients with constipation have prolonged oro-cecal transit time[27] and constipation can significantly increase the incidence of colorectal polyps[28]. However, colonoscopy for patients with constipation as the sole indication had fewer neoplastic lesions than for those undergoing routine screening colonoscopy[29].



DOI: 10.4240/wjgs.v15.i6.1138 Copyright ©The Author(s) 2023.

Figure 2 The receiver operation characteristic curves for age and gender adjusted performance of prediction models. A-G: Each model is for a subset of patient population defined by the cutoffs and the size of the subpopulation showing at the top of each box.

Additional randomized controlled double-blind studies with a larger sample size are needed to confirm the findings of the present study.

CONCLUSION

The current study identified key associations among colorectal polyps, SIBO and IBFD while demonstrating the moderate potential of LBT as an alternative noninvasive screening tool for colorectal polyps. SIBO was more prevalent in patients with colorectal polyps than those without and IBFD was more severe in patients with colorectal polyp than those without only when SIBO has present. This study also found that colorectal polyps were more common in older individuals and men. Moreover,

patients with polyps tended to have metabolic disorders such as diabetes and hyperlipidemia and were less likely to have non-organic abnormalities such as functional gastrointestinal disease and IBS.

ARTICLE HIGHLIGHTS

Research background

Polyps can often be found and removed by colonoscopy; however, this is an invasive and expensive test. Due to its safety, accessibility, and affordability, the lactulose breath test (LBT) is widely used as a noninvasive method of diagnosing small intestine bacterial overgrowth (SIBO). SIBO suggests dysbiosis of the intestinal flora, which is associated with the pathogenesis of adenomatous polyps and colorectal cancer.

Research motivation

New methods that can identify precancerous colorectal lesions can play an important role in early-stage colorectal cancer treatment and prevention.

Research objectives

To identify a potential association between colorectal polyps and SIBO or other relevant factors in a patient cohort with LBT results.

Research methods

This retrospective analysis of data from a total of 382 patients who had received an LBT were collected. SIBO was diagnosed by measuring LBT-derived hydrogen (H) and methane (M) levels according to 2017 North American Consensus recommendations. Logistic regression was used to assess the ability of LBT to predict colorectal polyps. Intestinal barrier function damage (IBFD) was determined by blood assays.

Research results

H and M levels revealed that the prevalence of SIBO was significantly higher in the polyp group than in the non-polyp group (41% vs 23%; 71% vs 59%, respectively). Within 90 min of lactulose ingestion, the peak H values in the adenomatous and inflammatory/hyperplastic polyp patients were significantly higher than those in the non-polyp group. In regression analysis with age and gender adjustment, colorectal polyps were most accurately predicted with models using M peak values or combined H and M values limited by North American Consensus recommendations for SIBO. These models had a sensitivity of ≥ 0.67 , a specificity of ≥ 0.64 , and an accuracy of ≥ 0.66 .

Research conclusions

The current study made key associations among colorectal polyps and SIBO and demonstrated that LBT has moderate potential as an alternative noninvasive screening tool for colorectal polyps.

Research perspectives

Due to its safety, accessibility, and affordability, the LBT has the potential to become one of the routine non-invasive screening methods for polyps and precancerous lesions. Furthermore, non-invasive tests such as fecal occult blood testing and LBT will help to improve the detection rate of precancerous lesions during colonoscopy screening.

ACKNOWLEDGEMENTS

We appreciate the participation of patients in the study.

FOOTNOTES

Author contributions: Lan Li and Zhang XY have contributed equally to this work. Lan Li participated the study design and wrote the manuscript draft; Zhang XY conducted clinical data collections and participated the writing of the draft; Zhou HM and Qin Y performed lactulose breath test tests; Xie WR participated statistical analysis of the data; Ding WJ interpreted the lactulose breath test results; Yu JS participated the study design, performed the statistical analysis of the data, and edited the final version of the manuscript; He XX conceived the study concept and design, supervised all the work, provided the study funding, and reviewed the final version of the manuscript. All authors listed have made a substantial, direct, and intellectual contribution to the work and approved it for publication.

Supported by the Key-Area Research and Development Program of Guangdong Province, No. 2022B1111070006; the Guangdong Innovation Research Team for Higher Education, No. 2021KCXTD025.

Institutional review board statement: The study protocol was approved by the institutional review board and the Ethics Committee of the First Affiliated Hospital of Guangdong Pharmaceutical University, No. 2019045.

Informed consent statement: All study participants or their legal guardian provided informed written consent about personal and medical data collection prior to study enrollment.

Conflict-of-interest statement: The authors have no conflicts of interest to declare. There are no ethical or legal conflicts involved in the article.

Data sharing statement: Statistical code, and dataset available from the corresponding author at hexingxiang@gdpu.edu.cn. Participants gave informed consent for data sharing.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Lan Li [0000-0002-3919-029X](https://orcid.org/0000-0002-3919-029X); Hui-Min Zhou [0000-0001-9740-2361](https://orcid.org/0000-0001-9740-2361); Wen-Rui Xie [0000-0001-7180-5090](https://orcid.org/0000-0001-7180-5090); Xing-Xiang He [0000-0003-0007-8513](https://orcid.org/0000-0003-0007-8513).

Corresponding Author's Membership in Professional Societies: Expert Committee of Guangzhou Regional Digestive Disease Medical Quality Control Center, No. [2022]1750; China Association for Promotion of Health Science and Technology, Deputy Director, No. 0050202210003.

S-Editor: Zhang H

L-Editor: A

P-Editor: Li X

REFERENCES

- 1 **Chinese Society of Gastroenterology**, Cancer Collaboration Group of Chinese Society of Gastroenterology, Chinese Medical Association. Chinese consensus on prevention of colorectal neoplasia (2021, Shanghai). *J Dig Dis* 2022; **23**: 58-90 [PMID: [34984819](https://pubmed.ncbi.nlm.nih.gov/34984819/) DOI: [10.1111/1751-2980.13079](https://doi.org/10.1111/1751-2980.13079)]
- 2 **Brethauer M**, Kalager M, Adami HO. Do's and don'ts in evaluation of endoscopic screening for gastrointestinal cancers. *Endoscopy* 2016; **48**: 75-80 [PMID: [26382306](https://pubmed.ncbi.nlm.nih.gov/26382306/) DOI: [10.1055/s-0034-1393094](https://doi.org/10.1055/s-0034-1393094)]
- 3 **Øines M**, Helsing LM, Brethauer M, Emilsson L. Epidemiology and risk factors of colorectal polyps. *Best Pract Res Clin Gastroenterol* 2017; **31**: 419-424 [PMID: [28842051](https://pubmed.ncbi.nlm.nih.gov/28842051/) DOI: [10.1016/j.bpg.2017.06.004](https://doi.org/10.1016/j.bpg.2017.06.004)]
- 4 **Hikino K**, Koido M, Otomo N, Tomizuka K, Ikegawa S, Matsuda K, Momozawa Y, Biobank Japan Project, Mushihiro T, Terao C. Genome-wide association study of colorectal polyps identified highly overlapping polygenic architecture with colorectal cancer. *J Hum Genet* 2022; **67**: 149-156 [PMID: [34671089](https://pubmed.ncbi.nlm.nih.gov/34671089/) DOI: [10.1038/s10038-021-00980-4](https://doi.org/10.1038/s10038-021-00980-4)]
- 5 **Pop OL**, Vodnar DC, Diaconeasa Z, Istrati M, Bințișan A, Bințișan VV, Suharoschi R, Gabbianelli R. An Overview of Gut Microbiota and Colon Diseases with a Focus on Adenomatous Colon Polyps. *Int J Mol Sci* 2020; **21** [PMID: [33028024](https://pubmed.ncbi.nlm.nih.gov/33028024/) DOI: [10.3390/ijms21197359](https://doi.org/10.3390/ijms21197359)]
- 6 **Brethauer M**. Colorectal cancer screening. *J Intern Med* 2011; **270**: 87-98 [PMID: [21575082](https://pubmed.ncbi.nlm.nih.gov/21575082/) DOI: [10.1111/j.1365-2796.2011.02399.x](https://doi.org/10.1111/j.1365-2796.2011.02399.x)]
- 7 **Montminy EM**, Jang A, Conner M, Karlitz JJ. Screening for Colorectal Cancer. *Med Clin North Am* 2020; **104**: 1023-1036 [PMID: [33099448](https://pubmed.ncbi.nlm.nih.gov/33099448/) DOI: [10.1016/j.mcna.2020.08.004](https://doi.org/10.1016/j.mcna.2020.08.004)]
- 8 **Kerckhoffs AP**, Visser MR, Samsom M, van der Rest ME, de Vogel J, Harmsen W, Akkermans LM. Critical evaluation of diagnosing bacterial overgrowth in the proximal small intestine. *J Clin Gastroenterol* 2008; **42**: 1095-1102 [PMID: [18936644](https://pubmed.ncbi.nlm.nih.gov/18936644/) DOI: [10.1097/MCG.0b013e31818474d7](https://doi.org/10.1097/MCG.0b013e31818474d7)]
- 9 **Pitcher CK**, Farmer AD, Haworth JJ, Treadway S, Hobson AR. Performance and Interpretation of Hydrogen and Methane Breath Testing Impact of North American Consensus Guidelines. *Dig Dis Sci* 2022; **67**: 5571-5579 [PMID: [35366119](https://pubmed.ncbi.nlm.nih.gov/35366119/) DOI: [10.1007/s10620-022-07487-8](https://doi.org/10.1007/s10620-022-07487-8)]
- 10 **Sánchez-Montes C**, Ortiz V, Bastida G, Rodríguez E, Yago M, Beltrán B, Aguas M, Iborra M, Garrigues V, Ponce J, Nos P. Small intestinal bacterial overgrowth in inactive Crohn's disease: influence of thiopurine and biological treatment. *World J Gastroenterol* 2014; **20**: 13999-14003 [PMID: [25320539](https://pubmed.ncbi.nlm.nih.gov/25320539/) DOI: [10.3748/wjg.v20.i38.13999](https://doi.org/10.3748/wjg.v20.i38.13999)]
- 11 **Ghoshal UC**, Shukla R, Ghoshal U. Small Intestinal Bacterial Overgrowth and Irritable Bowel Syndrome: A Bridge between Functional Organic Dichotomy. *Gut Liver* 2017; **11**: 196-208 [PMID: [28274108](https://pubmed.ncbi.nlm.nih.gov/28274108/) DOI: [10.5009/gnl16126](https://doi.org/10.5009/gnl16126)]
- 12 **Choi CH**, Chang SK. Role of Small Intestinal Bacterial Overgrowth in Functional Gastrointestinal Disorders. *J*

- Neurogastroenterol Motil* 2016; **22**: 3-5 [PMID: 26717927 DOI: 10.5056/jnm15196]
- 13 **Fialho A**, Fialho A, Thota P, McCullough AJ, Shen B. Small Intestinal Bacterial Overgrowth Is Associated with Non-Alcoholic Fatty Liver Disease. *J Gastrointest Liver Dis* 2016; **25**: 159-165 [PMID: 27308646 DOI: 10.15403/jgld.2014.1121.252.iwg]
 - 14 **Rana SV**, Malik A, Bhadada SK, Sachdeva N, Morya RK, Sharma G. Malabsorption, Orocecal Transit Time and Small Intestinal Bacterial Overgrowth in Type 2 Diabetic Patients: A Connection. *Indian J Clin Biochem* 2017; **32**: 84-89 [PMID: 28149017 DOI: 10.1007/s12291-016-0569-6]
 - 15 **Feng X**, Li X, Zhang X, Chen W, Tian Y, Yang Q, Yang Y, Pan H, Jiang Z. Hepatic Encephalopathy in Cirrhotic Patients and Risk of Small Intestinal Bacterial Overgrowth: A Systematic Review and Meta-Analysis. *Biomed Res Int* 2022; **2022**: 2469513 [PMID: 36303585 DOI: 10.1155/2022/2469513]
 - 16 **Hale VL**, Chen J, Johnson S, Harrington SC, Yab TC, Smyrk TC, Nelson H, Boardman LA, Druliner BR, Levin TR, Rex DK, Ahnen DJ, Lance P, Ahlquist DA, Chia N. Shifts in the Fecal Microbiota Associated with Adenomatous Polyps. *Cancer Epidemiol Biomarkers Prev* 2017; **26**: 85-94 [PMID: 27672054 DOI: 10.1158/1055-9965.EPI-16-0337]
 - 17 **Rezasoltani S**, Asadzadeh Aghdaei H, Dabiri H, Akhavan Sepahi A, Modarressi MH, Nazemalhosseini Mojarad E. The association between fecal microbiota and different types of colorectal polyp as precursors of colorectal cancer. *Microb Pathog* 2018; **124**: 244-249 [PMID: 30142468 DOI: 10.1016/j.micpath.2018.08.035]
 - 18 **Gao R**, Kong C, Li H, Huang L, Qu X, Qin N, Qin H. Dysbiosis signature of mycobiota in colon polyp and colorectal cancer. *Eur J Clin Microbiol Infect Dis* 2017; **36**: 2457-2468 [PMID: 28821976 DOI: 10.1007/s10096-017-3085-6]
 - 19 **Rezaie A**, Buresi M, Lembo A, Lin H, McCallum R, Rao S, Schmulson M, Valdovinos M, Zakko S, Pimentel M. Hydrogen and Methane-Based Breath Testing in Gastrointestinal Disorders: The North American Consensus. *Am J Gastroenterol* 2017; **112**: 775-784 [PMID: 28323273 DOI: 10.1038/ajg.2017.46]
 - 20 **Watson KM**, Gaulke CA, Tsikitis VL. Understanding the microbiome: a primer on the role of the microbiome in colorectal neoplasia. *Ann Gastroenterol* 2020; **33**: 223-236 [PMID: 32382225 DOI: 10.20524/aog.2020.0467]
 - 21 **Pimentel M**, Saad RJ, Long MD, Rao SSC. ACG Clinical Guideline: Small Intestinal Bacterial Overgrowth. *Am J Gastroenterol* 2020; **115**: 165-178 [PMID: 32023228 DOI: 10.14309/ajg.0000000000000501]
 - 22 **Newberry C**, Tierney A, Pickett-Blakely O. Lactulose Hydrogen Breath Test Result Is Associated with Age and Gender. *Biomed Res Int* 2016; **2016**: 1064029 [PMID: 27073800 DOI: 10.1155/2016/1064029]
 - 23 **Brenner H**, Altenhofen L, Stock C, Hoffmeister M. Incidence of colorectal adenomas: birth cohort analysis among 4.3 million participants of screening colonoscopy. *Cancer Epidemiol Biomarkers Prev* 2014; **23**: 1920-1927 [PMID: 25012996 DOI: 10.1158/1055-9965.EPI-14-0367]
 - 24 **Waldmann E**, Heinze G, Ferlitsch A, Gessl I, Sallinger D, Jeschek P, Britto-Arias M, Salzl P, Fasching E, Jilma B, Kundi M, Trauner M, Ferlitsch M. Risk factors cannot explain the higher prevalence rates of precancerous colorectal lesions in men. *Br J Cancer* 2016; **115**: 1421-1429 [PMID: 27764840 DOI: 10.1038/bjc.2016.324]
 - 25 **Milano A**, Bianco MA, Buri L, Cipolletta L, Grossi E, Rotondano G, Tessari F, Efthymakis K, Neri M. Metabolic syndrome is a risk factor for colorectal adenoma and cancer: a study in a White population using the harmonized criteria. *Therap Adv Gastroenterol* 2019; **12**: 1756284819867839 [PMID: 31523276 DOI: 10.1177/1756284819867839]
 - 26 **Yu Y**, Wu J. Presence of Metabolic Syndrome and Thyroid Nodules in Subjects with Colorectal Polyps. *Med Sci Monit* 2021; **27**: e927935 [PMID: 33518699 DOI: 10.12659/MSM.927935]
 - 27 **Bin Waqar SH**, Rehan A. Methane and Constipation-predominant Irritable Bowel Syndrome: Entwining Pillars of Emerging Neurogastroenterology. *Cureus* 2019; **11**: e4764 [PMID: 31363445 DOI: 10.7759/cureus.4764]
 - 28 **Liu B**. [Correlation between chronic constipation and colorectal neoplasms]. *Zhonghua Wei Chang Wai Ke Za Zhi* 2017; **20**: 255-257 [PMID: 28338155]
 - 29 **Obusez EC**, Lian L, Kariv R, Burke CA, Shen B. Diagnostic yield of colonoscopy for constipation as the sole indication. *Colorectal Dis* 2012; **14**: 585-591 [PMID: 21689337 DOI: 10.1111/j.1463-1318.2011.02664.x]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

