

World Journal of *Gastrointestinal Surgery*

World J Gastrointest Surg 2023 July 27; 15(7): 1262-1558



Contents

Monthly Volume 15 Number 7 July 27, 2023

REVIEW

- 1262 Pathophysiological consequences and treatment strategy of obstructive jaundice
Liu JJ, Sun YM, Xu Y, Mei HW, Guo W, Li ZL

MINIREVIEWS

- 1277 Carbon footprints in minimally invasive surgery: Good patient outcomes, but costly for the environment
Chan KS, Lo HY, Shelat VG

ORIGINAL ARTICLE

Basic Study

- 1286 Primary animal experiment to test the feasibility of a novel Y-Z magnetic hepatic portal blocking band
Zhang MM, Li CG, Xu SQ, Mao JQ, Ren YX, Zhang YH, Ma J, Shi AH, Lyu Y, Yan XP
- 1294 Magnetic compression anastomosis for reconstruction of digestive tract after total gastrectomy in beagle model
Zhang MM, Li CG, Xu SQ, Mao JQ, Zhang YH, Shi AH, Li Y, Lyu Y, Yan XP
- 1304 Differences in metabolic improvement after metabolic surgery are linked to the gut microbiota in non-obese diabetic rats
Luo X, Tan C, Tao F, Xu CY, Zheng ZH, Pang Q, He XA, Cao JQ, Duan JY
- 1317 Intervention effects and related mechanisms of glycyrrhizic acid on zebrafish with Hirschsprung-associated enterocolitis
Liu MK, Chen YJ, Chen F, Lin ZX, Zhu ZC, Lin Y, Fang YF, Wu DM
- 1331 Histological study of the structural layers around the esophagus in the lower mediastinum
Saito T, Muro S, Fujiwara H, Umebayashi Y, Sato Y, Tokunaga M, Akita K, Kinugasa Y

Case Control Study

- 1340 Liver transplantation for combined hepatocellular carcinoma and cholangiocarcinoma: A multicenter study
Kim J, Joo DJ, Hwang S, Lee JM, Ryu JH, Nah YW, Kim DS, Kim DJ, You YK, Yu HC
- 1354 Optimal choice of stapler and digestive tract reconstruction method after distal gastrectomy for gastric cancer: A prospective case-control study
Wu Z, Zhou ZG, Li LY, Gao WJ, Yu T

Retrospective Cohort Study

- 1363 Impact of perioperative blood transfusion on oncological outcomes in ampullary carcinoma patients underwent pancreaticoduodenectomy
Fei H, Zhang XJ, Sun CY, Li Z, Li ZF, Guo CG, Zhao DB

Retrospective Study

- 1375** Nomogram based on clinical characteristics for predicting overall survival in gastric cancer patients with preoperative anemia
Long Y, Zhou XL, Zhang CL, Wang YN, Pan WS
- 1388** Major complications after ultrasound-guided liver biopsy: An annual audit of a Chinese tertiary-care teaching hospital
Chai WL, Lu DL, Sun ZX, Cheng C, Deng Z, Jin XY, Zhang TL, Gao Q, Pan YW, Zhao QY, Jiang TA
- 1397** Different percutaneous transhepatic biliary stent placements and catheter drainage in the treatment of middle and low malignant biliary obstruction
Yang YB, Yan ZY, Jiao Y, Yang WH, Cui Q, Chen SP
- 1405** Utilization of deep neuromuscular blockade combined with reduced abdominal pressure in laparoscopic radical gastrectomy for gastric cancer: An academic perspective
Zhang YW, Li Y, Huang WB, Wang J, Qian XE, Yang Y, Huang CS
- 1416** Efficacy of peritoneal drainage in very-low-birth-weight neonates with Bell's stage II necrotizing enterocolitis: A single-center retrospective study
Shen Y, Lin Y, Fang YF, Wu DM, He YB
- 1423** Emergency exploratory laparotomy and radical gastrectomy in patients with gastric cancer combined with acute upper gastrointestinal bleeding
Kuang F, Wang J, Wang BQ
- 1434** Correlation of serum albumin level on postoperative day 2 with hospital length of stay in patients undergoing emergency surgery for perforated peptic ulcer
Xie D, Lu PL, Xu W, You JY, Bi XG, Xian Y

Clinical Trials Study

- 1442** Laboratory scoring system to predict hepatic indocyanine green clearance ability during fluorescence imaging-guided laparoscopic hepatectomy
Chen ZR, Zeng QT, Shi N, Han HW, Chen ZH, Zou YP, Zhang YP, Wu F, Xu LQ, Jin HS

Observational Study

- 1454** Incidence, characteristics and risk factors for alveolar recruitment maneuver-related hypotension in patients undergoing laparoscopic colorectal cancer resection
Zhang NR, Zheng ZN, Wang K, Li H
- 1465** New classification system for radical rectal cancer surgery based on membrane anatomy
Jiang HH, Ni ZZ, Chang Y, Li AJ, Wang WC, Lv L, Peng J, Pan ZH, Liu HL, Lin MB

Randomized Controlled Trial

- 1474** Transcutaneous electrical acupoint stimulation in adult patients receiving gastrectomy/colorectal resection: A randomized controlled trial
Hou YT, Pan YY, Wan L, Zhao WS, Luo Y, Yan Q, Zhang Y, Zhang WX, Mo YC, Huang LP, Dai QX, Jia DY, Yang AM, An HY, Wu AS, Tian M, Fang JQ, Wang JL, Feng Y

SYSTEMATIC REVIEWS

- 1485 Combined and intraoperative risk modelling for oesophagectomy: A systematic review
Grantham JP, Hii A, Shenfine J
- 1501 Spleen-preserving distal pancreatectomy from multi-port to reduced-port surgery approach
Hsieh CL, Tsai TS, Peng CM, Cheng TC, Liu YJ
- 1512 Resection of isolated liver oligometastatic disease in pancreatic ductal adenocarcinoma: Is there a survival benefit? A systematic review
Halle-Smith JM, Powell-Brett S, Roberts K, Chatzizacharias NA

META-ANALYSIS

- 1522 Outcome of split liver transplantation *vs* living donor liver transplantation: A systematic review and meta-analysis
Garzali IU, Akbulut S, Aloun A, Naffa M, Aksoy F

CASE REPORT

- 1532 Idiopathic hypereosinophilic syndrome with hepatic sinusoidal obstruction syndrome: A case report and literature review
Xu XT, Wang BH, Wang Q, Guo YJ, Zhang YN, Chen XL, Fang YF, Wang K, Guo WH, Wen ZZ
- 1542 Reoperation for heterochronic intraductal papillary mucinous neoplasm of the pancreas after bile duct neoplasm resection: A case report
Xiao G, Xia T, Mou YP, Zhou YC
- 1549 Successful resection of colonic metastasis of lung cancer after colonic stent placement: A case report and review of the literature
Nakayama Y, Yamaguchi M, Inoue K, Hamaguchi S, Tajima Y

ABOUT COVER

Editorial Board Member of *World Journal of Gastrointestinal Surgery*, Georgios Tsoulfas, AGAF, FACS, FICS, MD, PhD, Professor, Transplant Surgery, Aristotle University of Thessaloniki School of Medicine, Thessaloniki 54124, Greece. tsoulfasg@gmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Gastrointestinal Surgery* (WJGS, *World J Gastrointest Surg*) is to provide scholars and readers from various fields of gastrointestinal surgery with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGS mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal surgery and covering a wide range of topics including biliary tract surgical procedures, biliopancreatic diversion, colectomy, esophagectomy, esophagostomy, pancreas transplantation, and pancreatectomy, etc.

INDEXING/ABSTRACTING

The WJGS is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports/Science Edition, PubMed, PubMed Central, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJGS as 2.0; IF without journal self cites: 1.9; 5-year IF: 2.2; Journal Citation Indicator: 0.52; Ranking: 113 among 212 journals in surgery; Quartile category: Q3; Ranking: 81 among 93 journals in gastroenterology and hepatology; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Rui-Rui Wu, Production Department Director: Xiang Li, Editorial Office Director: Jia-Ru Fan.

NAME OF JOURNAL

World Journal of Gastrointestinal Surgery

ISSN

ISSN 1948-9366 (online)

LAUNCH DATE

November 30, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Peter Schemmer

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/1948-9366/editorialboard.htm>

PUBLICATION DATE

July 27, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Retrospective Study

Different percutaneous transhepatic biliary stent placements and catheter drainage in the treatment of middle and low malignant biliary obstruction

Yao-Bo Yang, Zhao-Yong Yan, Yang Jiao, Wei-Hao Yang, Qi Cui, Si-Pan Chen

Specialty type: Surgery

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Praktijnjo M, Germany; Reddy KR, United States

Received: March 13, 2023

Peer-review started: March 13, 2023

First decision: March 28, 2023

Revised: April 21, 2023

Accepted: May 6, 2023

Article in press: May 6, 2023

Published online: July 27, 2023



Yao-Bo Yang, Zhao-Yong Yan, Yang Jiao, Wei-Hao Yang, Qi Cui, Si-Pan Chen, Department of Interventional Radiology, Shaanxi Provincial People's Hospital, Xi'an 710068, Shaanxi Province, China

Corresponding author: Si-Pan Chen, Doctor, Professor, Department of Interventional Radiology, Shaanxi Provincial People's Hospital, No. 256 Youyi West Road, Beilin District, Xi'an 710068, Shaanxi Province, China. panda_sp@163.com

Abstract

BACKGROUND

For cases of middle and low biliary obstruction with left and right hepatic duct dilatation, the type of approach and whether different approaches affect the difficulty of puncture operation and intraoperative and postoperative complications have not been discussed in detail.

AIM

To compare the efficacy of different percutaneous transhepatic biliary stent placements and catheter drainage in treating middle and low biliary obstruction.

METHODS

A retrospective analysis was performed on the medical records of 424 patients with middle and low biliary obstruction who underwent percutaneous liver puncture biliary stent placement and catheter drainage at the Department of Interventional Radiology, Shaanxi Provincial People's Hospital between March 2016 and March 2022. Based on the puncture path, patients were categorized into two groups: Subxiphoid left hepatic lobe approach group (Group A, 224 cases) and right intercostal, right hepatic lobe approach group (Group B, 200 cases). Liver function improvement, postoperative biliary bleeding incidence, postoperative pain duration, and abdominal effusion leakage around the drainage tube were compared between the two groups at 3 d and 1 wk after the surgery. Patient survival time was recorded during follow-up.

RESULTS

All 424 surgeries were successful without adverse events. Group A comprised 224 cases, and Group B had 200 cases. There was no statistically significant difference in basic data between Group A and Group B ($P > 0.05$). No significant difference

in postoperative biliary bleeding incidence was observed between the groups ($P > 0.05$). The decreased rates for total bilirubin (Group A: 69.23 ± 4.50 , Group B: 63.79 ± 5.65), direct bilirubin (Group A: 79.30 ± 11.19 , Group B: 63.62 ± 5.64), and alkaline phosphatase (Group A: 60.51 ± 12.23 , Group B: 42.68 ± 23.56) in the 1st wk after surgery were significantly faster in Group A than in Group B. The decreased rate of gamma-glutamyl transpeptidase was also significantly faster in Group A at both 3 d (Group A: 40.56 ± 10.32 , Group B: 32.22 ± 5.12) and 1 wk (Group A: 73.19 ± 7.05 , Group B: 58.81 ± 18.98) after surgery ($P < 0.05$). Group A experienced significantly less peritoneal effusion leakage around the drainage tube than Group B ($P < 0.05$). The patient survival rate was higher in Group A compared to Group B ($P < 0.05$).

CONCLUSION

In treating jaundice patients with middle and low biliary obstruction, a percutaneous left liver puncture demonstrated better clinical efficacy than a percutaneous right liver puncture.

Key Words: Biliary obstruction; Puncture route; Stent placement; Survival rate; Malignant tumor; Digestive tract

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: For patients with unresectable middle to low level malignant biliary obstruction, percutaneous transhepatic biliary drainage has been widely used in clinical practice. The selection of the approach and whether different approaches will affect the difficulty of puncture surgery, as well as intraoperative and postoperative complications, have not been discussed in detail. This study compared the clinical efficacy of two different puncture pathways in the treatment of middle and low level biliary obstruction. The authors found that the clinical efficacy of patients undergoing percutaneous left liver puncture was superior to that of patients undergoing percutaneous right liver puncture.

Citation: Yang YB, Yan ZY, Jiao Y, Yang WH, Cui Q, Chen SP. Different percutaneous transhepatic biliary stent placements and catheter drainage in the treatment of middle and low malignant biliary obstruction. *World J Gastrointest Surg* 2023; 15(7): 1397-1404

URL: <https://www.wjgnet.com/1948-9366/full/v15/i7/1397.htm>

DOI: <https://dx.doi.org/10.4240/wjgs.v15.i7.1397>

INTRODUCTION

In patients with unresectable middle and low malignant biliary obstruction, the primary cause is often extrahepatic biliary obstruction and obstruction below the cystic duct opening. Common symptoms include yellowing of the skin and sclera, high fever, abdominal pain, and decreased appetite. In cases of complete obstruction, the stool color may become pale or even clay-like[1]. Percutaneous transhepatic cholangial drainage (PTCD), a widely used palliative care method[2, 3], offers advantages such as a straightforward procedure, effective jaundice relief, and convenient bile drainage, making it a popular choice in clinical settings.

For cases of middle and low biliary obstruction with left and right hepatic duct dilatation, the type of approach and whether different approaches affect the difficulty of puncture operation and intraoperative and postoperative complications have not been discussed in detail. This study compared the clinical efficacy of two different puncture paths in treating middle and low biliary obstruction.

MATERIALS AND METHODS

General data

Between March 2016 and March 2022, 424 medical records of patients with inoperable middle and low malignant biliary obstruction were analyzed. Patients were divided into two groups based on the puncture approach: Group A, consisting of 224 cases using the subxiphoid left hepatic lobe approach; and Group B, consisting of 200 cases using the right intercostal, right hepatic lobe approach. Inclusion criteria[4,5] were as follows: (1) Clinical presentation, magnetic resonance cholangiopancreatography, computed tomography, ultrasound, and other imaging and laboratory examination results indicating inoperable middle and low malignant biliary obstruction; (2) Anticipated survival time of more than 3 mo; and (3) Successful percutaneous puncture biliary drainage and metal stent implantation. Exclusion criteria included: (1) Incomplete clinical and follow-up data; (2) Patients with multiple organ failure who could not tolerate percutaneous transhepatic biliary drainage and stent implantation; (3) Patients who underwent chemotherapy, radiotherapy, or Billroth II subtotal gastrectomy; and (4) Abnormal coagulation function. All patients received liver protection treatment after the operation. Preoperative and postoperative total bilirubin, direct bilirubin, aspartate aminotransferase, and gamma-glutamyl transpeptidase values were recorded and calculated at 1 wk and 3 d after the procedure, and their rates were determined. Follow-ups were conducted every 3 mo.

Intervention operation

Traditional disinfection and local puncture anesthesia were employed. The puncture site was typically situated at the lower xiphoid process on the right abdominal wall (left bile duct approach) or between the 8th and 9th intercostal spaces of the right seasonal rib (right bile duct approach). The puncture direction and angle were designed for digital subtraction angiography. A Cook 22 G/15 cm biliary puncture needle was used to access the bile duct of the liver, and the needle core was removed. The successful puncture was confirmed by observing the outflow of pale-yellow bile, and a compatible 0.018-inch short guidewire was introduced through the puncture needle. After removing the puncture needle, a matching coaxial introducer sheath (4 F) was inserted along the guidewire. A contrast medium was administered through the introducer sheath to assess the location, extent, and severity of the obstruction of the bile duct. A 0.035-inch ultra-smooth guidewire was inserted through the introducer sheath, and the obstruction site was adjusted.

A memory alloy stent was introduced into the biliary system using a pusher along the guidewire, ensuring that the upper end of the stent was above the obstruction, and its lower end was within the obstructed bile duct and duodenum. Both ends needed to extend approximately 1 cm beyond the narrowed section. After the stent was deployed, the metal stent could open smoothly. An 8.5 F external biliary drainage tube was placed through the guidewire, and contrast medium was injected again to verify the resolution of the bile duct obstruction, adequate bile drainage, absence of clotting in the biliary tract, firm fixation of the drainage tube, and proper connection with the drainage bag. **Figure 1** displays the imaging data for percutaneous stent placement and catheter drainage of the left/right hepatic ducts. Prior to the procedure, all patients received an intramuscular injection of 10 mg diazepam, 70 mg pethidine hydrochloride, and 10 mg racemolamine hydrochloride.

Statistical analysis

SPSS23.0 statistical software was used for data analysis. Measurement data were expressed as mean \pm SD, and count data were expressed as numbers (percentage). An independent sample *t*-test was used for comparison between the two groups. The χ^2 test was used for comparison between groups. The survival rate was analyzed with the Kaplan-Meier survival curve, and $P < 0.05$ was considered statistically significant.

RESULTS

Basic data of patients

A total of 424 patients participated in the study, with no severe adverse events (such as death and critical cardiac or cerebrovascular incidents) observed during the follow-up period. The cohort included 227 males (53.5%) and 197 females (46.5%), with an average age of 68.92 ± 0.36 years. Among the patients, 46 cases (10.8%) had diabetes, 79 cases (18.6%) were smokers, 101 cases (23.8%) had coronary heart disease, and 45 cases (10.6%) experienced hypertension. Additionally, 33 cases (7.8%) presented with hyperlipidemia. No significant differences in the baseline characteristics were identified between the two groups ($P > 0.05$) (**Table 1**).

Comparison of clinical efficacy between the two groups

No significant differences were observed between the two groups regarding the incidence of postoperative biliary bleeding and the duration of postoperative pain ($P > 0.05$). A comparison of postoperative liver function indices revealed that the total bilirubin reduction rate, direct bilirubin reduction rate, and alkaline phosphatase reduction rate were significantly higher in Group A during the 1st wk after surgery. Moreover, the gamma-glutamyl transpeptidase reduction rate in Group A was considerably faster than in Group B at both 3 d and 1 wk post-surgery ($P < 0.05$). Group A also exhibited significantly lower leakage of peritoneal effusion around the drainage tube compared to Group B ($P < 0.05$). The survival rate for patients in Group A surpassed that of Group B ($P < 0.05$) (**Table 2** and **Figure 2**).

For patients with jaundice undergoing treatment for middle and low level malignant biliary obstruction, those receiving percutaneous left hepatic puncture demonstrated significantly better liver function improvement, reduced peritoneal effusion leakage around the drainage tube, and enhanced survival compared to patients undergoing percutaneous right hepatic puncture.

DISCUSSION


Pancreatic head cancer and periampullary cancer (including ampullary cancer, lower common bile duct cancer, and duodenal papilla cancer) are frequent causes of middle and low biliary obstruction as well as a group of gastrointestinal malignancies characterized by insidious onset, rapid progression, and poor treatment outcomes and prognosis[6-10]. Many patients are diagnosed at advanced stages, missing the opportunity for curative surgery. Hence, interventions to alleviate biliary obstruction are necessary to extend survival and improve patient quality of life[11,12]. Kubo *et al*[13] demonstrated that interventional surgery for malignant obstructive jaundice has broad applicability, minimal invasiveness, high success rates, and rapid postoperative recovery, with PTCD being a common procedure[14-16].

Currently, there is some debate surrounding the advantages and disadvantages of percutaneous left hepatic biliary stent placement and catheter drainage *vs* percutaneous right hepatic biliary stent placement and catheter drainage for treating low malignant obstructive jaundice. Dumonceau *et al*[17] investigated the benefits of PTCD and endoscopic retrograde cholangiopancreatography in treating biliary obstruction, concluding that PTCD is a simpler procedure with

Table 1 Comparison of baseline characteristics between the two groups

Clinical features	Group A, n = 224	Group B, n = 200	t value	P value
Male sex	124	103	0.632	0.427
Age in yr	69.76 ± 12.24	68.46 ± 14.45	0.816	0.675
Diabetes	21	25	1.060	0.302
Smoking	45	34	0.665	0.415
Coronary heart disease	53	48	0.070	0.935
Hypertension	23	22	0.060	0.870
Hyperlipidemia	14	19	1.555	0.212

Group A: Subxiphoid left hepatic lobe approach group; Group B: Right intercostal, right hepatic lobe approach group.



DOI: 10.4240/wjgs.v15.i7.1397 Copyright ©The Author(s) 2023.

Figure 1 Images of cholangiography and fixation of skin drainage tube. A and B: Cholangiography and fixation of skin drainage tube after subxiphoid percutaneous puncture biliary stent placement and drainage of the left liver; C and D: Fixation of skin drainage tube in cholangiography after percutaneous puncture biliary stent placement and catheter drainage under xiphoid process.

shorter transhepatic access and broader drainage coverage, making it more effective than endoscopic retrograde cholangiopancreatography for patients with biliary obstruction. However, Inoue *et al*[18] reported that improper puncture during percutaneous liver biliary drainage and stent implantation might result in vascular damage and biliary bleeding, with percutaneous right hepatic puncture causing more liver parenchyma damage and liver injury. Additionally, patients carrying a drainage bag on their right side experience a significant reduction in their quality of life. Due to the limited selection of percutaneous left hepatic puncture routes and scarce literature, many believe this approach is challenging and exposes patients to high radiation levels. Thus, few studies in China have investigated the treatment of low malignant obstructive jaundice with biliary stent placement and catheter drainage *via* percutaneous left hepatic puncture.


In line with a humanistic care approach and aiming to enhance long-term survival, our hospital predominantly opted for percutaneous left hepatic biliary stent placement and catheter drainage in treating middle-low malignant obstructive jaundice. We compared the clinical efficacy of two distinct puncture paths in addressing low biliary obstruction. Our study revealed that the reduction rates of total bilirubin, direct bilirubin, aspartate aminotransferase, and gamma-glutamyl transpeptidase in Group A were significantly faster than those in Group B presurgery and post-surgery ($P < 0.05$). Liver function recovery was notably superior in comparison to percutaneous right hepatic puncture. Factors contributing to these results include the slender left hepatic duct, the small left hepatic lobe, and minimal liver damage. Furthermore, stent implantation ensures an unobstructed common bile duct.

Owing to the intact bile duct in the right hepatic lobe, increased bile duct tension accelerates bile excretion. After the right hepatic puncture, the bile duct was exposed to the external environment, leading to a decrease in intrahepatic bile

Table 2 Comparison of liver function levels and clinical efficacy before and after operation between the two groups

	Group A, <i>n</i> = 224	Group B, <i>n</i> = 200	<i>t</i> value	<i>P</i> value
Biliary tract bleeding	2	1	0.232	0.630
Duration of postoperative pain in h	2.10 ± 0.99	2.40 ± 1.89	-0.443	0.663
Peritoneal fluid leaking from around the drainage tube	13	31	10.681	0.001
Survival time in mo	26.2 ± 7.39	16.1 ± 2.72	4.054	0.001
Total bilirubin in μmol/L				
Preoperative	208.23 ± 81.81	233.52 ± 92.41	-0.648	0.525
After 3 d	152.11 ± 86.21	141.15 ± 46.55	0.354	0.728
After 1 wk	64.09 ± 26.89	88.29 ± 45.40	-1.451	0.164
Rate of decline 3 d after surgery	30.40 ± 11.84	37.59 ± 13.65	-1.259	0.224
Rate of decline 1 wk after surgery	69.23 ± 4.50	63.79 ± 5.65	2.379	0.029
Direct bilirubin in μmol/L				
Preoperative	170.73 ± 69.82	194.38 ± 72.66	-0.742	0.468
After 3 d	106.62 ± 61.87	114.37 ± 38.77	-0.336	0.741
After 1 wk	34.26 ± 23.37	72.83 ± 33.03	-3.014	0.007
Rate of decline 3 d after surgery	39.96 ± 12.05	40.05 ± 12.78	-0.015	0.988
Rate of decline 1 wk after surgery	79.30 ± 11.19	63.62 ± 5.64	3.956	0.001
Alanine aminotransferase (U/L)				
Preoperative	127.20 ± 95.72	170.10 ± 109.80	-0.931	0.364
After 3 d	64.60 ± 24.35	102.20 ± 51.81	-2.077	0.052
After 1 wk	31.60 ± 14.72	43.00 ± 13.40	-10.810	0.087
Rate of decline 3 d after surgery	35.99 ± 33.13	34.46 ± 9.89	0.140	0.890
Rate of decline 1 wk after surgery	67.84 ± 22.83	60.38 ± 25.41	0.691	0.498
Aspartate aminotransferase (U/L)				
Preoperative	254.00 ± 192.84	144.90 ± 42.79	1.746	0.098
After 3 d	51.60 ± 13.14	71.40 ± 12.17	-3.495	0.003
After 1 wk	36.60 ± 15.12	38.60 ± 16.04	-0.287	0.778
Rate of decline 3 d after surgery	72.31 ± 15.13	46.81 ± 18.41	3.383	0.003
Rate of decline 1 wk after surgery	80.45 ± 13.15	68.51 ± 18.28	1.676	0.111
Alkaline phosphatase in IU/L				
Preoperative	995.00 ± 398.24	587.70 ± 199.03	2.893	0.010
After 3 d	798.00 ± 161.08	467.00 ± 155.36	4.677	0.001
After 1 wk	371.60 ± 129.25	305.80 ± 115.78	1.199	0.246
Rate of decline 3 d after surgery	14.24 ± 17.92	19.27 ± 15.73	-0.666	0.514
Rate of decline 1 wk after surgery	60.51 ± 12.23	42.68 ± 23.56	2.125	0.048
Gamma-glutamyl transpeptidase in IU/L				
Preoperative	704.30 ± 364.56	434.80 ± 111.44	2.236	0.038
After 3 d	417.00 ± 202.68	293.80 ± 71.79	1.812	0.087
After 1 wk	181.40 ± 78.62	161.70 ± 52.93	0.657	0.519
Rate of decline 3 d after surgery	40.56 ± 10.32	32.22 ± 5.12	2.286	0.035
Rate of decline 1 wk after surgery	73.19 ± 7.05	58.81 ± 18.98	2.246	0.038

Group A: Subxiphoid left hepatic lobe approach group; Group B: Right intercostal, right hepatic lobe approach group.



DOI: 10.4240/wjgs.v15.i7.1397 Copyright ©The Author(s) 2023.

Figure 2 Kaplan-Meier survival curve analysis of the two groups. The survival rate for patients who were in the subxiphoid left hepatic lobe approach group surpassed that of the patients who were in the right intercostal, right hepatic lobe approach group ($P < 0.05$). Group A: Subxiphoid left hepatic lobe approach group; Group B: Right intercostal, right hepatic lobe approach group.

duct tension and a slower bile excretion rate. Consequently, patients experienced a gradual decrease in bilirubin levels and a prolonged recovery period from liver damage in the short term. We observed and analyzed the significantly lower peritoneal effusion leakage around the drainage tube in Group A compared to Group B, considering the diminished liver function and protein synthesis in patients with advanced tumor stages, which contributed to ascites development. The puncture location for patients who underwent percutaneous right hepatic puncture resulted in a relatively low puncture point in both supine and lateral positions. Excessive abdominal pressure frequently accompanies abdominal fluid leakage from the puncture site, substantially diminishing quality of life. In contrast, the elevated puncture point in patients receiving percutaneous left hepatic puncture led to minimal fluid exudation at the puncture site, whether standing, lying laterally, or supine, resulting in a negligible impact on the quality of life. Group A displayed a marginally higher survival rate, which we attributed to the reduced effect on quality of life, improved mood and comfort, and consequently better mental state, dietary intake, and nutrition, ultimately leading to a longer patient survival time.

Both domestic and international findings combined with our study revealed that percutaneous left hepatic biliary stenting and catheter drainage for treating low biliary obstruction substantially enhanced liver function, quality of life, postoperative pain duration, leakage of peritoneal effusion around the drainage tube, and overall patient survival.

This research was carried out retrospectively and reflected the experiences of a single institution. Additionally, our study acknowledges the constraint posed by the limited sample size, which may result in variations among some of the documented findings.

CONCLUSION

In managing low biliary obstruction, percutaneous left hepatic biliary stent placement and catheter drainage demonstrated superior therapeutic efficacy compared to percutaneous right hepatic stent placement and catheter drainage. Consequently, for patients experiencing low biliary obstruction, prioritizing percutaneous left hepatic biliary stenting and catheter drainage is recommended.

ARTICLE HIGHLIGHTS

Research background

For cases of middle and low biliary obstruction with left and right hepatic duct dilatation, the type of approach and whether different approaches affect the difficulty of puncture operation and intraoperative and postoperative complications have not been discussed in detail.

Research motivation

This study compared the clinical efficacy of two different puncture paths in treating middle and low biliary obstruction. The study prioritized the optimal puncture pathway.

Research objectives

This study compared the efficacy of different pathways to find the best improvement in patient quality of life and survival rate.

Research methods

A retrospective analysis was performed on the medical records of 424 patients with middle and low biliary obstruction who underwent percutaneous liver puncture biliary stent placement and catheter drainage between March 2016 and March 2022. Based on the puncture path, patients were categorized into two groups: subxiphoid left hepatic lobe approach group (Group A) and right intercostal, right hepatic lobe approach group (Group B). Liver function improvement, postoperative biliary bleeding incidence, postoperative pain duration, and abdominal effusion leakage around the drainage tube were compared between the two groups at 3 d and 1 wk after the operation.

Research results

The decreased rates for total bilirubin, direct bilirubin, and alkaline phosphatase 1 wk after surgery were significantly faster in Group A than in Group B. The decreased rate of gamma-glutamyl transpeptidase was also significantly faster in Group A at both 3 d and 1 wk after surgery. Group A experienced significantly less peritoneal effusion leakage around the drainage tube than Group B. The patient survival rate was higher in Group A compared to Group B.

Research conclusions

The study proposed a humanistic care perspective, improving patient survival treatment in the later stage and prioritized the optimal puncture pathway.

Research perspectives

This research was carried out retrospectively and reflected the experiences of a single institution. More studies should be performed in the future.

FOOTNOTES

Author contributions: Chen SP initiated the project and designed the experiment; Yang YB conducted clinical data collection; Yan ZY, Yang WH, and Cui Q performed postoperative follow-up and recorded data; Jiao Y conducted data collation and statistical analysis; Yang YB wrote the original manuscript; Chen SP revised the paper; all authors reviewed and approved the final manuscript.

Institutional review board statement: The Ethics Committee of Shaanxi Provincial People's Hospital approved the study.

Informed consent statement: Due to the retrospective nature of this study, the requirement for informed consent of patients was waived.

Conflict-of-interest statement: The author(s) declare having no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Data sharing statement: All data generated or analyzed during this study are included in this published article.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Si-Pan Chen 0000-0001-9313-796X.

S-Editor: Wang JL

L-Editor: Filipodia

P-Editor: Chen YL

REFERENCES

- Kang MJ**, Yun EH, Jung KW, Park SJ. Incidence, mortality and survival of gallbladder, extrahepatic bile duct, and pancreatic cancer using Korea central cancer registry database: 1999-2019. *Ann Hepatobiliary Pancreat Surg* 2022; **26**: 220-228 [PMID: 35909086 DOI: 10.14701/ahbps.22-041]

- 2 **Jung JH**, Yoon SJ, Lee OJ, Shin SH, Han IW, Heo JS. Surgical outcomes and prognostic factors of distal common bile duct adenocarcinoma: chronological analysis in a single high-volume institutional experience. *BMC Surg* 2022; **22**: 258 [PMID: [35787702](#) DOI: [10.1186/s12893-022-01649-1](#)]
- 3 **Zhang H**, Li F, Huang J, Huo C. Fishing line assisted endoscopic placement of multiple plastic biliary stents for unresectable malignant hilar biliary obstruction: a retrospective study. *BMC Gastroenterol* 2021; **21**: 435 [PMID: [34798836](#) DOI: [10.1186/s12876-021-02014-x](#)]
- 4 **Yamashita Y**, Tachikawa A, Shimokawa T, Yamazaki H, Itonaga M, Sakai Y, Sugiyama H, Nakai Y, Tanaka K, Isayama H, Kitano M. Covered versus uncovered metal stent for endoscopic drainage of a malignant distal biliary obstruction: Meta-analysis. *Dig Endosc* 2022; **34**: 938-951 [PMID: [35114036](#) DOI: [10.1111/den.14260](#)]
- 5 **Camacho JC**, Brody LA, Covey AM. Treatment of Malignant Bile Duct Obstruction: What the Interventional Radiologist Needs to Know. *Semin Intervent Radiol* 2021; **38**: 300-308 [PMID: [34393340](#) DOI: [10.1055/s-0041-1731269](#)]
- 6 **Kurita A**, Uza N, Asada M, Yoshimura K, Takemura T, Yazumi S, Kodama Y, Seno H. Stent placement above the sphincter of Oddi is a useful option for patients with inoperable malignant hilar biliary obstruction. *Surg Endosc* 2022; **36**: 2869-2878 [PMID: [34076770](#) DOI: [10.1007/s00464-021-08576-x](#)]
- 7 **Gao DJ**, Yang JF, Ma SR, Wu J, Wang TT, Jin HB, Xia MX, Zhang YC, Shen HZ, Ye X, Zhang XF, Hu B. Endoscopic radiofrequency ablation plus plastic stent placement versus stent placement alone for unresectable extrahepatic biliary cancer: a multicenter randomized controlled trial. *Gastrointest Endosc* 2021; **94**: 91-100.e2 [PMID: [33359435](#) DOI: [10.1016/j.gie.2020.12.016](#)]
- 8 **Bernon MM**, Shaw J, Burmeister S, Chinnery G, Hofmeyr S, Kloppers JC, Jonas E, Krige JE. Distal malignant biliary obstruction: a prospective randomized trial comparing plastic and uncovered self-expanding metal stents in the palliation of symptomatic jaundice. *S Afr J Surg* 2018; **56**: 30-34 [PMID: [29638090](#) DOI: [10.17159/2078-5151/2018/v56n1a2208](#)]
- 9 **Coronel M**, Lee JH, Coronel E. Endoscopic Ultrasound for the Diagnosis and Staging of Biliary Malignancy. *Clin Liver Dis* 2022; **26**: 115-125 [PMID: [34802657](#) DOI: [10.1016/j.cld.2021.08.010](#)]
- 10 **Jin H**, Pang Q, Liu H, Li Z, Wang Y, Lu Y, Zhou L, Pan H, Huang W. Prognostic value of inflammation-based markers in patients with recurrent malignant obstructive jaundice treated by reimplantation of biliary metal stents: A retrospective observational study. *Medicine (Baltimore)* 2017; **96**: e5895 [PMID: [28099348](#) DOI: [10.1097/MD.0000000000005895](#)]
- 11 **Felici C**, Mannavola F, Stucci LS, Duda L, Cafforio P, Porta C, Tucci M. Circulating tumor cells from melanoma patients show phenotypic plasticity and metastatic potential in xenograft NOD.CB17 mice. *BMC Cancer* 2022; **22**: 754 [PMID: [35820816](#) DOI: [10.1186/s12885-022-09829-1](#)]
- 12 **Sakai Y**, Iwai T, Shimura K, Gon K, Koizumi K, Ijima M, Chiba K, Nakatani S, Sugiyama H, Tsuyuguchi T, Kamisawa T, Maetani I, Kida M. Safety and efficacy of metallic stent for unresectable distal malignant biliary obstruction in elderly patients. *World J Gastroenterol* 2018; **24**: 69-75 [PMID: [29358883](#) DOI: [10.3748/wjg.v24.i1.69](#)]
- 13 **Kubo N**, Harimoto N, Shibuya K, Ishii N, Tsukagoshi M, Igarashi T, Watanabe A, Araki K, Miyazaki M, Kuwano H, Shirabe K. Successful treatment of isolated bile leakage after hepatectomy combination therapy with percutaneous transhepatic portal embolization and bile duct ablation with ethanol: a case report. *Surg Case Rep* 2018; **4**: 61 [PMID: [29915920](#) DOI: [10.1186/s40792-018-0463-y](#)]
- 14 **Hyun H**, Choi SY, Kim KA, Ko SB. Safety and Efficacy of Percutaneous Biliary Covered Stent Placement in Patients with Malignant Biliary Hilar Obstruction; Correlation with Liver Function. *Cardiovasc Intervent Radiol* 2016; **39**: 1298-1305 [PMID: [27224987](#) DOI: [10.1007/s00270-016-1375-5](#)]
- 15 **Melenhorst MC**, Scheffer HJ, Vroomen LG, Kazemier G, van den Tol MP, Meijerink MR. Percutaneous Irreversible Electroporation of Unresectable Hilar Cholangiocarcinoma (Klatskin Tumor): A Case Report. *Cardiovasc Intervent Radiol* 2016; **39**: 117-121 [PMID: [25994516](#) DOI: [10.1007/s00270-015-1126-z](#)]
- 16 **Park JK**, Woo YS, Noh DH, Yang JI, Bae SY, Yun HS, Lee JK, Lee KT, Lee KH. Efficacy of EUS-guided and ERCP-guided biliary drainage for malignant biliary obstruction: prospective randomized controlled study. *Gastrointest Endosc* 2018; **88**: 277-282 [PMID: [29605722](#) DOI: [10.1016/j.gie.2018.03.015](#)]
- 17 **Dumonceau JM**, Tringali A, Papanikolaou IS, Blero D, Mangiavillano B, Schmidt A, Vanbiervliet G, Costamagna G, Devière J, Garcia-Cano J, Gyökeres T, Hassan C, Prat F, Siersema PD, van Hooft JE. Endoscopic biliary stenting: indications, choice of stents, and results: European Society of Gastrointestinal Endoscopy (ESGE) Clinical Guideline – Updated October 2017. *Endoscopy* 2018; **50**: 910-930 [PMID: [30086596](#) DOI: [10.1055/a-0659-9864](#)]
- 18 **Inoue T**, Naitoh I, Okumura F, Ozeki T, Anbe K, Iwasaki H, Nishie H, Mizushima T, Sano H, Nakazawa T, Yoneda M, Joh T. Reintervention for stent occlusion after bilateral self-expandable metallic stent placement for malignant hilar biliary obstruction. *Dig Endosc* 2016; **28**: 731-737 [PMID: [27088888](#) DOI: [10.1111/den.12657](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

