

World Journal of *Gastrointestinal Surgery*

World J Gastrointest Surg 2023 September 27; 15(9): 1841-2097



Contents

Monthly Volume 15 Number 9 September 27, 2023

REVIEW

- 1841 Indocyanine green dye and its application in gastrointestinal surgery: The future is bright green
Lim ZY, Mohan S, Balasubramaniam S, Ahmed S, Siew CCH, Shelat VG
- 1858 Hepatic ischemia-reperfusion syndrome and its effect on the cardiovascular system: The role of treprostinil, a synthetic prostacyclin analog
Mouratidou C, Pavlidis ET, Katsanos G, Kotoulas SC, Mouloudi E, Tsoulfas G, Galanis IN, Pavlidis TE

MINIREVIEWS

- 1871 Advances and challenges of gastrostomy insertion in children
Bitar R, Azaz A, Rawat D, Hobeldin M, Miqdady M, Abdelsalam S
- 1879 Surgical decompression for the management of abdominal compartment syndrome with severe acute pancreatitis: A narrative review
Nasa P, Chanchalani G, Juneja D, Malbrain ML

ORIGINAL ARTICLE

Retrospective Cohort Study

- 1892 Excision of malignant and pre-malignant rectal lesions by transanal endoscopic microsurgery in patients under 50 years of age
Shilo Yaacobi D, Berger Y, Shaltiel T, Bekhor EY, Khalifa M, Issa N
- 1901 Safety and feasibility of modified duct-to-mucosa pancreaticojejunostomy during pancreaticoduodenectomy: A retrospective cohort study
Sun Y, Yu XF, Yao H, Xu S, Ma YQ, Chai C

Retrospective Study

- 1910 Application of early enteral nutrition nursing based on enhanced recovery after surgery theory in patients with digestive surgery
Shao YR, Ke X, Luo LH, Xu JD, Xu LQ
- 1919 Autologous bone marrow infusion *via* portal vein combined with splenectomy for decompensated liver cirrhosis: A retrospective study
Liu BC, Cheng MR, Lang L, Li L, Si YH, Li AJ, Xu Q, Zhang H
- 1932 Application of multidisciplinary collaborative nursing with family care for enhanced recovery after surgery in children with inguinal hernia
Wang XM, Hou Q
- 1941 Preoperative and postoperative complications as risk factors for delayed gastric emptying following pancreaticoduodenectomy: A single-center retrospective study
Xie FL, Ren LJ, Xu WD, Xu TL, Ge XQ, Li W, Ge XM, Zhou WK, Li K, Zhang YH, Wang Z

- 1950** Efficacy of ileus tube combined with meglumine diatrizoate in treating postoperative inflammatory bowel obstruction after surgery
Yang W, Pu J
- 1959** Effect of internet multiple linkage mode-based extended care combined with in-hospital comfort care on colorectal cancer patients undergoing colostomy
Xu L, Zhou MZ
- 1969** Short- and long-term results of open *vs* laparoscopic multisegmental resection and anastomosis for synchronous colorectal cancer located in separate segments
Quan JC, Zhou XJ, Mei SW, Liu JG, Qiu WL, Zhang JZ, Li B, Li YG, Wang XS, Chang H, Tang JQ
- 1978** Prediction model of stress ulcer after laparoscopic surgery for colorectal cancer established by machine learning algorithm
Yu DM, Wu CX, Sun JY, Xue H, Yuwen Z, Feng JX
- 1986** Effect of two surgical approaches on the lung function and prognosis of patients with combined esophago-gastric cancer
Sun CB, Han XQ, Wang H, Zhang YX, Wang MC, Liu YN
- 1995** Clinical significance of serum oxidative stress and serum uric acid levels before surgery for hepatitis B-related liver cancer
Hou JX, Wang YB, Wu J, Ding GS, Wu Y, Wei LH, Wang F, Zhang ZM
- 2003** Multifactor analysis of the technique in total laparoscopic gastric cancer
Shi JK, Wang B, Zhang XS, Lv P, Chen YL, Ren SY
- 2012** Value of enhanced computed tomography in differentiating small mesenchymal tumours of the gastrointestinal from smooth muscle tumours
Nie WJ, Jing Z, Hua M
- 2021** Risk factors for myocardial injury during living donor liver transplantation in pediatric patients with biliary atresia
Wu YL, Li TY, Gong XY, Che L, Sheng MW, Yu WL, Weng YQ

Observational Study

- 2032** Comparative detection of syndecan-2 methylation in preoperative and postoperative stool DNA in patients with colorectal cancer
Song JH, Oh TJ, An S, Lee KH, Kim JY, Kim JS

Prospective Study

- 2042** Preoperative prediction of microvascular invasion in hepatocellular carcinoma using ultrasound features including elasticity
Jiang D, Qian Y, Tan BB, Zhu XL, Dong H, Qian R
- 2052** Quantitative evaluation of colorectal tumour vasculature using contrast-enhanced ultrasound: Correlation with angiogenesis and prognostic significance
Li MH, Li WW, He L, Li JF, Zhang SY

CASE REPORT

- 2063** Laparoscopy-assisted gastrectomy for advanced gastric cancer patients with situs inversus totalis: Two case reports and review of literature
Liu HB, Cai XP, Lu Z, Xiong B, Peng CW
- 2074** Acute flare of systemic lupus erythematosus with extensive gastrointestinal involvement: A case report and review of literature
Huang H, Li P, Zhang D, Zhang MX, Yu K
- 2083** Surgical management of gallstone ileus after one anastomosis gastric bypass: A case report
El Feghali E, Akel R, Chamaa B, Kazan D, Chakhtoura G
- 2089** Dual transformation therapy for giant hepatocellular carcinoma: Two case reports and review of literature
Gao Q, Zhu GZ, Han CY, Ye XP, Huang HS, Mo ST, Peng T

ABOUT COVER

Editorial Board Member of *World Journal of Gastrointestinal Surgery*, Sung Uk Bae, MD, PhD, Associate Professor, Department of Surgery, Keimyung University Dongsan Hospital, Daegu 42601, South Korea.
sabiston0000@hanmail.net

AIMS AND SCOPE

The primary aim of *World Journal of Gastrointestinal Surgery* (WJGS, *World J Gastrointest Surg*) is to provide scholars and readers from various fields of gastrointestinal surgery with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGS mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal surgery and covering a wide range of topics including biliary tract surgical procedures, biliopancreatic diversion, colectomy, esophagectomy, esophagostomy, pancreas transplantation, and pancreatectomy, etc.

INDEXING/ABSTRACTING

The WJGS is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports/Science Edition, PubMed, PubMed Central, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJGS as 2.0; IF without journal self cites: 1.9; 5-year IF: 2.2; Journal Citation Indicator: 0.52; Ranking: 113 among 212 journals in surgery; Quartile category: Q3; Ranking: 81 among 93 journals in gastroenterology and hepatology; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Rui-Rui Wu, Production Department Director: Xiang Li, Editorial Office Director: Jia-Ru Fan.

NAME OF JOURNAL

World Journal of Gastrointestinal Surgery

ISSN

ISSN 1948-9366 (online)

LAUNCH DATE

November 30, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Peter Schemmer

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/1948-9366/editorialboard.htm>

PUBLICATION DATE

September 27, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Retrospective Study

Effect of two surgical approaches on the lung function and prognosis of patients with combined esophagogastric cancer

Chong-Bing Sun, Xiao-Qing Han, Hao Wang, Yi-Xuan Zhang, Meng-Chun Wang, Yong-Ning Liu

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B, B
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Terashima M, Japan; Wei W, China

Received: May 24, 2023

Peer-review started: May 24, 2023

First decision: June 13, 2023

Revised: July 4, 2023

Accepted: July 31, 2023

Article in press: July 31, 2023

Published online: September 27, 2023



Chong-Bing Sun, Hao Wang, Meng-Chun Wang, Yong-Ning Liu, Department of General Surgery, Weifang People's Hospital, Weifang 261041, Shandong Province, China

Xiao-Qing Han, Department of Spinal Surgery, Weifang People's Hospital, Weifang 261041, Shandong Province, China

Yi-Xuan Zhang, Department of Medical, Weifang People's Hospital, Weifang 261041, Shandong Province, China

Corresponding author: Yong-Ning Liu, MSc, Attending Doctor, Department of General Surgery, Weifang People's Hospital, No. 151 Kuiwen District, Weifang 261041, Shandong Province, China. rmyyliuyn@wfmc.edu.cn

Abstract

BACKGROUND

Adenocarcinoma of the esophagogastric junction has a center of origin within 5 cm of the esophagogastric junction. Surgical resection remains the main treatment. A transthoracic approach is recommended for Siewert I adenocarcinoma of the esophagogastric junction and a transabdominal approach is recommended for Siewert III adenocarcinoma of the esophagogastric junction. However, there is a need to determine the optimal surgical approach for Siewert II adenocarcinoma of the esophagogastric junction to improve lung function and the prognosis of patients.

AIM

To investigate and compare the surgical effects, postoperative changes in pulmonary function, and prognoses of two approaches to treating combined esophagogastric cancer.

METHODS

One hundred and thirty-eight patients with combined esophagogastric cancer treated by general and thoracic surgeries in our hospital were selected. They were divided into group A comprising 70 patients (transabdominal approach) and group B comprising 68 patients (transthoracic approach) based on the surgical approach. The indexes related to surgical trauma, number of removed lymph nodes, indexes of lung function before and after surgery, survival rate, and survival duration of the two groups were compared 3 years after surgery.

RESULTS

The duration of surgery, length of hospital stay, and postoperative drainage duration of the patients in group A were shorter than those of the patients in group B, and the volume of blood loss caused by surgery was lower for group A than for group B ($P < 0.05$). At the one-month postoperative review, the first second, maximum ventilation volume, forceful lung volume, and lung volume values were higher for group A than for group B ($P < 0.05$). Preoperatively, the QLQ-OES18 scale scores of the patients in group A were higher than those in group B on re-evaluation at 3 mo postoperatively ($P < 0.05$). The surgical complication rate of the patients in group A was 10.00%, which was lower than that of patients in group B, which was 23.53% ($P < 0.05$).

CONCLUSION

Transabdominal and transthoracic surgical approaches are comparable in treating combined esophagogastric cancer; however, the former results in lesser surgical trauma, milder changes in pulmonary function, and fewer complications.

Key Words: Transabdominal approach; Transthoracic approach; Esophagogastric Junction cancer; Pulmonary function; Prognosis; Adenocarcinoma

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Surgical resection remains the main treatment for adenocarcinoma of the esophagogastric junction. The transthoracic approach is recommended for Siewert I, and the transabdominal approach for Siewert III adenocarcinomas of the esophagogastric junction. However, the optimal surgical approach for Siewert II adenocarcinoma of the esophagogastric junction remains inconclusive. We found that the transabdominal approach has the advantage of lesser surgical trauma, lesser impact on patients' pulmonary function, and fewer complications.

Citation: Sun CB, Han XQ, Wang H, Zhang YX, Wang MC, Liu YN. Effect of two surgical approaches on the lung function and prognosis of patients with combined esophagogastric cancer. *World J Gastrointest Surg* 2023; 15(9): 1986-1994

URL: <https://www.wjgnet.com/1948-9366/full/v15/i9/1986.htm>

DOI: <https://dx.doi.org/10.4240/wjgs.v15.i9.1986>

INTRODUCTION

Adenocarcinoma of the esophagogastric junction has a center of origin within 5 cm of the esophagogastric junction. It is classified as a separate disease because of its anatomical location and biological characteristics, and its incidence is increasing significantly globally[1]. Surgical resection remains the main treatment. The transthoracic and transabdominal approaches are recommended for Siewert I and III adenocarcinomas of the esophagogastric junction, respectively, for complete resection[2]. However, the optimal surgical approach for Siewert II adenocarcinoma of the esophagogastric junction has not been established[3]. The three common approaches for the surgical treatment of Siewert II adenocarcinoma of the esophagogastric junction are transthoracic, transabdominal, and combined transthoracic and abdominal[4]. However, the transthoracic and transabdominal approaches are more commonly used. This study aimed to investigate and compare the surgical effects, changes in postoperative pulmonary function, and differences in prognosis associated with the transthoracic and transabdominal epigastric approaches for the treatment of adenocarcinoma of the esophagogastric junction.

MATERIALS AND METHODS

Information

One hundred and thirty-eight patients with combined esophagogastric cancer surgically treated by general and thoracic surgeries in our hospital were recruited between July 2015 and June 2017 for this study. Based on the surgical approach, they were divided into group A comprising 70 patients (transabdominal approach) and group B comprising 68 patients (transthoracic approach). The inclusion criteria were as follows: (1) Esophageal cancer diagnosed according to the criteria in the NCCN Esophageal Cancer Guidelines 2015 V3 edition[5]; (2) age of 19 to 75 years; (3) esophageal cancer confirmed by biopsies taken by fiberoptic esophagoscopy before surgery and Siewert type II adenocarcinoma confirmed by pathology; and (4) absence of distant metastasis during preoperative examination. The exclusion criteria were as follows: (1) Patients requiring surgery due to emergencies, such as obstruction, perforation, or bleeding; (2) coagulation disorders; (3) palliative tumor resection; (4) previous history of open-heart or open abdominal surgery; (5) preoperative history of radiotherapy; and (6) missing data. The study protocol did not violate the relevant medical ethics requirements.

The characteristics of group A were as follows: Age range of 42–75 years with an average of 58.3 ± 7.0 years; gender distribution: 40 males and 30 females; TNM tumor stages: 13 cases of stage I, 30 cases of stage II, and 27 cases of stage III; tumor lesion diameter of 5.27 ± 1.40 cm; and Lauren's staging: 50 cases of intestinal type, 6 cases of diffuse type, 14 cases of mixed type.

The characteristics of group B were as follows: Age range of 40–75 years with an average of 56.8 ± 6.9 years; gender distribution: 34 males and 34 females; tumor TNM stages: 16 cases of stage I, 30 cases of stage II, and 22 cases of stage III; tumor lesion diameter of 5.10 ± 1.35 cm; Lauren typing: 52 cases of intestinal type, 4 cases of diffuse type, and 12 cases of mixed type. The baseline information of the two patient groups was above that of the baseline comparison, and the difference was not statistically significant ($P > 0.05$).

Surgical method

All patients were diagnosed preoperatively using gastroscopy and underwent cardiac color Doppler ultrasonography, pulmonary function test, chest computed tomography (CT), abdominal CT, routine electrocardiogram, routine biochemical tests, and coagulation function tests (Figure 1).

Group A: Transabdominal approach to surgical treatment (Figure 2). After the successful induction of general anesthesia, the patient was placed on his back and disinfected routinely. An incision was made in the middle of the upper abdomen, the abdomen was opened in layers, and the transverse colon was elevated. The gastrosplenic ligament was opened along its upper edge, on the left near the splenic flexure of the colon, and on the right at the hepatic flexure of the colon, exposing the greater omentum and the anterior lobe of the transverse mesocolon. There was an upward separation along the greater curvature of the stomach. The left vessel of the gastric omentum was ligated, and the lymph nodes were swept. The primordial band of the liver and stomach was cut along the lesser curvature of the stomach on the right side of the cardia, the stomach was turned upward, and the gastric artery and vein were severed. The right and left diaphragmatic angles were opened, the anterior and posterior vagus nerve trunks were separated, and the lower end of the esophagus was freed and cut 3–5 cm from the cardia. The anastomotic stapler was placed in the esophagus, and the anastomosis was fixed in a purse string. The closing apparatus was placed close to the gastric body and sutured, and the esophagogastric anastomosis was strengthened intermittently. After the abdominal cavity was irrigated to stop bleeding, an abdominal drain was placed, and the abdomen was closed layer-by-layer. The abdominal and lower mediastinal lymph nodes were dissected.

Group B: Transthoracic approach to surgical treatment. After the successful induction of general anesthesia for transthoracic surgical access, the patient was placed in the supine position and routinely disinfected. An incision was made in the middle of the upper abdomen, and the abdomen was opened in layers. The transverse colon was elevated. The gastrosplenic ligament was opened along its upper edge, on the left near the splenic flexure of the colon, and on the right to the hepatic flexure of the colon, exposing the greater omentum and the anterior lobe of the transverse mesocolon. There was an upward separation along the greater curvature of the stomach, the left vessel of the gastric omentum was ligated, and the lymph nodes were swept. The primordial band of the liver and stomach was separated along the lesser curvature of the stomach on the right side of the cardia, the stomach was turned upward, and the gastric artery and vein were severed. The right and left diaphragmatic angles were opened, the anterior and posterior vagus nerve trunks were separated, and the lower end of the esophagus was freed and cut 3–5 cm from the cardia. The anastomotic stapler was placed in the esophagus, and the anastomosis was secured in a purse string. The closing apparatus was placed close to the gastric body and sutured, and the esophagogastric anastomosis was strengthened intermittently. After the abdominal cavity was irrigated to stop bleeding, an abdominal drain was placed, and the abdomen was closed layer-by-layer. The mediastinal and perigastric lymph nodes were dissected.


The tumor center, number of lymph nodes, and positive status were confirmed based on the postoperative pathological findings of the specimen. R staging of the residual disease was used to determine the rate of radical resection: r0 indicated no signs of a residual tumor to the naked eye and under the microscope; r1 indicated a residual tumor that was not visible to the naked eye but was visible at the margin under the microscope; and r2 indicated a tumor visible to the naked eye at the cut edge.

Indices for observation and detection methods

We compared the duration of surgery, bleeding volume, number of cleared lymph nodes, number of positive lymph nodes, positive margin rate, and length of hospital stay. We also compared the two groups based on the four indicators of forced expiratory volume within the first second (FEV1), maximum ventilation volume (MVV), forceful lung volume (FVC), and lung volume (VC) as percentages of their expected values; surgical complications; and 3-year survival rates and durations before and after surgery.


Pulmonary function was measured preoperatively and 1 mo postoperatively. A JAEGER Flowscreen pulmonary function tester (Jaeger, Germany) was used to examine the patients while fully awake and in the sitting position[6]. The main indices included VC, FVC, FEV, and MVV, and the data are expressed as percentages of the actual value to the desired value.

The quality of survival was assessed preoperatively and at 3 mo postoperatively. The QLQ-OES18 scale for esophageal cancer[7] was used to evaluate the quality of survival; it contains 19 questionnaire items, 18 of which have a score range of 0–3, and one which has a score range of 0–4, with a total score of 58. Higher scores indicated higher quality of survival for the patient.



DOI: 10.4240/wjgs.v15.i9.1986 Copyright ©The Author(s) 2023.

Figure 1 The approaches. A: Transabdominal approach; B: Transthoracic approach.



DOI: 10.4240/wjgs.v15.i9.1986 Copyright ©The Author(s) 2023.

Figure 2 Survival function diagram of the two patient groups.

Statistical analysis

Statistical analysis was performed using SPSS 21.0 software. Measurement data such as duration of surgery, length of hospital stay, and bleeding volume were expressed as mean and standard deviation ($\bar{x} \pm s$) for both groups. The *t*-test was used to compare the two groups. The χ^2 test was used to compare the counting data of the groups. The Kaplan-Meier method was used for survival analysis. Statistical significance was set at $P < 0.05$.

RESULTS

Comparison of the durations of surgery of the patient groups

The durations of surgery, lengths of hospital stay, and postoperative drainage durations of patients in group A were shorter than those in group B. Bleeding caused by surgery was lower in group A than in group B ($P < 0.05$), and there was no statistically significant difference in the positive incision margin rate between the patients in groups A and B ($P > 0.05$) (Table 1).

Comparison of lymph node dissection in the two groups

There were no statistically significant differences in the number of cleared lymph nodes, positive lymph nodes, lower mediastinal lymph nodes, subdiaphragmatic lymph nodes, or abdominal lymph nodes between groups A and B ($P > 0.05$) (Table 2).

Table 1 Comparison of operation time between two groups of patients

Group	n	Operation time (min)	Bleeding volume (mL)	Positive rate of cutting edge (%)	Postoperative drainage time (d)	Postoperative landing time (h)	Length of stay (d)
A group	70	168.1 ± 15.7	136.8 ± 36.1	1 (1.43)	3.71 ± 0.84	27.81 ± 7.51	12.30 ± 2.13
B group	68	188.0 ± 19.3	188.2 ± 43.7	3 (4.41)	4.40 ± 1.22	29.40 ± 7.82	13.54 ± 2.35
t/ χ^2		-6.653	-7.542	1.091	-3.879	-1.218	-3.250
P value		0.000	0.000	0.296	0.000	0.225	0.001

Table 2 Comparison of lymph node dissection effect between two groups of patients (mean ± SD, number)

Group	n	Number of lymph nodes cleaned	Number of positive lymph nodes	Inferior mediastinal lymph nodes	Subphrenic lymph node	Number of abdominal lymph nodes
A group	70	33.87 ± 3.82	3.65 ± 1.20	3.81 ± 1.20	2.56 ± 0.72	27.50 ± 2.95
B group	68	34.33 ± 2.90	3.92 ± 1.53	4.03 ± 1.15	2.74 ± 0.75	27.56 ± 3.02
t value		-0.795	-1.155	-1.099	-1.438	-0.118
P value		0.428	0.250	0.274	0.153	0.906

Comparison of changes in pulmonary function in the two groups

The preoperative values of FEV1, MVV, FVC, and VC for the patients in groups A and B were not significantly different ($P > 0.05$). However, the values obtained during the review conducted one month after surgery showed higher percentages of FEV1, MVV, FVC, and VC relative to their expected values for group A than for group B ($P < 0.05$) (Table 3).

Comparison of the quality of the survival scores for the two groups

Before surgery, the QLQ-OES18 scores of patients in groups A and B were compared, and the difference was not statistically significant ($P > 0.05$). When re-evaluated 3 mo after surgery, the QLQ-OES18 scores for the patients in group A were higher than those for the patients in group B ($P < 0.05$) (Table 4).

Comparison of the complication rates of the two groups

The surgical complication rate of 10.00% for group A was lower than that of 23.53% for group B ($P < 0.05$; Table 5).

Comparison of the prognoses of the two groups of patients

After 3 years of postoperative follow-up, there was no statistically significant difference in the survival rate of 35.71% for group A relative to 29.41% for group B ($P > 0.05$) (Table 6).

The median duration of survival was 30.0 mo for the patients in group A and 29.0 mo for those in group B. The difference between the two groups was not statistically significant ($P > 0.05$) (Figure 2).

DISCUSSION

Adenocarcinoma of the esophagogastric junction is considered a special type of tumor because of its anatomical location and physiological function. It is independent of esophageal and gastric cancers and is more common in patients with Siewert II and III types[8]. Surgery for adenocarcinoma of the esophagogastric junction is usually performed by a gastrointestinal or thoracic surgeon or both. However, the choice of surgical approach for the Siewert II type has not been established. The advantage of combined thoracoabdominal therapy for adenocarcinoma of the esophagogastric junction is that it allows for complete dissection of the abdominal and mediastinal lymph nodes. However, this surgical approach is more invasive, increases the risk of surgery, and does not significantly improve the long-term survival of patients; therefore, most scholars recommend the transthoracic or transabdominal approach[9]. The advantage of the transthoracic approach is that it can completely expose the structures and tissues of the esophagus and cardia; however, exposure of the tissues near the distal stomach and spleen is poor. The transabdominal approach can fully expose the abdomen and facilitate the dissection of the abdominal lymph node; however, exposure to the distal esophagus is poor[9-11].

The duration of surgery and the severity of intraoperative blood loss are lateral reflections of surgical trauma and are interrelated[12]. Less intraoperative bleeding ensures less obstruction of the surgical field and a shorter duration of

Table 3 Comparison of pulmonary function parameters between two groups of patients (mean \pm SD)

Group	n	Preoperative	1 mo after operation	t value	P value	Preoperative	1 mo after operation	t value	P value
		FEV1 (%)				MVV (%)			
A group	70	95.66 \pm 8.64	90.21 \pm 8.50	3.734	0.000	98.16 \pm 9.26	93.48 \pm 9.11	2.992	0.003
B group	68	97.03 \pm 8.11	86.30 \pm 7.76	7.937	0.000	96.32 \pm 8.58	88.75 \pm 8.36	5.248	0.000
t value		-0.960	2.820			1.210	3.175		
P value		0.339	0.006			0.228	0.002		
		FVC (%)				VC (%)			
A group	70	97.34 \pm 8.14	92.36 \pm 6.06	4.067	0.000	93.06 \pm 4.85	90.01 \pm 4.43	3.854	0.000
B group	68	99.03 \pm 7.93	89.51 \pm 7.24	7.359	0.000	94.41 \pm 5.00	88.26 \pm 5.25	7.048	0.000
t value		-1.235	2.510			-1.610	2.119		
P value		0.219	0.013			0.11	0.036		

FEV1: First second; MVV: Maximum ventilation volume; FVC: Forceful lung volume; VC: Lung volume.

Table 4 Comparison of quality of life scores between the two groups (mean \pm SD, scores)

Group	n	Preoperative	3 mo after operation	t value	P value
A group	70	31.83 \pm 6.60	43.09 \pm 5.57	-10.816	0.000
B group	68	30.50 \pm 5.78	40.14 \pm 5.42	-10.100	0.000
t value		1.258	3.152		
P value		0.211	0.002		

Table 5 Comparison of complication rates between the two groups

Group	n	Pulmonary infection	Abdominal infection	Anastomotic fistula	Pyothorax	Incision infection	Pleural effusion	Complication rate (%)
A group	70	1	2	1	0	1	2	7 (10.00)
B group	68	7	0	3	2	1	3	16 (23.53)
χ^2								4.546
P value								0.033

Table 6 Survival comparison

Group	n	Subsist	Die
A group	70	25 (35.71)	45 (64.29)
B group	68	20 (29.41)	48 (70.59)
χ^2		0.623	
P value		0.430	

bleeding. The results of this study showed that the duration of surgery, length of hospital stay, and postoperative drainage of the patients in group A were shorter than those of the patients in group B. The blood loss during surgery was significantly lower for Group A than for Group B. This indicates that the transabdominal is less traumatic than the transthoracic approach. The pectoral muscle or the rib cage of the patient needs to be severed to expose the left side of the chest cavity, which is rich in intercostal vessels, and bleeding can easily occur when opening the chest. The transabdominal approach, which involves entering the abdominal cavity through the white line of the abdomen, results in less severe blood loss and a shorter duration of surgery. However, it is difficult to expose the organs and lymph nodes in the

abdominal cavity, the scope of surgery is relatively small, the duration of surgery is prolonged, and intraoperative bleeding and duration of surgery may be further prolonged if adhesions and anatomical abnormalities are present in the abdominal cavity during exposure. The results of this study showed no statistically significant difference in the positive margin rate between the two groups, indicating that the outcomes of the two surgical approaches were comparable. Complete removal of regional lymph nodes is a key factor for the long-term survival of patients with Siewert type II adenocarcinoma of the esophagogastric junction and helps improve the long-term survival of patients during the progressive stage[13]. Studies have shown[14-18] that lymph node metastases are more likely to metastasize to the abdominal cavity than to the thoracic cavity in patients with type II esophagogastric junction cancer. Therefore, more attention should be paid to the removal of abdominal lymph nodes during surgery for type II esophagogastric junction cancer. In this study, there were no statistically significant differences in the number of removed lymph nodes, positive lymph nodes, lower mediastinal lymph nodes, subdiaphragmatic lymph nodes, or abdominal lymph nodes between the two groups. Increased attention of the surgeon combined with the use of laparoscopy and other techniques has resulted in no significant difference in the number of positive lymph node dissections associated with the current transabdominal and transthoracic approaches.

FEV1, MVV, FVC, and VC, expressed as percentages of the expected values, were higher for group A than for group B. Both groups showed different degrees of decline. Owing to the residual effects of general anesthesia, early postoperative pain, and the use of a chest strap, chest compliance decreased, and respiratory function decreased. The differences in postoperative pulmonary function associated with the two surgical approaches in this study were mainly attributed to the transabdominal approach through the right anterolateral incision, which maintained the integrity of the diaphragm during open thoracotomy with relatively little damage to the chest wall muscles. In addition, a shorter duration of surgery of the thoracic cavity resulted in less interference with the lung tissue. The transthoracic approach caused more severe damage to the chest wall muscles, impairment of the diaphragmatic integrity, greater interference with the lung tissue, and a greater deterioration of lung function. Group A had a significantly lower rate of surgical complications than group B. The transthoracic approach may have disrupted the normal muscles of the chest during the left thoracic incision, requiring incision and re-suturing of the diaphragm and damaging the respiratory muscles; this was detrimental to postoperative sputum expulsion and affected respiratory function. The intraoperative collapse of the left lung; ventilation of the right lung; intra-thoracic surgical involvement and compression of the lung tissue, heart, and blood vessels; and postoperative chest tube placement increase the inflammatory response in the thoracic cavity. Transthoracic surgery destroys the intercostal nerves of the patient, results in more severe postoperative pain than abdominal surgery, and is more likely to result in sputum accumulation. As a result, the incidence of pulmonary complications was higher for group B, which may have led to longer postoperative hospital stays. Patients in group A had higher QLQ-OES18 scores than those in group B when re-evaluated 3 mo postoperatively. Patients who underwent surgery with the transabdominal approach had a better prognosis. The 3-year postoperative follow-up evaluation revealed no statistically significant differences between the survival rates and median durations of survival of the two groups. The results of this study are consistent with those of previous studies[3,19-23], showing that the treatment of Siewert II esophagogastric junction cancer through the left thorax and abdomen is equally reliable.

Current clinical studies on the treatment of combined esophagogastric cancer mainly compare the efficacies and adverse effects of the different approaches[24-27]; they less frequently focus on the changes in the parameters of pulmonary function. In this study, the clinical outcomes, lymph node removal, and lung function recovery of patients with combined esophagogastric cancer treated using transabdominal and transthoracic surgical approaches were studied comprehensively. This was more conducive to finding a more advantageous surgical approach. There were some limitations and shortcomings in the design of this study. It was a single-center retrospective study with a short duration of follow-up. Therefore, the results need to be confirmed by prospective, multicenter, randomized controlled clinical studies.

CONCLUSION

Transabdominal and transthoracic surgical approaches for the treatment of combined esophagogastric cancer are comparable. However, the former has the advantages of milder surgical trauma, less impact on pulmonary function, and fewer complications. Thus, it is suitable for older patients with frailty, cardiopulmonary insufficiency, or more complicated diseases.

ARTICLE HIGHLIGHTS

Research background

Different types of esophagogastric junction adenocarcinoma have different operation methods.

Research motivation

We need to determine the optimal surgical approach for Siewert II adenocarcinoma of the esophagogastric junction to improve lung function and the prognosis of patients.

Research objectives

To investigate and compare the surgical effects, postoperative changes in pulmonary function, and prognoses of two approaches to treating combined esophagogastric cancer.

Research methods

Patients with esophageal gastric cancer who received combined treatment in our hospital were selected, and the relevant indicators were compared after grouping.

Research results

The transabdominal approach has the advantages of less trauma, less impact on lung function and fewer complications.

Research conclusions

Transabdominal surgical approaches is suitable for older patients with frailty, cardiopulmonary insufficiency, or more complicated diseases.

Research perspectives

It was a single-center retrospective study with a short duration of follow-up. Therefore, the results need to be confirmed by prospective, multicenter, randomized controlled clinical studies.

FOOTNOTES

Author contributions: Sun CB and Liu YN designed the research study; Sun CB, Han XQ and Wang H performed the research; Zhang YX contributed new reagents and analytic tools; Wang MC and Sun CB analyzed the data and wrote the manuscript; All authors have read and approve the final manuscript.

Institutional review board statement: The study was reviewed and approved by the Weifang People's Hospital Institutional Review Board, No. AF/SQ-01/01.0.

Informed consent statement: All study participants or their legal guardian provided informed written consent about personal and medical data collection prior to study enrolment.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

Data sharing statement: No additional data are available.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Chong-Bing Sun 0009-0008-5472-1168; Xiao-Qing Han 0009-0003-9739-8317; Hao Wang 0009-0006-3007-9516; Yi-Xuan Zhang 0009-0002-5501-965X; Meng-Chun Wang 0009-0009-9127-8318; Yong-Ning Liu 0009-0000-7926-3265.

S-Editor: Fan JR

L-Editor: A

P-Editor: Cai YX

REFERENCES

- Chevallay M**, Bollschweiler E, Chandramohan SM, Schmidt T, Koch O, Demanzoni G, Mönig S, Allum W. Cancer of the gastroesophageal junction: a diagnosis, classification, and management review. *Ann N Y Acad Sci* 2018; **1434**: 132-138 [PMID: 30138540 DOI: 10.1111/nyas.13954]
- Türeci O**, Sahin U, Schulze-Bergkamen H, Zvirbulis Z, Lordick F, Koeberle D, Thuss-Patience P, Ettrich T, Arnold D, Bassermann F, Al-Batran SE, Wiechen K, Dhaene K, Maurus D, Gold M, Huber C, Krivoschik A, Arozullah A, Park JW, Schuler M. A multicentre, phase IIa study of zolbetuximab as a single agent in patients with recurrent or refractory advanced adenocarcinoma of the stomach or lower oesophagus: the MONO study. *Ann Oncol* 2019; **30**: 1487-1495 [PMID: 31240302 DOI: 10.1093/annonc/mdz199]
- Kumamoto T**, Kurahashi Y, Niwa H, Nakanishi Y, Okumura K, Ozawa R, Ishida Y, Shinohara H. True esophagogastric junction adenocarcinoma: background of its definition and current surgical trends. *Surg Today* 2020; **50**: 809-814 [PMID: 31278583 DOI: 10.1007/s00595-019-01843-4]
- Urabe M**, Matsusaka K, Ushiku T, Fukuyo M, Rahmutulla B, Yamashita H, Seto Y, Fukayama M, Kaneda A. Adenocarcinoma of the stomach and esophagogastric junction with low DNA methylation show poor prognoses. *Gastric Cancer* 2023; **26**: 95-107 [PMID: 36224483 DOI: 10.1007/s10120-022-01111-1]

- 10.1007/s10120-022-01344-3]
- 5 **Manabe N**, Matsueda K, Haruma K. Epidemiological Review of Gastroesophageal Junction Adenocarcinoma in Asian Countries. *Digestion* 2022; **103**: 29-36 [PMID: [34718236](#) DOI: [10.1159/000519602](#)]
- 6 **Treese C**, Hartl K, Pötzsch M, Dahlmann M, von Winterfeld M, Berg E, Hummel M, Timm L, Rau B, Walther W, Daum S, Kobelt D, Stein U. S100A4 Is a Strong Negative Prognostic Marker and Potential Therapeutic Target in Adenocarcinoma of the Stomach and Esophagus. *Cells* 2022; **11** [PMID: [35326507](#) DOI: [10.3390/cells11061056](#)]
- 7 **Zhang CD**, Takeshima H, Sekine S, Yamashita S, Liu YY, Hattori N, Abe H, Yamashita H, Fukuda M, Imamura Y, Ushiku T, Katai H, Makino H, Watanabe M, Seto Y, Ushijima T. Prediction of tissue origin of adenocarcinomas in the esophagogastric junction by DNA methylation. *Gastric Cancer* 2022; **25**: 336-345 [PMID: [34557982](#) DOI: [10.1007/s10120-021-01252-y](#)]
- 8 **Jung MK**, Schmidt T, Chon SH, Chevally M, Berlth F, Akiyama J, Gutschow CA, Mönig SP. Current surgical treatment standards for esophageal and esophagogastric junction cancer. *Ann N Y Acad Sci* 2020; **1482**: 77-84 [PMID: [32798235](#) DOI: [10.1111/nyas.14454](#)]
- 9 **Suhara H**, Hirooka Y, Kawashima H, Ohno E, Ishikawa T, Nakamura M, Miyahara R, Ishigami M, Hashimoto S, Goto H. Transabdominal ultrasound elastography of the esophagogastric junction predicts reflux esophagitis. *J Med Ultrason (2001)* 2019; **46**: 99-104 [PMID: [30043241](#) DOI: [10.1007/s10396-018-0890-1](#)]
- 10 **Dai Z**, Lang W, Yang H, Tian J, Sun W, Pekbay B, Lin Y, Wang M, Cui B, Yang S, Li H, Luo L, Guo H, Zhang L. Validation of EORTC QLQ-OES18 for Chinese patients with esophageal cancer. *Dis Esophagus* 2017; **30**: 1-7 [PMID: [28859386](#) DOI: [10.1093/dote/dox046](#)]
- 11 **DeCarlo C**, Manxhari C, Boitano LT, Mohebbi J, Schwartz SI, Eagleton MJ, Conrad MF. Transabdominal approach associated with increased long-term laparotomy complications after open abdominal aortic aneurysm repair. *J Vasc Surg* 2021; **73**: 1603-1610 [PMID: [33080323](#) DOI: [10.1016/j.jvs.2020.08.154](#)]
- 12 **Trevellin E**, Pirozzolo G, Fassan M, Vettor R. Prognostic value of stem cell markers in esophageal and esophagogastric junction cancer: a meta-analysis. *J Cancer* 2020; **11**: 4240-4249 [PMID: [32368307](#) DOI: [10.7150/jca.33699](#)]
- 13 **Ajani JA**, D'Amico TA, Bentrem DJ, Chao J, Corvera C, Das P, Denlinger CS, Enzinger PC, Fanta P, Farjah F, Gerdes H, Gibson M, Glasgow RE, Hayman JA, Hochwald S, Hofstetter WL, Ilson DH, Jaroszewski D, Johung KL, Keswani RN, Kleinberg LR, Leong S, Ly QP, Matkowskyj KA, McNamara M, Mulcahy MF, Paluri RK, Park H, Perry KA, Pimiento J, Poultides GA, Roses R, Strong VE, Wiesner G, Willett CG, Wright CD, McMillian NR, Pluchino LA. Esophageal and Esophagogastric Junction Cancers, Version 2.2019, NCCN Clinical Practice Guidelines in Oncology. *J Natl Compr Canc Netw* 2019; **17**: 855-883 [PMID: [31319389](#) DOI: [10.6004/jnccn.2019.0033](#)]
- 14 **van Workum F**, Versteegen MHP, Klarenbeek BR, Bouwense SAW, van Berge Henegouwen MI, Daams F, Gisbertz SS, Hannink G, Haveman JW, Heisterkamp J, Jansen W, Kouwenhoven EA, van Lanschot JJB, Nieuwenhuijzen GAP, van der Peet DL, Polat F, Ubels S, Wijnhoven BPL, Rovers MM, Rosman C; ICAN collaborative research group. Intrathoracic vs Cervical Anastomosis After Totally or Hybrid Minimally Invasive Esophagectomy for Esophageal Cancer: A Randomized Clinical Trial. *JAMA Surg* 2021; **156**: 601-610 [PMID: [33978698](#) DOI: [10.1001/jamasurg.2021.1555](#)]
- 15 **Lee Y**, Min SH, Park KB, Park YS, Ahn SH, Park DJ, Kim HH. Long-term Outcomes of Laparoscopic Versus Open Transhiatal Approach for the Treatment of Esophagogastric Junction Cancer. *J Gastric Cancer* 2019; **19**: 62-71 [PMID: [30944759](#) DOI: [10.5230/jgc.2019.19.e1](#)]
- 16 **Berlth F**, Hoelscher AH. History of Esophagogastric Junction Cancer Treatment and Current Surgical Management in Western Countries. *J Gastric Cancer* 2019; **19**: 139-147 [PMID: [31245158](#) DOI: [10.5230/jgc.2019.19.e18](#)]
- 17 **Hoshino I**, Gunji H, Ishige F, Iwatate Y, Takiguchi N, Ikeda A, Soda H, Tonooka T, Sato N, Kawahara K, Nabeya Y. Surgical treatment strategy for esophagogastric junction cancers based on the tumor diameter. *BMC Surg* 2019; **19**: 152 [PMID: [31651313](#) DOI: [10.1186/s12893-019-0614-5](#)]
- 18 **Shinohara H**, Haruki S, Ohata Y, Ueda H, Arita K, Ito K, Matsumoto A, Takiguchi N. [A Case of Advanced Esophagogastric Junction Cancer Responding to Combined Modality Therapy]. *Gan To Kagaku Ryoho* 2020; **47**: 466-468 [PMID: [32381917](#)]
- 19 **Pericay C**, Macías-Declara I, Arrazubi V, Vilà L, Marín M. Treatment in esophagogastric junction cancer: Past, present and future. *Cir Esp (Engl Ed)* 2019; **97**: 459-464 [PMID: [31155142](#) DOI: [10.1016/j.ciresp.2019.03.016](#)]
- 20 **Hosoda K**, Yamashita K, Tsuruta H, Moriya H, Mieno H, Ema A, Washio M, Watanabe M. Prognoses of advanced esophago-gastric junction cancer may be modified by thoracotomy and splenectomy. *Oncol Lett* 2018; **15**: 1200-1210 [PMID: [29399174](#) DOI: [10.3892/ol.2017.7441](#)]
- 21 **Takada K**, Yabuuchi Y, Yamamoto Y, Yoshida M, Kawata N, Takizawa K, Kishida Y, Ito S, Imai K, Hotta K, Ishiwatari H, Matsubayashi H, Kawabata T, Ono H. Predicting the depth of superficial adenocarcinoma of the esophagogastric junction. *J Gastroenterol Hepatol* 2022; **37**: 363-370 [PMID: [34820917](#) DOI: [10.1111/jgh.15741](#)]
- 22 **Thompson ED**, Zahurak M, Murphy A, Cornish T, Cuka N, Abdelfatah E, Yang S, Duncan M, Ahuja N, Taube JM, Anders RA, Kelly RJ. Patterns of PD-L1 expression and CD8 T cell infiltration in gastric adenocarcinomas and associated immune stroma. *Gut* 2017; **66**: 794-801 [PMID: [26801886](#) DOI: [10.1136/gutjnl-2015-310839](#)]
- 23 **Chen Y**, Hu L, Lin H, Yu H, You J. Serum metabolomic profiling for patients with adenocarcinoma of the esophagogastric junction. *Metabolomics* 2022; **18**: 26 [PMID: [35441991](#) DOI: [10.1007/s11306-022-01883-x](#)]
- 24 **Zheng YH**, Zhao EH. Recent advances in multidisciplinary therapy for adenocarcinoma of the esophagus and esophagogastric junction. *World J Gastroenterol* 2022; **28**: 4299-4309 [PMID: [36159003](#) DOI: [10.3748/wjg.v28.i31.4299](#)]
- 25 **Tang Z**, Wang Y, Liu D, Wang X, Xu C, Yu Y, Cui Y, Tang C, Li Q, Sun J, Zhang Q, Ji Y, Ma G, Li H, Shen Z, Shen K, Zheng R, Hou Z, Liu T, Wang J, Sun Y. The Neo-PLANET phase II trial of neoadjuvant camrelizumab plus concurrent chemoradiotherapy in locally advanced adenocarcinoma of stomach or gastroesophageal junction. *Nat Commun* 2022; **13**: 6807 [PMID: [36357415](#) DOI: [10.1038/s41467-022-34403-5](#)]
- 26 **Rice TW**, Lu M, Ishwaran H, Blackstone EH; Worldwide Esophageal Cancer Collaboration Investigators. Precision Surgical Therapy for Adenocarcinoma of the Esophagus and Esophagogastric Junction. *J Thorac Oncol* 2019; **14**: 2164-2175 [PMID: [31442498](#) DOI: [10.1016/j.jtho.2019.08.004](#)]
- 27 **Kumar NA**, Desouza A, Bhandare MS, Murugan JR, Khandelwal G, Chaudhari V, Ostwal V, Shrikhande SV. Curative resection for adenocarcinoma of the gastro-esophageal junction following neo-adjuvant chemotherapy-thoraco-abdominal vs. trans-abdominal approach. *Langenbecks Arch Surg* 2021; **406**: 613-621 [PMID: [33242137](#) DOI: [10.1007/s00423-020-02020-9](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

