


World Journal of *Gastrointestinal Surgery*

World J Gastrointest Surg 2024 January 27; 16(1): 1-259



EDITORIAL

- 1 Novel prognostic factors after radical resection of hepatocellular carcinoma: Updating an old issue
Bencini L
- 6 Prospects in the application of ultrasensitive chromosomal aneuploidy detection in precancerous lesions of gastric cancer
Qian ST, Xie FF, Zhao HY, Liu QS, Cai DL

MINIREVIEWS

- 13 Prognostic value of ultrasound in early arterial complications post liver transplant
Zhao NB, Chen Y, Xia R, Tang JB, Zhao D

ORIGINAL ARTICLE**Case Control Study**

- 21 Added value of ratio of cross diameters of the appendix in ultrasound diagnosis of acute appendicitis
Gu FW, Wu SZ

Retrospective Cohort Study

- 29 Oncological features and prognosis of colorectal cancer in human immunodeficiency virus-positive patients: A retrospective study
Yang FY, He F, Chen DF, Tang CL, Woraiikat S, Li Y, Qian K

Retrospective Study

- 40 Laparoscopic *vs* open surgery for gastric cancer: Assessing time, recovery, complications, and markers
Lu YY, Li YX, He M, Wang YL
- 49 Single-incision laparoscopic transabdominal preperitoneal repair in the treatment of adult female patients with inguinal hernia
Zhu XJ, Jiao JY, Xue HM, Chen P, Qin CF, Wang P
- 59 Computerized tomography-guided therapeutic percutaneous puncture catheter drainage-combined with somatostatin for severe acute pancreatitis: An analysis of efficacy and safety
Zheng XL, Li WL, Lin YP, Huang TL
- 67 Impact of open hepatectomy on postoperative bile leakage in patients with biliary tract cancer
Wu G, Li WY, Gong YX, Lin F, Sun C
- 76 Clinical observation of gastrointestinal function recovery in patients after hepatobiliary surgery
Zeng HJ, Liu JJ, Yang YC

- 85 Predictive value of machine learning models for lymph node metastasis in gastric cancer: A two-center study
Lu T, Lu M, Wu D, Ding YY, Liu HN, Li TT, Song DQ
- 95 Post-operative morbidity after neoadjuvant chemotherapy and resection for gallbladder cancer: A national surgical quality improvement program analysis
Kim M, Stroever S, Aploks K, Ostapenko A, Dong XD, Seshadri R
- 103 Risk factors for recurrence of common bile duct stones after surgical treatment and effect of ursodeoxycholic acid intervention
Yuan WH, Zhang Z, Pan Q, Mao BN, Yuan T
- 113 Clinical efficacy of modified Kamikawa anastomosis in patients with laparoscopic proximal gastrectomy
Wu CY, Lin JA, Ye K
- 124 Clinical effect of laparoscopic radical resection of colorectal cancer based on propensity score matching
Liu Y, Wang XX, Li YL, He WT, Li H, Chen H
- 134 Different timing for abdominal paracentesis catheter placement and drainage in severe acute pancreatitis complicated by intra-abdominal fluid accumulation
Chen R, Chen HQ, Li RD, Lu HM
- 143 Comparison of different preoperative objective nutritional indices for evaluating 30-d mortality and complications after liver transplantation
Li C, Chen HX, Lai YH
- 155 Predictive value of NLR, Fib4, and APRI in the occurrence of liver failure after hepatectomy in patients with hepatocellular carcinoma
Kuang TZ, Xiao M, Liu YF
- 166 Practical effect of different teaching modes in teaching gastrointestinal surgery nursing
Rong XJ, Ning Z
- Observational Study**
- 173 Predictive factors and model validation of post-colon polyp surgery *Helicobacter pylori* infection
Zhang ZS
- Randomized Controlled Trial**
- 186 Micro-power negative pressure wound technique reduces risk of incision infection following loop ileostomy closure
Xu DY, Bai BJ, Shan L, Wei HY, Lin DF, Wang Y, Wang D
- 196 Paravertebral block's effect on analgesia and inflammation in advanced gastric cancer patients undergoing transarterial chemoembolization and microwave ablation
Xiong YF, Wei BZ, Wang YF, Li XF, Liu C

META-ANALYSIS

- 205 Unraveling the efficacy network: A network meta-analysis of adjuvant external beam radiation therapy methods after hepatectomy
Yang GY, He ZW, Tang YC, Yuan F, Cao MB, Ren YP, Li YX, Su XR, Yao ZC, Deng MH
- 215 Estimation of Physiologic Ability and Surgical Stress scoring system for predicting complications following abdominal surgery: A meta-analysis spanning 2004 to 2022
Pang TS, Cao LP
- 228 Role of Oncostatin M in the prognosis of inflammatory bowel disease: A meta-analysis
Yang Y, Fu KZ, Pan G

CASE REPORT

- 239 Endoscopic treatment of extreme esophageal stenosis complicated with esophagotracheal fistula: A case report
Fang JH, Li WM, He CH, Wu JL, Guo Y, Lai ZC, Li GD
- 248 Intestinal tuberculosis with small bowel stricture and hemorrhage as the predominant manifestation: Three case reports
Huang G, Wu KK, Li XN, Kuai JH, Zhang AJ

LETTER TO THE EDITOR

- 257 Sarcopenia in cirrhotic patients: Does frailty matter while waiting for a liver transplant?
Li XJ, He K

ABOUT COVER

Editorial Board Member of *World Journal of Gastrointestinal Surgery*, Renato Pietroletti, PhD, Associate Professor, Professor, Department of Applied Clinical and Biotechnological Sciences, University of L'Aquila, L'Aquila 67100, AQ, Italy. renato.pietroletti@univaq.it

AIMS AND SCOPE

The primary aim of *World Journal of Gastrointestinal Surgery* (*WJGS, World J Gastrointest Surg*) is to provide scholars and readers from various fields of gastrointestinal surgery with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGS mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal surgery and covering a wide range of topics including biliary tract surgical procedures, biliopancreatic diversion, colectomy, esophagectomy, esophagostomy, pancreas transplantation, and pancreatectomy, *etc.*

INDEXING/ABSTRACTING

The *WJGS* is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports/Science Edition, PubMed, PubMed Central, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for *WJGS* as 2.0; IF without journal self cites: 1.9; 5-year IF: 2.2; Journal Citation Indicator: 0.52; Ranking: 113 among 212 journals in surgery; Quartile category: Q3; Ranking: 81 among 93 journals in gastroenterology and hepatology; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Zi-Hang Xu, Production Department Director: Xiang Li, Editorial Office Director: Jia-Ru Fan.

NAME OF JOURNAL

World Journal of Gastrointestinal Surgery

ISSN

ISSN 1948-9366 (online)

LAUNCH DATE

November 30, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Peter Schemmer

POLICY OF CO-AUTHORS

<https://www.wjgnet.com/1948-9366/editorialboard.htm>

PUBLICATION DATE

January 27, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

<https://www.wjgnet.com/bpg/GerInfo/310>

ARTICLE PROCESSING CHARGE


<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Observational Study

Predictive factors and model validation of post-colon polyp surgery *Helicobacter pylori* infection

Zheng-Sen Zhang

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): C, C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Bhansali A, India;
Rastogi A, India

Received: November 6, 2023

Peer-review started: November 6, 2023

First decision: November 16, 2023

Revised: December 3, 2023

Accepted: December 20, 2023

Article in press: December 20, 2023

Published online: January 27, 2024



Zheng-Sen Zhang, Health Management Center, Affiliated Hospital of Shaoxing University (Shaoxing Municipal Hospital), Shaoxing 312000, Zhejiang Province, China

Corresponding author: Zheng-Sen Zhang, BMed, Doctor, Health Management Center, Affiliated Hospital of Shaoxing University (Shaoxing Municipal Hospital), No. 999 Zhongxing South Road, Yuecheng District, Shaoxing 312000, Zhejiang Province, China.
sxzs02@126.com

Abstract

BACKGROUND

Recently, research has linked *Helicobacter pylori* (*H. pylori*) stomach infection to colonic inflammation, mediated by toxin production, potentially impacting colorectal cancer occurrence.

AIM

To investigate the risk factors for post-colon polyp surgery, *H. pylori* infection, and its correlation with pathologic type.

METHODS

Eighty patients who underwent colon polypectomy in our hospital between January 2019 and January 2023 were retrospectively chosen. They were then randomly split into modeling ($n = 56$) and model validation ($n = 24$) sets using R. The modeling cohort was divided into an *H. pylori*-infected group ($n = 37$) and an *H. pylori*-uninfected group ($n = 19$). Binary logistic regression analysis was used to analyze the factors influencing the occurrence of *H. pylori* infection after colon polyp surgery. A roadmap prediction model was established and validated. Finally, the correlation between the different pathological types of colon polyps and the occurrence of *H. pylori* infection was analyzed after colon polyp surgery.

RESULTS

Univariate results showed that age, body mass index (BMI), literacy, alcohol consumption, polyp pathology type, high-risk adenomas, and heavy diet were all influential factors in the development of *H. pylori* infection after intestinal polypectomy. Binary multifactorial logistic regression analysis showed that age, BMI, and type of polyp pathology were independent predictors of the occurrence of *H. pylori* infection after intestinal polypectomy. The area under the receiver operating characteristic curve was 0.969 [95% confidence interval (95% CI): 0.928–1.000] and 0.898 (95% CI: 0.773–1.000) in the modeling and validation sets,

respectively. The slope of the calibration curve of the graph was close to 1, and the goodness-of-fit test was $P > 0.05$ in the two sets. The decision analysis curve showed a high rate of return in both sets. The results of the correlation analysis between different pathological types and the occurrence of *H. pylori* infection after colon polyp surgery showed that hyperplastic polyps, inflammatory polyps, and the occurrence of *H. pylori* infection were not significantly correlated. In contrast, adenomatous polyps showed a significant positive correlation with the occurrence of *H. pylori* infection.

CONCLUSION

Age, BMI, and polyps of the adenomatous type were independent predictors of *H. pylori* infection after intestinal polypectomy. Moreover, the further constructed column-line graph prediction model of *H. pylori* infection after intestinal polypectomy showed good predictive ability.

Key Words: Colon polyps; *Helicobacter pylori*; Risk factors; Pathologic type; Columnar graphic modeling

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: *Helicobacter pylori* (*H. pylori*) infection is reportedly a risk factor for the development of colonic adenomas, especially progressive or multiple adenomas. However, few studies have examined the risk factors for *H. pylori* infection after therapeutic colon polypectomy and the type of polyp pathology associated with its occurrence. This randomized study evaluated the risk factors for the development of *H. pylori* infections in patients with colon polyps and the relationship between their pathology and the development of *H. pylori* infections.

Citation: Zhang ZS. Predictive factors and model validation of post-colon polyp surgery *Helicobacter pylori* infection. *World J Gastrointest Surg* 2024; 16(1): 173-185

URL: <https://www.wjgnet.com/1948-9366/full/v16/i1/173.htm>

DOI: <https://dx.doi.org/10.4240/wjgs.v16.i1.173>

INTRODUCTION

Colonic polyps are tumor-like lesions that grow on the mucosal surface of the colon, usually in the form of protruding or elevated masses or fleshy lesions[1]. They do not invade surrounding tissues and are clinically classified based on their histologic features and susceptibility to malignant transformation. Mainly, there are colorectal adenomatous polyps, inflammatory polyps, and hyperplastic polyps[2]. Colonic adenomatous polyps are abnormal tissues that may develop into colorectal cancer (CC); colonic adenomatous polyps are the most common in clinical practice and carry a higher risk of developing cancer[3]. According to the 2016 United States Guidelines for Follow-up after Colorectal Polypectomy[4], the presence of adenomas that are more than 1 cm in diameter, three or more in number, or that exhibit villous or high-grade intraepithelial neoplasia, along with the occurrence of any of the aforementioned criteria, suggests a high risk of cancer development in colorectal polyp case. To reduce the risk of CC, further development of colonic adenomatous polyps must be inhibited to the greatest extent possible, through prevention and early treatment[5]. In the last decade, *Helicobacter pylori* (*H. pylori*) infection of the stomach has been demonstrated to induce an inflammatory response in the colon through the production of toxins, thereby promoting the development of CC, to some extent[6]. Considering the increasing number of patients with colonic polyps in our country and the large number of *H. pylori* infections, an in-depth understanding of the current status and risk factors for *H. pylori* infections in these patients is essential[7]. A previous study[8] revealed that the development of colon tumors is significantly associated with *H. pylori* infection. Simultaneously, *H. pylori* infection is also identified as a risk factor for the development of colon adenomas, especially progressive or multiple adenomas. Therefore, this study aimed to analyze the risk factors for the development of *H. pylori* infection in patients with colonic polyps, and the relationship between their pathological type and the development of *H. pylori* infection.

MATERIALS AND METHODS

Objects of study

Eighty patients who underwent colon polypectomy at our hospital between January 2019 and January 2023 were retrospectively selected as participants. They were randomly divided into a modeling cohort ($n = 56$) and a model validation cohort ($n = 24$) at a ratio of 7:3 using the R language.

Inclusion and exclusion criteria

Inclusion criteria: (1) Participants who met the indications for colonoscopic polypectomy; (2) those who underwent the 14C-urea breath test; (3) had no immune system disease or immune dysfunction; (4) no psychiatric disorders and were able to communicate and interact normally; and (5) had complete clinical data. Participants that met all the above criteria were included in this study.

Exclusion criteria: Participants who met any one of the following criteria were excluded from the study: (1) Participants with a previous history of gastrointestinal disease or colon tumor; (2) those who presented with coagulation disorders after discontinuing oral anticoagulant medication for < 1 wk; and (3) those who were on medication prior to *H. pylori* screening.

Observation of grouping and *H. pylori*-infection

The modeling cohort was divided into an *H. pylori*-infected group ($n = 37$) and an *H. pylori*-uninfected group ($n = 19$) according to whether the patients developed an *H. pylori* infection. Patients were monitored for the occurrence of *H. pylori* infection, which served as the endpoint.

Observation indicators

The general information of the patients was collected through electronic medical records. This included general information [sex, age, body mass index (BMI), exercise, education, smoking and drinking habits, history of hypertension and diabetes mellitus, and heavy diet consumptions] and speciality information (number, size, location, and pathological type of the polyps, and whether they were high-risk adenomas).

We analyzed the risk factors for developing *H. pylori* infection after colon polyp surgery by observing the age, sex, BMI, and exercise of patients in the modeling cohort (*H. pylori*-infected and *H. pylori*-uninfected groups). In addition, we assessed whether or not they smoked, consumed alcohol, suffered from high blood pressure, consumed a heavy diet, and had diabetes mellitus. The number, size, location, and the pathological type of polyps, and the presence of high-risk adenomas, were also assessed. All of the information was used to develop and validate a roadmap prediction model. Finally, the correlation between the different pathological types and the occurrence of *H. pylori* after colon polyp surgery was analyzed.

Statistical analysis

SPSS 26.0 software and R software were used to analyze the data. The collected count data were expressed as cases (%); χ^2 or Fisher exact test was used for unordered data, and the Mann-Whitney *U* test was used for ordered data. Univariate and multivariate binary logistic regression analyses were used to analyze the factors influencing the development of *H. pylori* infection after colon polyp surgery and to develop a column-line graph prediction model. The discriminative power of the validation set and calibration graphs were used to assess the accuracy of the column-line graphs. The area under the receiver operating characteristic (ROC) curve (AUC) was used to evaluate the discriminative ability of the column diagram. Calibration curves for the model were calculated, and the consistency of the model was verified using the Hosmer-Lemeshow test. Decision curve analysis (DCA) was performed to evaluate the discriminative ability of the model. $P < 0.05$ was considered statistically significant, and correlations were tested using Spearman's test.

RESULTS

Baseline clinical characteristics

A total of 93 patients with severe traumatic brain injury were included in the study: 56 in the modeling cohort and 24 in the validation cohort. All patients were aged 30–72 years at the time of diagnosis; 42 (52.50%) were male and 38 (47.50%) were female. Other baseline information regarding the modeling and validation cohorts is shown in [Table 1](#).

Comparison of clinical data between *H. pylori*-infected and *H. pylori*-uninfected groups of patients in the model cohort

There were no statistically significant differences in sex composition, exercise and smoking status, history of hypertension and diabetes mellitus, number of polyps, polyp size, or polyp site in the model cohort ($P > 0.05$). The differences in age, BMI, literacy level, alcohol consumption, polyp pathological type, high-risk adenomas, and heavy diet consumption in the *H. pylori*-infected group were statistically significant when compared with the *H. pylori*-uninfected group ($P < 0.05$; [Table 2](#)).

Univariate analysis of the occurrence of *H. pylori* infection in the model cohort of patients

In the model cohort, *H. pylori* infection was the dependent variable and assigned 1, and its absence was assigned 0. Variables with $P < 0.05$ in the clinical data were included in the univariate analysis. The univariate results showed that age, BMI, literacy level, alcohol consumption, type of polyp pathology, high-risk adenomas, and a heavy diet consumption were all influential factors in the occurrence of *H. pylori* infections after intestinal polypectomy ([Table 3](#)).

Multifactorial analysis of the occurrence of *H. pylori* infection

Variables with $P < 0.05$ in the univariate analysis were included in the binary multivariate logistic regression analysis, which showed that age, BMI, and pathologic type of polyp were independent predictors of the development of *H. pylori*

Table 1 Baseline information for modeling and validation cohorts

Sports event		Total population (n = 80)	Modeling queues (n = 56)	Validation queue (n = 24)
Gender [n (%)]	Male	42 (52.50)	30 (53.57)	12 (50.00)
	Female	38 (47.50)	26 (46.43)	12 (50.00)
Age (yr, mean ± SD)		36.15 ± 11.79	36.67 ± 10.00	35.92 ± 13.58
BMI [n (%)]		22.42 ± 3.44	22.20 ± 2.32	22.64 ± 4.55
Movement [n (%)]	< 1 h/wk	38 (47.50)	27 (48.21)	11 (45.83)
	≥ 1 h/wk	42 (52.50)	29 (51.79)	13 (54.17)
Literacy [n (%)]	Primary and below	30 (37.50)	20 (35.71)	10 (41.67)
	Junior high school and secondary school	29 (36.25)	21 (37.50)	8 (33.33)
	Junior college or above	21 (26.25)	15 (26.79)	6 (25.00)
Smoking [n (%)]	Be	33 (41.25)	24 (42.86)	9 (37.50)
	Clogged	45 (56.25)	32 (57.14)	13 (54.17)
Alcohol consumption [n (%)]	Be	44 (55.00)	31 (55.36)	13 (54.17)
	Clogged	34 (42.50)	25 (44.64)	9 (37.50)
History of hypertension [n (%)]	Be	32 (40.00)	22 (39.29)	10 (41.67)
	Clogged	48 (60.00)	34 (60.71)	14 (58.33)
History of diabetes [n (%)]	Be	14 (17.50)	10 (17.86)	4 (16.67)
	Clogged	66 (82.50)	46 (82.14)	20 (83.33)
Number of polyps [n (%)]	An odd one	38 (47.50)	27 (48.21)	11 (45.83)
	Multi- (faceted, ethnic etc.)	42 (52.50)	29 (51.79)	13 (54.17)
Polyp size [n (%)]	< 1 cm	33 (41.25)	23 (41.07)	10 (41.67)
	≥ 1 cm	47 (58.75)	33 (58.93)	14 (58.33)
Polyp site [n (%)]	Proximal	27 (33.75)	19 (33.93)	8 (33.33)
	Far end	25 (31.25)	18 (32.14)	7 (29.17)
	Whole colon	28 (35.00)	19 (33.93)	9 (37.50)
Type of polyp pathology [n (%)]	Adenomatous polyp	35 (43.75)	24 (42.86)	11 (45.83)
	Non-adenomatous polyp	45 (56.25)	32 (57.14)	13 (54.17)
High-risk adenomas [n (%)]	Be	23 (28.75)	16 (28.57)	7 (29.17)
	Clogged	57 (71.25)	40 (71.43)	17 (70.83)
Heavy diet [n (%)]	Be	46 (57.50)	32 (57.14)	14 (58.33)
	Clogged	34 (42.50)	24 (42.86)	10 (41.67)

BMI: Body mass index.

infection after intestinal polypectomy, with the model equation: Logistic = $-3.798 - 0.342 \times \text{age} + 1.222 \times \text{BMI} - 3.760 \times \text{type of polyp pathology}$ (Table 4).

Modeling of a column-line diagram to predict the occurrence of *H. pylori* infection in patients after colon polyp surgery

The resulting three independent risk factors (age, BMI, and polyp pathology type) were used to construct a prediction model using R software, and subsequent column-line graph model, as shown in Figure 1. The C-statistic of the model was calculated using the R language software as 0.809, with a 95% confidence interval (95%CI) of 0.761–0.890 and a standard error of 0.030 ($P < 0.001$). The C-statistic was calculated using the R language software as 0.818, with a standard error of 0.030 ($P < 0.001$), and the 10000 Bootstrap calculated a C statistic of 0.818. The slope of the generated column-line graph calibration curve was close to 1 (Figure 2), with a goodness-of-fit test of $P > 0.05$ and a high degree of consistency between the predicted and actual events. The area under the ROC curve of the column-line diagram prediction model was 0.969 (95%CI: 0.928–1.000) (Figure 3). The decision analysis curve is shown in Figure 4, where the X-axis indicates the threshold

Table 2 Comparison of clinical data between *Helicobacter pylori*-infected and *Helicobacter pylori*-uninfected groups of patients in the model cohort

Sports event		Hp infection group (n = 37)	Hp uninfected group (n = 19)	χ^2 value	P value
Gender [n (%)]	Male	20 (54.05)	10 (52.63)	0.010	0.920
	Female	17 (45.95)	9 (47.37)		
Age (yr, mean \pm SD)			46.58 \pm 3.50	4.788	0.000
BMI [n (%)]			20.37 \pm 1.65	5.114	0.000
Movement [n (%)]	< 1 h/wk	18 (48.65)	9 (47.37)	0.008	0.928
	\geq 1 h/wk	19 (51.35)	10 (52.63)		
Literacy [n (%)]	Primary and below	16 (43.24)	4 (21.05)	2.348	0.019
	Junior high school and secondary school	15 (40.54)	6 (31.58)		
	Junior college or above	6 (16.22)	9 (47.37)		
Smoking [n (%)]	Be	16 (43.24)	8 (42.11)	0.007	0.935
	Clogged	21 (56.76)	11 (57.89)		
Alcohol consumption [n (%)]	Be	24 (64.86)	7 (36.84)	3.989	0.046
	Clogged	13 (35.14)	12 (63.16)		
History of hypertension [n (%)]	Be	15 (40.54)	7 (36.84)	0.072	0.788
	Clogged	22 (59.46)	12 (63.16)		
History of diabetes [n (%)]	Be	9 (24.32)	1 (5.26)	1.946	0.163
	Clogged	28 (75.68)	18 (94.74)		
Number of polyps [n (%)]	An odd one	17 (45.95)	10 (52.63)	0.225	0.636
	Multi- (faceted, ethnic etc.)	20 (54.05)	9 (47.37)		
Polyp size [n (%)]	< 1 cm	15 (40.54)	8 (42.11)	0.013	0.910
	\geq 1 cm	22 (59.46)	11 (57.89)		
Polyp site [n (%)]	Proximal	10 (27.03)	9 (47.37)	2.326	0.313
	Far end	13 (35.14)	5 (26.32)		
	Whole colon	14 (37.84)	5 (26.32)		
Type of polyp pathology [n (%)]	Adenomatous polyp	20 (56.76)	4 (21.05)	5.583	0.018
	Non-adenomatous polyp	17 (27.03)	15 (52.63)		
High-risk adenomas [n (%)]	Be	14 (37.84)	2 (10.53)	4.588	0.032
	Clogged	23 (62.16)	17 (89.47)		
Heavy diet [n (%)]	Be	25 (67.57)	7 (36.84)	4.839	0.028
	Clogged	12 (32.43)	12 (63.16)		

BMI: Body mass index.

probability, the Y-axis indicates the net return, and the black solid line indicates the net return using the column-line diagram prediction model, which shows a higher return and further confirms the effectiveness of the column-line diagram prediction model.

Validation of the column-line diagram model

Based on the validation cohort ($n = 24$), which was divided into *H. pylori*-infected ($n = 16$) and *H. pylori*-uninfected ($n = 8$) groups, the column-line diagram of the risk of *H. pylori* infection was externally validated using an ROC curve, and the lower product of the ROC curve was 0.898 (95%CI: 0.773–1.000) (Figure 5). The slope of the generated calibration curve for the column-line diagram was close to 1 (Figure 6), and the result of the Hosmer-Lemeshow test was, $\chi^2 = 10.609$, $P = 0.157 > 0.05$. The decision curve showed a higher net benefit of the model (Figure 7), suggesting that the calibration of the

Table 3 Univariate analysis of *Helicobacter pylori*-infected and *Helicobacter pylori*-uninfected groups

Considerations	B	SE	Wals	P value	OR	95%CI
Age	-0.169	0.049	12.137	0.000	0.844	0.768-0.929
BMI	0.738	0.216	11.708	0.001	2.093	1.371-3.194
Educational attainment	-0.912	0.393	5.383	0.020	0.402	0.186~0.868
Drinking wine	1.152	0.587	3.850	0.050	3.165	1.001-10.004
Types of polyp pathology	1.484	0.652	5.178	0.023	0.227	0.063-0.814
High-risk adenoma	1.644	0.821	4.010	0.045	5.174	1.035-25.852
Heavy diet	1.399	0.596	5.506	0.019	4.052	1.259-13.038

OR: Odds ratio; 95%CI: 95% confidence interval; BMI: Body mass index.

Table 4 Multifactorial analysis of *Helicobacter pylori*-infected and *Helicobacter pylori*-uninfected groups

considerations	B	SE	Wals	P value	OR	95%CI
Age	-0.342	0.145	5.574	0.018	0.710	0.535-0.944
BMI	1.222	0.446	7.524	0.006	3.395	1.418-8.130
Types of polyp pathology	-3.760	1.772	4.505	0.034	0.023	0.001-0.750
Constant	-3.798	7.216	0.277	0.599	0.022	-




Figure 1 Column-line diagram.


column-line diagram model in the validation group was better.

Analysis of the correlation between different pathology types and the occurrence of *H. pylori* infection after colon polyps

Three pathologic types were identified in the model cohort of patients with colon polyps. In the *H. pylori*-infected group, these included seven cases of inflammatory polyps, 10 cases of hyperplastic polyps, and 20 cases of adenomatous polyps. In the *H. pylori*-uninfected group, these included five cases of inflammatory polyps, 10 cases of hyperplastic polyps, and four cases of adenomatous polyps. Correlation analysis of the occurrence of different pathological types and *H. pylori* infection after colon polyp surgery was performed. The results of the correlation analysis showed no significant correlation between hyperplastic polyps, inflammatory polyps, and the occurrence of *H. pylori* infection. In contrast, adenomatous polyps showed a significant positive correlation with the occurrence of *H. pylori* infection (Table 5).

Table 5 Correlation between different types of pathology and occurrence of *Helicobacter pylori* infection after colon polyp surgery

		Inflammatory polyp	Hyperplastic polyp	Adenomatous polyp
<i>Helicobacter pylori</i> infection	Correlation coefficient	-0.085	-0.253	0.316
	Sig. (bilateral)	0.532	0.060	0.018

**Figure 2** Calibration curves of the column graph prediction model.

DISCUSSION

H. pylori is a bacterium that parasitizes areas such as the stomach or duodenum, and can survive for long periods of time under conditions of little oxygen. It not only has the ability to secrete toxic substances, which contribute to gastrointestinal disorders such as gastric ulcers and gastritis, but also has an impact on the rate of infection[9,10]. *H. pylori* can be detected not only by non-invasive methods such as the urease breath test and serum *H. pylori* antibodies, but also by invasive methods such as biopsy of the gastric mucosa with special staining of the tissue biopsies. Moreover, its important role in the digestive system has been widely recognized[11]. However, few studies have addressed the exact relationship between the three pathological types of colon polyps, namely, adenomatous, inflammatory, and hyperplastic colon polyps, and *H. pylori* infection. Further research is needed to clarify this relationship. Therefore, the present study developed a column-line diagram model focusing on analyzing the factors influencing the occurrence of *H. pylori* infection in patients with gastrointestinal polyps and exploring the correlation between the pathological types of polyps and the occurrence of *H. pylori* infection.

By comparing the clinical data of patients with gastrointestinal polyps, we found that there were statistically significant differences between the *H. pylori*-infected and *H. pylori*-uninfected groups in terms of age, BMI, literacy, alcohol consumption, polyp pathology type, presence of high-risk adenomas, and heavy diets consumption. There were no significant differences in the other indicators. The results of the binary logistic one-way regression analysis assigned a value of 1 to the occurrence of *H. pylori* infection and a value of 2 to the non-occurrence of *H. pylori* infection as the dependent variables. Moreover, the factors with significant differences in the aforementioned clinical data as the covariates, showed that age, BMI, literacy, alcohol consumption, polyp pathology type, high-risk adenomas, and heavy diets consumption were the factors influencing the occurrence of *H. pylori* infection after intestinal polypectomy. Subsequently, we performed a binary logistic regression analysis, of the factors with significant differences in the univariate analysis as covariates and found that age, BMI, and polyp pathology type were independent predictors of the occurrence of *H. pylori* infection after intestinal polypectomy. Among them, younger age is associated with a greater likelihood of developing *H. pylori* infection after intestinal polypectomy. This may be because younger patients, with continuous changes in their social environment, are presented with increasing work and life pressures, which tend to result in lower resistance of their bodies and thus are more susceptible to *H. pylori* infection[12]. Additionally, adolescents tend to favor convenient diets, such as high-fat, high-sugar, and high-salt foods, which subsequently increase the risk of *H. pylori* infection[13]. In contrast, older people have a more regular lifestyle, pay more attention to healthy eating and




Figure 3 Receiver operating characteristic curve of the column chart prediction model.




Figure 4 Decision analysis curve.

living habits, have frequent medical checkups, and follow their doctors' advice. This reduces their likelihood of becoming infected with *H. pylori*[14,15].

In contrast to the age trend, regarding BMI and polyp pathology type, we found that higher BMI is associated with a greater likelihood of *H. pylori* infection after intestinal polypectomy. Thus patients with adenomatous polyps on polyp pathology had a greater likelihood of *H. pylori* infection. It has been reported in the literature[16,17] that this can be because there is an association between BMI and *H. pylori* infection, and that the two factors can interact with each other. Due to the long-term intake of excessively high calories, the immune environment of their organs is changed, which leads




Figure 5 Verification queue receiver operating characteristic curve.




Figure 6 Verification queue calibration curve.

to the expansion of adipose tissues and the activation of macrophages through the secretion of chemokines, subsequently causing a localized inflammatory response. Consequently, the immune microenvironment of obese patients creates favorable conditions for the survival of *H. pylori*; thus, obese people are more likely to be infected with *H. pylori*. This is similar to the findings of Xie *et al*[18] Additionally, changes in the intragastric microenvironment due to *H. Pylori* may lead to intestinal microecological disorders, further affecting the intestinal microecology of the patients. This may lead to intestinal tumor-like lesions and adenomatous polyps[19]. Thus, adenomatous polyps in patients are often accompanied by *H. pylori* infection. The results of a study by Zhang *et al*[20] showed that the proportion of adenomatous polyps occurring in *H. pylori*-infected populations was significantly higher than that in *H. pylori*-uninfected populations. This is




Figure 7 Verification queue decision analysis curve.

similar to the results of the present study and further supports the findings of the present study.

Additionally, to further clarify the predictive value of age, BMI, and polyp pathology type in the occurrence of *H. pylori* infection after intestinal polypectomy, we utilized the R software to establish a column-line graph model. The C statistic of this model was calculated using the R language software as 0.809, which indicated that the model had a stronger discriminatory ability and was able to distinguish patients with high likelihood to develop *H. pylori* infection. The slope of the calibration curve of the column-line graph it generated was close to 1, and the test of goodness of fit was $P > 0.05$, which showed that the model had a strong calibration ability. The consistency between the predicted events and the actual events was high, and the area under the ROC curve was 0.969 (95%CI: 0.928-1.000). This indicated that the model was more efficacious in predicting the risk of *H. pylori* infection. Furthermore, the AUC value was closer to 1, indicating that the model is more capable of discriminating risk. The decision analysis curve showed a higher yield, further confirming the validity of the column-line graph prediction model. Further external validation ROC curve product under the curve was 0.898 (95%CI: 0.773-1.000), which indicated that the model also performed well in the external validation cohort and had good generalization ability. The slope of the generated column-line graph calibration curve was close to 1, with a Hosmer-Lemeshow test result of $P > 0.05$. Moreover, the decision curve showed a higher net gain of the model, suggesting that the column-line graph model had a better calibration ability in the validation cohort. The column-line diagram model of *H. pylori* infection risk obtained in this study showed good predictive and calibration abilities for both in-sample and out-of-sample validations. According to the visualized form of the column-line diagram, age ≤ 50 years, lower education level, and higher BMI are associated with higher risk of *H. pylori* infection after intestinal polypectomy. Moreover, patients with adenomatous polyps often have *H. pylori* infection. This showed effective clinical discrimination of the high-risk group of *H. pylori* infection after intestinal polypectomy, based on the information of patients in the aforementioned key factors. Therefore, the present study illustrated simple predictors that are favorable for the early prevention of *H. pylori* infection.

CONCLUSION

In conclusion, age, BMI, and polyp pathology of the adenomatous type were independent predictors of *H. pylori* infection after intestinal polypectomy. In addition, the columnar graph prediction model of *H. pylori* infection after intestinal polypectomy showed good predictive ability, which provided assistance in the clinical identification of high-risk groups of *H. pylori* infection after intestinal polypectomy. This is beneficial for the timely prevention of *H. pylori* infection. However, because this study was a retrospective analysis, the sample size was limited, and more clinical indicators should be added for further comprehensive assessment and establishment of a more comprehensive prediction model.

ARTICLE HIGHLIGHTS

Research background

Colon polyps are tumor-like lesions that grow on the surface of the colonic mucosa, usually in the form of a protruding or bulging mass, or meaty lesion. They are abnormal tissue that can develop into colorectal cancer. Considering that the number of patients with colon polyps in our country has been rising and that a large number of *Helicobacter pylori* (*H. pylori*) infections also exist, an in-depth understanding of the current status of *H. pylori* infections in patients with colonic polyps in our country and the risk factors for these infections is necessary.

Research motivation

The development of colon tumors is significantly associated with *H. pylori* infection, of which colonic adenomatous polyps may develop into colon cancer. It is also a risk factor for the development of colonic adenomas, especially progressive or multiple adenomas. However, few clinical studies have investigated the correlation between the pathological types of colonic polyps and *H. pylori* infection.

Research objectives

To investigate the risk factors for the development of *H. pylori* infection after colon polyp surgery, and to establish the relationship between the type of pathology and its occurrence.

Research methods

Eighty patients who underwent colon polypectomy in our hospital from January 2019 to January 2023 were retrospectively selected as participants, and randomly divided into a modeling cohort ($n = 56$) and a model validation cohort ($n = 24$) at a ratio of 7:3 using R. Simultaneously, based on whether the patients were infected with *H. pylori*, the modeling cohort was divided into an *H. pylori*-infected group ($n = 37$) and an *H. pylori*-uninfected group ($n = 19$). The risk factors for *H. pylori* after colon polyp surgery were analyzed by comparing the age, sex, body mass index (BMI), and exercise status of patients in the modeling cohort (*H. pylori*-infected and *H. pylori*-uninfected groups). In addition, whether or not they smoked, consumed alcohol, suffered from hypertension and diabetes mellitus, and had heavy diets, and the number, size, location, and the pathological type of the polyps, and whether or not they were high-risk adenomas, were also analyzed. A binary logistic regression analysis was used to analyze the factors influencing the occurrence of *H. pylori* infection after colon polyp surgery. A roadmap prediction model was therefore established and validated; receiver operating characteristic was used to evaluate the predictive efficacy of the model; calibration curves were used to assess the consistency between predicted and actual events. DCA curves were also used to evaluate the validity of the model; and finally, the correlation between the different pathological types of colon polyps and the occurrence of *H. pylori* infection was analyzed after colon polyp surgery.

Research results

Age, BMI, and polyp pathology type were independent predictors of *H. pylori* infection after intestinal polypectomy. Additionally, the *H. pylori* infection risk column-line diagram model obtained in this study demonstrated good predictive and calibration abilities for both in-sample and out-of-sample validations. The visualized form of the column-line diagram showed that for age ≤ 50 years, the lower the education level, the higher the risk of *H. pylori* infection after intestinal polypectomy, the higher the BMI, the higher the risk of *H. pylori* infection, and that patients with adenomatous polyps often have *H. pylori* infection. This is conducive to the effective clinical discrimination of patients at high risk of *H. pylori* infection, after intestinal polypectomy, based on the information of the above mentioned key factors. Moreover, the predictors obtained in this study are favorable for the early prevention of *H. pylori* infection.

Research conclusions

Age, BMI, and polyp pathology of the adenomatous type were all independent predictors of *H. pylori* infection after intestinal polypectomy, and the column-line graph prediction model of *H. pylori* infection after intestinal polypectomy showed good predictive ability. This provides assistance in the clinical identification of high-risk groups for *H. pylori* infection after intestinal polypectomy and is conducive to timely prevention.

Research perspectives

This study was a retrospective analysis with a limited sample size, and additional clinical indicators need to be added for further comprehensive assessment and predictive modeling.

FOOTNOTES

Author contributions: Zhang ZS was responsible for the methodology, investigation, software, data curation, formal analysis, writing the original draft and editing, conceptualization, validation.

Institutional review board statement: The study was reviewed and approved by the Ethics Committee of Affiliated Hospital of Shaoxing University (Shaoxing Municipal Hospital (NO.2023-072-01)).

Informed consent statement: All study participants, or their legal guardian, provided informed written consent prior to study enrollment.

Conflict-of-interest statement: The authors have no conflicts of interest to declare.

Data sharing statement: No additional data are available.

STROBE statement: The authors have read the STROBE Statement—checklist of items, and the manuscript was prepared and revised according to the STROBE Statement—checklist of items.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Zheng-Sen Zhang [0009-0003-6473-1451](https://orcid.org/0009-0003-6473-1451).

S-Editor: Lin C

L-Editor: A

P-Editor: Yu HG

REFERENCES

- 1 Yan Z, Sun YM, Gao F, Lang HB, Zhang J. Analysis of risk factors for delayed bleeding after endoscopic mucosal resection of colon polyps. *Zhongguo Yiyao Daobao* 2023; **20**: 132-135 [DOI: [10.20047/j.issn1673-7210.2023.17.30](https://doi.org/10.20047/j.issn1673-7210.2023.17.30)]
- 2 Han L, Jiang SL. Observation on the efficacy of endoscopic mucosal resection in patients with colonic polyps. *Xiandai Yixueyujiankang Yanjiu (Dianziban)* 2023; **7**: 56-59 [DOI: [10.3969/j.issn.2096-3718.2023.03.017](https://doi.org/10.3969/j.issn.2096-3718.2023.03.017)]
- 3 Sninsky JA, Shore BM, Lupu GV, Crockett SD. Risk Factors for Colorectal Polyps and Cancer. *Gastrointest Endosc Clin N Am* 2022; **32**: 195-213 [PMID: [35361331](https://pubmed.ncbi.nlm.nih.gov/35361331/) DOI: [10.1016/j.giec.2021.12.008](https://doi.org/10.1016/j.giec.2021.12.008)]
- 4 Shuai XW, Xie PY. Guidelines for colonoscopic follow-up protocols after resection of colon cancer--Updated consensus of the U.S. Multicenter Task Force on Colorectal Cancer and the American Cancer Society (2006). *Zhongguo Neijing Zazhi* 2007; **889-891**
- 5 Chao G, Zhu Y, Fang L. Retrospective study of risk factors for colorectal adenomas and non-adenomatous polyps. *Transl Cancer Res* 2020; **9**: 1670-1677 [PMID: [35117515](https://pubmed.ncbi.nlm.nih.gov/35117515/) DOI: [10.21037/tcr.2020.01.69](https://doi.org/10.21037/tcr.2020.01.69)]
- 6 Chen CC, Liou JM, Lee YC, Hong TC, El-Omar EM, Wu MS. The interplay between *Helicobacter pylori* and gastrointestinal microbiota. *Gut Microbes* 2021; **13**: 1-22 [PMID: [33938378](https://pubmed.ncbi.nlm.nih.gov/33938378/) DOI: [10.1080/19490976.2021.1909459](https://doi.org/10.1080/19490976.2021.1909459)]
- 7 Hou Y M, Li H, Wang L, Xiang X H, Yang S G. Association of colon polyps with *Helicobacter pylori* infection and gastric polyps. *Zhongguo Neijing Zazhi* 2023; **29**: 73-80 [DOI: [10.12235/E20220103](https://doi.org/10.12235/E20220103)]
- 8 Yang QJ, Zheng J, Yang J, Luo R, Leng J, Jin Q, Ma HL. Study on the effect of *Helicobacter pylori* infection on nonalcoholic fatty liver disease and its associated colorectal polyps. *Jiating Yixue* 2021; **4**: 3855-3862 [DOI: [10.12114/j.issn.1007-9572.2021.00.549](https://doi.org/10.12114/j.issn.1007-9572.2021.00.549)]
- 9 Yang Z P, Tian Y J, Wang Y Z, Zhao Y D. Relationship between oral *Helicobacter pylori* infection and *Helicobacter pylori* gastritis. *Beihua Daxue Xuebao(Ziranxueban)* 2022; **23**: 352-356 [DOI: [10.11713/j.issn.1009-4822.2022.03.013](https://doi.org/10.11713/j.issn.1009-4822.2022.03.013)]
- 10 de Brito BB, da Silva FAF, Soares AS, Pereira VA, Santos MLC, Sampaio MM, Neves PHM, de Melo FF. Pathogenesis and clinical management of *Helicobacter pylori* gastric infection. *World J Gastroenterol* 2019; **25**: 5578-5589 [PMID: [31602159](https://pubmed.ncbi.nlm.nih.gov/31602159/) DOI: [10.3748/wjg.v25.i37.5578](https://doi.org/10.3748/wjg.v25.i37.5578)]
- 11 Qiu E, Li Z, Han S. Methods for detection of *Helicobacter pylori* from stool sample: current options and developments. *Braz J Microbiol* 2021; **52**: 2057-2062 [PMID: [34392499](https://pubmed.ncbi.nlm.nih.gov/34392499/) DOI: [10.1007/s42770-021-00589-x](https://doi.org/10.1007/s42770-021-00589-x)]
- 12 Liu L, Zhu ZH. Influencing factors of *Helicobacter pylori* infection in gastroenterology patients and its relationship with gastrointestinal diseases. *Xiandai Yangsheng (Xiabanyueban)* 2020; **20**: 52-55 [DOI: [10.3969/j.issn.1671-0223\(x\).2020.11.016](https://doi.org/10.3969/j.issn.1671-0223(x).2020.11.016)]
- 13 Cao YF, Li XL, Wang YF, Wu WZ, Xu C, Wang SS, Huang HX. Study on the influencing factors of *Helicobacter pylori* infection based on age stratification. *Zhejiang Yixue* 2023; **45**: 140-144 [DOI: [10.12056/j.issn.1006-2785.2023.45.2.2022-2609](https://doi.org/10.12056/j.issn.1006-2785.2023.45.2.2022-2609)]
- 14 LI XY, GAO CG. Analysis of the results of 102 cases of 13C-urea breath test for detecting Hp infection in patients and related influencing factors in Xuchang Central Hospital. *Yixue Yanjiu* 2021; **30**: 1010-1012 [DOI: [10.3969/j.issn.1004-437X.2021.06.014](https://doi.org/10.3969/j.issn.1004-437X.2021.06.014)]
- 15 Li X, Wang J. Survey on factors affecting compliance with standardized treatment of *Helicobacter pylori* infection patients in digestive system and intervention countermeasures. *Guizhou Yixue* 2023; **47**: 549-550 [DOI: [10.3969/j.issn.1000-744X.2023.04.024](https://doi.org/10.3969/j.issn.1000-744X.2023.04.024)]
- 16 Yusuf Tohti, Li K. Research progress on the correlation between *Helicobacter pylori* infection and obesity. *Zhonghua Feipangyudaixiebing Zazhi* 2020; **6**: 196-199 [DOI: [10.3877/cma.j.issn.2095-9605.2020.03.011](https://doi.org/10.3877/cma.j.issn.2095-9605.2020.03.011)]
- 17 AlAli MN, Bamehriz F, Arishi H, Aldeghaither MK, Alabdullatif F, Alnaeem KA, Alzamil AF, AlHashim IR, Alhaizan S, Aljuhani T, Aldohayan A. Trends in bariatric surgery and incidentalomas at a single institution in Saudi Arabia: a retrospective study and literature review. *Ann Saudi Med* 2020; **40**: 389-395 [PMID: [33007169](https://pubmed.ncbi.nlm.nih.gov/33007169/) DOI: [10.5144/0256-4947.2020.389](https://doi.org/10.5144/0256-4947.2020.389)]
- 18 Xie Q, He Y, Zhou D, Jiang Y, Deng Y, Li R. Recent research progress on the correlation between metabolic syndrome and *Helicobacter pylori* infection. *Peer J* 2023; **11**: e15755 [PMID: [37483988](https://pubmed.ncbi.nlm.nih.gov/37483988/) DOI: [10.7717/peerj.15755](https://doi.org/10.7717/peerj.15755)]
- 19 Ouyang Y, Zhang W, Huang Y, Wang Y, Shao Q, Wu X, Lu N, Xie C. Effect of *Helicobacter pylori* eradication on hyperplastic gastric polyps: A systematic review and meta-analysis. *Helicobacter* 2021; **26**: e12838 [PMID: [34333811](https://pubmed.ncbi.nlm.nih.gov/34333811/) DOI: [10.1111/hel.12838](https://doi.org/10.1111/hel.12838)]

- 20 **Zhang P**, Wang P, Hong R, Huang ML Q. Clinical characteristics of Hp infection in patients with colorectal polyps and its relationship with G-17, sIL-2R and COX-2. *Zhonghua Yiyuan Ganranxue Zazhi* 2023; **33**: 81-85 [DOI: [10.11816/cn.ni.2023-212986](https://doi.org/10.11816/cn.ni.2023-212986)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: office@baishideng.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

