


World Journal of *Gastrointestinal Surgery*

World J Gastrointest Surg 2024 January 27; 16(1): 1-259



EDITORIAL

- 1 Novel prognostic factors after radical resection of hepatocellular carcinoma: Updating an old issue
Bencini L
- 6 Prospects in the application of ultrasensitive chromosomal aneuploidy detection in precancerous lesions of gastric cancer
Qian ST, Xie FF, Zhao HY, Liu QS, Cai DL

MINIREVIEWS

- 13 Prognostic value of ultrasound in early arterial complications post liver transplant
Zhao NB, Chen Y, Xia R, Tang JB, Zhao D

ORIGINAL ARTICLE**Case Control Study**

- 21 Added value of ratio of cross diameters of the appendix in ultrasound diagnosis of acute appendicitis
Gu FW, Wu SZ

Retrospective Cohort Study

- 29 Oncological features and prognosis of colorectal cancer in human immunodeficiency virus-positive patients: A retrospective study
Yang FY, He F, Chen DF, Tang CL, Woraiikat S, Li Y, Qian K

Retrospective Study

- 40 Laparoscopic *vs* open surgery for gastric cancer: Assessing time, recovery, complications, and markers
Lu YY, Li YX, He M, Wang YL
- 49 Single-incision laparoscopic transabdominal preperitoneal repair in the treatment of adult female patients with inguinal hernia
Zhu XJ, Jiao JY, Xue HM, Chen P, Qin CF, Wang P
- 59 Computerized tomography-guided therapeutic percutaneous puncture catheter drainage-combined with somatostatin for severe acute pancreatitis: An analysis of efficacy and safety
Zheng XL, Li WL, Lin YP, Huang TL
- 67 Impact of open hepatectomy on postoperative bile leakage in patients with biliary tract cancer
Wu G, Li WY, Gong YX, Lin F, Sun C
- 76 Clinical observation of gastrointestinal function recovery in patients after hepatobiliary surgery
Zeng HJ, Liu JJ, Yang YC

- 85 Predictive value of machine learning models for lymph node metastasis in gastric cancer: A two-center study
Lu T, Lu M, Wu D, Ding YY, Liu HN, Li TT, Song DQ
- 95 Post-operative morbidity after neoadjuvant chemotherapy and resection for gallbladder cancer: A national surgical quality improvement program analysis
Kim M, Stroever S, Aploks K, Ostapenko A, Dong XD, Seshadri R
- 103 Risk factors for recurrence of common bile duct stones after surgical treatment and effect of ursodeoxycholic acid intervention
Yuan WH, Zhang Z, Pan Q, Mao BN, Yuan T
- 113 Clinical efficacy of modified Kamikawa anastomosis in patients with laparoscopic proximal gastrectomy
Wu CY, Lin JA, Ye K
- 124 Clinical effect of laparoscopic radical resection of colorectal cancer based on propensity score matching
Liu Y, Wang XX, Li YL, He WT, Li H, Chen H
- 134 Different timing for abdominal paracentesis catheter placement and drainage in severe acute pancreatitis complicated by intra-abdominal fluid accumulation
Chen R, Chen HQ, Li RD, Lu HM
- 143 Comparison of different preoperative objective nutritional indices for evaluating 30-d mortality and complications after liver transplantation
Li C, Chen HX, Lai YH
- 155 Predictive value of NLR, Fib4, and APRI in the occurrence of liver failure after hepatectomy in patients with hepatocellular carcinoma
Kuang TZ, Xiao M, Liu YF
- 166 Practical effect of different teaching modes in teaching gastrointestinal surgery nursing
Rong XJ, Ning Z
- Observational Study**
- 173 Predictive factors and model validation of post-colon polyp surgery *Helicobacter pylori* infection
Zhang ZS
- Randomized Controlled Trial**
- 186 Micro-power negative pressure wound technique reduces risk of incision infection following loop ileostomy closure
Xu DY, Bai BJ, Shan L, Wei HY, Lin DF, Wang Y, Wang D
- 196 Paravertebral block's effect on analgesia and inflammation in advanced gastric cancer patients undergoing transarterial chemoembolization and microwave ablation
Xiong YF, Wei BZ, Wang YF, Li XF, Liu C

META-ANALYSIS

- 205** Unraveling the efficacy network: A network meta-analysis of adjuvant external beam radiation therapy methods after hepatectomy
Yang GY, He ZW, Tang YC, Yuan F, Cao MB, Ren YP, Li YX, Su XR, Yao ZC, Deng MH
- 215** Estimation of Physiologic Ability and Surgical Stress scoring system for predicting complications following abdominal surgery: A meta-analysis spanning 2004 to 2022
Pang TS, Cao LP
- 228** Role of Oncostatin M in the prognosis of inflammatory bowel disease: A meta-analysis
Yang Y, Fu KZ, Pan G

CASE REPORT

- 239** Endoscopic treatment of extreme esophageal stenosis complicated with esophagotracheal fistula: A case report
Fang JH, Li WM, He CH, Wu JL, Guo Y, Lai ZC, Li GD
- 248** Intestinal tuberculosis with small bowel stricture and hemorrhage as the predominant manifestation: Three case reports
Huang G, Wu KK, Li XN, Kuai JH, Zhang AJ

LETTER TO THE EDITOR

- 257** Sarcopenia in cirrhotic patients: Does frailty matter while waiting for a liver transplant?
Li XJ, He K

ABOUT COVER

Editorial Board Member of *World Journal of Gastrointestinal Surgery*, Renato Pietroletti, PhD, Associate Professor, Professor, Department of Applied Clinical and Biotechnological Sciences, University of L'Aquila, L'Aquila 67100, AQ, Italy. renato.pietroletti@univaq.it

AIMS AND SCOPE

The primary aim of *World Journal of Gastrointestinal Surgery* (*WJGS, World J Gastrointest Surg*) is to provide scholars and readers from various fields of gastrointestinal surgery with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGS mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal surgery and covering a wide range of topics including biliary tract surgical procedures, biliopancreatic diversion, colectomy, esophagectomy, esophagostomy, pancreas transplantation, and pancreatectomy, *etc.*

INDEXING/ABSTRACTING

The *WJGS* is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports/Science Edition, PubMed, PubMed Central, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for *WJGS* as 2.0; IF without journal self cites: 1.9; 5-year IF: 2.2; Journal Citation Indicator: 0.52; Ranking: 113 among 212 journals in surgery; Quartile category: Q3; Ranking: 81 among 93 journals in gastroenterology and hepatology; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Zi-Hang Xu, Production Department Director: Xiang Li, Editorial Office Director: Jia-Ru Fan.

NAME OF JOURNAL

World Journal of Gastrointestinal Surgery

ISSN

ISSN 1948-9366 (online)

LAUNCH DATE

November 30, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Peter Schemmer

POLICY OF CO-AUTHORS

<https://www.wjgnet.com/1948-9366/editorialboard.htm>

PUBLICATION DATE

January 27, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

<https://www.wjgnet.com/bpg/GerInfo/310>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Role of Oncostatin M in the prognosis of inflammatory bowel disease: A meta-analysis

Yue Yang, Kan-Zuo Fu, Gu Pan

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Matsui T, Japan

Received: November 14, 2023

Peer-review started: November 14, 2023

First decision: December 8, 2023

Revised: December 19, 2023

Accepted: December 29, 2023

Article in press: December 29, 2023

Published online: January 27, 2024



Yue Yang, Gu Pan, Department of Gastroenterology III, Heilongjiang Provincial Hospital, Harbin 150036, Heilongjiang Province, China

Kan-Zuo Fu, Department of Nursing, The Second Hospital of Harbin, Harbin 150056, Heilongjiang Province, China

Corresponding author: Gu Pan, BSc, Nurse, Department of Gastroenterology III, Heilongjiang Provincial Hospital, No. 405 Guogoli Street, Nangang District, Harbin 150036, Heilongjiang Province, China. pangu6196@126.com

Abstract

BACKGROUND

Oncostatin M (OSM) is a pleiotropic cytokine which is implicated in the pathogenesis of inflammatory bowel disease (IBD).

AIM

To evaluate the prognostic role of OSM in IBD patients.

METHODS

Literature search was conducted in electronic databases (Google Scholar, Embase, PubMed, Science Direct, Springer, and Wiley). Studies were selected if they reported prognostic information about OSM in IBD patients. Outcome data were synthesized, and meta-analyses were performed to estimate standardized mean differences (SMDs) in OSM levels between treatment responders and non-responders and to seek overall correlations of OSM with other inflammatory biomarkers.

RESULTS

Sixteen studies (818 Crohn's disease and 686 ulcerative colitis patients treated with anti-tumor necrosis factor-based therapies) were included. OSM levels were associated with IBD severity. A meta-analysis found significantly higher OSM levels in non-responders than in responders to therapy [SMD 0.80 (0.33, 1.27); $P = 0.001$], in non-remitters than in remitters [SMD 0.75 (95%CI: 0.35 to 1.16); $P < 0.0001$] and in patients with no mucosal healing than in those with mucosal healing [SMD 0.63 (0.30, 0.95); $P < 0.0001$]. Area under receiver operator curve values showed considerable variability between studies but in general higher OSM levels were associated with poor prognosis. OSM had significant correlations with Simple Endoscopic Score of Crohn's disease [$r = 0.47$ (95%CI: 0.25 to 0.64); $P < 0.0001$], Mayo Endoscopic Score [$r = 0.35$ (95%CI: 0.28 to 0.41); $P < 0.0001$], fecal

calprotectin [$r = 0.19$ (95% CI: 0.08 to 0.3); $P = 0.001$], C-reactive protein [$r = 0.25$ (95% CI: 0.11 to 0.39); $P < 0.0001$], and platelet count [$r = 0.28$ (95% CI: 0.17 to 0.39); $P < 0.0001$].

CONCLUSION

OSM is a potential candidate for determining the severity of disease and predicting the outcomes of anti-tumor necrosis factor-based therapies in IBD patients.

Key Words: Inflammatory bowel disease; Crohn's disease; Ulcerative colitis; Oncostatin M; Prognosis

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Higher Oncostatin M (OSM) expression/levels are found to be associated with worse disease outcomes which shows that OSM can be used as a surrogate marker of poor prognosis in inflammatory bowel disease patients treated with anti-tumor necrosis factor based therapies. Thus, OSM appears to be an attractive biomarker for patient selection and clinical decision-making. However, owing to the presence of heterogeneity in included studies, this evidence should be refined in future studies.

Citation: Yang Y, Fu KZ, Pan G. Role of Oncostatin M in the prognosis of inflammatory bowel disease: A meta-analysis. *World J Gastrointest Surg* 2024; 16(1): 228-238

URL: <https://www.wjgnet.com/1948-9366/full/v16/i1/228.htm>

DOI: <https://dx.doi.org/10.4240/wjgs.v16.i1.228>

INTRODUCTION

Inflammatory bowel disease (IBD) is a disease of the gastrointestinal tract with two main types: Crohn's disease, and ulcerative colitis. Crohn's disease can affect any part of the gastrointestinal tract, whereas ulcerative colitis mainly affects the colon. IBD may arise at any age but usually onsets at early adulthood[1]. Primary surgery is required for approximately 32%, 55%, 70%, and 82% of Crohn's disease patients after 5, 10, 15, and 20 years of diagnosis, respectively[2]. It is speculated that an altered immune response to gut flora depending on individual's hereditary variability and environmental influences may be involved in the etiology of IBD. Age at onset, location, behavior, perianal disease in Crohn's disease and disease extent in ulcerative colitis are important determinants of disease condition[3].

There is an increasing trend in the prevalence of IBD. In the United States, the prevalence of IBD has increased from 0.8% (1.8 million) in 1999 to 1.3% (3 million) in 2015[4]. Globally, the prevalence of IBD is rising in newly industrialized countries[5]. Concomitantly, the prevalence rates of pediatric-onset IBD are also increasing even in regions where this disease was not previously reported[6]. IBD is clinically difficult-to-treat disease that affects younger individuals and leads to long-term morbidity. Resistance to therapeutic agents is a hallmark of its management that necessitates personalized medicine research and development. Use of alternative drugs is frequent, but prediction of disease course and response to a particular therapy can profoundly benefit to patients.

IBD is a lifelong incurable disease that alternates with remission and relapse. Management may require 5-aminosalicylates, thiopurines, steroids, and biologics such as antibodies against tumor necrosis factor alpha (TNF α), vedolizumab, ustekinumab *etc.*[7]. TNF α is involved in IBD onset and progression. Anti-TNF α antibody-based drugs including infliximab, adalimumab, certolizumab, and golimumab are the mainstay in the treatment of IBD[8]. However, about 40% of patients do not respond to anti-TNF therapies, and among those who initially respond, several develop resistance to treatment. This necessitates IBD research to focus not only on the development of newer drugs, but also to identify biomarkers that can predict response to a therapy in advance[9].

Oncostatin M (OSM) is a proinflammatory cytokine belonging to the interleukin-6 family. OSM is produced mostly in hematopoietic tissues including T-lymphocytes, monocytes, macrophages, dendritic cells, neutrophils, eosinophils, and mast cells[10]. It is a pleiotropic factor that participates in several organismic processes including hematopoiesis, differentiation, regeneration, and inflammation. On the other hand, several pathological processes including arthritis, ossification, dermatitis, fibrosis, gingivitis, and carcinogenesis are found to have OSM mediation[11]. In colorectal cancer, higher OSM levels are associated with advanced disease and metastasis[12]. One of the major pathological processes in which the involvement of OSM has been found critical is the inflammation of various parts including the joints, skin, lungs, and intestine[10].

The role of OSM in the pathogenesis of IBD was first described in a discovery of single nucleotide polymorphism in OSM receptors[13]. OSM mediates its effects by binding to a glycoprotein called gp130 and this complex then activates the OSM receptor for signaling[14]. OSM is highly expressed in inflamed mucosa of IBD patients in comparison with normal individuals. Elevated OSM levels are also found in serum of IBD patients. Moreover, higher OSM levels are observed in first-degree relatives of multiple-affected families in comparison with normal families[15]. Several studies have evaluated the prognostic role of OSM in IBD patients. However, there are variabilities in the degree of associations between OSM and disease or interventional outcomes. This necessitates a systematic review of this area. The aim of the

present study was to identify studies that evaluated the prognostic role of OSM in IBD patients in order to synthesize the reported outcomes and to perform meta-analyses of statistical indices for seeking up-to-date evidence of the prognostic role of OSM in IBD prognosis.

MATERIALS AND METHODS

Inclusion and exclusion criteria

Studies were included in this meta-analysis if they: (1) Evaluated IBD patients receiving a therapy who were subjected to OSM measurements in serum or tissue; (2) evaluated prognostic role of OSM in predicting disease outcomes and reported statistical indices of this relationship; and (3) reported correlations between baseline OSM and other important indicators of disease. Exclusion criteria were: (1) Studies involving the prognostic role of OSM in combination with other biomarkers; (2) molecular studies not providing any prognostic outcome; (3) molecular studies evaluating a possible role of OSM in IBD therapeutics; (4) preclinical studies; and (5) reviews and congress abstracts.

Literature search

The literature search was conducted in electronic databases (Google Scholar, Ebsco, PubMed, Science Direct, Springer, and Wiley) using the most relevant keywords. Primary search strategy was: Inflammatory bowel disease OR Crohn's disease OR ulcerative colitis AND oncostatin M AND prognosis OR prognostic OR predictor. The literature search encompassed original research articles published in English language from the date of inception of the database till September 2023.

Data analysis

The quality assessment of the included studies was performed with the Newcastle-Ottawa Scale for the Quality Assessment of Observational Studies. Demographic information, disease pathological indices, previous treatments, study design and conduct features, and study outcome data including OSM levels at baseline, OSM levels in association with response, remission, and mucosal healing rates, statistical data depicting the relationship between baseline OSM levels and outcomes of disease, and correlation coefficients between OSM and other variables of IBD etiology were extracted from research articles of the included studies and were organized in datasheets. Important characteristics of the included studies were tabulated, and outcome data were synthesized for use in analyses.

Area under the receiver-operator curve (AUC) values depicting the relationship between OSM and disease indicators including response rate, remission rate, and mucosal healing rate reported by individual studies were tabulated. A meta-analysis of standardized mean differences (SMDs) in OSM levels between responders and non-responders, remitters and non-remitters, and in patients with mucosal healing and no mucosal healing was performed. Correlation coefficients between OSM and other variables of disease etiology reported by the individual studies were first converted to z-scores and were pooled under random-effects model by deriving variance from respective sample sizes. Overall estimates were back transformed into correlation coefficients.

RESULTS

Sixteen studies[16-31] were included in this review (Figure 1). In these studies, 1353 IBD (818 Crohn's disease and 686 ulcerative colitis) patients were evaluated. Important characteristics of the included studies are presented in Supplementary Tables 1 and 2. The quality of these studies was generally good. The lack of unexposed cohort was the main constraint which was observed for 7 studies. One of the included studies was double-blind, randomized, placebo-controlled with high quality. An assessment of the quality of other included studies with the Newcastle-Ottawa Scale is presented in Supplementary Table 3.

OSM levels in patients with Crohn's disease and ulcerative colitis were generally similar. In Cao *et al*[18], fecal OSM levels (mean \pm SD; pg/mL) were 7 ± 3 in Crohn's disease and 11 ± 4 in ulcerative colitis patients. In Cao *et al*[19], serum OSM levels (mean \pm SD; pg/mL) were 119.4 (range: 34.8 - 240.6) in Crohn's disease and 122.1 (range: 58.7 - 294.9) in ulcerative colitis patients. In Verstockt *et al*[28], OSM expression levels (NPX; OLINK proximity extension technology values) were 6.4 (IQR: 5.5, 7) in Crohn's disease and 6.5 (IQR: 4.35, 4.9) in UC patients. In the study of West *et al*[29], log 2 OSM mRNA expression levels relative to control were 5 (IQR: 4, 8) in Crohn's disease and 5.2 (IQR: 4.5, 7) in ulcerative colitis patients.

However, OSM levels were associated with disease severity. In Cao *et al*[18], fecal OSM levels (mean \pm SD; pg/mL) were 7 ± 2 in mild, 8 ± 5 in moderate, and 14 ± 4 in severe IBD cases whereas in Cao *et al*[19], serum OSM levels (mean \pm SD; pg/mL) were 10 ± 27 in mild, 220 ± 240 in moderate, and 340 ± 150 in severe IBD cases. Mohamed *et al*[24] reported serum OSM levels to be 109.5 ± 25.5 in mild, 116.2 ± 27.6 in moderate, and 144.8 ± 33.5 in severe IBD cases. In West *et al* [29], OSM expression relative to control was 2 (IQR: 0, 2.2) in mild, 5 (IQR: 4, 5.2) in moderate, and 4 (IQR: 3, 4.5) in severe IBD cases.

A meta-analysis found significantly higher OSM levels in non-responders than in responders to therapy [SMD 0.80 (95%CI: 0.33 to 1.27); $P = 0.001$]. OSM levels were also significantly higher in non-remitters in comparison with remitters [SMD 0.75 (95%CI: 0.35 to 1.16); $P < 0.0001$] and in patients with no mucosal healing than in those with mucosal healing [SMD 0.63 (95%CI: 0.30 to 0.95); $P < 0.0001$; Figure 2]. OSM levels and tissue expression data of all included studies are




Figure 1 A flowchart of study screening and selection process. OSM: Oncostatin M; IBD: Inflammatory bowel disease.

presented in [Supplementary Table 4](#).

A synthesis of AUC values of treatment outcomes revealed that anti-TNF treatment had poor outcomes in patients with higher OSM levels in most studies (Tables 1 and 2). AUC values of OSM predicting the response of anti-TNF treatment in IBD patients ranged from 0.56 [95%CI: 0.31 to 0.82] to 0.91 [95%CI: 0.81 to 1.0] whereas the AUC values of OSM in distinguishing between responders and non-responders (including remission and mucosal healing) to anti-TNF therapy ranged from 0.52 to 0.9. Two studies did not report numeric data. Among these, O'Connell *et al*[26], who studied 21 patients with ulcerative colitis, did not find an association of pretreatment colonic OSM expression with the outcomes of infliximab therapy, and Mateos *et al*[21], who studied 22 patients with Crohn's disease, reported that OSM levels measured before induction therapy predicted response to infliximab treatment.

The correlation coefficients between OSM and Simple Endoscopic Score and Mayo Endoscopic Score were 0.47 [95%CI: 0.25 to 0.64] ($P < 0.0001$) and 0.35 [95%CI: 0.28 to 0.41] ($P < 0.0001$) respectively. The correlation coefficients between OSM and fecal calprotectin, C-reactive protein, and platelet count were 0.19 [95%CI: 0.08 to 0.3] ($P = 0.001$), 0.25 [95%CI: 0.11 to 0.39] ($P < 0.0001$), and 0.28 [95%CI: 0.17 to 0.39] ($P < 0.0001$) respectively (Figure 3). The correlation coefficients between OSM and other inflammatory/hematological markers observed in the included studies are given in [Supplementary Table 5](#).

DISCUSSION

OSM has emerged as an important biomarker for determining disease condition and response to anti-TNF therapies in IBD patients. Higher OSM levels are found to be associated with disease severity and therapeutic non-response. AUC values reported by the individual studies showed that higher OSM levels predicted poor response and could be used to distinguish responders from non-responders of anti-TNF therapy. OSM had significant correlations with Simple Endoscopic Score, Mayo Endoscopic Score, fecal calprotectin, C-reactive protein, and platelet count.

Where there is always a need to search for newer drugs, there is also a need to identify markers which can predict the effectiveness of a therapy in advance. Although C-reactive protein is a commonly used marker for predicting response to a therapy, it is non-specific to IBD[16,32,33]. Fecal calprotectin is more important for IBD outcome prediction. Higher FC levels are found to be associated with no response to therapy[34-36]. Cao *et al*[18] found higher AUC value for fecal calprotectin (0.834) than fecal OSM (0.763) in predicting response to anti-TNF therapy. In the meta-analysis of correlation coefficients, we have found a significant correlation between OSM and fecal calprotectin in IBD patients.

It has been observed that OSM predicts therapeutic response better to anti-TNF than to other pharmacological treatments. Bertani *et al*[17] found OSM to be a useful biomarker for predicting response to anti-TNF therapy (AUC 0.91) but not to vedolizumab (AUC 0.56). Verstockt *et al*[28] also found low AUC values for distinguishing remitters from non-remitters after vedolizumab therapy both by serum OSM (0.51) and colonic OSM (0.685). Nishioka *et al*[25] found a relatively higher AUC value for mucosal OSM expression in distinguishing resistant from sensitive patients to anti-TNF therapy (0.83) compared to ustekinumab (0.77).

Table 1 Area under receiver operator curve values for the prediction of treatment outcomes by the Oncostatin M

Ref.	Prognostic association	OSM cutoff	AUC [95%CI]	Sensitivity [95%CI]	Specificity [95%CI]
Bertani <i>et al</i> [16], 2020	Prediction of no mucosal healing after anti-TNF α therapy at week 54 by baseline serum OSM	14	0.91 [0.81, 1]	96% [82, 100]	89% [67, 97]
Bertani <i>et al</i> [16], 2020	Prediction of no mucosal healing after anti-TNF α therapy at week 54 by serum OSM at week 14		0.83 [0.7, 0.95]		
Bertani <i>et al</i> [17], 2022	Prediction of no mucosal healing after anti-TNF α therapy at week 54 by baseline serum OSM	14	0.91 [0.84, 0.99]	91% [78, 97]	90% [75, 97]
Bertani <i>et al</i> [17], 2022	Prediction of non-response to vedolizumab therapy at week 54 by baseline serum OSM		0.56 [0.42, 0.7]		
Cao <i>et al</i> [18], 2021	Prediction of non-response to infliximab at week 54 by baseline fecal OSM		0.638		
Cao <i>et al</i> [18], 2021	Prediction of non-response to infliximab at week 28 by baseline fecal OSM	132	0.763	66.7%	92.5%
Ezirike Ladipo <i>et al</i> [21], 2021	Prediction of response to anti-TNF therapy by OSM expression in biopsies		OSM expression in pre-treatment biopsies did not predict response to anti-TNF in a pediatric population		
Mateos <i>et al</i> [22], 2021	Prediction of response to infliximab in a calprotectin log drop measurement model		OSM was found to have predicting ability to infliximab response		
Minar <i>et al</i> [23], 2019	Prediction of no remission after anti-TNF α therapy at week 12 by baseline serum OSM	144	0.71 [0.52, 0.89]	71%	78%
Minar <i>et al</i> [23], 2019	Prediction of non-response to anti-TNF α therapy at week 12 by baseline serum OSM	117	0.69 [0.5, 0.89]		
Mohamed <i>et al</i> [24], 2022	Prediction of no remission after anti-TNF α therapy by baseline serum OSM	119	0.56 [0.31, 0.82]	66.7%	54.2%
O'connell <i>et al</i> [26], 2022	Prediction of response to infliximab by colonic OSM expression		No association of pretreatment colonic OSM expression with outcomes of Infliximab therapy		
Zhou <i>et al</i> [31], 2019	Prediction of the response to PF-00547659 (anti-human mucosal addressin cell adhesion molecule-1) therapy		Baseline OSM expression/levels were unable to predict response		

AUC: Area under receiver operator curve; OSM: Oncostatin M; TNF: Tumor necrosis factor.

Minar *et al*[23] found no association between clinical remission and OSM 3 months after anti-TNF therapy but observed a significant association between OSM and clinical response one year after treatment. They suggested that duration of response evaluation may affect the outcomes. However, Cao *et al*[18] found higher AUC value (0.76) of serum OSM to predict nonresponse at week 28 in comparison with AUC value observed at week 54 of treatment (0.64). Verstockt *et al* [28] found no significant association between serum OSM and endoscopic remission after 6 months of anti-TNF therapy. On the other hand, Bertani *et al*[16,17] and Guo *et al*[20] found significant associations between serum OSM and response to anti-TNF therapy after one year of treatment.

In a transcriptomic gene expression study, Zhou *et al*[31] found that on week 12 of PF-00547659 (anti-human mucosal addressin cell adhesion molecule-1 antibody) treatment the OSM expression and serum levels decreased profoundly from baseline in patients with ulcerative colitis who achieved response, remission, or mucosal healing. Whereas the change in serum OSM levels was 1.4-fold among responders, the change in OSM expression among responders and those achieving mucosal healing was 6.1-fold and 7.4-fold respectively[31]. Verstockt *et al*[28] who found mucosal OSM to predict response to anti-TNF therapy, did not find serum OSM to do the same. In the study of Zhou *et al*[31], baseline OSM expression did not predict therapeutic outcomes. Whether this difference can be attributed to the mechanism of action of drug (PF-00547659 *vs* anti-TNF based therapies) remains to be evaluated. OSM acts synergistically with TNF to promote inflammation in stromal cells and this phenomenon may not be exhibited by the other modulators such as human mucosal addressin cell adhesion molecule-1.

Verstockt *et al*[28] performed immunohistochemical staining on resected tissues and found OSM expression in the macrophages lying in superficial lamina propria as well as in the epithelial granulomas and multinucleated giant cells. In this study, macrophagic OSM expression had a strong correlation with mucosal OSM. OSM expression is found consistently higher in inflamed parts of intestine where it promotes inflammation in gut stromal cells in response to microbial challenges[22]. O'Connell *et al*[26] who studied 21 acute severe ulcerative colitis patients observed a greater degree of immunostaining in the mucosal epithelial cells rather than stromal cells which provides impetus for studying OSM immunostaining in different IBD phenotypes. This study did not find an association between OSM expression levels and response to infliximab used as rescue therapy.

We found that OSM levels were not much different between patients with Crohn's disease and those with ulcerative colitis[19,28,29]. However, Cao *et al*[18] found fecal OSM levels to be significantly higher in patients with ulcerative colitis

Table 2 Area under receiver operator curve values for identifying/distinguishing treatment response by OSM

Ref.	Prognostic association	OSM cutoff	AUC [95%CI]	Sensitivity [95%CI]	Specificity [95%CI]
Cao <i>et al</i> [18], 2021	Identification of mucosal healing after infliximab therapy by fecal OSM		0.702		
Cao <i>et al</i> [18], 2021	Identification of clinical remission after infliximab therapy by fecal OSM		0.674		
Cao <i>et al</i> [19], 2022	Identification of mucosal healing after infliximab therapy by serum OSM	64.1	0.84 [0.75, 0.91]	81.8%	80.8%
Cao <i>et al</i> [19], 2022	Identification of clinical response to infliximab therapy by serum OSM	83	0.90 [0.8, 0.96]	86.4%	87%
Cao <i>et al</i> [19], 2022	Identification of clinical remission after infliximab therapy by serum OSM	98.9	0.9 [0.83, 0.95]	82.1%	86.4%
Guo <i>et al</i> [20], 2022	Distinction between remitters and non-remitters after 1 year of anti-TNF α therapy in CD patients by serum OSM	169	0.88 [0.79, 0.96]	76% [58, 88]	91% [80, 96]
Guo <i>et al</i> [20], 2022	Distinction between remitters and non-remitters after 1 year of anti-TNF α therapy in UC patients by serum OSM	234	0.94 [0.87, 1]	80% [55, 93]	96% [79, 99]
Nishioka <i>et al</i> [25], 2021	Distinction between anti-TNF α resistant and sensitive patients by mucosal OSM mRNA		0.83		
Nishioka <i>et al</i> [25], 2021	Distinction between ustekinumab resistant and sensitive patients by mucosal OSM mRNA		0.77		
Verstockt <i>et al</i> [28], 2021	Distinction between remitters and non-remitters 6 months after surgery by serum OSM		0.80 [0.68, 0.92]		
Verstockt <i>et al</i> [28], 2021	Distinction between remitters and non-remitters after anti-TNF α therapy by serum OSM		0.52 [0.44, 0.61]		
Verstockt <i>et al</i> [28], 2021	Distinction between remitters and non-remitters after anti-TNF α therapy by colonic OSM		0.74 [0.54, 0.94]		
Verstockt <i>et al</i> [28], 2021	Distinction between remitters and non-remitters after vedolizumab therapy by serum OSM		0.51 [0.43, 0.59]		
Verstockt <i>et al</i> [28], 2021	Distinction between remitters and non-remitters after vedolizumab therapy by colonic OSM		0.69 [0.53, 0.84]		
West <i>et al</i> [29], 2017	Distinction between responders and non-responders to infliximab by mucosal OSM mRNA		0.99	100	91.7
Yokoyama <i>et al</i> [30], 2023	Distinction between responders and non-responders to anti-TNF α by mucosal OSM mRNA		0.94		
Yokoyama <i>et al</i> [30], 2023	Distinction between CORT-dependent vs non-dependent remission by mucosal OSM mRNA		0.79		
Zhou <i>et al</i> [31], 2019	Distinction between responders and non-responders to F-00547659 by change in OSM expression in inflamed tissue		0.88		
Zhou <i>et al</i> [31], 2019	Distinction between remitters and non-remitters to F-00547659 by change in OSM expression in inflamed tissue		0.81		
Zhou <i>et al</i> [31], 2019	Distinction between mucosal healing and no mucosal healing by F-00547659 therapy by the change in OSM expression during treatment in inflamed tissue		0.83		

AUC: Area under receiver operator curve; OSM: Oncostatin M; TNF: Tumor necrosis factor.

than in Crohn's disease. On the other hand, OSM levels were associated with disease severity as there was an increasing trend of OSM levels from mild, to moderate and severe disease[18,19,24,29]. OSM levels are also found higher in IBD patients in comparison with healthy controls[18,21,24,28]. These outcomes are similar to a study that characterized serum inflammatory protein profile and found a differential regulation of OSM between patients with ulcerative colitis and healthy controls but not between patients with Crohn's disease and ulcerative colitis[37].

Although, most of the studies included herein identified OSM as a potential biomarker of IBD severity and predictor of response to anti-TNF therapies, some studies could not find so. Ezirike Ladipo *et al*[21] who studied 98 children with IBD reported that OSM or OSM receptor expression did not predict response to anti-TNF treatment, although OSM was associated with disease severity. Mohamed *et al*[24] reported that OSM did not have an appreciable ability to predict the response to therapy. O'Connell *et al*[26] also reported that colonic OSM expression was unable to predict infliximab treatment outcomes. Verstockt *et al*[28] reported that serum OSM levels had an AUC value of 0.52 in distinguishing




Figure 2 A forest graph showing the outcomes of a meta-analysis of standardized mean differences between responders and non-responders, remitters and non-remitters, and mucosal healers and non-healers of anti-tumor necrosis factor-based therapies in patients with inflammatory bowel disease. OSM: Oncostatin M; TNF: Tumor necrosis factor; SMD: Standardized mean difference.

between remitters and non-remitters after anti-TNF therapy. In the study of Zhou *et al*[31], baseline OSM levels were unable to predict response to PF-00547659.

West *et al*[29] reported that among the 64 cytokines evaluated, the OSM and its receptor were most intensely overexpressed in the inflamed mucosa of IBD patients. They suggested that OSM may also be involved in developing resistance to anti-TNF therapies. According to West *et al*[29], haematopoietically derived OSM appears to mediate intestinal pathology by promoting inflammatory behavior in gut-resident stromal cells which is a novel system of leukocyte-stromal cell cross talk that may have relevance in multiple mucosal tissues. Because of its stabilizing interactions with extracellular matrix components, OSM may play a critical role in the etiology of disease. Thus, OSM may act as an inflammatory amplifier and driver of disease chronicity by promoting chemokines, cytokines, and adhesion factor production from intestinal stromal cells.

There are some limitations of this review. There were inconsistencies in the outcome data of individual studies in measuring OSM levels/expression and their numerical presentations due to which not all data could be meta-analyzed. Moreover, in some studies, numerical outcome data were not accompanied by the variance. To account for such constraints, we either performed a meta-analysis of SMDs or tabulated the outcomes systematically. This constraint also precluded us from having a generalized estimate of OSM levels/concentrations. Moreover, for data where meta-analyses were possible, we observed higher statistical heterogeneity. Methodological differences of individual studies could have also played a role in the variabilities observed in this review.

CONCLUSION

Most of the studies attempting to seek relationship between OSM and disease or treatment outcomes have found that higher OSM expression/levels to be associated with worse disease outcomes which shows that OSM can be used as a surrogate marker of poor prognosis in IBD patients treated with anti-TNF treatments. This makes OSM an attractive biomarker for patient selection and clinical decision-making. However, some studies could not recognize such associations and others found non-significant correlations between OSM and other indicators such as fecal calprotectin, C-reactive protein, and platelets. Therefore, more studies are required to validate present day evidence and to explore the behavior of OSM for IBD treatments other than anti-TNF drugs.




Figure 3 A forest graph showing the outcomes of a meta-analysis of correlation coefficients between baseline Oncostatin M and other baseline inflammatory biomarkers/disease indicators. OSM: Oncostatin M; TNF: Tumor necrosis factor.

ARTICLE HIGHLIGHTS

Research background

Oncostatin M (OSM) is a pleiotropic factor that participates in several physiological processes such as hematopoiesis, differentiation, regeneration, and inflammation, and pathological processes such as arthritis, ossification, dermatitis, fibrosis, gingivitis, and carcinogenesis.

Research motivation

Higher OSM levels in patients with inflammatory bowel disease (IBD) provided impetus for reviewing the outcomes of studies that evaluated the prognostic role of OSM in IBD patients.

Research objectives

The objective of this research was to systematically review relevant studies and perform meta-analyses of statistical indices for seeking current evidence about the role of OSM in IBD prognosis.

Research methods

After a literature search in electronic databases, studies were identified for synthesis. Meta-analyses were performed to estimate standardized mean differences in OSM levels between responders and non-responders, and to pool correlations of OSM with other inflammatory biomarkers.

Research results

OSM levels were associated with disease severity and were significantly higher in non-responders, in non-remitters, and

in patients with no mucosal healing after anti-tumor necrosis factor (anti-TNF) therapy. Area under receiver operator curve values showed considerable variability between studies but in general higher OSM levels were associated with poor prognosis. OSM had significant correlations with Simple Endoscopic Score of Crohn's disease, Mayo Endoscopic Score, fecal calprotectin, C-reactive protein, and platelet count.

Research conclusions

OSM can potentially determine IBD severity and can predict the outcomes of anti-tumor necrosis factor-based therapies.

Research perspectives

Future studies may refine the outcomes reported herein. It could also be interesting to explore the role of OSM in achieving response to non-anti-TNF therapies.

FOOTNOTES

Co-first authors: Yue Yang and Kan-Zuo Fu.

Author contributions: Yang Y and Fu KZ wrote the manuscript; Yang Y, Fu KZ, and Pan G collected and analyzed the data; all authors read and approved the final manuscript.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

PRISMA 2009 Checklist statement: The authors have read the PRISMA 2009 Checklist, and the manuscript was prepared and revised according to the PRISMA 2009 Checklist.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Gu Pan [0009-0009-9248-5714](https://orcid.org/0009-0009-9248-5714).

S-Editor: Gong ZM

L-Editor: A

P-Editor: Xu ZH

REFERENCES

- 1 **Wehkamp J**, Götz M, Herrlinger K, Steurer W, Stange EF. Inflammatory Bowel Disease. *Dtsch Arztebl Int* 2016; **113**: 72-82 [PMID: 26900160 DOI: [10.3238/arztebl.2016.0072](https://doi.org/10.3238/arztebl.2016.0072)]
- 2 **Sato Y**, Matsui T, Yano Y, Tsurumi K, Okado Y, Matsushima Y, Koga A, Takahashi H, Ninomiya K, Ono Y, Takatsu N, Beppu T, Nagahama T, Hisabe T, Takaki Y, Hirai F, Yao K, Higashi D, Futami K, Washio M. Long-term course of Crohn's disease in Japan: Incidence of complications, cumulative rate of initial surgery, and risk factors at diagnosis for initial surgery. *J Gastroenterol Hepatol* 2015; **30**: 1713-1719 [PMID: 26094852 DOI: [10.1111/jgh.13013](https://doi.org/10.1111/jgh.13013)]
- 3 **Burisch J**, Jess T, Martinato M, Lakatos PL; ECCO -EpiCom. The burden of inflammatory bowel disease in Europe. *J Crohns Colitis* 2013; **7**: 322-337 [PMID: 23395397 DOI: [10.1016/j.crohns.2013.01.010](https://doi.org/10.1016/j.crohns.2013.01.010)]
- 4 **Dahlhamer JM**, Zammitti EP, Ward BW, Wheaton AG, Croft JB. Prevalence of Inflammatory Bowel Disease Among Adults Aged ≥ 18 Years - United States, 2015. *MMWR Morb Mortal Wkly Rep* 2016; **65**: 1166-1169 [PMID: 27787492 DOI: [10.15585/mmwr.mm6542a3](https://doi.org/10.15585/mmwr.mm6542a3)]
- 5 **Ng SC**, Shi HY, Hamidi N, Underwood FE, Tang W, Benchimol EI, Panaccione R, Ghosh S, Wu JCY, Chan FKL, Sung JY, Kaplan GG. Worldwide incidence and prevalence of inflammatory bowel disease in the 21st century: a systematic review of population-based studies. *Lancet* 2017; **390**: 2769-2778 [PMID: 29050646 DOI: [10.1016/S0140-6736\(17\)32448-0](https://doi.org/10.1016/S0140-6736(17)32448-0)]
- 6 **Kuenzig ME**, Fung SG, Marderfeld L, Mak JWY, Kaplan GG, Ng SC, Wilson DC, Cameron F, Henderson P, Kotze PG, Bhatti J, Fang V, Gerber S, Guay E, Kotteduwa Jayawarden S, Kadota L, Maldonado D F, Osei JA, Sandarage R, Stanton A, Wan M; InsightScope Pediatric IBD Epidemiology Group, Benchimol EI. Twenty-first Century Trends in the Global Epidemiology of Pediatric-Onset Inflammatory Bowel Disease: Systematic Review. *Gastroenterology* 2022; **162**: 1147-1159.e4 [PMID: 34995526 DOI: [10.1053/j.gastro.2021.12.282](https://doi.org/10.1053/j.gastro.2021.12.282)]
- 7 **Wang LF**, Chen PR, He SK, Duan SH, Zhang Y. Predictors and optimal management of tumor necrosis factor antagonist nonresponse in inflammatory bowel disease: A literature review. *World J Gastroenterol* 2023; **29**: 4481-4498 [PMID: 37621757 DOI: [10.3748/wjg.v29.i29.4481](https://doi.org/10.3748/wjg.v29.i29.4481)]
- 8 **Guerra I**, Bermejo F. Management of inflammatory bowel disease in poor responders to infliximab. *Clin Exp Gastroenterol* 2014; **7**: 359-367 [PMID: 25258548 DOI: [10.2147/CEG.S45297](https://doi.org/10.2147/CEG.S45297)]
- 9 **Thomas H**. IBD: Oncostatin M promotes inflammation in IBD. *Nat Rev Gastroenterol Hepatol* 2017; **14**: 261 [PMID: 28400625 DOI: [10.1038/nrgastro.2017.47](https://doi.org/10.1038/nrgastro.2017.47)]
- 10 **West NR**, Owens BMJ, Hegazy AN. The oncostatin M-stromal cell axis in health and disease. *Scand J Immunol* 2018; **88**: e12694 [PMID: 30111111 DOI: [10.1111/sji.12694](https://doi.org/10.1111/sji.12694)]

- 29926972 DOI: [10.1111/sji.12694](https://doi.org/10.1111/sji.12694)]
- 11 **Wolf CL**, Pruett C, Lighter D, Jorczyk CL. The clinical relevance of OSM in inflammatory diseases: a comprehensive review. *Front Immunol* 2023; **14**: 1239732 [PMID: [37841259](https://pubmed.ncbi.nlm.nih.gov/37841259/) DOI: [10.3389/fimmu.2023.1239732](https://doi.org/10.3389/fimmu.2023.1239732)]
 - 12 **Gurluler E**, Tumay LV, Guner OS, Kucukmetin NT, Hizli B, Zorluoglu A. Oncostatin-M as a novel biomarker in colon cancer patients and its association with clinicopathologic variables. *Eur Rev Med Pharmacol Sci* 2014; **18**: 2042-2047 [PMID: [25027345](https://pubmed.ncbi.nlm.nih.gov/25027345/)]
 - 13 **Mirkov MU**, Verstockt B, Cleynen I. Genetics of inflammatory bowel disease: beyond NOD2. *Lancet Gastroenterol Hepatol* 2017; **2**: 224-234 [PMID: [28404137](https://pubmed.ncbi.nlm.nih.gov/28404137/) DOI: [10.1016/S2468-1253\(16\)30111-X](https://doi.org/10.1016/S2468-1253(16)30111-X)]
 - 14 **Adrian-Segarra JM**, Schindler N, Gajawada P, Lörchner H, Braun T, Pöling J. The AB loop and D-helix in binding site III of human Oncostatin M (OSM) are required for OSM receptor activation. *J Biol Chem* 2018; **293**: 7017-7029 [PMID: [29511087](https://pubmed.ncbi.nlm.nih.gov/29511087/) DOI: [10.1074/jbc.RA118.001920](https://doi.org/10.1074/jbc.RA118.001920)]
 - 15 **Verstockt S**, Verstockt B, Vermeire S. Oncostatin M as a new diagnostic, prognostic and therapeutic target in inflammatory bowel disease (IBD). *Expert Opin Ther Targets* 2019; **23**: 943-954 [PMID: [31587593](https://pubmed.ncbi.nlm.nih.gov/31587593/) DOI: [10.1080/14728222.2019.1677608](https://doi.org/10.1080/14728222.2019.1677608)]
 - 16 **Bertani L**, Fornai M, Fornili M, Antonioli L, Benvenuti L, Tapete G, Baiano Svizzero G, Ceccarelli L, Mumolo MG, Baglietto L, de Bortoli N, Bellini M, Marchi S, Costa F, Blandizzi C. Serum oncostatin M at baseline predicts mucosal healing in Crohn's disease patients treated with infliximab. *Aliment Pharmacol Ther* 2020; **52**: 284-291 [PMID: [32506635](https://pubmed.ncbi.nlm.nih.gov/32506635/) DOI: [10.1111/apt.15870](https://doi.org/10.1111/apt.15870)]
 - 17 **Bertani L**, Barberio B, Fornili M, Antonioli L, Zanzi F, Casadei C, Benvenuti L, Facchin S, D'Antongiovanni V, Lorenzon G, Ceccarelli L, Baglietto L, de Bortoli N, Bellini M, Costa F, Savarino EV, Fornai M. Serum oncostatin M predicts mucosal healing in patients with inflammatory bowel diseases treated with anti-TNF, but not vedolizumab. *Dig Liver Dis* 2022; **54**: 1367-1373 [PMID: [35393259](https://pubmed.ncbi.nlm.nih.gov/35393259/) DOI: [10.1016/j.dld.2022.03.008](https://doi.org/10.1016/j.dld.2022.03.008)]
 - 18 **Cao Y**, Dai Y, Zhang L, Wang D, Hu W, Yu Q, Wang X, Yu P, Liu W, Ping Y, Sun T, Sang Y, Liu Z, Chen Y, Tao Z. Combined Use of Fecal Biomarkers in Inflammatory Bowel Diseases: Oncostatin M and Calprotectin. *J Inflamm Res* 2021; **14**: 6409-6419 [PMID: [34880643](https://pubmed.ncbi.nlm.nih.gov/34880643/) DOI: [10.2147/JIR.S342846](https://doi.org/10.2147/JIR.S342846)]
 - 19 **Cao Y**, Dai Y, Zhang L, Wang D, Yu Q, Hu W, Wang X, Yu P, Ping Y, Sun T, Sang Y, Liu Z, Chen Y, Tao Z. Serum oncostatin M is a potential biomarker of disease activity and infliximab response in inflammatory bowel disease measured by chemiluminescence immunoassay. *Clin Biochem* 2022; **100**: 35-41 [PMID: [34843732](https://pubmed.ncbi.nlm.nih.gov/34843732/) DOI: [10.1016/j.clinbiochem.2021.11.011](https://doi.org/10.1016/j.clinbiochem.2021.11.011)]
 - 20 **Guo A**, Ross C, Chande N, Gregor J, Ponich T, Khanna R, Sey M, Beaton M, Yan B, Kim RB, Wilson A. High oncostatin M predicts lack of clinical remission for patients with inflammatory bowel disease on tumor necrosis factor α antagonists. *Sci Rep* 2022; **12**: 1185 [PMID: [35075155](https://pubmed.ncbi.nlm.nih.gov/35075155/) DOI: [10.1038/s41598-022-05208-9](https://doi.org/10.1038/s41598-022-05208-9)]
 - 21 **Ezirike Ladipo J**, He Z, Chikwava K, Robbins K, Beri J, Molle-Rios Z. Oncostatin-M Does Not Predict Treatment Response in Inflammatory Bowel Disease in a Pediatric Cohort. *J Pediatr Gastroenterol Nutr* 2021; **73**: 352-357 [PMID: [34117193](https://pubmed.ncbi.nlm.nih.gov/34117193/) DOI: [10.1097/MPG.0000000000003201](https://doi.org/10.1097/MPG.0000000000003201)]
 - 22 **Mateos B**, Sáez-González E, Moret I, Hervás D, Iborra M, Cerrillo E, Tortosa L, Nos P, Beltrán B. Plasma Oncostatin M, TNF- α , IL-7, and IL-13 Network Predicts Crohn's Disease Response to Infliximab, as Assessed by Calprotectin Log Drop. *Dig Dis* 2021; **39**: 1-9 [PMID: [32325460](https://pubmed.ncbi.nlm.nih.gov/32325460/) DOI: [10.1159/000508069](https://doi.org/10.1159/000508069)]
 - 23 **Minar P**, Lehn C, Tsai YT, Jackson K, Rosen MJ, Denson LA. Elevated Pretreatment Plasma Oncostatin M Is Associated With Poor Biochemical Response to Infliximab. *Crohn's Colitis* 2019; **1**: otz026 [PMID: [31667468](https://pubmed.ncbi.nlm.nih.gov/31667468/) DOI: [10.1093/crocol/otz026](https://doi.org/10.1093/crocol/otz026)]
 - 24 **Mohamed GA**, Mohamed HAE-L, Abo Halima AS, Elshaarawy MEA, Khedr A. Changes in serum oncostatin M levels during treatment of inflammatory bowel disease. *Egypt J Hosp Med* 2022; **89**: 7217-7225 [DOI: [10.21608/ejhm.2022.273068](https://doi.org/10.21608/ejhm.2022.273068)]
 - 25 **Nishioka K**, Ogino H, Chinen T, Ihara E, Tanaka Y, Nakamura K, Ogawa Y. Mucosal IL23A expression predicts the response to Ustekinumab in inflammatory bowel disease. *J Gastroenterol* 2021; **56**: 976-987 [PMID: [34448069](https://pubmed.ncbi.nlm.nih.gov/34448069/) DOI: [10.1007/s00535-021-01819-7](https://doi.org/10.1007/s00535-021-01819-7)]
 - 26 **O'Connell J**, Doherty J, Buckley A, Cormican D, Dunne C, Hartery K, Larkin J, MacCarthy F, McCormick P, McKiernan S, Mehigan B, Muldoon C, Ryan C, O'Sullivan J, Kevans D. Colonic oncostatin M expression evaluated by immunohistochemistry and infliximab therapy outcome in corticosteroid-refractory acute severe ulcerative colitis. *Intest Res* 2022; **20**: 381-385 [PMID: [35263959](https://pubmed.ncbi.nlm.nih.gov/35263959/) DOI: [10.5217/ir.2021.00073](https://doi.org/10.5217/ir.2021.00073)]
 - 27 **Verstockt B**, Verstockt S, Dehairs J, Ballet V, Blevi H, Wollants WJ, Breynaert C, Van Assche G, Vermeire S, Ferrante M. Low TREM1 expression in whole blood predicts anti-TNF response in inflammatory bowel disease. *EBioMedicine* 2019; **40**: 733-742 [PMID: [30685385](https://pubmed.ncbi.nlm.nih.gov/30685385/) DOI: [10.1016/j.ebiom.2019.01.027](https://doi.org/10.1016/j.ebiom.2019.01.027)]
 - 28 **Verstockt S**, Verstockt B, Machiels K, Vancamelbeke M, Ferrante M, Cleynen I, De Hertogh G, Vermeire S. Oncostatin M Is a Biomarker of Diagnosis, Worse Disease Prognosis, and Therapeutic Nonresponse in Inflammatory Bowel Disease. *Inflamm Bowel Dis* 2021; **27**: 1564-1575 [PMID: [33624092](https://pubmed.ncbi.nlm.nih.gov/33624092/) DOI: [10.1093/ibd/izab032](https://doi.org/10.1093/ibd/izab032)]
 - 29 **West NR**, Hegazy AN, Owens BMJ, Bullers SJ, Linggi B, Buonocore S, Coccia M, Görtz D, This S, Stockenhuber K, Pott J, Friedrich M, Ryzhakov G, Baribaud F, Brodmerkel C, Cieluch C, Rahman N, Müller-Newen G, Owens RJ, Kühl AA, Maloy KJ, Plevy SE; Oxford IBD Cohort Investigators, Keshav S, Travis SPL, Powrie F. Oncostatin M drives intestinal inflammation and predicts response to tumor necrosis factor-neutralizing therapy in patients with inflammatory bowel disease. *Nat Med* 2017; **23**: 579-589 [PMID: [28368383](https://pubmed.ncbi.nlm.nih.gov/28368383/) DOI: [10.1038/nm.4307](https://doi.org/10.1038/nm.4307)]
 - 30 **Yokoyama Y**, Yamakawa T, Miyake T, Kazama T, Hayashi Y, Hirayama D, Nakase H. Mucosal gene expression of inflammatory cytokines as biomarkers for predicting treatment response in patients with inflammatory bowel disease. *Research Square* 2023 [DOI: [10.21203/rs.3.rs-2841128/v1](https://doi.org/10.21203/rs.3.rs-2841128/v1)]
 - 31 **Zhou H**, Xi L, Ziemek D, O'Neil S, Lee J, Stewart Z, Zhan Y, Zhao S, Zhang Y, Page K, Huang A, Maciejewski M, Zhang B, Gorelick KJ, Fitz L, Pradhan V, Cataldi F, Vincent M, Von Schack D, Hung K, Hassan-Zahrae M. Molecular Profiling of Ulcerative Colitis Subjects from the TURANDOT Trial Reveals Novel Pharmacodynamic/Efficacy Biomarkers. *J Crohn's Colitis* 2019; **13**: 702-713 [PMID: [30901380](https://pubmed.ncbi.nlm.nih.gov/30901380/) DOI: [10.1093/ecco-jcc/jjy217](https://doi.org/10.1093/ecco-jcc/jjy217)]
 - 32 **Kiss LS**, Papp M, Lovasz BD, Vegh Z, Golovics PA, Janka E, Varga E, Szathmari M, Lakatos PL. High-sensitivity C-reactive protein for identification of disease phenotype, active disease, and clinical relapses in Crohn's disease: a marker for patient classification? *Inflamm Bowel Dis* 2012; **18**: 1647-1654 [PMID: [22081542](https://pubmed.ncbi.nlm.nih.gov/22081542/) DOI: [10.1002/ibd.21933](https://doi.org/10.1002/ibd.21933)]
 - 33 **Oh K**, Oh EH, Baek S, Song EM, Kim GU, Seo M, Hwang SW, Park SH, Yang DH, Kim KJ, Byeon JS, Myung SJ, Yang SK, Ye BD. Elevated C-reactive protein level during clinical remission can predict poor outcomes in patients with Crohn's disease. *PLoS One* 2017; **12**: e0179266 [PMID: [28622356](https://pubmed.ncbi.nlm.nih.gov/28622356/) DOI: [10.1371/journal.pone.0179266](https://doi.org/10.1371/journal.pone.0179266)]
 - 34 **Dai C**, Cao Q, Jiang M. Clinical Utility of Fecal Calprotectin Monitoring in Asymptomatic Patients with Inflammatory Bowel Disease.

- Inflamm Bowel Dis* 2017; **23**: E46-E47 [PMID: 28816761 DOI: 10.1097/MIB.0000000000001225]
- 35 **Beltrán B**, Iborra M, Sáez-González E, Marqués-Miñana MR, Moret I, Cerrillo E, Tortosa L, Bastida G, Hinojosa J, Poveda-Andrés JL, Nos P. Fecal Calprotectin Pretreatment and Induction Influximab Levels for Prediction of Primary Nonresponse to Influximab Therapy in Crohn's Disease. *Dig Dis* 2019; **37**: 108-115 [PMID: 30149385 DOI: 10.1159/000492626]
- 36 **Petryszyn P**, Staniak A, Wolosińska A, Ekk-Cierniakowski P. Faecal calprotectin as a diagnostic marker of inflammatory bowel disease in patients with gastrointestinal symptoms: meta-analysis. *Eur J Gastroenterol Hepatol* 2019; **31**: 1306-1312 [PMID: 31464777 DOI: 10.1097/MEG.0000000000001509]
- 37 **Andersson E**, Bergemalm D, Kruse R, Neumann G, D'Amato M, Repsilber D, Halfvarson J. Subphenotypes of inflammatory bowel disease are characterized by specific serum protein profiles. *PLoS One* 2017; **12**: e0186142 [PMID: 28982144 DOI: 10.1371/journal.pone.0186142]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: office@baishideng.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

