


World Journal of *Gastrointestinal Surgery*

World J Gastrointest Surg 2024 January 27; 16(1): 1-259



EDITORIAL

- 1 Novel prognostic factors after radical resection of hepatocellular carcinoma: Updating an old issue
Bencini L
- 6 Prospects in the application of ultrasensitive chromosomal aneuploidy detection in precancerous lesions of gastric cancer
Qian ST, Xie FF, Zhao HY, Liu QS, Cai DL

MINIREVIEWS

- 13 Prognostic value of ultrasound in early arterial complications post liver transplant
Zhao NB, Chen Y, Xia R, Tang JB, Zhao D

ORIGINAL ARTICLE**Case Control Study**

- 21 Added value of ratio of cross diameters of the appendix in ultrasound diagnosis of acute appendicitis
Gu FW, Wu SZ

Retrospective Cohort Study

- 29 Oncological features and prognosis of colorectal cancer in human immunodeficiency virus-positive patients: A retrospective study
Yang FY, He F, Chen DF, Tang CL, Woraiikat S, Li Y, Qian K

Retrospective Study

- 40 Laparoscopic *vs* open surgery for gastric cancer: Assessing time, recovery, complications, and markers
Lu YY, Li YX, He M, Wang YL
- 49 Single-incision laparoscopic transabdominal preperitoneal repair in the treatment of adult female patients with inguinal hernia
Zhu XJ, Jiao JY, Xue HM, Chen P, Qin CF, Wang P
- 59 Computerized tomography-guided therapeutic percutaneous puncture catheter drainage-combined with somatostatin for severe acute pancreatitis: An analysis of efficacy and safety
Zheng XL, Li WL, Lin YP, Huang TL
- 67 Impact of open hepatectomy on postoperative bile leakage in patients with biliary tract cancer
Wu G, Li WY, Gong YX, Lin F, Sun C
- 76 Clinical observation of gastrointestinal function recovery in patients after hepatobiliary surgery
Zeng HJ, Liu JJ, Yang YC

- 85 Predictive value of machine learning models for lymph node metastasis in gastric cancer: A two-center study
Lu T, Lu M, Wu D, Ding YY, Liu HN, Li TT, Song DQ
- 95 Post-operative morbidity after neoadjuvant chemotherapy and resection for gallbladder cancer: A national surgical quality improvement program analysis
Kim M, Stroever S, Aploks K, Ostapenko A, Dong XD, Seshadri R
- 103 Risk factors for recurrence of common bile duct stones after surgical treatment and effect of ursodeoxycholic acid intervention
Yuan WH, Zhang Z, Pan Q, Mao BN, Yuan T
- 113 Clinical efficacy of modified Kamikawa anastomosis in patients with laparoscopic proximal gastrectomy
Wu CY, Lin JA, Ye K
- 124 Clinical effect of laparoscopic radical resection of colorectal cancer based on propensity score matching
Liu Y, Wang XX, Li YL, He WT, Li H, Chen H
- 134 Different timing for abdominal paracentesis catheter placement and drainage in severe acute pancreatitis complicated by intra-abdominal fluid accumulation
Chen R, Chen HQ, Li RD, Lu HM
- 143 Comparison of different preoperative objective nutritional indices for evaluating 30-d mortality and complications after liver transplantation
Li C, Chen HX, Lai YH
- 155 Predictive value of NLR, Fib4, and APRI in the occurrence of liver failure after hepatectomy in patients with hepatocellular carcinoma
Kuang TZ, Xiao M, Liu YF
- 166 Practical effect of different teaching modes in teaching gastrointestinal surgery nursing
Rong XJ, Ning Z
- Observational Study**
- 173 Predictive factors and model validation of post-colon polyp surgery *Helicobacter pylori* infection
Zhang ZS
- Randomized Controlled Trial**
- 186 Micro-power negative pressure wound technique reduces risk of incision infection following loop ileostomy closure
Xu DY, Bai BJ, Shan L, Wei HY, Lin DF, Wang Y, Wang D
- 196 Paravertebral block's effect on analgesia and inflammation in advanced gastric cancer patients undergoing transarterial chemoembolization and microwave ablation
Xiong YF, Wei BZ, Wang YF, Li XF, Liu C

META-ANALYSIS

- 205 Unraveling the efficacy network: A network meta-analysis of adjuvant external beam radiation therapy methods after hepatectomy
Yang GY, He ZW, Tang YC, Yuan F, Cao MB, Ren YP, Li YX, Su XR, Yao ZC, Deng MH
- 215 Estimation of Physiologic Ability and Surgical Stress scoring system for predicting complications following abdominal surgery: A meta-analysis spanning 2004 to 2022
Pang TS, Cao LP
- 228 Role of Oncostatin M in the prognosis of inflammatory bowel disease: A meta-analysis
Yang Y, Fu KZ, Pan G

CASE REPORT

- 239 Endoscopic treatment of extreme esophageal stenosis complicated with esophagotracheal fistula: A case report
Fang JH, Li WM, He CH, Wu JL, Guo Y, Lai ZC, Li GD
- 248 Intestinal tuberculosis with small bowel stricture and hemorrhage as the predominant manifestation: Three case reports
Huang G, Wu KK, Li XN, Kuai JH, Zhang AJ

LETTER TO THE EDITOR

- 257 Sarcopenia in cirrhotic patients: Does frailty matter while waiting for a liver transplant?
Li XJ, He K

ABOUT COVER

Editorial Board Member of *World Journal of Gastrointestinal Surgery*, Renato Pietroletti, PhD, Associate Professor, Professor, Department of Applied Clinical and Biotechnological Sciences, University of L'Aquila, L'Aquila 67100, AQ, Italy. renato.pietroletti@univaq.it

AIMS AND SCOPE

The primary aim of *World Journal of Gastrointestinal Surgery* (*WJGS, World J Gastrointest Surg*) is to provide scholars and readers from various fields of gastrointestinal surgery with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGS mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal surgery and covering a wide range of topics including biliary tract surgical procedures, biliopancreatic diversion, colectomy, esophagectomy, esophagostomy, pancreas transplantation, and pancreatectomy, *etc.*

INDEXING/ABSTRACTING

The *WJGS* is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports/Science Edition, PubMed, PubMed Central, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for *WJGS* as 2.0; IF without journal self cites: 1.9; 5-year IF: 2.2; Journal Citation Indicator: 0.52; Ranking: 113 among 212 journals in surgery; Quartile category: Q3; Ranking: 81 among 93 journals in gastroenterology and hepatology; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Zi-Hang Xu, Production Department Director: Xiang Li, Editorial Office Director: Jia-Ru Fan.

NAME OF JOURNAL

World Journal of Gastrointestinal Surgery

ISSN

ISSN 1948-9366 (online)

LAUNCH DATE

November 30, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Peter Schemmer

POLICY OF CO-AUTHORS**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/1948-9366/editorialboard.htm>

PUBLICATION DATE

January 27, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

<https://www.wjgnet.com/bpg/GerInfo/310>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Prospects in the application of ultrasensitive chromosomal aneuploidy detection in precancerous lesions of gastric cancer

Su-Ting Qian, Fei-Fei Xie, Hao-Yu Zhao, Qing-Sheng Liu, Dan-Li Cai

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Kotelevets SM, Russia

Received: November 18, 2023

Peer-review started: November 18, 2023

First decision: December 6, 2023

Revised: December 11, 2023

Accepted: December 26, 2023

Article in press: December 26, 2023

Published online: January 27, 2024



Su-Ting Qian, Fei-Fei Xie, Hao-Yu Zhao, Department of Digestive, Hangzhou Hospital of Traditional Chinese Medicine, Hangzhou 310007, Zhejiang Province, China

Qing-Sheng Liu, Science and Education Section, Hangzhou Hospital of Traditional Chinese Medicine, Hangzhou 310007, Zhejiang Province, China

Dan-Li Cai, Intensive Care Unit, The First Affiliated Hospital of Zhejiang Chinese Medical University, Hangzhou 311122, Zhejiang Province, China

Corresponding author: Dan-Li Cai, MM, Assistant Chief Physician, Intensive Care Unit, The First Affiliated Hospital of Zhejiang Chinese Medical University, No. 1 Gaojiao Road, Yuhang District, Hangzhou 311122, Zhejiang Province, China. 13858159603@139.com

Abstract

Gastric cancer (GC) is a prevalent malignant tumor within the digestive system, with over 40% of new cases and deaths related to GC globally occurring in China. Despite advancements in treatment modalities, such as surgery supplemented by adjuvant radiotherapy or chemotherapeutic agents, the prognosis for GC remains poor. New targeted therapies and immunotherapies are currently under investigation, but no significant breakthroughs have been achieved. Studies have indicated that GC is a heterogeneous disease, encompassing multiple subtypes with distinct biological characteristics and roles. Consequently, personalized treatment based on clinical features, pathologic typing, and molecular typing is crucial for the diagnosis and management of precancerous lesions of gastric cancer (PLGC). Current research has categorized GC into four subtypes: Epstein-Barr virus-positive, microsatellite instability, genome stability, and chromosome instability (CIN). Technologies such as multi-omics analysis and gene sequencing are being employed to identify more suitable novel testing methods in these areas. Among these, ultrasensitive chromosomal aneuploidy detection (UCAD) can detect CIN at a genome-wide level in subjects using low-depth whole genome sequencing technology, in conjunction with bioinformatics analysis, to achieve qualitative and quantitative detection of chromosomal stability. This editorial reviews recent research advancements in UCAD technology for the diagnosis and management of PLGC.

Key Words: Gastric cancer; Precancerous lesions of gastric cancer; Molecular typing; Ultrasensitive chromosomal aneuploidy detection; Adjuvant therapy; Application prospects

Core Tip: The purpose of this editorial is to provide an overview of the current diagnostic and therapeutic guidelines for gastric precancerous lesions, and to explore the potential clinical application of ultrasensitive chromosomal aneuploidy detection in the field of gastric cancer prevention and control. By doing so, this article aims to advance future research in this area.

Citation: Qian ST, Xie FF, Zhao HY, Liu QS, Cai DL. Prospects in the application of ultrasensitive chromosomal aneuploidy detection in precancerous lesions of gastric cancer. *World J Gastrointest Surg* 2024; 16(1): 6-12

URL: <https://www.wjgnet.com/1948-9366/full/v16/i1/6.htm>

DOI: <https://dx.doi.org/10.4240/wjgs.v16.i1.6>

INTRODUCTION

Gastric cancer (GC) is the third leading cause of cancer-related deaths globally, with the highest incidence particularly in East Asia, Central, and Eastern Europe[1]. The number of incidences of GC will likely increase in the future due to higher socioeconomic status and aging populations[2]. In recent times, the prevalence of pan-cancer screening and gastroscopy has led to early identification and treatment of a growing number of patients in the initial (gastric pre-cancer) phase. The use of endoscopic mucosal resection (EMR) and endoscopic submucosal dissection (ESD) has significantly enhanced patient survival. However, the majority of cases are still diagnosed at an advanced stage, resulting in the majority of patients succumbing to GC[3]. Despite the advancements in medical technology, a significant number of patients are reluctant to undergo endoscopy. Thus, researchers are still actively investigating the stages of precancerous lesions of gastric cancer (PLGC) and searching for more expedited and streamlined diagnostic approaches. These remain pressing issues in current PLGC research.

Due to the subtle symptoms of GC, the condition is typically discovered at an advanced stage in patients who seek medical attention, leading to higher mortality rates and a poorer prognosis. Advancements in next-generation sequencing and other genomic technologies have spurred new research into the molecular characteristics of GC[4]. Gene mutations, chromosomal aberrations, differential gene expression, and epigenetic alterations are some of the genetic and epigenetic factors that influence the pathogenesis of GC. Thus, The Cancer Genome Atlas (TCGA) network proposes a four-subtype classification scheme for GC based on the molecular biology of potential tumors for each subtype, including Epstein-Barr virus (EBV)-positive, microsatellite instability (MSI), genome stability (GS) type, and chromosome instability (CIN) type [5]. One of the most frequent genetic alterations is CIN, often referred to as chromosomal copy number aberration (CNA). The ultrasensitive chromosomal aneuploidy detection (UCAD) technology relies on low-coverage whole-genome sequencing (LC-WGS) to diagnose or predict a patient's tumor risk by identifying chromosomal CNA in a sample.

The purpose of this editorial is to offer an updated overview of the established diagnostic and treatment criteria for gastric precancerous lesions. It covers emerging diagnostic and treatment techniques, and explores the potential for deploying UCAD technology as an early diagnostic tool.

PLGC: EARLY PREVENTION AND CURRENT STATUS IN CLINICAL DIAGNOSIS

At this stage, the clinical consensus of PLGC can be categorized into atrophy (loss of gastric glands), intestinal metaplasia (replacement of gastric epithelium by intestinal epithelium), and dysplasia (intraepithelial neoplasia, including low-grade intraepithelial neoplasia and high-grade intraepithelial neoplasia)[6].

Helicobacter pylori (*H. pylori*) infection, the most common chronic bacterial infection in humans, is the strongest known risk factor for gastric carcinogenesis. Chronic and persistent inflammation resulting from *H. pylori* in the stomach lining may give rise to progressive atrophic gastritis and intestinal metaplasia[7].

H. pylori strains have a cytotoxin-associated gene A (CagA) that encodes a 120-140 kDa CagA protein, an oncoprotein that affects tumor cells. It also contains additional virulence factors, including vacuolating cytotoxin A, duodenal ulcer-promoting gene A protein, outer inflammatory protein A, and gamma-glutamyl transpeptidase. Most individuals infected with *H. pylori* are asymptomatic, yet have an increased risk of developing peptic ulcers or gastric adenocarcinoma[8,9].

In addition to *H. pylori*, age, tobacco and alcohol consumption, high salt intake and a diet low in fruits and vegetables, familial susceptibility, previous gastric surgery, and pernicious anemia are all relevant risk factors. It is clear that control of the above risk factors is an important basis for the prevention of PLGC[10].

Screening for cancer risk factors is a straightforward process; however, precisely identifying at-risk groups for early screening poses a challenge. To mitigate this, it is essential to implement effective strategies that optimize the selection of individuals who are most likely to benefit from early screening. The primary diagnostic method currently used is endoscopic biopsy, with the challenge being the identification of individuals at risk of developing PLGC in the absence of positive signs[11]. It is apparent that endoscopy, as an invasive medical examination, receives limited acceptance from

certain demographics who may also be restricted by underlying diseases and age. The development and implementation of novel early cancer diagnostic tools must adhere to the reliability, reproducibility, and cost control standards set by the World Health Organization. Furthermore, they must prioritize ease of use and patient comfort[12].

Available studies have shown a significant level of heterogeneity in the histopathology and molecular biology of GC. Screening for hallmark molecules of PLGC involves identifying genetic or protein markers that can be used for early diagnosis and prognostic determination. Most of the screened molecules such as oncogenes, intercellular adhesion molecules, growth factors, and certain hormone receptors exhibit deficiencies in sensitivity, specificity, and reliability, with only a few being recognized.

A paper published in Nature a few years ago by TCGA proposed dividing GC into four subtypes based on molecular characteristics[13].

EBV-positive phenotype is frequently associated with PIK3CA mutations, DNA hypermethylation, and amplification of JAK2, CD274, and PDCD1LG2. MSI is a type characterized by a high mutation rate, including mutations in genes that activate oncogene signaling pathways. The GS type, which occurs most often in the histologically diffuse form, is caused by mutations in RHOA or fusions of GTPase-activating protein genes in the THO family. CIN type, which has the hallmark heterozygous chromosomes and in situ amplification of the receptor tyrosine kinase.

The conventional method of devising treatment plans based on tumor phenotypic characteristics will be replaced by a mode that considers gene alterations instead[14]. Utilizing single gene expression changes can direct targeted medication therapy, which is a more rational, efficient and personalized mode of treatment compared to chemotherapy for the same morphology type. Additionally, multi-gene detection will become an essential research area[15].

GUIDELINES FOR THE TREATMENT OF PLGC: STANDARDS AND DEVELOPMENTS

Early GC without lymph node metastasis may be treated through endoscopic or surgical means, depending on the extent of tumor invasion. Adjuvant radiotherapy or chemotherapy is not necessary after surgery.

Locally advanced GC or early GC with lymph node metastasis should be managed using comprehensive surgical treatment. The decision of direct radical surgery *vs* preoperative neoadjuvant chemotherapy, before radical surgery, depends on the depth of tumor invasion and whether or not lymph node metastasis is present. For patients with locally advanced GC who have undergone successful radical surgery, adjuvant treatment planning (including adjuvant chemotherapy and, if necessary, adjuvant chemoradiotherapy) should be based on the postoperative pathological staging.

Recurrent or metastatic GC requires comprehensive treatment consisting mainly of medication. Palliative surgery, radiotherapy, interventional therapy, and local therapies should also be administered as needed, along with optimal supportive therapies, including analgesia, stenting, and nutrition support. It is essential to give supportive therapies actively at the appropriate time[16].

Endoscopic techniques

EMR is the endoscopic removal of mucosal lesions, in whole or in part, for the diagnosis and treatment of superficial tumors of the gastrointestinal tract[17].

ESD is a new technology developed on the basis of EMR. According to the lesions with different parts, sizes and infiltration depths, special electrocautery knives such as IT knives, Dual knives, Hook knives, *etc.* are selected for endoscopic gradual separation of tissues between the mucous membrane layer and the intrinsic muscular layer, and finally the lesion mucous membrane and submucosal layer are completely removed[18].

Other endoscopic treatments include laser therapy, argon knife, microwave therapy, *etc.* They can only remove the tumor, but cannot obtain complete pathological specimens or be sure that the tumor is completely removed. Therefore, they are mostly used for the treatment of precancerous gastric lesions and require close follow-up after treatment, and are not recommended as the first choice of treatment for early GC[19,20].

Surgery and chemotherapy

PLGC surgery is divided into radical surgery and non-radical surgery. Radical surgery should completely remove the primary lesion and thoroughly clear the regional lymph nodes, mainly including standard surgery, modified surgery and extended surgery; non-radical surgery mainly includes palliative surgery and tumor reduction surgery[21,22].

Chemotherapy is divided into palliative chemotherapy, adjuvant chemotherapy, neoadjuvant chemotherapy, and conversion therapy.

Neoadjuvant chemotherapy is recommended for locally advanced GC without distant metastases (T3/4, N+). The regimen should consist of a two-drug combination of platinum and fluorouracil or a three-drug combination based on the two-drug regimen combined with paclitaxel. It should not be used as a single agent[11].

The GC management process is shown in [Figure 1](#).

UCAD APPLICATION PROSPECTS

In the article "*Hallmarks of cancer: The next generation*" in *Cell*, the authors systematically summarized various hallmarks of cancer development. They highlighted genomic instability (or chromosomal instability) as the fundamental cause of cancer development, present in almost all malignant tumors. Thus, genomic instability is considered the root cause of




Figure 1 The gastric cancer management process. EMR: Endoscopic mucosal resection; ESD: Endoscopic submucosal dissection; GC: Gastric cancer; PLGC: Precancerous lesions of gastric cancer; UCAD: Ultrasensitive chromosomal aneuploidy detection.

cancer[23].

CIN is a feature of certain cells that results in the production of daughter cells that have an altered chromosome number or structure. This is a common feature in cancer cells. The fundamental aspects of chromosomal instability include changes in chromosome copy number, which leads to the emergence of cells with aneuploidy, as well as alterations in chromosome structure. These changes can include polyploidy, chromosomal translocations, genomic chaos, and non-clonal chromosomal aberrations[24].

The mechanism behind cancer cells is the irregular distribution of chromosomes during mitosis in zygote cells and the persistence of errors in chromosome segregation. This leads to modifications in chromosome copy number, as well as intrachromosomal segment amplification or deletion[25].

The substantial loss, gain, and rearrangement of DNA accumulate, resulting in chromosomal instability. This instability leads to extensive genomic complexity, which is a hallmark of cancer that is either present or expected. Chromosomal instability is strongly associated with tumor staging, metastasis, poor prognosis, and treatment resistance. This includes the loss or amplification of driver genes, focal rearrangements, extrachromosomal DNA, micronucleus formation, and the activation of innate immune signaling. These mechanisms lead to the diversification of tumor subclones in terms of space and time. This facilitates metastasis, accelerates the phenotypic adaptation of tumors, promotes cellular immortality, the escape from immune surveillance, and the development of resistance to drug treatments, among other processes[26].

CIN has been extensively studied as an innovative test. Using patient-derived tumor organoids to imitate colorectal cancer, researchers confirmed the existence of anaphase chromatin bridges in colorectal cancer cell lines. Causes consisted of insufficient disassembly of sister chromatids, telomere fusion producing bicentric chromosomes, and incomplete replication. These manifestations of CIN are prevalent in organs resembling colorectal cancer and have a significant impact on tumor evolution and therapeutic response. The levels of CIN and tolerance to mitotic errors play a crucial role in shaping the aneuploid landscapes and karyotypic heterogeneity[27].

Copy number and structural changes due to CIN are common features of multiple myeloma (MM). Primary and secondary genetic events caused by CIN lead to an increase in malignant plasma cell genomic instability by interfering with cell cycle checkpoints, thus accelerating proliferation. Thus, an assessment of CIN in MM and its precursor states may help to mitigate the progression of symptomatic disease and the risk of relapse[28].

CIN plays a role in the development of some subtypes of diffuse infiltrating gliomas. In isocitrate dehydrogenase-mutant astrocytomas, copy number variation (CNV) levels rise overall with tumor grade, duration, and proximity of infiltrating cells within the same tumor. The detection of CIN through CNV, DNA methylation, and/or gene expression profiling can effectively identify gliomas influenced by this molecular mechanism[29].

UCAD is a cutting-edge approach for identifying cellular chromosomal instability. This technique leverages body fluid free DNA or tissue genomic DNA as a template and utilizes LC-WGS, together with bioinformatics analysis, to identify chromosomal instability at the genome-wide scale. This technique has the potential to aid in the detection of tumors, monitor the effectiveness of treatment, and assess the likelihood of recurrence and metastasis. Samples including peripheral blood, exfoliated cells, menstrual blood, fresh and frozen tissues, formalin-fixed paraffin-embedded, urine, and others, can be analyzed for diagnostic purposes.

UCAD has been applied in clinical studies at this stage. Ye *et al*[30] collected samples of pancreatic cystic fluid from 102 patients with pancreatic cystic neoplasms (PCN) and employed UCAD to examine distinct CIN characteristics among different types of PCN. The deletion of chr3p and chr6p was used to define subtypes of serous cystadenoma. Meanwhile, the gain of chr1q and chr8q was associated with latent malignant PCN and facilitated the detection of high-risk intraductal papillary mucinous neoplasm.

Wang *et al*[31] collected plasma samples from 47 patients suspected of having lesions in the biliary tract. They utilized UCAD to analyze free DNA for CNV analysis through low-coverage whole genome sequencing. The results demonstrated the most frequent copy number gains in chr3q (7/29) and chr8q (6/29), with the most prevalent copy number losses noted in chr7p (6/29), chr17p (6/29), and chr19p (6/29). The sensitivity and specificity of the plasma CNV assay for the diagnosis of biliary tract cancer were 89.7% and 88.9%, respectively.

Feng *et al*[32] obtained 196 plasma samples from two groups of patients: a discovery cohort of individuals with PLC who were not eligible for surgery, and a validation cohort of patients who underwent pathologically confirmed hepatectomies. Of the 172 individuals, 22 (95.7%) surgically ineligible hepatocellular carcinomas were identified as having CNV in at least 1 of the 29 segments. 54 (69.4%) of the hepatocellular carcinomas eligible for surgery received positive screenings, and subsequently, confirmed to be cancerous by pathologic examination. Additionally, 26/27 non-cancers were identified with negative screenings.

Ye *et al*[33] gathered mucosal samples from 40 patients with GC from the hospital. 20 were microbiome-enriched, 5 tested positive for EBV DNA, and 15 had *H. pylori* DNA. Meanwhile, 20 of the samples were found to have CIN. UCAD can be used to identify 3 distinct subtypes of GC in the Chinese population, providing valuable guidance for future research on GC treatment and prevention. UCAD can identify three distinct subtypes of GC in the Chinese population. This finding could provide valuable guidance for further research on the prevention and treatment of GC.

CONCLUSION

In recent years, there has been an acceleration in precision treatment for PLGC, however, obstacles remain. The evolving endoscopic technology and surgical treatment have not sufficiently addressed the high recurrence rate after surgery. While there is a clinical benefit trend in anti-HER2 therapy and anti-CLDN18.2 monoclonal antibody therapy, additional research is necessary to optimize treatment and maximize patient benefit[34,35]. Early diagnostic treatment remains a pressing and essential need for improvement with a significant demand for less invasive or non-invasive testing and highly specific biomarkers[36]. Research efforts must focus on developing more convenient, comfortable, and highly accurate diagnostic tests, like tests on body fluids such as peripheral blood, urine or saliva, and gastric fluids, for early assessment of the likelihood of patient progression. Early treatment is critical for reducing the risk of developing GC.

In this context, UCAD has been researched and applied as a new test for early prevention in patients with positive early sequencing results or for gastroscopy and treatment to timely remove lesions at a stage when PLGC has not progressed. With the gradual decrease in the cost of second-generation sequencing, UCAD assays are expected to become more cost-effective in the near future. Meanwhile, the UCAD assay can capture both human and microbiome DNA while detecting specimens from mucosal, blood, and urine sources. This convenience, comfort, and information make it a superior technology for GC subtyping compared to other methods. Overall, this new approach may improve diagnostic and therapeutic strategies for PLGC in a cost-effective manner, achieving true precision.

ACKNOWLEDGEMENTS

We gratefully acknowledge the kind cooperation of all authors in the preparation of this paper.

FOOTNOTES

Author contributions: Cai DL designed the overall concept and outline of the manuscript; Qian ST contributed to the design and writing of the manuscript; Xie FF, Zhao HY, and Liu QS contributed to the editing the manuscript, illustrations, and review of literature.

Conflict-of-interest statement: All authors have no conflict-of-interest to disclose.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Su-Ting Qian 0009-0003-3308-7560; Fei-Fei Xie 0009-0005-1455-6215; Hao-Yu Zhao 0009-0005-0599-3851; Qing-Sheng Liu 0000-0003-1351-1443; Dan-Li Cai 0000-0003-1299-2273.

S-Editor: Zhang H

L-Editor: A

P-Editor: Zhang YL

REFERENCES

- 1 Röcken C. Predictive biomarkers in gastric cancer. *J Cancer Res Clin Oncol* 2023; **149**: 467-481 [PMID: 36260159 DOI: 10.1007/s00432-022-04408-0]
- 2 Smyth EC, Nilsson M, Grabsch HI, van Grieken NC, Lordick F. Gastric cancer. *Lancet* 2020; **396**: 635-648 [PMID: 32861308 DOI: 10.1016/S0140-6736(20)31288-5]
- 3 Pimentel-Nunes P, Libânio D, Marcos-Pinto R, Areia M, Leja M, Esposito G, Garrido M, Kikuste I, Megraud F, Matysiak-Budnik T, Annibale B, Dumonceau JM, Barros R, Fléjou JF, Carneiro F, van Hooft JE, Kuipers EJ, Dinis-Ribeiro M. Management of epithelial precancerous conditions and lesions in the stomach (MAPS II): European Society of Gastrointestinal Endoscopy (ESGE), European Helicobacter and Microbiota Study Group (EHMSG), European Society of Pathology (ESP), and Sociedade Portuguesa de Endoscopia Digestiva (SPED) guideline update 2019. *Endoscopy* 2019; **51**: 365-388 [PMID: 30841008 DOI: 10.1055/a-0859-1883]
- 4 Zeng Y, Jin RU. Molecular pathogenesis, targeted therapies, and future perspectives for gastric cancer. *Semin Cancer Biol* 2022; **86**: 566-582 [PMID: 34933124 DOI: 10.1016/j.semcancer.2021.12.004]
- 5 Yuen ST, Leung SY. Genomics Study of Gastric Cancer and Its Molecular Subtypes. *Adv Exp Med Biol* 2016; **908**: 419-439 [PMID: 27573784 DOI: 10.1007/978-3-319-41388-4_21]
- 6 Gullo I, Grillo F, Mastracci L, Vanoli A, Carneiro F, Saragoni L, Limarzi F, Ferro J, Parente P, Fassan M. Precancerous lesions of the stomach, gastric cancer and hereditary gastric cancer syndromes. *Pathologica* 2020; **112**: 166-185 [PMID: 33179620 DOI: 10.32074/1591-951X-166]
- 7 Toyoshima O, Nishizawa T, Koike K. Endoscopic Kyoto classification of Helicobacter pylori infection and gastric cancer risk diagnosis. *World J Gastroenterol* 2020; **26**: 466-477 [PMID: 32089624 DOI: 10.3748/wjg.v26.i5.466]
- 8 de Brito BB, da Silva FAF, Soares AS, Pereira VA, Santos MLC, Sampaio MM, Neves PHM, de Melo FF. Pathogenesis and clinical management of Helicobacter pylori gastric infection. *World J Gastroenterol* 2019; **25**: 5578-5589 [PMID: 31602159 DOI: 10.3748/wjg.v25.i37.5578]
- 9 FitzGerald R, Smith SM. An Overview of Helicobacter pylori Infection. *Methods Mol Biol* 2021; **2283**: 1-14 [PMID: 33765303 DOI: 10.1007/978-1-0716-1302-3_1]
- 10 Eusebi LH, Telese A, Marasco G, Bazzoli F, Zagari RM. Gastric cancer prevention strategies: A global perspective. *J Gastroenterol Hepatol* 2020; **35**: 1495-1502 [PMID: 32181516 DOI: 10.1111/jgh.15037]
- 11 Hoshi H. Management of Gastric Adenocarcinoma for General Surgeons. *Surg Clin North Am* 2020; **100**: 523-534 [PMID: 32402298 DOI: 10.1016/j.suc.2020.02.004]
- 12 Hu Y, Lv X, Wei W, Li X, Zhang K, Zhu L, Gan T, Zeng H, Yang J, Rao N. Quantitative Analysis on Molecular Characteristics Evolution of Gastric Cancer Progression and Prognosis. *Adv Biol (Weinh)* 2023; **7**: e2300129 [PMID: 37357148 DOI: 10.1002/adbi.202300129]
- 13 Cancer Genome Atlas Research Network. Comprehensive molecular characterization of gastric adenocarcinoma. *Nature* 2014; **513**: 202-209 [PMID: 25079317 DOI: 10.1038/nature13480]
- 14 Wadowska K, Bil-Lula I, Trembecki L, Śliwińska-Mossoń M. Genetic Markers in Lung Cancer Diagnosis: A Review. *Int J Mol Sci* 2020; **21** [PMID: 32604993 DOI: 10.3390/ijms21134569]
- 15 Bartlett J, Amemiya Y, Arts H, Bayani J, Eng B, Grafodatskaya D, Kamel Reid S, Lariviere M, Lo B, McClure R, Mittal V, Sadikovic B, Sadis S, Seth A, Smith J, Zhang X, Feilletter H. Multisite verification of the accuracy of a multi-gene next generation sequencing panel for detection of mutations and copy number alterations in solid tumours. *PLoS One* 2021; **16**: e0258188 [PMID: 34597339 DOI: 10.1371/journal.pone.0258188]
- 16 Ajani JA, D'Amico TA, Brentem DJ, Chao J, Cooke D, Corvera C, Das P, Enzinger PC, Enzler T, Fanta P, Farjah F, Gerdes H, Gibson MK, Hochwald S, Hofstetter WL, Ilson DH, Keswani RN, Kim S, Kleinberg LR, Klemperer SJ, Lacy J, Ly QP, Matkowskyj KA, McNamara M, Mulcahy MF, Outlaw D, Park H, Perry KA, Pimiento J, Poultsides GA, Reznik S, Roses RE, Strong VE, Su S, Wang HL, Wiesner G, Willett CG, Yakoub D, Yoon H, McMillian N, Pluchino LA. Gastric Cancer, Version 2.2022, NCCN Clinical Practice Guidelines in Oncology. *J Natl Compr Canc Netw* 2022; **20**: 167-192 [PMID: 35130500 DOI: 10.6004/jncn.2022.0008]
- 17 Ahmed Y, Othman M. EMR/ESD: Techniques, Complications, and Evidence. *Curr Gastroenterol Rep* 2020; **22**: 39 [PMID: 32542462 DOI: 10.1007/s11894-020-00777-z]
- 18 Liu Q, Ding L, Qiu X, Meng F. Updated evaluation of endoscopic submucosal dissection versus surgery for early gastric cancer: A systematic review and meta-analysis. *Int J Surg* 2020; **73**: 28-41 [PMID: 31783166 DOI: 10.1016/j.ijssu.2019.11.027]
- 19 Niknam N, Obanor S, Lee LA. Endoscopic methods for the detection and treatment of gastric cancer. *Curr Opin Gastroenterol* 2022; **38**: 436-442 [PMID: 35881962 DOI: 10.1097/MOG.0000000000000867]
- 20 Young E, Philpott H, Singh R. Endoscopic diagnosis and treatment of gastric dysplasia and early cancer: Current evidence and what the future may hold. *World J Gastroenterol* 2021; **27**: 5126-5151 [PMID: 34497440 DOI: 10.3748/wjg.v27.i31.5126]
- 21 Johnston FM, Beckman M. Updates on Management of Gastric Cancer. *Curr Oncol Rep* 2019; **21**: 67 [PMID: 31236716 DOI: 10.1007/s11912-019-0820-4]
- 22 Wang Y, Zhang L, Yang Y, Lu S, Chen H. Progress of Gastric Cancer Surgery in the era of Precision Medicine. *Int J Biol Sci* 2021; **17**: 1041-1049 [PMID: 33867827 DOI: 10.7150/ijbs.56735]
- 23 Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. *Cell* 2011; **144**: 646-674 [PMID: 21376230 DOI: 10.1016/j.cell.2011.02.013]
- 24 Kuang X, Li J. Chromosome instability and aneuploidy as context-dependent activators or inhibitors of antitumor immunity. *Front Immunol* 2022; **13**: 895961 [PMID: 36003402 DOI: 10.3389/fimmu.2022.895961]
- 25 Bach DH, Zhang W, Sood AK. Chromosomal Instability in Tumor Initiation and Development. *Cancer Res* 2019; **79**: 3995-4002 [PMID: 31350294 DOI: 10.1158/0008-5472.CAN-18-3235]
- 26 Drews RM, Hernando B, Tarabichi M, Haase K, Lesluyes T, Smith PS, Morrill Gavarró L, Couturier DL, Liu L, Schneider M, Brenton JD, Van Loo P, Macintyre G, Markowitz F. A pan-cancer compendium of chromosomal instability. *Nature* 2022; **606**: 976-983 [PMID: 35705807]

- DOI: [10.1038/s41586-022-04789-9](https://doi.org/10.1038/s41586-022-04789-9)]
- 27 **Bolhaqueiro ACF**, Ponsioen B, Bakker B, Klaasen SJ, Kucukkose E, van Jaarsveld RH, Vivié J, Verlaan-Klink I, Hami N, Spierings DCJ, Sasaki N, Dutta D, Boj SF, Vries RGJ, Lansdorp PM, van de Wetering M, van Oudenaarden A, Clevers H, Kranenburg O, Foijer F, Snippert HJG, Kops GJPL. Ongoing chromosomal instability and karyotype evolution in human colorectal cancer organoids. *Nat Genet* 2019; **51**: 824-834 [PMID: [31036964](https://pubmed.ncbi.nlm.nih.gov/31036964/) DOI: [10.1038/s41588-019-0399-6](https://doi.org/10.1038/s41588-019-0399-6)]
 - 28 **Neuse CJ**, Lomas OC, Schliemann C, Shen YJ, Manier S, Bustoros M, Ghobrial IM. Genome instability in multiple myeloma. *Leukemia* 2020; **34**: 2887-2897 [PMID: [32651540](https://pubmed.ncbi.nlm.nih.gov/32651540/) DOI: [10.1038/s41375-020-0921-y](https://doi.org/10.1038/s41375-020-0921-y)]
 - 29 **Richardson TE**, Walker JM, Abdullah KG, McBrayer SK, Viapiano MS, Mussa ZM, Tsankova NM, Snuderl M, Hatanpaa KJ. Chromosomal instability in adult-type diffuse gliomas. *Acta Neuropathol Commun* 2022; **10**: 115 [PMID: [35978439](https://pubmed.ncbi.nlm.nih.gov/35978439/) DOI: [10.1186/s40478-022-01420-w](https://doi.org/10.1186/s40478-022-01420-w)]
 - 30 **Ye M**, Zhang B, Han X, Wei X, Wang Y, Cao W, Wu J, Chen C, Sun X, Sun K, Li H, Zhang Q, Liang T. Low-Pass Genomic Sequencing Reveals Novel Subtypes of Pancreatic Cystic Neoplasms. *Ann Surg Oncol* 2023; **30**: 5804-5812 [PMID: [37249723](https://pubmed.ncbi.nlm.nih.gov/37249723/) DOI: [10.1245/s10434-023-13676-0](https://doi.org/10.1245/s10434-023-13676-0)]
 - 31 **Wang X**, Fu XH, Qian ZL, Zhao T, Duan AQ, Ruan X, Zhu B, Yin L, Zhang YJ, Yu WL. Non-invasive detection of biliary tract cancer by low-coverage whole genome sequencing from plasma cell-free DNA: A prospective cohort study. *Transl Oncol* 2021; **14**: 100908 [PMID: [33059123](https://pubmed.ncbi.nlm.nih.gov/33059123/) DOI: [10.1016/j.tranon.2020.100908](https://doi.org/10.1016/j.tranon.2020.100908)]
 - 32 **Feng S**, Ding Z, Wang J, Qian Z, Li S, Zhang C, Xin H, Liu S, Ding G, Hu M, Meng Y, Li N. Investigation of Plasma cell-free cancer genome chromosomal instability as a tool for targeted minimally invasive biomarkers for primary liver cancer diagnoses. *Cancer Med* 2020; **9**: 5075-5085 [PMID: [32458568](https://pubmed.ncbi.nlm.nih.gov/32458568/) DOI: [10.1002/cam4.3142](https://doi.org/10.1002/cam4.3142)]
 - 33 **Ye LP**, Mao XL, Zhou XB, Wang Y, Xu SW, He SQ, Qian ZL, Zhang XG, Zhai LJ, Peng JB, Gu BB, Jin XX, Song YQ, Li SW. Cost-effective low-coverage whole-genome sequencing assay for the risk stratification of gastric cancer. *World J Gastrointest Oncol* 2022; **14**: 690-702 [PMID: [35321281](https://pubmed.ncbi.nlm.nih.gov/35321281/) DOI: [10.4251/wjgo.v14.i3.690](https://doi.org/10.4251/wjgo.v14.i3.690)]
 - 34 **Janjigian YY**, Kawazoe A, Yañez P, Li N, Lonardi S, Kolesnik O, Barajas O, Bai Y, Shen L, Tang Y, Wyrwicz LS, Xu J, Shitara K, Qin S, Van Cutsem E, Tabernero J, Li L, Shah S, Bhagia P, Chung HC. The KEYNOTE-811 trial of dual PD-1 and HER2 blockade in HER2-positive gastric cancer. *Nature* 2021; **600**: 727-730 [PMID: [34912120](https://pubmed.ncbi.nlm.nih.gov/34912120/) DOI: [10.1038/s41586-021-04161-3](https://doi.org/10.1038/s41586-021-04161-3)]
 - 35 **Zhong W**, Lu Y, Ma Z, He Y, Ding Y, Yao G, Zhou Z, Dong J, Fang Y, Jiang W, Wang W, Huang Y. Development of a Humanized VHH Based Recombinant Antibody Targeting Claudin 18.2 Positive Cancers. *Front Immunol* 2022; **13**: 885424 [PMID: [35837391](https://pubmed.ncbi.nlm.nih.gov/35837391/) DOI: [10.3389/fimmu.2022.885424](https://doi.org/10.3389/fimmu.2022.885424)]
 - 36 **Necula L**, Matei L, Dragu D, Neagu AI, Mambet C, Nedeianu S, Bleotu C, Diaconu CC, Chivu-Economescu M. Recent advances in gastric cancer early diagnosis. *World J Gastroenterol* 2019; **25**: 2029-2044 [PMID: [31114131](https://pubmed.ncbi.nlm.nih.gov/31114131/) DOI: [10.3748/wjg.v25.i17.2029](https://doi.org/10.3748/wjg.v25.i17.2029)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: office@baishideng.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

