# World Journal of Gastrointestinal Surgery

World J Gastrointest Surg 2024 March 27; 16(3): 635-973





Published by Baishideng Publishing Group Inc

GS WŮ

## World Journal of Gastrointestinal Surgery

### Contents

### Monthly Volume 16 Number 3 March 27, 2024

### **EDITORIAL**

635 Ex vivo liver resection and auto-transplantation and special systemic therapy in perihilar cholangiocarcinoma treatment

Tchilikidi KY

641 Indocyanine green: The guide to safer and more effective surgery Fransvea P, Chiarello MM, Fico V, Cariati M, Brisinda G

### **MINIREVIEWS**

- 650 Alcohol associated liver disease and bariatric surgery: Current perspectives and future directions Cooper KM, Colletta A, Hebda N, Devuni D
- 658 Applications of gastric peroral endoscopic myotomy in the treatment of upper gastrointestinal tract disease

Chang SY, Jin GH, Sun HB, Yang D, Tang TY

### **ORIGINAL ARTICLE**

### **Retrospective Cohort Study**

670 Evaluation of bacterial contamination and medium-term oncological outcomes of intracorporeal anastomosis for colon cancer: A propensity score matching analysis

Kayano H, Mamuro N, Kamei Y, Ogimi T, Miyakita H, Nakagohri T, Koyanagi K, Mori M, Yamamoto S

681 Rescue from complications after pancreaticoduodenectomies at a low-volume Caribbean center: Value of tailored peri-pancreatectomy protocols

Cawich SO, Dixon E, Shukla PJ, Shrikhande SV, Deshpande RR, Mohammed F, Pearce NW, Francis W, Johnson S, Bujhawan J

689 Comparison of prognosis and postoperative morbidities between standard pancreaticoduodenectomy and the TRIANGLE technique for resectable pancreatic ductal adenocarcinoma

Hang HX, Cai ZH, Yang YF, Fu X, Qiu YD, Cheng H

700 Analysis of the impact of immunotherapy efficacy and safety in patients with gastric cancer and liver metastasis

Liu K, Wu CX, Liang H, Wang T, Zhang JY, Wang XT

### **Retrospective Study**

710 Clinical observation of extraction-site incisional hernia after laparoscopic colorectal surgery Fan BH, Zhong KL, Zhu LJ, Chen Z, Li F, Wu WF

717 Predicting short-term major postoperative complications in intestinal resection for Crohn's disease: A machine learning-based study

Wang FT, Lin Y, Yuan XQ, Gao RY, Wu XC, Xu WW, Wu TQ, Xia K, Jiao YR, Yin L, Chen CQ



| Contor | World Journal of Gastrointestinal Surgery  |
|--------|--|
| Conter | Monthly Volume 16 Number 3 March 27, 2024  |
| 731    | Analysis of factors impacting postoperative pain and quality of life in patients with mixed hemorrhoids: A retrospective study   |
|        | Sun XW, Xu JY, Zhu CZ, Li SJ, Jin LJ, Zhu ZD   |
| 740    | Pre-operative visceral adipose tissue radiodensity is a potentially novel prognostic biomarker for early endoscopic post-operative recurrence in Crohn's disease   |
|        | Gu P, Dube S, Gellada N, Choi SY, Win S, Lee YJ, Yang S, Haritunians T, Melmed GY, Vasiliauskas EA, Bonthala N, Syal<br>G, Yarur AJ, Ziring D, Rabizadeh S, Fleshner P, Kallman C, Devkota S, Targan SR, Li D, McGovern DP |
| 751    | Clinical study on the relationship between liver cirrhosis, ascites, and hyponatremia  |
|        | Li XJ, Meng HH   |
| 759    | Comparison of the clinical effects of dual-modality endoscopy and traditional laparotomy for the treatment of intra- and extrahepatic bile duct stones   |
|        | Wang W, Xia H, Dai B   |
| 768    | Role of ablation therapy in conjunction with surgical resection for neuroendocrine tumors involving the liver  |
|        | Ostapenko A, Stroever S, Eyasu L, Kim M, Aploks K, Dong XD, Seshadri R   |
| 777    | Feasibility and safety of minimally invasive multivisceral resection for T4b rectal cancer: A 9-year review  |
|        | Chan KS, Liu B, Tan MNA, How KY, Wong KY   |
| 790    | MH-STRALP: A scoring system for prognostication in patients with upper gastrointestinal bleeding   |
|        | Hu JN, Xu F, Hao YR, Sun CY, Wu KM, Lin Y, Zhong L, Zeng X   |
|        | Clinical Trials Study  |
| 807    | Early postoperative complications after transverse colostomy closure, a retrospective study  |
|        | Liu F, Luo XJ, Li ZW, Liu XY, Liu XR, Lv Q, Shu XP, Zhang W, Peng D  |
| 816    | Clinical study of enhanced recovery after surgery in laparoscopic appendectomy for acute appendicitis  |
|        | Li ZL, Ma HC, Yang Y, Chen JJ, Wang ZJ   |
|        | Observational Study  |
| 823    | Reinforced tissue matrix to strengthen the abdominal wall following reversal of temporary ostomies or to treat incisional hernias  |
|        | Lake SP, Deeken CR, Agarwal AK   |
|        | Randomized Controlled Trial  |
| 833    | Whole-process case management effects on mental state and self-care ability in patients with liver cancer  |
|        | Ju MD, Qin Q, Li M   |
|        | Clinical and Translational Research  |
| 842    | Construction and validation of somatic mutation-derived long non-coding RNAs signatures of genomic instability to predict prognosis of hepatocellular carcinoma  |
|        | Duan BT, Zhao XK, Cui YY, Liu DZ, Wang L, Zhou L, Zhang XY   |



Contents

World Journal of Gastrointestinal Surgery

### Monthly Volume 16 Number 3 March 27, 2024

### **Basic Study**

- 860 Influence of different magnetic forces on the effect of colonic anastomosis in rats Tian BY, Zhang MM, Ma J, Lyu Y, Yan XP
- 871 Inflammatory responses in esophageal mucosa before and after laparoscopic antireflux surgery Ergun P, Kipcak S, Selvi Gunel N, Yildirim Sozmen E, Bor S
- 882 Etanercept-synthesizing adipose-derived stem cell secretome: A promising therapeutic option for inflammatory bowel disease

Kim SJ, Kim OH, Hong HE, Ju JH, Lee DS

### SYSTEMATIC REVIEWS

893 Impact of frailty on short-term postoperative outcomes in patients undergoing colorectal cancer surgery: A systematic review and meta-analysis

Zhou Y, Zhang XL, Ni HX, Shao TJ, Wang P

### **META-ANALYSIS**

907 Endoscopic-ultrasound-guided biliary drainage with placement of electrocautery-enhanced lumenapposing metal stent for palliation of malignant biliary obstruction: Updated meta-analysis

Peng ZX, Chen FF, Tang W, Zeng X, Du HJ, Pi RX, Liu HM, Lu XX

921 Clinical efficacy and safety of erlotinib combined with chemotherapy in the treatment of advanced pancreatic cancer: A meta-analysis

Liu XY. Pan HN. Yu Y

### **CASE REPORT**

932 Link between mutations in ACVRL1 and PLA2G4A genes and chronic intestinal ulcers: A case report and review of literature

Tang YJ, Zhang J, Wang J, Tian RD, Zhong WW, Yao BS, Hou BY, Chen YH, He W, He YH

- 944 Mucinous neoplasm of the appendix: A case report and review of literature Chang HC, Kang JC, Pu TW, Su RY, Chen CY, Hu JM
- 955 Abdominal cocoon syndrome-a rare culprit behind small bowel ischemia and obstruction: Three case reports

Vipudhamorn W, Juthasilaparut T, Sutharat P, Sanmee S, Supatrakul E

966 Endoscopic ultrasound-guided lauromacrogol injection for treatment of colorectal cavernous hemangioma: Two case reports

Zhu HT, Chen WG, Wang JJ, Guo JN, Zhang FM, Xu GQ, Chen HT



### Contents

World Journal of Gastrointestinal Surgery

Monthly Volume 16 Number 3 March 27, 2024

### **ABOUT COVER**

Editorial Board Member of World Journal of Gastrointestinal Surgery, Jia-Gang Han, MD, Professor, Department of General Surgery, Beijing Chaoyang Hospital, Capital Medical University, Beijing 100020, China. hjg211@163.com

### **AIMS AND SCOPE**

The primary aim of World Journal of Gastrointestinal Surgery (WJGS, World J Gastrointest Surg) is to provide scholars and readers from various fields of gastrointestinal surgery with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGS mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal surgery and covering a wide range of topics including biliary tract surgical procedures, biliopancreatic diversion, colectomy, esophagectomy, esophagostomy, pancreas transplantation, and pancreatectomy, etc.

### **INDEXING/ABSTRACTING**

The WJGS is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports/Science Edition, PubMed, PubMed Central, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJGS as 2.0; IF without journal self cites: 1.9; 5-year IF: 2.2; Journal Citation Indicator: 0.52; Ranking: 113 among 212 journals in surgery; Quartile category: Q3; Ranking: 81 among 93 journals in gastroenterology and hepatology; and Quartile category: Q4.

### **RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: Zi-Hang Xu; Production Department Director: Xiang Li; Editorial Office Director: Jia-Ru Fan.

| NAME OF JOURNAL                                     | INSTRUCTIONS TO AUTHORS                       |
|---|---|
| World Journal of Gastrointestinal Surgery           | https://www.wjgnet.com/bpg/gerinfo/204        |
| <b>ISSN</b>   | GUIDELINES FOR ETHICS DOCUMENTS               |
| ISSN 1948-9366 (online)                             | https://www.wjgnet.com/bpg/GerInfo/287        |
| LAUNCH DATE   | GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH |
| November 30, 2009                                   | https://www.wjgnet.com/bpg/gerinfo/240        |
| FREQUENCY   | PUBLICATION ETHICS                            |
| Monthly   | https://www.wjgnet.com/bpg/GerInfo/288        |
| EDITORS-IN-CHIEF                                    | PUBLICATION MISCONDUCT                        |
| Peter Schemmer                                      | https://www.wjgnet.com/bpg/gerinfo/208        |
| EDITORIAL BOARD MEMBERS                             | ARTICLE PROCESSING CHARGE                     |
| https://www.wjgnet.com/1948-9366/editorialboard.htm | https://www.wjgnet.com/bpg/gerinfo/242        |
| PUBLICATION DATE                                    | STEPS FOR SUBMITTING MANUSCRIPTS              |
| March 27, 2024                                      | https://www.wjgnet.com/bpg/GerInfo/239        |
| COPYRIGHT   | ONLINE SUBMISSION                             |
| © 2024 Baishideng Publishing Group Inc              | https://www.f6publishing.com                  |

© 2024 Baishideng Publishing Group Inc. All rights reserved. 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA E-mail: office@baishideng.com https://www.wjgnet.com



S WU

### World Journal of Gastrointestinal Surgery

Submit a Manuscript: https://www.f6publishing.com

World J Gastrointest Surg 2024 March 27; 16(3): 650-657

DOI: 10.4240/wjgs.v16.i3.650

ISSN 1948-9366 (online)

MINIREVIEWS

### Alcohol associated liver disease and bariatric surgery: Current perspectives and future directions

Katherine M Cooper, Alessandro Colletta, Nicholas Hebda, Deepika Devuni

Specialty type: Transplantation

Provenance and peer review: Invited article; Externally peer reviewed.

Peer-review model: Single blind

### Peer-review report's scientific quality classification

Grade A (Excellent): 0 Grade B (Very good): 0 Grade C (Good): C Grade D (Fair): 0 Grade E (Poor): 0

P-Reviewer: Skok P, Slovenia

Received: November 23, 2023 Peer-review started: November 23, 2023

First decision: January 24, 2024 Revised: January 27, 2024 Accepted: February 26, 2024 Article in press: February 26, 2024 Published online: March 27, 2024



Katherine M Cooper, Alessandro Colletta, Nicholas Hebda, Deepika Devuni, Department of Medicine, UMass Chan Medical School, Worcester, MA 01655, United States

Deepika Devuni, Division of Gastroenterology, UMass Chan Medical School, Worcester, MA 01655, United States

Corresponding author: Katherine M Cooper, MD, Doctor, Department of Medicine, UMass Chan Medical School, 55 Lake Ave North, Worcester, MA 01655, United States. katherine.cooper@umassmed.edu

### Abstract

Bariatric surgery is a routinely performed procedure and is associated with a reduction in all-cause mortality in patients with obesity. However, bariatric surgery has also been linked to increased alcohol use with up to 30% of these patients developing alcohol use disorder (AUD). The mechanism of AUD after bariatric surgery is multifactorial and includes anatomic, metabolic, and neurohumoral changes associated with post-surgical anatomy. These patients are at increased risk of alcohol associated liver disease and, in some cases, require liver transplantation. In this article, we provide a scoping review of epidemiology, pathophysiology, and clinical outcomes of alcohol-related health conditions after bariatric surgery.

Key Words: Obesity medicine; Alcohol use disorder; Roux-en-Y gastric bypass; Vertical sleeve gastrectomy; Liver transplant; Simultaneous liver transplant and bariatric surgery

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** Bariatric surgery improves obesity related health co-morbidities but also increases the risk of developing alcohol related health problems. Up to 33% of bariatric surgery patients develop an alcohol use disorder (AUD) with peak onset 2 years after surgery and up to 5% develop alcohol associated liver disease (ALD) and cirrhosis. Further research is needed to develop clinical recommendations for AUD screening and to define risk factors for chronic ALD after bariatric surgery.

WJGS https://www.wjgnet.com

Citation: Cooper KM, Colletta A, Hebda N, Devuni D. Alcohol associated liver disease and bariatric surgery: Current perspectives and future directions. World J Gastrointest Surg 2024; 16(3): 650-657 URL: https://www.wjgnet.com/1948-9366/full/v16/i3/650.htm DOI: https://dx.doi.org/10.4240/wjgs.v16.i3.650

### INTRODUCTION

Over 500000 individuals undergo bariatric surgery annually worldwide[1]. Although bariatric surgery reduces the development of obesity-related diseases and improves all-cause mortality[2], it may also increase the risk of harmful drinking[3] and alter the liver's susceptibility to injury in the setting of alcohol use[4]. In recent years, the practice of simultaneous bariatric surgery and liver transplantation has gained favor in patients with metabolic dysfunctionassociated steatotic liver disease (MASLD) due to the potential improved peri-operative outcomes[5]. Thus, understanding alcohol-related disorders after bariatric surgery will be increasingly important in the clinical setting. In this article, we provide a scoping review of epidemiology, pathophysiology, and clinical outcomes of alcohol use disorder (AUD) and alcohol associated liver disease (ALD) after bariatric surgery and discuss opportunities to enhance postoperative care for these patients.

### EPIDEMIOLOGY OF ALCOHOL USE AFTER BARIATRIC SURGERY

AUD is characterized by a pattern of alcohol consumption that persists despite negative personal and health consequences[6]. The prevalence of alcohol use has been steadily increasing with some data suggesting over 40% of adults engage in potentially harmful drinking[6]. An expanding body of literature suggests that bariatric surgery increases the risk of alcohol misuse. Estimates suggest that between 2%-33% of adult bariatric surgery patients develop AUD, with the peak incidence occurring in the second post-operative year [3,4,7,8]. This risk may be higher in adolescent bariatric surgery recipients with a recent multi-center study reporting an AUD prevalence of approximately 45% eight years postsurgery[9].

Multiple studies have identified the following as risk factors for harmful alcohol use after bariatric surgery: younger age, tobacco use, pre-surgical alcohol use, and reduced social support[8,10,11]. Additionally, one large prospective study identified higher household income and lower-level education as risk factors for AUD[7]. Patient sex influences risk as well: female sex is a risk factor for AUD in adolescent bariatric surgery recipients<sup>[12]</sup>, while male sex is a risk factor in adult bariatric surgery recipients [10]. Although adult males have a higher risk of AUD, most bariatric surgery recipients with AUD are female, reflecting the overall higher proportion of females in the bariatric surgery population[3]. The risk of post-surgical AUD may also depend on the type of surgery performed. Some studies indicate a higher risk following gastric bypass procedures as compared to restrictive procedures[11,13], although this relationship is not yet fully elucidated[7,11,13].

Although pre-surgical alcohol use is associated with AUD following bariatric surgery, new onset alcohol use is also common[7,14-16]. In a recently published study, Kim et al[14] utilized a large commercial insurance database to assess alcohol use in patients who underwent bariatric surgery compared to patients who underwent cholecystectomy. They found that 2.8% of the post-bariatric surgery group developed de novo alcohol use or dependence compared to 1.7% of the post-cholecystectomy group. On subgroup analyses, Roux-en-Y gastric bypass (RYGB) was associated with a 150% higher risk of acquiring an alcohol-associated diagnosis (e.g., alcohol use, alcohol use, and alcohol-associated hepatitis) compared to post-cholecystectomy controls (95% CI 1.40-1.62, P < 0.001). In contrast, vertical sleeve gastrectomy (VSG) was associated with a slightly reduced risk of acquiring an alcohol-associated diagnosis. Notably, since this study was conducted using International Classification of Diseases, Tenth Revision (ICD-10) codes, the context of these diagnoses was not available. A multi-center retrospective study within the United States Veterans Affairs Healthcare System used the Alcohol Use Disorder Identification Test (AUDIT-C) to assess for harmful alcohol use among bariatric surgery patients (n = 1539 VSG, n = 854 RYGB) compared to non-surgical controls[15]. Among patients without pre-surgical alcohol use, new unhealthy alcohol use was significantly more common in the surgical groups at post operative years 3, 5, and 8. At the end of the study period, the risk of unhealthy alcohol use was 7.9% in VSG patients, 9.2% in RYGB patients, and 4.4%-4.5% in non-surgical control patients. Similarly, a recent prospective study observed a quadratic relationship between alcohol use and time after bariatric surgery, with annual increases in both quantity and frequency of alcohol consumption from years 1-7 after bariatric surgery[16]. This discrepancy in findings may be due to differing follow-up periods, as Kim et al[14] followed patients for a mean of 2.7 years whereas the latter two studies had an 8 year follow up and noted a significant increase in harmful alcohol use between 3-8 years post-surgery.

### PATHOPHYSIOLOGY OF ALCOHOL USE DISORDER AFTER BARIATRIC SURGERY

The mechanisms by which bariatric surgery increases the risk of AUD have been studied but are not fully understood. Several factors are likely to contribute, including the metabolic, anatomic, and downstream neurologic changes resulting from the surgery[3]. A basic understanding of each surgical procedure is important when considering post-operative



WJGS | https://www.wjgnet.com

alcohol use. Briefly, in RYGB, the stomach is divided into a small gastric pouch (using the proximal stomach) and a larger distal remnant. The gastric pouch is then connected directly to the mid-jejunum via the Roux - or alimentary - limb, bypassing the distal stomach and duodenum[17]. As a result, nutrients quickly pass from the stomach to the distal small intestine. Vertical sleeve gastrectomy involves stapling the stomach along a vertical line from the pylorus to the angle of His and removing the occluded portion which includes the fundus, body, and part of the antrum[18]. This results in a narrow tube-like stomach that is about 70%-80% smaller in size (Figure 1).

### Pharmacokinetics

The anatomic changes from both RYGB and VSG result in faster gastric emptying and reduced first-past metabolism [3,19, 20]. As a result, blood alcohol concentration peaks faster and at higher concentrations after any given ingestion of alcohol. For instance, the oral bioavailability of alcohol has been shown to increase by more than 30% in women who undergo VSG[21]. Likewise, a study found that women with a history of RYGB experience more brisk and larger amplitude changes in blood alcohol concentration. Specifically, after consuming 26 g of alcohol (equivalent to about two shots of hard liquor), blood alcohol concentration increased to a level greater than the United States legal driving limit in less than 10 min[22].

In addition to objective measurements, patients report subjectively increased response to alcohol. A qualitative study evaluating the addictive properties of alcohol following bariatric surgery reported that patients experienced the feeling of intoxication faster than they did prior to surgery [23]. Other studies have used patient-facing questionnaires including the Addiction Research Center Inventory and the Alcohol Sensitivity Questionnaire to quantify subjective response to alcohol. In these studies, patients reported increased feelings of drunkenness and higher sensitivity to alcohol after bariatric surgery [24,25]. Because fast onset drugs are known to have higher addictive potential [26], this brisk increase in response to alcohol has been thought to play a major role in the development of post-bariatric surgery AUD. It was thought that reduced first-pass metabolism after bariatric surgery was due a relative reduction in gastric alcohol dehydrogenase activity. However, studies evaluating alcohol dehydrogenase have been inconsistent, and currently there is insufficient evidence to support that reduced alcohol dehydrogenase activity is the primary factor behind the rapid increases in blood alcohol concentration observed after bariatric surgery [22,27,28].

### Hormonal and reward pathways

Metabolic changes influencing alcohol use after bariatric surgery arise from alterations in the secretion of multiple gut hormones that occur after both RYGB and VSG[3,19,29]. Two key hormones linking alcohol use and bariatric surgery are ghrelin and glucagon-like peptide-1 (GLP-1).

Ghrelin: The role of ghrelin in the development of AUD after bariatric surgery has been explored extensively. Early theories proposed that changes in ghrelin levels directly influence alcohol intake after RYGB. However, it is now thought that hormonal changes affect reward-seeking pathways in the brain[30-32]. Pre-clinical models have demonstrated that RYGB surgery is associated with a heightened response to alcohol in central reward centers[30,31] and that this heightened response can be attenuated by antagonizing the ghrelin receptor: growth hormone secretagogue receptor (GHSR) 1a[31]. In fact, a recent study found that GHSR activity regulates the tonic rate of neuronal firing in a key reward center in the brain, the ventral tegmental area (VTA). In this study, researchers treated mice with a GHSR-antagonist, and observed a reduction in the action potential firing rate of VTA dopaminergic neurons in non-RYGB mice but observed no effect in mice with RYGB[32]. They concluded that alcohol misuse after RYGB may be related to reduced GHSR control of the VTA dopaminergic system. Similar animal models have been developed for VSG[33,34]. In contrast to RYGB models, mice with VSG demonstrated reduced alcohol seeking behavior compared to control mice. This finding was initially attributed to the removal of the ghrelin producing part of the stomach, but this was refuted when administration of ghrelin at different concentrations failed to impact ethanol consumption. Interestingly, administration of low dose GHSR antagonist induced a significant reduction in alcohol consumption in the VSG mice but not in control mice[35]. Collectively these data demonstrate that GHSR signaling is linked to changes in brain reward pathways including enhanced dopamine sensitivity and increased reward response to alcohol after bariatric surgery. Differences observed between preclinical models of VSG and RYGB may help explain differences in the prevalence of AUD in these groups.

GLP-1: GLP-1 is a gut-brain peptide that regulates appetite and food intake, and it has been increasingly recognized for its role in alcohol-mediated behaviors. For instance, animal studies have shown that administration of liraglutide, a GLP-1 agonist, attenuates the rewarding properties of alcohol[36,37]. Specifically, liraglutide has been shown to impair the release of dopamine in the nucleus accumbens of mice after acute injections of ethanol. After treatment with liraglutide, mice were less likely to demonstrate preference for areas where ethanol was in their environment. Data from these preclinical models suggest that GLP-1 receptor agonists may attenuate the ability of alcohol to activate the mesolimbic dopamine system and may reduce alcohol intake. Interestingly, a nation-wide cohort study in Denmark found that GLP-1 agonists were associated with a lower incidence of alcohol-related health events, though this protective effect was limited to the first 3 months of therapy [38]. While these findings highlight the potential clinical relevance of GLP-1 pathways in treating AUD, the use of GLP-1 receptor agonists in reducing alcohol intake following weight reduction surgery has not been studied.

### Psychology and addiction

When discussing potential mechanisms of increased alcohol use after bariatric surgery, it is worth mentioning the "addiction transfer" hypothesis. This theory assumes that patients pursuing bariatric surgery have addictive eating habits that are transferred to alcohol consumption post-operatively when food intake is limited by appetite[39]. However, there





Figure 1 Mechanisms of alcohol related liver disease after bariatric surgery. Several anatomic, metabolic, and neurohumoral changes occur after bariatric surgery. A: Changes in the gastric anatomy affect the metabolism and pharmacokinetics properties of alcohol and lead to enhanced alcohol sensitivity and reduced tolerance; B: These alterations are linked to changes in brain reward pathways that can increase the risk of developing alcohol use disorders; C: Further, reduced first pass metabolism results in increased delivery of alcohol to the liver; D: Together, these may result in earlier onset alcohol associated liver injury and chronic alcohol associated liver disease. MASLD: Metabolic dysfunction-associated steatotic liver disease; BAC: Blood alcohol concentration; GHSR, or ghrelin receptor: Growth hormone secretagogue receptor; RYGB: Roux-en-Y gastric bypass. Image created using Biorender.com.

are several lines of evidence which demonstrate that addiction transfer alone is insufficient to explain observed trends in post-bariatric surgery alcohol use.

First, studies have shown that pre-operative addictive eating behaviors and binge eating do not increase the likelihood of developing AUD[40,41] which directly opposes the idea that post-surgical AUD originates from food-related addictive behaviors. Furthermore, increased alcohol use has been observed in patients following prophylactic total gastrectomy, even in the absence of obesity or history of addictive eating behaviors[42,43]. Second, the tendency for alcohol use to increase one to two years after bariatric surgery contradicts the addiction transfer theory as the reduction in food intake occurs almost immediately post-operatively[44]. Third, the incidence of AUD after bariatric surgery varies depending on the type of surgery, suggesting that changes inherent to each surgical procedure play a role in the development of AUD [7].

Additionally, the anatomic changes resulting from these surgeries alter the metabolism and pharmacokinetics of alcohol[45]. Post-surgery weight loss combined with reduced gastric volume and accelerated gastric emptying results in higher blood alcohol concentrations and leads to increased alcohol sensitivity and lower tolerance[46]. These changes are likely critical in the development of AUD after weight loss surgeries. While the addiction transfer hypothesis may explain a behavioral component in a minority of patients, its general application can increase stigma and discourage care seeking amongst the majority[45]. Therefore, healthcare providers should be cautious when referencing this hypothesis to avoid perpetuating stigma in a disorder that has a clear biological basis.

### ALCOHOL-RELATED OUTCOMES AFTER BARIATRIC SURGERY

Beyond AUD, bariatric surgery is associated with an increased risk of developing ALD. There are likely several reasons for this relationship. Patients undergoing bariatric surgery likely have some baseline hepatic disease, commonly in the form of steatotic liver disease[47]. Obesity and alcohol consumption have a synergistic effect on the liver; patients with a BMI over 35 are more likely to develop liver damage from any given amount of alcohol compared to patients with a BMI under 35[48,49]. While bariatric surgery can reduce hepatic fibrosis and/or eliminate steatohepatitis in 70%-80% of patients achieving a minimum 5-10-kilogram weight loss[50], it remains unclear if the liver continues to be predisposed to alcohol-associated damage, particularly when exposed to relatively higher concentrations of alcohol after surgery.

The exact prevalence of ALD after bariatric surgery is unknown. Two recent studies estimate the incidence of ALD to be 0.7% in bariatric surgery recipients[4] compared to 0.3% in the general population[51,52]. Mellinger *et al*[4] compared the risk of alcohol-associated cirrhosis (AC) in obese patients who underwent bariatric surgery and those who did not in

two time periods: before 2008 and from 2008 to 2016. While initial unadjusted analyses suggested no increased risk of AC in the bariatric surgery group, further analyses revealed a significant interaction between the time period and patient gender. Adjusted analyses demonstrated that women who underwent bariatric surgery had a 2.1-times greater risk of AC (95%CI 1.8-2.4) and men had a 1.3-times greater risk of AC (95%CI 1.1-1.6) compared to their counterparts who did not undergo the surgery. In another study, Alvarado *et al*[52] assessed the risk of ALD following bariatric surgery recipients were more likely to develop both ALD (95%CI 1.22-1.37) and AC (95%CI 1.37-1.42). Additionally, they observed a significant increase in ALD and AC cases among bariatric surgery patients from 2005 to 2015, which aligns with Mellinger *et al*'s findings about the long-term risk of AC[4].

Bariatric surgery status has also been linked to poorer outcomes in patients with ALD. In a study of 2,634 patients with alcohol-associated hepatitis, those with prior RYGB had a higher 30-day readmission rate (P < 0.01) and increased overall mortality (P = 0.03) compared to non-surgical patients[53]. Conversely, an analysis of the National Inpatient Sample found that prior RYGB was associated with an increased risk of hepatic encephalopathy and infection but was not associated with higher mortality among patients hospitalized for ALD[54]. It is possible that the higher rate of adverse outcomes, despite a relatively younger patient population, is due to the delivery of higher concentrations of alcohol to the liver for any given ingestion of alcohol. Ultimately, the mortality benefits of bariatric surgery are attenuated by the development of alcohol-associated co-morbidities in patients who develop AUD[55].

### BARIATRIC SURGERY AND LIVER TRANSPLANTATION

In line with the trends for ALD, there has been an increase in liver transplants for AC among post-bariatric surgery patients. Patients with a history of bariatric surgery that progress to needing a liver transplant in the setting of ALD are typically younger, female, and are more likely to be diagnosed with mood disorders compared to those without a history of bariatric surgery[56]. This cohort tends to present with more severe hepatic decompensation and progresses to liver transplant significantly faster than those who have not undergone bariatric surgery[57]. Post-liver transplant outcomes in this cohort have not been extensively studied, though a small study found that post-transplant biliary complications and rejection rates were similar in patients with and without a history of RYGB.

Over the last decade, there has been a trend towards performing simultaneous liver transplant and bariatric surgery (usually VSG) in eligible patients. Although limited, a few small studies reporting patient outcomes have emerged in recent years[58-62]. Generally, simultaneous bariatric surgery has not shown to increase liver transplant related mortality. Two prospective studies showed that patients with simultaneous surgeries had lower degrees of hepatic steatosis and fewer weight-related co-morbidities three years post-surgery compared to patients who had liver transplant alone[60,61]. It is important to note that post-surgical alcohol use is not addressed in these reports. Therefore, it remains unclear whether the risk of AUD in this cohort is similar to that in the general population. For those who do develop AUD, the risk of recurrent ALD is likely very different. While the pre-surgical fibrosis is obviated by transplantation, post-transplant patients are at risk of complications like strictures, chronic cholestasis, and rejection. These complications may interact with alcohol and result in liver dysfunction. To-date, no studies have addressed these clinical questions or assessed the risk of AUD or ALD in patients who have undergone simultaneous liver transplant-bariatric surgery.

### FUTURE DIRECTIONS: CLINICAL CARE FOR AUD AFTER BARIATRIC SURGERY

Currently, there are no screening guidelines for AUD following bariatric surgery. Providers should counsel patients that there is no known "safe" level of alcohol intake and that standard recommendations for alcohol consumption (less than 7 drinks per week for women, less than 14 drinks per week for men) generally do not apply after the procedure. When concerns arise, screening questionnaires used in the general population, such as the AUDIT-C, can be used to identify potentially harmful alcohol use. The AUDIT-C is a three-question survey that can help categorize alcohol use behaviors over the past 12 months[63]. The AUDIT-C has proven effective in detecting AUD and predicting long-term outcomes, including hospitalizations for alcohol-related diagnoses. A 2019 study by White *et al*[16] assessed the use of the AUDIT-C in women with a history of bariatric surgery and found that a score  $\geq$  3 had the highest combined sensitivity (76.4%) and specificity (81.6%) for identifying alcohol-related health problems. Using validated questionnaires may be more helpful than serum markers of alcohol consumption (*e.g.*, phosphatidylethanol), which may vary unpredictably in patients with a history of bariatric surgery[64].

While early screening is important for early diagnosis, bariatric surgery centers may also consider more intensive evaluation later, particularly in the second post-operative year[7]. Notably, patients undergoing simultaneous liver transplant and bariatric surgery may experience compound risk for AUD when considering the potential for psychological distress after liver transplantation. Therefore, these patients should receive counseling on risk reduction strategies and undergo similar, if not more stringent, post-operative monitoring for alcohol use compared to individuals with other liver disease etiologies. In patients who develop liver disease, identifying the primary etiology may be challenging for clinicians. While non-invasive markers exist to identify non-alcohol associated liver diseases, they often lack specificity. In patients with suspected alcohol use, liver biopsy may be required to assess the relative contribution of alcohol to liver damage[65-67].

Zaishidena® WJGS | https://www.wjgnet.com

### CONCLUSION

Patients who have undergone bariatric surgery represent a high-risk cohort for alcohol-related health conditions including alcohol-associated liver disease. More information is needed to develop clinical recommendations for AUD and ALD in this cohort, though it is essential to advise patients that there is no "safe" level of alcohol consumption after bariatric surgery. Future studies should focus on identifying risk factors for hepatic fibrosis and ALD development, evaluating the effectiveness of medication-associated therapies for AUD in bariatric surgery recipients, and assessing alcohol-related outcomes in patients who undergo simultaneous liver transplant-bariatric surgery.

### FOOTNOTES

Author contributions: Cooper KM conceptualized this review article; Cooper KM, Colletta A, Hebda N, and Devuni D wrote the original draft, reviewed and edited the final draft and gave final approval; Devuni D provided supervision for the writing of this manuscript.

Conflict-of-interest statement: Deepika Devuni has received grant funding from Sequana Medical for a clinical trial unrelated to the present work and grant funding from the NIAAA for research unrelated to this work. All other authors have no conflicts of interest to report.

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: https://creativecommons.org/Licenses/by-nc/4.0/

Country/Territory of origin: United States

**ORCID** number: Katherine M Cooper 0000-0002-6030-4773; Alessandro Colletta 0000-0002-3865-1015; Deepika Devuni 0000-0002-1011-9414.

S-Editor: Gong ZM L-Editor: A P-Editor: Xu ZH

### REFERENCES

- 1 Angrisani L, Santonicola A, Iovino P, Vitiello A, Zundel N, Buchwald H, Scopinaro N. Bariatric Surgery and Endoluminal Procedures: IFSO Worldwide Survey 2014. Obes Surg 2017; 27: 2279-2289 [PMID: 28405878 DOI: 10.1007/s11695-017-2666-x]
- 2 Wiggins T, Guidozzi N, Welbourn R, Ahmed AR, Markar SR. Association of bariatric surgery with all-cause mortality and incidence of obesity-related disease at a population level: A systematic review and meta-analysis. PLoS Med 2020; 17: e1003206 [PMID: 32722673 DOI: 10.1371/journal.pmed.1003206]
- Ivezaj V, Benoit SC, Davis J, Engel S, Lloret-Linares C, Mitchell JE, Pepino MY, Rogers AM, Steffen K, Sogg S. Changes in Alcohol Use 3 after Metabolic and Bariatric Surgery: Predictors and Mechanisms. Curr Psychiatry Rep 2019; 21: 85 [PMID: 31410716 DOI: 10.1007/s11920-019-1070-8]
- Mellinger JL, Shedden K, Winder GS, Fernandez AC, Lee BP, Waljee J, Fontana R, Volk ML, Blow FC, Lok ASF. Bariatric surgery and the 4 risk of alcohol-related cirrhosis and alcohol misuse. Liver Int 2021; 41: 1012-1019 [PMID: 33529460 DOI: 10.1111/liv.14805]
- Chierici A, Alromayan M, DEFATICO S, Céline D, Vinci D, Rodolphe A, Schiavo L, Iannelli A. Is bariatric surgery safer before, during, or 5 after liver transplantation? A systematic review and meta-analysis. J Liver Transpl 2023; 9: 100139 [DOI: 10.1016/j.liver.2023.100139]
- Nehring SM, Chen RJ, Freeman AM. Alcohol Use Disorder. StatPearls. Treasure Island (FL): StatPearls Publishing Copyright © 2023, 6 StatPearls Publishing LLC., 2023
- Ibrahim N, Alameddine M, Brennan J, Sessine M, Holliday C, Ghaferi AA. New onset alcohol use disorder following bariatric surgery. Surg 7 Endosc 2019; 33: 2521-2530 [PMID: 30350107 DOI: 10.1007/s00464-018-6545-x]
- King WC, Chen JY, Courcoulas AP, Dakin GF, Engel SG, Flum DR, Hinojosa MW, Kalarchian MA, Mattar SG, Mitchell JE, Pomp A, Pories 8 WJ, Steffen KJ, White GE, Wolfe BM, Yanovski SZ. Alcohol and other substance use after bariatric surgery: prospective evidence from a U.S. multicenter cohort study. Surg Obes Relat Dis 2017; 13: 1392-1402 [PMID: 28528115 DOI: 10.1016/j.soard.2017.03.021]
- 9 White GE, Boles RE, Courcoulas AP, Yanovski SZ, Zeller MH, Jenkins TM, Inge TH. A Prospective Cohort of Alcohol-Use and Alcoholrelated Problems Before and After Metabolic and Bariatric Surgery in Adolescents. Ann Surg 2023; 278: e519-e525 [PMID: 36538630 DOI: 10.1097/SLA.00000000005759
- King WC, Chen JY, Mitchell JE, Kalarchian MA, Steffen KJ, Engel SG, Courcoulas AP, Pories WJ, Yanovski SZ. Prevalence of alcohol use 10 disorders before and after bariatric surgery. JAMA 2012; 307: 2516-2525 [PMID: 22710289 DOI: 10.1001/jama.2012.6147]
- Svensson PA, Anveden Å, Romeo S, Peltonen M, Ahlin S, Burza MA, Carlsson B, Jacobson P, Lindroos AK, Lönroth H, Maglio C, Näslund I, 11 Sjöholm K, Wedel H, Söderpalm B, Sjöström L, Carlsson LM. Alcohol consumption and alcohol problems after bariatric surgery in the Swedish obese subjects study. Obesity (Silver Spring) 2013; 21: 2444-2451 [PMID: 23520203 DOI: 10.1002/oby.20397]
- Zeller MH, Washington GA, Mitchell JE, Sarwer DB, Reiter-Purtill J, Jenkins TM, Courcoulas AP, Peugh JL, Michalsky MP, Inge TH; Teen-12 LABS Consortium and in collaboration with the TeenView Study Group. Alcohol use risk in adolescents 2 years after bariatric surgery. Surg Obes Relat Dis 2017; 13: 85-94 [PMID: 27567561 DOI: 10.1016/j.soard.2016.05.019]
- Conason A, Teixeira J, Hsu CH, Puma L, Knafo D, Geliebter A. Substance use following bariatric weight loss surgery. JAMA Surg 2013; 148: 13



WJGS | https://www.wjgnet.com

145-150 [PMID: 23560285 DOI: 10.1001/2013.jamasurg.265]

- Kim HP, Jiang Y, Farrell TM, Peat CM, Hayashi PH, Barritt AS 4th. Roux-en-Y Gastric Bypass Is Associated With Increased Hazard for De 14 Novo Alcohol-related Complications and Liver Disease. J Clin Gastroenterol 2022; 56: 181-185 [PMID: 33780222 DOI: 10.1097/MCG.000000000001506]
- Maciejewski ML, Smith VA, Berkowitz TSZ, Arterburn DE, Mitchell JE, Olsen MK, Liu CF, Livingston EH, Funk LM, Adeyemo A, Bradley 15 KA. Association of Bariatric Surgical Procedures With Changes in Unhealthy Alcohol Use Among US Veterans. JAMA Netw Open 2020; 3: e2028117 [PMID: 33346846 DOI: 10.1001/jamanetworkopen.2020.28117]
- White GE, Courcoulas AP, Richardson GA, Mair C, King WC. Alcohol Use Thresholds for Identifying Alcohol-related Problems Before and 16 Following Roux-en-Y Gastric Bypass. Ann Surg 2019; 269: 1001-1009 [PMID: 31082893 DOI: 10.1097/SLA.000000000003078]
- Pucci A, Batterham RL. Mechanisms underlying the weight loss effects of RYGB and SG: similar, yet different. J Endocrinol Invest 2019; 42: 17 117-128 [PMID: 29730732 DOI: 10.1007/s40618-018-0892-2]
- 18 Seeras K, Sankararaman S, Lopez PP. Sleeve Gastrectomy. StatPearls: StatPearls Publishing Copyright © 2023, StatPearls Publishing LLC., 2018
- Csendes A, Braghetto I. Changes in the anatomy and physiology of the distal esophagus and stomach after sleeve gastrectomy. J Obes Weight 19 Loss Ther 2016; 6: 297 [DOI: 10.4172/2165-7904.1000297]
- Oneta CM, Simanowski UA, Martinez M, Allali-Hassani A, Parés X, Homann N, Conradt C, Waldherr R, Fiehn W, Coutelle C, Seitz HK. 20 First pass metabolism of ethanol is strikingly influenced by the speed of gastric emptying. Gut 1998; 43: 612-619 [PMID: 9824340 DOI: 10.1136/gut.43.5.612
- Seyedsadjadi N, Acevedo MB, Alfaro R, Ramchandani VA, Plawecki MH, Rowitz B, Pepino MY. Site of Alcohol First-Pass Metabolism 21 Among Women. JAMA Netw Open 2022; 5: e223711 [PMID: 35315921 DOI: 10.1001/jamanetworkopen.2022.3711]
- Steffen KJ, Engel SG, Pollert GA, Li C, Mitchell JE. Blood alcohol concentrations rise rapidly and dramatically after Roux-en-Y gastric 22 bypass. Surg Obes Relat Dis 2013; 9: 470-473 [PMID: 23507629 DOI: 10.1016/j.soard.2013.02.002]
- Yoder R, MacNeela P, Conway R, Heary C. How Do Individuals Develop Alcohol Use Disorder After Bariatric Surgery? A Grounded Theory 23 Exploration. Obes Surg 2018; 28: 717-724 [PMID: 29032488 DOI: 10.1007/s11695-017-2936-7]
- 24 Acevedo MB, Eagon JC, Bartholow BD, Klein S, Bucholz KK, Pepino MY. Sleeve gastrectomy surgery: when 2 alcoholic drinks are converted to 4. Surg Obes Relat Dis 2018; 14: 277-283 [PMID: 29305304 DOI: 10.1016/j.soard.2017.11.010]
- Acevedo MB, Teran-Garcia M, Bucholz KK, Eagon JC, Bartholow BD, Burd NA, Khan N, Rowitz B, Pepino MY. Alcohol sensitivity in 25 women after undergoing bariatric surgery: a cross-sectional study. Surg Obes Relat Dis 2020; 16: 536-544 [PMID: 32075778 DOI: 10.1016/j.soard.2020.01.014]
- Allain F, Minogianis EA, Roberts DC, Samaha AN. How fast and how often: The pharmacokinetics of drug use are decisive in addiction. 26 Neurosci Biobehav Rev 2015; 56: 166-179 [PMID: 26116543 DOI: 10.1016/j.neubiorev.2015.06.012]
- Parikh M, Johnson JM, Ballem N; American Society for Metabolic and Bariatric Surgery Clinical Issues Committee. ASMBS position 27 statement on alcohol use before and after bariatric surgery. Surg Obes Relat Dis 2016; 12: 225-230 [PMID: 26968500 DOI: 10.1016/j.soard.2015.10.085]
- Changchien EM, Woodard GA, Hernandez-Boussard T, Morton JM. Normal alcohol metabolism after gastric banding and sleeve 28 gastrectomy: a case-cross-over trial. J Am Coll Surg 2012; 215: 475-479 [PMID: 22770864 DOI: 10.1016/j.jamcollsurg.2012.06.008]
- Akkary E, Duffy A, Bell R. Deciphering the sleeve: technique, indications, efficacy, and safety of sleeve gastrectomy. Obes Surg 2008; 18: 29 1323-1329 [PMID: 18535867 DOI: 10.1007/s11695-008-9551-6]
- Thanos PK, Subrize M, Delis F, Cooney RN, Culnan D, Sun M, Wang GJ, Volkow ND, Hajnal A. Gastric bypass increases ethanol and water 30 consumption in diet-induced obese rats. Obes Surg 2012; 22: 1884-1892 [PMID: 22976430 DOI: 10.1007/s11695-012-0749-2]
- Hajnal A, Zharikov A, Polston JE, Fields MR, Tomasko J, Rogers AM, Volkow ND, Thanos PK. Alcohol reward is increased after Roux-en-Y 31 gastric bypass in dietary obese rats with differential effects following ghrelin antagonism. PLoS One 2012; 7: e49121 [PMID: 23145091 DOI: 10.1371/journal.pone.0049121]
- Sirohi S, Richardson BD, Lugo JM, Rossi DJ, Davis JF. Impact of Roux-en-Y gastric bypass surgery on appetite, alcohol intake behaviors, and 32 midbrain ghrelin signaling in the rat. Obesity (Silver Spring) 2017; 25: 1228-1236 [PMID: 28500684 DOI: 10.1002/oby.21839]
- 33 Coluzzi I, Iossa A, Spinetti E, Silecchia G. Alcohol consumption after laparoscopic sleeve gastrectomy: 1-year results. Eat Weight Disord 2019; 24: 1131-1136 [PMID: 29411323 DOI: 10.1007/s40519-018-0486-1]
- Orellana E, Horvath N, Farokhnia M, Leggio L, Hajnal A. Changes in plasma ghrelin levels following surgical and non-surgical weight-loss in 34 female rats predict alcohol use. Brain Res Bull 2022; 188: 179-186 [PMID: 35901985 DOI: 10.1016/j.brainresbull.2022.07.019]
- Orellana ER, Jamis C, Horvath N, Hajnal A. Effect of vertical sleeve gastrectomy on alcohol consumption and preferences in dietary obese 35 rats and mice: A plausible role for altered ghrelin signaling. Brain Res Bull 2018; 138: 26-36 [PMID: 28802901 DOI: 10.1016/j.brainresbull.2017.08.004]
- Vallöf D, Maccioni P, Colombo G, Mandrapa M, Jörnulf JW, Egecioglu E, Engel JA, Jerlhag E. The glucagon-like peptide 1 receptor agonist 36 liraglutide attenuates the reinforcing properties of alcohol in rodents. Addict Biol 2016; 21: 422-437 [PMID: 26303264 DOI: 10.1111/adb.12295]
- Jerlhag E. Alcohol-mediated behaviours and the gut-brain axis; with focus on glucagon-like peptide-1. Brain Res 2020; 1727: 146562 [PMID: 37 31759971 DOI: 10.1016/j.brainres.2019.146562]
- Wium-Andersen IK, Wium-Andersen MK, Fink-Jensen A, Rungby J, Jørgensen MB, Osler M. Use of GLP-1 receptor agonists and 38 subsequent risk of alcohol-related events. A nationwide register-based cohort and self-controlled case series study. Basic Clin Pharmacol Toxicol 2022; 131: 372-379 [PMID: 35968738 DOI: 10.1111/bcpt.13776]
- 39 McFadden KM. Cross-addiction: from morbid obesity to substance abuse. Bariat Nurs Surg Pat 2010; 5: 145-178 [DOI: 10.1089/bar.2010.9922
- Wong E, Fleishman A, Brem A, Jones DB, Wee CC. High-Risk Alcohol Use and Disordered Eating Behavior Before and 1 Year After Sleeve 40 Gastrectomy. Obes Surg 2022; 32: 593-598 [PMID: 35088252 DOI: 10.1007/s11695-021-05847-3]
- Dickhut C, Hase C, Gruner-Labitzke K, Mall JW, Köhler H, de Zwaan M, Müller A. No addiction transfer from preoperative food addiction to 41 other addictive behaviors during the first year after bariatric surgery. Eur Eat Disord Rev 2021; 29: 924-936 [PMID: 34460134 DOI: 10.1002/erv.2857]
- Yokoyama A, Takagi T, Ishii H, Wada N, Maruyama K, Takagi S, Hayashida M. Gastrectomy enhances vulnerability to the development of 42 alcoholism. *Alcohol* 1995; **12**: 213-216 [PMID: 7639953 DOI: 10.1016/0741-8329(94)00096-V]



- 43 Ito S, Izumi T, Arakawa M. Gastrectomy is a risk factor for alcoholism. *Intern Med* 1999; **38**: 751 [PMID: 10480310 DOI: 10.2169/internalmedicine.38.751]
- 44 Azam H, Shahrestani S, Phan K. Alcohol use disorders before and after bariatric surgery: a systematic review and meta-analysis. *Ann Transl* Med 2018; 6: 148 [PMID: 29862237 DOI: 10.21037/atm.2018.03.16]
- 45 Engel SG, Schaefer LM, Kerver GA, Leone LM, Smith G, Mitchell JE, Gunstad J, Crosby RD, Steffen K. The rewarding effects of alcohol after bariatric surgery: do they change and are they associated with pharmacokinetic changes? *Surg Obes Relat Dis* 2022; 18: 190-195 [PMID: 34583891 DOI: 10.1016/j.soard.2021.08.011]
- 46 Blackburn AN, Hajnal A, Leggio L. The gut in the brain: the effects of bariatric surgery on alcohol consumption. Addict Biol 2017; 22: 1540-1553 [PMID: 27578259 DOI: 10.1111/adb.12436]
- 47 Newsome PN, Cramb R, Davison SM, Dillon JF, Foulerton M, Godfrey EM, Hall R, Harrower U, Hudson M, Langford A, Mackie A, Mitchell-Thain R, Sennett K, Sheron NC, Verne J, Walmsley M, Yeoman A. Guidelines on the management of abnormal liver blood tests. *Gut* 2018; 67: 6-19 [PMID: 29122851 DOI: 10.1136/gutjnl-2017-314924]
- 48 Åberg F, Helenius-Hietala J, Puukka P, Färkkilä M, Jula A. Interaction between alcohol consumption and metabolic syndrome in predicting severe liver disease in the general population. *Hepatology* 2018; 67: 2141-2149 [PMID: 29164643 DOI: 10.1002/hep.29631]
- 49 Ventura-Cots M, Watts AE, Bataller R. Binge drinking as a risk factor for advanced alcoholic liver disease. *Liver Int* 2017; 37: 1281-1283 [PMID: 28845617 DOI: 10.1111/liv.13482]
- 50 Lassailly G, Caiazzo R, Ntandja-Wandji LC, Gnemmi V, Baud G, Verkindt H, Ningarhari M, Louvet A, Leteurtre E, Raverdy V, Dharancy S, Pattou F, Mathurin P. Bariatric Surgery Provides Long-term Resolution of Nonalcoholic Steatohepatitis and Regression of Fibrosis. *Gastroenterology* 2020; 159: 1290-1301.e5 [PMID: 32553765 DOI: 10.1053/j.gastro.2020.06.006]
- 51 Han S, Yang Z, Zhang T, Ma J, Chandler K, Liangpunsakul S. Epidemiology of Alcohol-Associated Liver Disease. *Clin Liver Dis* 2021; 25: 483-492 [PMID: 34229835 DOI: 10.1016/j.cld.2021.03.009]
- 52 Alvarado-Tapias E, Marti-Aguado D, Kennedy K, Fernández-Carrillo C, Ventura-Cots M, Morales-Arraez D, Atkinson SR, Clemente-Sanchez A, Argemi J, Bataller R. Bariatric Surgery Is Associated with Alcohol-Related Liver Disease and Psychiatric Disorders Associated with AUD. Obes Surg 2023; 33: 1494-1505 [PMID: 36881347 DOI: 10.1007/s11695-023-06490-w]
- 53 Anugwom C, Thomson M, Freese RL, Lake JR, Lim N. Lower survival and higher rates of cirrhosis in patients with ROUX-EN-Y gastric bypass hospitalised with alcohol-associated hepatitis. *BMJ Open Gastroenterol* 2023; 10 [PMID: 36808070 DOI: 10.1136/bmjgast-2022-001083]
- 54 Yarra P, Dunn W, Younossi Z, Kuo YF, Singal AK. Association of Previous Gastric Bypass Surgery and Patient Outcomes in Alcohol-Associated Cirrhosis Hospitalizations. *Dig Dis Sci* 2023; **68**: 1026-1034 [PMID: 35788931 DOI: 10.1007/s10620-022-07591-9]
- 55 Mahmud N, Panchal S, Abu-Gazala S, Serper M, Lewis JD, Kaplan DE. Association Between Bariatric Surgery and Alcohol Use-Related Hospitalization and All-Cause Mortality in a Veterans Affairs Cohort. JAMA Surg 2023; 158: 162-171 [PMID: 36515960 DOI: 10.1001/jamasurg.2022.6410]
- 56 Fipps DC, Goetze RE, Clark MM, Mara K, Watt KD, Jowsey-Gregoire SG, Heimbach JK, Grothe K. Liver Transplantation After Bariatric Surgery: a Clinical Cohort Study. Obes Surg 2021; 31: 3700-3706 [PMID: 34021883 DOI: 10.1007/s11695-021-05483-x]
- 57 Lefere S, Stroobant L, Verhelst X, Vanlander A, Berrevoet F, Troisi RI, Van Nieuwenhove Y, Rogiers X, Van Vlierberghe H, Geerts A. Bariatric Surgery Patients Are at Risk for Alcoholic Liver Disease with Need for Liver Transplantation. *Obes Surg* 2020; **30**: 4659-4664 [PMID: 32594466 DOI: 10.1007/s11695-020-04806-8]
- 58 Wijarnpreecha K, Pungpapong S, Aby ES, Croome KP, Taner CB, Thompson CC, Kröner PT. Simultaneous liver transplant and sleeve gastrectomy not associated with worse index admission outcomes compared to liver transplant alone - a retrospective cohort study. *Transpl Int* 2020; 33: 1447-1452 [PMID: 32772404 DOI: 10.1111/tri.13713]
- 59 Yemini R, Nesher E, Braun M, Cohen M, Carmeli I, Mor E, Keidar A. Long-term outcomes of Roux-en-Y gastric bypass or sleeve gastrectomy in patients with cirrhosis; before, during or after liver transplantation: A single center's experience. *Clin Transplant* 2021; 35: e14374 [PMID: 34043840 DOI: 10.1111/ctr.14374]
- 60 Zamora-Valdes D, Watt KD, Kellogg TA, Poterucha JJ, Di Cecco SR, Francisco-Ziller NM, Taner T, Rosen CB, Heimbach JK. Long-term outcomes of patients undergoing simultaneous liver transplantation and sleeve gastrectomy. *Hepatology* 2018; 68: 485-495 [PMID: 29457842 DOI: 10.1002/hep.29848]
- 61 **Gunturu NS**, Castillo-Larios R, Bowers S, Edwards M, Burns J, Perry D, Elli EF. Combined Sleeve Gastrectomy with Liver Transplant in Patients with Obesity: a Feasibility Study. *Obes Surg* 2022; **32**: 3600-3604 [PMID: 36169908 DOI: 10.1007/s11695-022-06289-1]
- 62 Tariq N, Saharia A, Nwokedi U, Hobeika MJ, Mobley CM, Hsu D, Potter LM, Moore LW, Elaileh A, Sherman V, Ghobrial RM. Combined liver transplantation and sleeve gastrectomy: Report of a brief-interval staged approach. *Liver Transpl* 2023; 29: 422-430 [PMID: 35976078 DOI: 10.1002/lt.26560]
- 63 **Bush K**, Kivlahan DR, McDonell MB, Fihn SD, Bradley KA. The AUDIT alcohol consumption questions (AUDIT-C): an effective brief screening test for problem drinking. Ambulatory Care Quality Improvement Project (ACQUIP). Alcohol Use Disorders Identification Test. *Arch Intern Med* 1998; **158**: 1789-1795 [PMID: 9738608 DOI: 10.1001/archinte.158.16.1789]
- 64 Walther L, Brodén CM, Isaksson A, Hedenbro JL. Alcohol Consumption in Obese Patients Before and After Gastric Bypass as Assessed with the Alcohol Marker Phosphatidylethanol (PEth). *Obes Surg* 2018; **28**: 2354-2360 [PMID: 29500677 DOI: 10.1007/s11695-018-3165-4]
- 65 **Tiniakos DG**. Liver biopsy in alcoholic and non-alcoholic steatohepatitis patients. *Gastroenterol Clin Biol* 2009; **33**: 930-939 [PMID: 19646834 DOI: 10.1016/j.gcb.2009.05.009]
- 66 Theise ND. Histopathology of alcoholic liver disease. *Clin Liver Dis (Hoboken)* 2013; 2: 64-67 [PMID: 30992826 DOI: 10.1002/cld.172]
- 67 Mueller S. Noninvasive assessment of patients with alcoholic liver disease. *Clin Liver Dis (Hoboken)* 2013; 2: 68-71 [PMID: 30992827 DOI: 10.1002/cld.186]

Raisbideng® WJGS | https://www.wjgnet.com



### Published by Baishideng Publishing Group Inc 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA Telephone: +1-925-3991568 E-mail: office@baishideng.com Help Desk: https://www.f6publishing.com/helpdesk https://www.wjgnet.com

