

World Journal of *Gastrointestinal Surgery*

World J Gastrointest Surg 2024 April 27; 16(4): 974-1217



EDITORIAL

- 974 How to identify early complications in patients undergoing distal gastrectomy?
Tropeano G, Chiarello MM, Fico V, Brisinda G
- 982 Quality assessment of surgery for colorectal cancer: Where do we stand?
Morarasu S, Livadaru C, Dimofte GM
- 988 Emerging molecules, tools, technology, and future of surgical knife in gastroenterology
Kumar A, Goyal A
- 999 Carcinoembryonic antigen in the diagnosis, treatment, and follow-up of focal liver lesions
Dilek ON, Arslan Kahraman Dİ, Kahraman G
- 1008 Relationship between *Helicobacter pylori* infection and colorectal polyp/colorectal cancer
Liu Y, Yang DQ, Jiang JN, Jiao Y

MINIREVIEWS

- 1017 Near-infrared cholangiography with intragallbladder indocyanine green injection in minimally invasive cholecystectomy
Symeonidis S, Mantzoros I, Anestiadou E, Ioannidis O, Christidis P, Bitsianis S, Bisbinas V, Zapsalis K, Karastergiou T, Athanasiou D, Apostolidis S, Angelopoulos S
- 1030 Blastomas of the digestive system in adults: A review
Liu Y, El Jabbour T, Somma J, Nakanishi Y, Ligato S, Lee H, Fu ZY

ORIGINAL ARTICLE**Retrospective Study**

- 1043 Single-center retrospective study of the diagnostic value of double-balloon enteroscopy in Meckel's diverticulum with bleeding
He T, Yang C, Wang J, Zhong JS, Li AH, Yin YJ, Luo LL, Rao CM, Mao NF, Guo Q, Zuo Z, Zhang W, Wan P
- 1055 Prognostic value of a nomogram model for postoperative liver metastasis of colon cancer
Cheng DX, Xu KD, Liu HB, Liu Y
- 1066 Computer-assisted three-dimensional individualized extreme liver resection for hepatoblastoma in proximity to the major liver vasculature
Xiu WL, Liu J, Zhang JL, Wang JM, Wang XF, Wang FF, Mi J, Hao XW, Xia N, Dong Q
- 1078 Research on the prognostic value of adjusting intraperitoneal three-dimensional quality evaluation mode in laparoscopic cholecystectomy patients
Zhou Y, Chen ZQ

- 1087** Construction of a predictive model for acute liver failure after hepatectomy based on neutrophil-to-lymphocyte ratio and albumin-bilirubin score
Li XP, Bao ZT, Wang L, Zhang CY, Yang W
- 1097** Predicting short-term thromboembolic risk following Roux-en-Y gastric bypass using supervised machine learning
Ali H, Inayat F, Moond V, Chaudhry A, Afzal A, Anjum Z, Tahir H, Anwar MS, Dahiya DS, Afzal MS, Nawaz G, Sohail AH, Aziz M
- 1109** Comparative analysis of two digestive tract reconstruction methods in total laparoscopic radical total gastrectomy
Dong TX, Wang D, Zhao Q, Zhang ZD, Zhao XF, Tan BB, Liu Y, Liu QW, Yang PG, Ding PA, Zheng T, Li Y, Liu ZJ
- 1121** Incidence of surgical site infection in minimally invasive colorectal surgery
Ni LT, Zhao R, Ye YR, Ouyang YM, Chen X
- Observational Study**
- 1130** Burden of gallstone disease in the United States population: Prepandemic rates and trends
Unalp-Arida A, Ruhl CE
- Prospective Study**
- 1149** Kuicolong-yu enema decoction retains traditional Chinese medicine enema attenuates inflammatory response ulcerative colitis through TLR4/NF- κ B signaling pathway
Han L, Tang K, Fang XL, Xu JX, Mao XY, Li M

SYSTEMATIC REVIEWS

- 1155** Quality-adjusted life years and surgical waiting list: Systematic review of the literature
de la Plaza Llamas R, Ortega Azor L, Hernández Yuste M, Gorini L, Latorre-Fragua RA, Díaz Candelas DA, Al Shwely Abduljabar F, Gemio del Rey IA

META-ANALYSIS

- 1165** Impact of different anastomosis methods on post-recurrence after intestinal resection for Crohn's disease: A meta-analysis
Wang ZZ, Zhao CH, Shen H, Dai GP

CASE REPORT

- 1176** Suspected coexistence of perianal necrotizing sweet syndrome in chronic myelomonocytic leukemia: A case report
Yu KQ, Li HX, Wu J
- 1184** Successful splenic artery embolization in a patient with Behçet's syndrome-associated splenic rupture: A case report
Zhu GZ, Ji DH
- 1189** Stercoral perforation of the cecum: A case report
Yu HC, Pu TW, Kang JC, Chen CY, Hu JM, Su RY

- 1195** Percutaneous transhepatic stenting for acute superior mesenteric vein stenosis after pancreaticoduodenectomy with portal vein reconstruction: A case report
Lin C, Wang ZY, Dong LB, Wang ZW, Li ZH, Wang WB
- 1203** Endoscopic treatment of bleeding gastric ulcer causing gastric wall necrosis: A case report
Li WF, Gao RY, Xu JW, Yu XQ
- 1208** Intermittent melena and refractory anemia due to jejunal cavernous lymphangioma: A case report
Liu KR, Zhang S, Chen WR, Huang YX, Li XG

LETTER TO THE EDITOR

- 1215** New frontiers in ectopic pancreatic tissue management
Covantsev S

ABOUT COVER

Editorial Board Member of *World Journal of Gastrointestinal Surgery*, Arunachalam Rathnaswami, FACS, MBBS, MS, MNAMS, MCh, Ex-Professor and Head, Department of Surgical Gastroenterology, SRM Medical College Hospital and Research Institute SRMIST, Kattankulathur, Chennai, TamilNadu 603203, India. arunarathna@gmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Gastrointestinal Surgery* (*WJGS, World J Gastrointest Surg*) is to provide scholars and readers from various fields of gastrointestinal surgery with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGS mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal surgery and covering a wide range of topics including biliary tract surgical procedures, biliopancreatic diversion, colectomy, esophagectomy, esophagostomy, pancreas transplantation, and pancreatectomy, etc.

INDEXING/ABSTRACTING

The *WJGS* is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports/Science Edition, PubMed, PubMed Central, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for *WJGS* as 2.0; IF without journal self cites: 1.9; 5-year IF: 2.2; Journal Citation Indicator: 0.52; Ranking: 113 among 212 journals in surgery; Quartile category: Q3; Ranking: 81 among 93 journals in gastroenterology and hepatology; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Zi-Hang Xu, Production Department Director: Xiang Li, Cover Editor: Jia-Ru Fan.

NAME OF JOURNAL

World Journal of Gastrointestinal Surgery

ISSN

ISSN 1948-9366 (online)

LAUNCH DATE

November 30, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Peter Schemmer

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/1948-9366/editorialboard.htm>

PUBLICATION DATE

April 27, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.fcpublishing.com>

Retrospective Study

Prognostic value of a nomogram model for postoperative liver metastasis of colon cancer

De-Xin Cheng, Kang-Di Xu, Han-Bo Liu, Yi Liu

Specialty type: Gastroenterology and hepatology**Provenance and peer review:**

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind**Peer-review report's scientific quality classification**Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): C
Grade D (Fair): 0
Grade E (Poor): 0**P-Reviewer:** Cano-Valderrama O, Spain; Tougeron D, France**Received:** December 19, 2023**Peer-review started:** December 19, 2023**First decision:** January 9, 2024**Revised:** January 18, 2024**Accepted:** March 7, 2024**Article in press:** March 7, 2024**Published online:** April 27, 2024**De-Xin Cheng, Han-Bo Liu**, Cancer Center, Department of Interventional Medicine, Zhejiang Provincial People's Hospital (Affiliated People's Hospital, Hangzhou Medical College), Hangzhou 310014, Zhejiang Province, China**Kang-Di Xu, Yi Liu**, General Surgery, Cancer Center, Department of Vascular Surgery, Zhejiang Provincial People's Hospital (Affiliated People's Hospital, Hangzhou Medical College), Hangzhou 310014, Zhejiang Province, China**Corresponding author:** Yi Liu, MM, Attending Doctor, General Surgery, Cancer Center, Department of Vascular Surgery, Zhejiang Provincial People's Hospital (Affiliated People's Hospital, Hangzhou Medical College), No. 158 Shangtang Road, Hangzhou 310014, Zhejiang Province, China. liuyi4917@126.com**Abstract****BACKGROUND**

Colon cancer is one of the most common malignant tumors of the digestive system. Liver metastasis after colon cancer surgery is the primary cause of death in patients with colon cancer.

AIM

To construct a novel nomogram model including various factors to predict liver metastasis after colon cancer surgery.

METHODS

We retrospectively analyzed 242 patients with colon cancer who were admitted and underwent radical resection for colon cancer in Zhejiang Provincial People's Hospital from December 2019 to December 2022. Patients were divided into liver metastasis and non-liver metastasis groups. Sex, age, and other general and clinicopathological data (preoperative blood routine and biochemical test indexes) were compared. The risk factors for liver metastasis were analyzed using single-factor and multifactorial logistic regression. A predictive model was then constructed and evaluated for efficacy.

RESULTSSystemic inflammatory index (SII), C-reactive protein/albumin ratio (CAR), red blood cell distribution width (RDW), alanine aminotransferase, preoperative carcinoembryonic antigen level, and lymphatic metastasis were different between groups ($P < 0.05$). SII, CAR, and RDW were risk factors for liver metastasis after

colon cancer surgery ($P < 0.05$). The area under the curve was 0.93 for the column-line diagram prediction model constructed based on these risk factors to distinguish whether liver metastasis occurred postoperatively. The actual curve of the column-line diagram predicting the risk of postoperative liver metastasis was close to the ideal curve, with good agreement. The prediction model curves in the decision curve analysis showed higher net benefits for a larger threshold range than those in extreme cases, indicating that the model is safer.

CONCLUSION

Liver metastases after colorectal cancer surgery could be well predicted by a nomogram based on the SII, CAR, and RDW.

Key Words: Systemic immunoinflammatory index; C-reactive protein/albumin ratio; Erythrocyte distribution width; Colon cancer; Liver metastasis; Novel nomogram model

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Colon cancer is a highly aggressive and migratory malignant tumor for which the liver is the most common target organ for postoperative metastasis. Herein, we analyzed the general and clinicopathological data of 242 patients with colon cancer who underwent radical resection for colon cancer. The results showed that the systemic inflammatory index, C-reactive protein/albumin ratio, and red blood cell distribution width were risk factors for postoperative liver metastasis in patients with colon cancer. A columnar graph prediction model was subsequently developed based on these three factors, and its predictive efficacy was evaluated.

Citation: Cheng DX, Xu KD, Liu HB, Liu Y. Prognostic value of a nomogram model for postoperative liver metastasis of colon cancer. *World J Gastrointest Surg* 2024; 16(4): 1055-1065

URL: <https://www.wjgnet.com/1948-9366/full/v16/i4/1055.htm>

DOI: <https://dx.doi.org/10.4240/wjgs.v16.i4.1055>

INTRODUCTION

Colon cancer is a highly aggressive and migratory malignant tumor, which is most commonly characterized by intestinal obstruction, loss of body mass, abdominal pain, and blood in the stool[1]. The liver is the most common metastatic site in patients with colon cancer[2], with liver metastasis occurring in 25% of patients after surgery[3]. Therefore, designing a convenient and efficient predictive index for liver metastasis after colon cancer surgery would not only be beneficial for the early screening of colon cancer but also play an important role in knowing whether liver metastasis occurs after colon cancer surgery.

Several studies have shown that tumor-associated inflammatory cells can directly or indirectly act on tumors and are closely related to tumorigenesis and migration[4]. Systemic inflammation can occur within the tumor immune microenvironment and is closely related to tumor development[5,6]. The systemic inflammatory index (SII) is a test that assesses the impact of peripheral blood neutrophil, lymphocyte, and platelet counts on the biological behavior of cells. By detecting these relative values from these cell types, the SII measures whether their function is normal and subsequently assesses the patient's status[7]. Studies have demonstrated that SII is independently associated with postoperative liver metastasis in colon cancer, and its predictive ability is superior to that of other inflammatory factors[8]. Therefore, the SII may be a marker for the prediction of liver metastasis after colon cancer surgery.

C-reactive protein (CRP) levels are influenced by tumor necrosis factor- α , interleukin (IL)-1 β , and IL-6, indicating a strong association with acute inflammation[9]. Conversely, albumin levels have been associated with chronic inflammation and can indicate nutritional status[10,11]. The CRP/albumin ratio (CAR) reflects the inflammatory status and nutritional level of the patient, is less susceptible to the influence of other factors, and has better stability. Therefore, studies have confirmed that CAR can be used as a prognostic marker for colon cancer in clinical practice[12]. CAR has also been shown to be a reliable marker for predicting survival in patients with metastatic colon cancer[13]. Therefore, we hypothesized that CAR could be used as a new parameter for predicting liver metastasis after colon cancer surgery.

Owing to the many advantages of blood sample analysis, including easy accessibility and low invasiveness, as well as its ability to allow long-term monitoring and assessment of systemic status, clinical blood tests are commonplace. Among these, red cell distribution width (RDW) is a standard blood test measurement item that reflects the homogeneity of red blood cells[14]. RDW has been used as a prognostic marker for various cancers, including lung, liver, esophagogastric, and breast cancers[15,16], and has also been used as a potential prognostic marker for colon cancer. However, the predictive value of RDW for the occurrence of postoperative liver metastasis in colon cancer remains unclear. Therefore, a more accurate and comprehensive assessment of the predictive value of postoperative liver metastasis in colon cancer through the continuous exploration of new blood indices would be crucial in guiding clinical work.

In recent years, relevant studies have reported the factors influencing the occurrence of liver metastasis following colon cancer surgery; however, a more accurate prediction model is still lacking. Columnar graphs can visualize the probability of patients presenting with corresponding diseases, meaning that they are commonly used for the rapid clinical screening of high-risk groups and the development of effective interventions. Therefore, this retrospective study aimed to investigate the predictive value of the column chart model based on SII, CAR, and RDW for postoperative liver metastasis of colon cancer to aid in the clinical screening of postoperative liver metastasis of colon cancer.

MATERIALS AND METHODS

Clinical data

The general and clinical data of 242 patients with colon cancer admitted to the Zhejiang Provincial People's Hospital between December 2019 and December 2022 were retrospectively analyzed. Patients were divided into the transfer ($n = 52$) and non-transfer ($n = 190$) groups based on whether they developed liver metastasis after surgery or not. The analysis process is shown in [Figure 1](#).

Inclusion and exclusion criteria

The inclusion criteria were as follows: (1) Preoperative diagnosis of colon cancer by imaging and histopathology; (2) postoperative pathologic examination confirming colon cancer; (3) radical resection of colon cancer; (4) availability of complete clinicopathological data; and (5) complete follow-up data.

The exclusion criteria were as follows: (1) Preoperative colon cancer with distant metastasis; (2) colon cancer with perforation or hemorrhage; (3) previous history of malignancy; and (4) underlying severe cardiac, hepatic, renal, pulmonary, or hematological diseases.

Observation indexes

General and clinicopathological data were collected for all patients, including age, sex, body mass index (BMI), history of alcohol consumption, smoking, and history of diseases (cardiac disease, diabetes mellitus, and hypertension). Clinicopathological data included SII, CAR, RDW, total bilirubin, albumin, platelet count, white blood cell (WBC) count, alanine aminotransferase (ALT), aspartate aminotransferase (AST), surgical modality, vascular invasion, tumor site, type of pathology, postoperative chemotherapy, preoperative carcinoembryonic antigen (CEA) levels, and lymph node metastasis status.

Statistical analysis

The Statistical Package for Social Science (SPSS, version 26.0) was used to analyze all data. For normally distributed data points, comparisons between two groups were performed using *t*-test, and results are expressed as mean \pm SD. Non-normally distributed data were analyzed by the Mann-Whitney *U*-test and expressed as M (P25-P75). Comparisons of counting data between the two groups were performed using the χ^2 test, expressed as *n* (%). Binary logistic regression was applied to analyze the relevant influencing factors of liver metastasis after surgery for colon cancer.

A predictive model was constructed using *R* software based on multifactorial logistic regression analysis, and the model was internally validated using the bootstrap method with 1000 repetitive samples. The accuracy of the model was assessed by the area under the curve (AUC) of the receiver operating characteristic (ROC) curve, the calibration curve, and the decision curve analysis (DCA).

RESULTS

Comparison of general data between the liver transfer and non-transfer groups after colon cancer surgery

General information such as age, BMI, sex, drinking history, smoking history, and medical history of the patients in the two groups were analyzed separately. There was no significant difference between the two groups ($P > 0.05$) ([Table 1](#)).

Clinical data comparison between the two patient groups

To investigate the predictive value of the SII, CAR, and RDW for the development of liver metastases after surgery in patients with colon cancer, a one-way analysis of all case information collected was performed. The results revealed no significant difference between the two groups in terms of total bilirubin, albumin, platelet count, WBC count, AST level, surgical method, vascular invasion, tumor site, pathology type, or postoperative chemotherapy ($P > 0.05$). However, there was a significant difference between the metastatic and non-metastatic groups in the comparison of SII, CAR, RDW, ALT, preoperative CEA, and lymphatic metastasis ($P < 0.05$) ([Table 2](#)).

Multifactorial analysis of liver metastasis after colon cancer surgery

Binary logistic regression analysis was performed with the occurrence of liver transfer as the dependent variable and SII, CAR, RDW, ALT, preoperative CEA, and lymphatic metastasis as the independent variables. The variable assignments are shown in [Table 3](#). Factors affecting the univariate analysis ($P < 0.05$) (SII, CAR, RDW, ALT, preoperative CEA, and lymphatic metastasis) in the transfer and non-transfer groups were included in the binary logistic regression analysis,

Table 1 Comparison of the general data in patients in the transfer and non-transfer groups, *n* (%)

Considerations	Transfer group (<i>n</i> = 52)	Non-transfer group (<i>n</i> = 190)	<i>t/z/χ</i> ²	<i>P</i> value
Age (yr)	58.96 ± 16.59	61.31 ± 14.00	0.93	0.35
BMI (kg/m ²)	22.16 ± 3.84	22.56 ± 2.92	0.69	0.50
Sex			1.01	0.32
Male	30 (57.70)	124 (65.30)		
Female	22 (42.30)	66 (34.70)		
Drinking history			1.41	0.24
Yes	40 (76.90)	130 (68.40)		
No	12 (23.10)	60 (31.60)		
Smoking history			1.13	0.29
Yes	44 (84.60)	148 (77.90)		
No	8 (15.40)	42 (22.10)		
Heart disease			0.01	0.93
Yes	14 (26.90)	50 (26.30)		
No	38 (73.10)	140 (73.70)		
Diabetes			1.58	0.21
Yes	14 (26.90)	36 (18.90)		
No	38 (73.10)	154 (81.10)		
Hypertension			1.84	0.18
Yes	8 (15.40)	46 (24.20)		
No	44 (84.60)	144 (75.80)		

BMI: Body mass index.

which showed that SII, CAR, and RDW were independent risk factors for liver transfer in the postoperative period, and the model excluded the three factors of ALT, preoperative CEA, and lymphatic metastasis ($P > 0.05$) (Table 4).

Predictive value of SII, CAR, and RDW for postoperative liver metastasis in patients

The ROC curves showed that the predicted AUCs of SII, CAR, and RDW were 0.88, 0.68, and 0.67, respectively, with sensitivities of 96.20%, 65.40%, and 50.00% and specificities of 71.60%, 67.40%, and 87.40%, respectively (Table 5 and Figure 2).

Construction and validation of the column-line diagram model

Based on the results of the multifactorial logistic regression analysis, three factors, namely SII, CAR, and RDW, were used in a columnar graphical prediction model constructed using the R software for the occurrence of liver metastasis after radical resection of colon cancer (Figure 3). The ROC curve (Figure 4A) assessed that the AUC (95%CI) of the column chart for distinguishing the occurrence of liver metastasis after radical resection of colon cancer was 0.93 (0.89-0.96), and the optimal cut-off value of the column chart was 0.16, with a sensitivity of 91.30% and a specificity of 81.10%. Assessment of the calibration curve (Figure 4B) showed that the column chart could predict the occurrence of liver metastasis after radical resection of colon cancer (Figure 4B). Further assessment of the curves showing the predicted risk of occurrence of postoperative liver metastases from colon cancer in the column-line diagram and the actual risk of occurrence of postoperative liver metastases from colon cancer showed high homology. This result indicates that the predicted risk of occurrence of postoperative liver metastases from colon cancer by the column-line diagram matched well with the risk of occurrence of liver metastases from actual postoperative liver metastases from colon cancer. Based on the predicted model curve in the DCA curve compared to that in the extreme case, the net benefit of the model is superior in the larger threshold range (above 15%), indicating that the model is safer (Figure 4C).

DISCUSSION

Colon cancer is a highly aggressive and migratory malignant tumor, and the liver is the most common organ site for

Table 2 Comparison of the clinical data in the transfer and non-transfer groups

Considerations	Transfer group (n = 52)	Non-transfer group (n = 190)	t/z/ χ^2	P value
SII	385.21 ± 78.63	264.29 ± 64.43	-10.19	< 0.01
CAR	1.02 ± 0.40	0.79 ± 0.38	-3.88	< 0.01
RDW (FL)	17.64 ± 1.98	16.28 ± 1.26	-4.72	< 0.01
Total bilirubin (μmol/L)	13.45 ± 2.51	13.75 ± 3.28	0.71	0.48
Albumin (g/L)	42.08 ± 3.81	42.22 ± 2.71	0.25	0.80
Blood platelet count (10 ⁹ /L)	167.13 ± 65.02	157.29 ± 40.59	-1.04	0.30
WBC (10 ⁹ /L)	7.48 ± 2.58	7.52 ± 3.12	0.08	0.93
ALT (U/L)	47.88 ± 1.84	48.50 ± 1.59	2.40	0.017
AST (U/L)	47.29 ± 1.25	47.14 ± 1.76	-0.61	0.54
Surgical procedures			0.01	0.94
Laparoscopy	32 (61.50)	118 (62.10)		
Open the abdomen	20 (38.50)	72 (37.90)	0.01	0.93
Vascular invasion			3.26	0.07
Yes	24 (46.20)	62 (32.60)		
No	28 (53.80)	128 (67.40)	1.58	0.21
Tumor site			1.01	0.32
Left	30 (57.70)	124 (65.30)		
Right	22 (42.30)	66 (34.70)		
Pathological type			1.05	0.31
Adenocarcinoma	48 (92.30)	182 (95.80)		
Other	4 (7.70)	8 (4.20)		
Postoperative chemotherapy			3.26	0.07
Yes	24 (46.20)	62 (32.60)		
No	28 (53.80)	128 (67.40)		
Preoperative CEA (μg/L)			9.59	< 0.01
< 5	21 (40.40)	122 (64.20)		
≥ 5	31 (59.60)	68 (35.80)		
Lymphatic node transfer			4.90	0.03
Yes	28 (53.80)	70 (36.80)		
No	24 (46.20)	120 (63.20)		

ALT: Alanine aminotransferase; AST: Aspartate aminotransferase; CAR: C-reactive protein/albumin ratio; CEA: Carcinoembryonic antigen; RDW: Red blood cell distribution width; SII: Systemic inflammatory index; WBC: White blood cell.

postoperative metastasis[17]. Liver metastasis occurs in approximately 50% of patients with colon cancer, with metastases found at diagnosis in 25% of patients[18]. Further, metastasis is the primary cause of death in patients with colon cancer [19]. Therefore, assessing the predictive value of liver metastases after surgery in patients with colon cancer is clinically important. Overall, we found that SII, CAR, and RDW alone had some assessment value for the postoperative prognosis of patients with colon cancer[12,20,21], and the SII had a high predictive ability for the occurrence of liver metastasis after surgery in these patients[8]. Therefore, we constructed a column-line graph prediction model for the occurrence of liver metastasis after radical resection of colon cancer using SII, CAR, and RDW to help clinicians identify patients at high risk of liver metastasis. Such predictions would allow timely intervention, which could help promote optimal treatment timing.

An impaired immune function, systemic inflammatory state, and chronic inflammatory response promote the development and progression of malignant tumors and are closely associated with patient prognosis. SII is a novel inflammatory index closely related to inflammatory and immune pathways. This relationship can be expressed by the

Table 3 Variable assignment table

Project	Assignment description
Dependent variable	Transfer group '1', Non-transfer group '0'
SII	Continuous variable
CAR	Continuous variable
RDW	Continuous variable
ALT	Continuous variable
Preoperative CEA	≥ 5 µg/L '1', < 5 µg/L '0'
Lymphatic node transfer	Yes '1', No '0'

ALT: Alanine aminotransferase; CAR: C-reactive protein/albumin ratio; CEA: Carcinoembryonic antigen; RDW: Red blood cell distribution width; SII: Systemic inflammatory index.

Table 4 Multifactorial analysis of liver metastases after surgery for colon cancer

Considerations	β	SE	Wald χ ²	P value	OR	95%CI
SII	0.26	< 0.01	36.78	< 0.01	1.03	1.02-1.03
CAR	1.57	0.56	7.74	0.01	4.79	1.59-14.44
RDW	0.57	0.15	13.77	< 0.01	1.76	1.31-2.38
ALT	-0.25	0.15	2.87	0.09	0.78	0.58-1.04
Preoperative CEA	0.85	0.48	3.09	0.08	2.34	0.91-6.04
Lymphatic node transfer	0.7	0.49	2.04	0.15	2.01	0.77-5.23
Constant	-9.17	7.71	1.41	0.24	< 0.01	-

ALT: Alanine aminotransferase; CAR: C-reactive protein/albumin ratio; Carcinoembryonic antigen; RDW: Red blood cell distribution width; SII: Systemic inflammatory index.

Table 5 Predictive value of systemic inflammatory index, lymphatic metastasis, C-reactive protein/albumin ratio, and red blood cell distribution width in patients with postoperative liver metastasis

Considerations	Truncation value	AUC	SE	P value	95%CI	Sensitivity (%)	Idiosyncrasy (%)
SII	290.52	0.88	0.03	< 0.01	0.83-0.94	96.2	71.6
CAR	0.86	0.68	0.04	< 0.01	0.60-0.76	65.4	67.2
RDW	16.76	0.67	0.04	< 0.01	0.59-0.76	50	87.4

CAR: C-reactive protein/albumin ratio; RDW: Red blood cell distribution width; SII: Systemic inflammatory index; AUC: Area under the curve.

formula $SII = \text{platelets} \times \text{neutrophils} / \text{Lymphocytes} (\times 10^9/L)$ [19]. Studies have shown that the SII has a high value in predicting the prognosis of malignant tumors[22]. In the present study, we reviewed related studies on postoperative liver transfer for colorectal cancer, which showed that the preoperative SII level of patients in the transfer group was higher than that of both the transfer and non-transfer groups, indicating that the preoperative SII level has some predictive value for liver metastasis of colorectal cancer after surgery. The results of the ROC curve analysis revealed an AUC of 0.88, which suggests that the efficacy of the SII in predicting liver metastasis of colon cancer after surgery improved, with a sensitivity of 96.20%. This result is consistent with the findings of Lu *et al*[8] (AUC = 0.882).

Related studies have shown that the prognoses of patients with malignant tumors such as colorectal cancer[23] and hepatocellular carcinoma[24] are closely related. Preoperative CAR (a systemic inflammatory marker) has further been found to serve as a prognostic marker for liver metastases of colorectal cancer treated with potential radical resection[25]. Meanwhile, other studies have shown that CAR reflects the inflammatory status and nutritional level of the patient, with low susceptibility and high stability[26]. In the present study, we showed that CAR levels were higher in the group with liver metastases than in those without liver metastases after colon cancer surgery; this is consistent with the findings of Liao *et al*[12]. These results suggest that the preoperative CAR level has some predictive value for postoperative liver




Figure 1 Flow chart of the study. SII: Systemic inflammatory index; CAR: C-reactive protein/albumin ratio; RDW: Red blood cell distribution width.




Figure 2 Receiver operating characteristic curves showing the prediction sensitivity of systemic inflammatory index, C-reactive protein/albumin ratio, and red blood cell distribution width in patients with postoperative liver metastases. SII: Systemic inflammatory index; CAR: C-reactive protein/albumin ratio; RDW: Red blood cell distribution width.

metastasis of colon cancer. In our study, the ROC results showed that the efficacy of CAR in predicting postoperative hepatic metastasis of colon cancer was average (AUC = 0.68), indicating that there is a limitation to the value of CAR alone for predicting postoperative hepatic metastasis of colon cancer.

Platelet secretion, migration, and endothelial cell proliferation of vascular endothelial growth factor have an inducing effect and increase vascular permeability, while tumor cells penetrate the machinery of blood vessels to metastasize and




Figure 3 Nomogram prediction model of the risk of postoperative liver metastases in patients with colon cancer. SII: Systemic inflammatory index; CAR: C-reactive protein/albumin ratio; RDW: Red blood cell distribution width.




Figure 4 Predictive performance of the column graph model on the training set. A: receiver operating characteristic curve of the training set; B: Calibration curve of the training set; C: The DCA curve of the training set. SII: Systemic inflammatory index; CAR: C-reactive protein/albumin ratio; RDW: Red blood cell distribution width.

invade. RDW reflects the heterogeneous parameters of erythrocyte size and peripheral blood volume[27]. Studies have shown that the prognosis of patients with solid tumors is closely related to RDW[16]. In the present study, we showed that the RDW level was higher in the group with liver metastasis than in the group without liver metastasis after colon cancer surgery. This result is consistent with the findings of Lu *et al*[28]. These results suggest that preoperative RDW has a certain predictive value for postoperative liver metastasis in colon cancer. The ROC results showed that the efficacy of RDW in predicting the postoperative incidence of liver metastasis in colon cancer was average (AUC = 0.67), suggesting that the use of RDW alone does not have a good predictive value for liver metastasis in postoperative colon cancer.

Pang *et al*[29] showed that column-line diagrams performed better in risk stratification of prognosis for patients undergoing radical colon cancer surgery. Therefore, in the present study, we further constructed a column-line diagram prediction model of the occurrence of liver metastasis after radical resection of colon cancer by SII, CAR, and RDW. The ROC results showed an AUC (95%CI) of 0.93 (0.89-0.96). In a prior study, Nagata *et al*[30] developed a prediction model for liver metastasis involving tumor differentiation, pathological staging, and preoperative CEA level. The AUC (95%CI) of this model was 0.78 (0.71-0.84), which was less effective than the model we constructed, suggesting that the models constructed by SII, CAR, and RDW can improve the prediction of liver metastasis after colon cancer surgery accuracy. In addition, we plotted calibration curves and DCA curves to further evaluate the efficacy of the model. The results of the calibration curve analysis showed that the actual curves of the column-line graph model for predicting the occurrence of postoperative liver metastasis in colon cancer largely agreed with the ideal curves, which indicated that the model had good consistency. In addition, the DCA curve suggested that the net benefit of the model was superior to the threshold value of 15%, indicating that the model was safer and had a higher clinical utility.

Overall, the results of the present study show that poor SII, CAR, and RDW results are risk factors for liver metastasis after colon cancer surgery. The prediction model established in this study based on these risk factors has good discriminatory power and a high calibrating ability for the occurrence of liver metastasis after radical resection of colon cancer. Thus, this model could help clinicians more intuitively identify high-risk patients prone to liver metastasis after surgery, take targeted screening measures, and formulate individualized medical treatment strategies. However, this study had certain limitations. First, only the data of patients who underwent radical resection for colon cancer in our hospital at a fixed time period were selected, meaning that selection bias may have been an issue. Second, as this study was a retrospective study, there is a possibility of retrospective bias. Third, the single-center design limits the generalizability of the results. Finally, only the bootstrap method was used for the internal validation of the predictive model of the risk of liver metastasis after surgery for colon cancer, and no other validation method was used. Therefore, external validation with large samples and multicenter data is needed in the future to assess the generalizability of the model and further validate and improve its prediction ability.

CONCLUSION

We identified SII, CAR, and RDW as risk factors affecting the occurrence of liver metastasis after colon cancer surgery. Further, we showed that the combination of these three factors could effectively predict the risk of liver metastasis, and our prediction model based on these three risk factors had good predictive efficacy. Therefore, in clinical practice, SII, CAR, and RDW can be beneficial for the early screening of colon cancer and predict the occurrence of liver metastasis after surgery in advance, which can help determine the best time for treatment.

ARTICLE HIGHLIGHTS

Research background

Colon cancer is one of the most common malignant tumors of the digestive system. Clinical treatment primarily includes surgery, and chemo- and radiotherapy are used as auxiliary therapies to ensure comprehensive treatment. Surgical resection is an effective radical treatment for colon cancer; however, liver metastasis still occurs in 25% of patients after surgery.

Research motivation

In the current study, several methods to predict and assess liver metastasis after colon cancer surgery were investigated. Therefore, the identification of simple and efficient predictors of liver metastasis after colon cancer surgery is clinically important.

Research objectives

This study is aimed to predict the efficacy of postoperative liver metastasis in colon cancer using nomogram models constructed from systemic inflammatory index (SII), C-reactive protein/albumin ratio (CAR), and red blood cell distribution width (RDW) to provide predictive value in the clinic.

Research methods

This study retrospectively analyzed the clinicopathological data of 242 patients who underwent radical resection for colon cancer; analyzed the risk factors affecting the development of liver metastases in these patients; assessed nomogram models constructed using the SII, CAR, and RDW; and evaluated the predictive efficacy of the models.

Research results

The SII, CAR, and RDW are risk factors for liver metastasis after colon cancer surgery. The area under the receiver operating characteristic curve of the column-line diagram model constructed based on these three risk factors to predict whether liver metastasis occurred after colon cancer surgery was 0.93 (95%CI: 0.89-0.96). The calibration curve of the column-line diagram predicting the risk of postoperative liver metastasis from colon cancer matched well with the actual

risk of occurrence, and the net benefit of the model was better, indicating that the model was safer.

Research conclusions

The SII, CAR, and RDW are independent risk factors for the development of liver metastases after colon cancer surgery, and the predictive efficacy of the column-line graph model constructed using the SII, CAR, and RDW was high.

Research perspectives

To analyze the factors influencing the occurrence of liver metastases in patients with colon cancer and construct a nomogram model using the SII, CAR, and RDW to evaluate the predictive efficacy of the model.

FOOTNOTES

Author contributions: Cheng DX wrote the manuscript; Liu Y reviewed the manuscript; Xu KD and Liu HB collected the data; and all authors annotated the manuscript.

Institutional review board statement: This study was reviewed and approved by the Institutional Review Board of Zhejiang Provincial People's Hospital (Approval No. 2023-338).

Informed consent statement: The Institutional Review Board waived the requirement for informed consent.

Conflict-of-interest statement: The authors declare no conflicts of interest.

Data sharing statement: Data used in this study are available from the corresponding author.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: De-Xin Cheng 0009-0007-8791-249X; Kang-Di Xu 0000-0003-2253-9982; Han-Bo Liu 0000-0002-6624-0432; Yi Liu 0009-0000-1346-5155.

S-Editor: Wang JL

L-Editor: A

P-Editor: Yuan YY

REFERENCES

- 1 Ruan H, Leibowitz BJ, Zhang L, Yu J. Immunogenic cell death in colon cancer prevention and therapy. *Mol Carcinog* 2020; **59**: 783-793 [PMID: 32215970 DOI: 10.1002/mc.23183]
- 2 Van Cutsem E, Nordlinger B, Adam R, Köhne CH, Pozzo C, Poston G, Ychou M, Rougier P; European Colorectal Metastases Treatment Group. Towards a pan-European consensus on the treatment of patients with colorectal liver metastases. *Eur J Cancer* 2006; **42**: 2212-2221 [PMID: 16904315 DOI: 10.1016/j.ejca.2006.04.012]
- 3 Saad AM, Abdel-Rahman O. Initial systemic chemotherapeutic and targeted therapy strategies for the treatment of colorectal cancer patients with liver metastases. *Expert Opin Pharmacother* 2019; **20**: 1767-1775 [PMID: 31314604 DOI: 10.1080/14656566.2019.1642324]
- 4 Marozzi M, Parnigoni A, Negri A, Viola M, Vigetti D, Passi A, Karousou E, Rizzi F. Inflammation, Extracellular Matrix Remodeling, and Proteostasis in Tumor Microenvironment. *Int J Mol Sci* 2021; **22** [PMID: 34360868 DOI: 10.3390/ijms22158102]
- 5 Proctor MJ, Morrison DS, Talwar D, Balmer SM, O'Reilly DS, Foulis AK, Horgan PG, McMillan DC. An inflammation-based prognostic score (mGPS) predicts cancer survival independent of tumour site: a Glasgow Inflammation Outcome Study. *Br J Cancer* 2011; **104**: 726-734 [PMID: 21266974 DOI: 10.1038/sj.bjc.6606087]
- 6 Nguyen AV, Wu YY, Lin EY. STAT3 and sphingosine-1-phosphate in inflammation-associated colorectal cancer. *World J Gastroenterol* 2014; **20**: 10279-10287 [PMID: 25132744 DOI: 10.3748/wjg.v20.i30.10279]
- 7 Walzik D, Joisten N, Zacher J, Zimmer P. Transferring clinically established immune inflammation markers into exercise physiology: focus on neutrophil-to-lymphocyte ratio, platelet-to-lymphocyte ratio and systemic immune-inflammation index. *Eur J Appl Physiol* 2021; **121**: 1803-1814 [PMID: 33787989 DOI: 10.1007/s00421-021-04668-7]
- 8 Lu Y, Xin D, Wang F. Predictive Significance Of Preoperative Systemic Immune-Inflammation Index Determination In Postoperative Liver Metastasis Of Colorectal Cancer. *Onco Targets Ther* 2019; **12**: 7791-7799 [PMID: 31571929 DOI: 10.2147/OTT.S223419]
- 9 Nakazaki H. Preoperative and postoperative cytokines in patients with cancer. *Cancer* 1992; **70**: 709-713 [PMID: 1320454 DOI: 10.1002/1097-0142(19920801)70:3<709::aid-cnrcr2820700328>3.0.co;2-o]
- 10 McMillan DC, Watson WS, O'Gorman P, Preston T, Scott HR, McArdle CS. Albumin concentrations are primarily determined by the body cell mass and the systemic inflammatory response in cancer patients with weight loss. *Nutr Cancer* 2001; **39**: 210-213 [PMID: 11759282 DOI: 10.1207/S15327914nc392_8]

- 11 **Chojkier M.** Inhibition of albumin synthesis in chronic diseases: molecular mechanisms. *J Clin Gastroenterol* 2005; **39**: S143-S146 [PMID: 15758650 DOI: 10.1097/01.mcg.0000155514.17715.39]
- 12 **Liao CK, Yu YL, Lin YC, Hsu YJ, Chern YJ, Chiang JM, You JF.** Prognostic value of the C-reactive protein to albumin ratio in colorectal cancer: an updated systematic review and meta-analysis. *World J Surg Oncol* 2021; **19**: 139 [PMID: 33933070 DOI: 10.1186/s12957-021-02253-y]
- 13 **Haruki K, Shiba H, Horiuchi T, Sakamoto T, Gocho T, Fujiwara Y, Furukawa K, Misawa T, Yanaga K.** Impact of the C-reactive protein to albumin ratio on long-term outcomes after hepatic resection for colorectal liver metastases. *Am J Surg* 2017; **214**: 752-756 [PMID: 28187858 DOI: 10.1016/j.amjsurg.2017.02.001]
- 14 **Jandial A, Kumar S, Bhalla A, Sharma N, Varma N, Varma S.** Elevated Red Cell Distribution Width as a Prognostic Marker in Severe Sepsis: A Prospective Observational Study. *Indian J Crit Care Med* 2017; **21**: 552-562 [PMID: 28970653 DOI: 10.4103/ijccm.IJCCM_208_17]
- 15 **Koma Y, Onishi A, Matsuoka H, Oda N, Yokota N, Matsumoto Y, Koyama M, Okada N, Nakashima N, Masuya D, Yoshimatsu H, Suzuki Y.** Increased red blood cell distribution width associates with cancer stage and prognosis in patients with lung cancer. *PLoS One* 2013; **8**: e80240 [PMID: 24244659 DOI: 10.1371/journal.pone.0080240]
- 16 **Hu L, Li M, Ding Y, Pu L, Liu J, Xie J, Cabanero M, Li J, Xiang R, Xiong S.** Prognostic value of RDW in cancers: a systematic review and meta-analysis. *Oncotarget* 2017; **8**: 16027-16035 [PMID: 27926498 DOI: 10.18632/oncotarget.13784]
- 17 **Polk N, Budai B, Hitre E, Patócs A, Mersich T.** Corrigendum: High Neutrophil-to-Lymphocyte Ratio (NLR) and Systemic Immune-Inflammation Index (SII) are Markers of Longer Survival After Metastasectomy of Patients With Liver-Only Metastasis of Rectal Cancer. *Pathol Oncol Res* 2022; **28**: 1610658 [PMID: 36277961 DOI: 10.3389/pore.2022.1610658]
- 18 **Chandra R, Karalis JD, Liu C, Murimwa GZ, Voth Park J, Heid CA, Reznik SI, Huang E, Minna JD, Brekken RA.** The Colorectal Cancer Tumor Microenvironment and Its Impact on Liver and Lung Metastasis. *Cancers (Basel)* 2021; **13** [PMID: 34944826 DOI: 10.3390/cancers13246206]
- 19 **Wang Q, Zhu D.** The prognostic value of systemic immune-inflammation index (SII) in patients after radical operation for carcinoma of stomach in gastric cancer. *J Gastrointest Oncol* 2019; **10**: 965-978 [PMID: 31602335 DOI: 10.21037/jgo.2019.05.03]
- 20 **Nøst TH, Alcalá K, Urbarova I, Byrne KS, Guida F, Sandanger TM, Johansson M.** Systemic inflammation markers and cancer incidence in the UK Biobank. *Eur J Epidemiol* 2021; **36**: 841-848 [PMID: 34036468 DOI: 10.1007/s10654-021-00752-6]
- 21 **Bernstein CN, Benchimol EI, Bitton A, Murthy SK, Nguyen GC, Lee K, Cooke-Lauder J, Kaplan GG.** The Impact of Inflammatory Bowel Disease in Canada 2018: Extra-intestinal Diseases in IBD. *J Can Assoc Gastroenterol* 2019; **2**: S73-S80 [PMID: 31294387 DOI: 10.1093/jcag/gwy053]
- 22 **Chen JH, Zhai ET, Yuan YJ, Wu KM, Xu JB, Peng JJ, Chen CQ, He YL, Cai SR.** Systemic immune-inflammation index for predicting prognosis of colorectal cancer. *World J Gastroenterol* 2017; **23**: 6261-6272 [PMID: 28974892 DOI: 10.3748/wjg.v23.i34.6261]
- 23 **Matsuoka H, Ando K, Hu Q, Zaitzu Y, Tsuda Y, Hisamatsu Y, Nakashima Y, Kimura Y, Oki E, Mori M.** Postoperative C-reactive protein/albumin ratio is a biomarker of risk of recurrence and need for adjuvant chemotherapy for stage III colorectal cancer. *Int J Clin Oncol* 2020; **25**: 1318-1326 [PMID: 32279124 DOI: 10.1007/s10147-020-01672-3]
- 24 **Lin N, Li J, Ke Q, Wang L, Cao Y, Liu J.** Clinical Significance of C-Reactive Protein to Albumin Ratio in Patients with Hepatocellular Carcinoma: A Meta-Analysis. *Dis Markers* 2020; **2020**: 4867974 [PMID: 32963635 DOI: 10.1155/2020/4867974]
- 25 **Shibutani M, Nagahara H, Fukuoka T, Iseki Y, Hirakawa K, Ohira M.** Efficacy of Adjuvant Chemotherapy According to the Classification of Recurrence Risk Based on Systemic Inflammatory Markers in Patients With Liver Metastases of Colorectal Cancer. *Anticancer Res* 2019; **39**: 5039-5045 [PMID: 31519612 DOI: 10.21873/anticancerres.13695]
- 26 **Ito T, Shinkawa H, Takemura S, Tanaka S, Nishioka T, Miyazaki T, Ishihara A, Kubo S.** Impact of the Preoperative C-reactive Protein to Albumin Ratio on the Long-Term Outcomes of Hepatic Resection for Intrahepatic Cholangiocarcinoma. *Asian Pac J Cancer Prev* 2020; **21**: 2373-2379 [PMID: 32856868 DOI: 10.31557/APJCP.2020.21.8.2373]
- 27 **Chen W, Xin S, Xu B.** Value Research of NLR, PLR, and RDW in Prognostic Assessment of Patients with Colorectal Cancer. *J Healthc Eng* 2022; **2022**: 7971415 [PMID: 35469225 DOI: 10.1155/2022/7971415]
- 28 **Lu X, Huang X, Xue M, Zhong Z, Wang R, Zhang W, Wang L, Qiao Y, Ling F, Zhang Q, Zhang Y.** Prognostic significance of increased preoperative red cell distribution width (RDW) and changes in RDW for colorectal cancer. *Cancer Med* 2023; **12**: 13361-13373 [PMID: 37143237 DOI: 10.1002/cam4.6036]
- 29 **Pang X, Xu B, Lian J, Wang R, Wang X, Shao J, Tang S, Lu H.** Real-world survival of colon cancer after radical surgery: A single-institutional retrospective analysis. *Front Oncol* 2022; **12**: 914076 [PMID: 36185216 DOI: 10.3389/fonc.2022.914076]
- 30 **Nagata H, Ishihara S, Oba K, Tanaka T, Hata K, Kawai K, Nozawa H.** Development and Validation of a Prediction Model for Organ-Specific Recurrences After Curative Resection of Colon Cancer. *Dis Colon Rectum* 2019; **62**: 1043-1054 [PMID: 31318776 DOI: 10.1097/DCR.0000000000001430]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: office@baishideng.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

