

World Journal of *Gastrointestinal Surgery*

World J Gastrointest Surg 2024 April 27; 16(4): 974-1217



EDITORIAL

- 974 How to identify early complications in patients undergoing distal gastrectomy?
Tropeano G, Chiarello MM, Fico V, Brisinda G
- 982 Quality assessment of surgery for colorectal cancer: Where do we stand?
Morarasu S, Livadaru C, Dimofte GM
- 988 Emerging molecules, tools, technology, and future of surgical knife in gastroenterology
Kumar A, Goyal A
- 999 Carcinoembryonic antigen in the diagnosis, treatment, and follow-up of focal liver lesions
Dilek ON, Arslan Kahraman Dİ, Kahraman G
- 1008 Relationship between *Helicobacter pylori* infection and colorectal polyp/colorectal cancer
Liu Y, Yang DQ, Jiang JN, Jiao Y

MINIREVIEWS

- 1017 Near-infrared cholangiography with intragallbladder indocyanine green injection in minimally invasive cholecystectomy
Symeonidis S, Mantzoros I, Anestiadou E, Ioannidis O, Christidis P, Bitsianis S, Bisbinas V, Zapsalis K, Karastergiou T, Athanasiou D, Apostolidis S, Angelopoulos S
- 1030 Blastomas of the digestive system in adults: A review
Liu Y, El Jabbour T, Somma J, Nakanishi Y, Ligato S, Lee H, Fu ZY

ORIGINAL ARTICLE

Retrospective Study

- 1043 Single-center retrospective study of the diagnostic value of double-balloon enteroscopy in Meckel's diverticulum with bleeding
He T, Yang C, Wang J, Zhong JS, Li AH, Yin YJ, Luo LL, Rao CM, Mao NF, Guo Q, Zuo Z, Zhang W, Wan P
- 1055 Prognostic value of a nomogram model for postoperative liver metastasis of colon cancer
Cheng DX, Xu KD, Liu HB, Liu Y
- 1066 Computer-assisted three-dimensional individualized extreme liver resection for hepatoblastoma in proximity to the major liver vasculature
Xiu WL, Liu J, Zhang JL, Wang JM, Wang XF, Wang FF, Mi J, Hao XW, Xia N, Dong Q
- 1078 Research on the prognostic value of adjusting intraperitoneal three-dimensional quality evaluation mode in laparoscopic cholecystectomy patients
Zhou Y, Chen ZQ

- 1087** Construction of a predictive model for acute liver failure after hepatectomy based on neutrophil-to-lymphocyte ratio and albumin-bilirubin score

Li XP, Bao ZT, Wang L, Zhang CY, Yang W

- 1097** Predicting short-term thromboembolic risk following Roux-en-Y gastric bypass using supervised machine learning

Ali H, Inayat F, Moond V, Chaudhry A, Afzal A, Anjum Z, Tahir H, Anwar MS, Dahiya DS, Afzal MS, Nawaz G, Sohail AH, Aziz M

- 1109** Comparative analysis of two digestive tract reconstruction methods in total laparoscopic radical total gastrectomy

Dong TX, Wang D, Zhao Q, Zhang ZD, Zhao XF, Tan BB, Liu Y, Liu QW, Yang PG, Ding PA, Zheng T, Li Y, Liu ZJ

- 1121** Incidence of surgical site infection in minimally invasive colorectal surgery

Ni LT, Zhao R, Ye YR, Ouyang YM, Chen X

Observational Study

- 1130** Burden of gallstone disease in the United States population: Prepandemic rates and trends

Unalp-Arida A, Ruhl CE

Prospective Study

- 1149** Kuicolong-yu enema decoction retains traditional Chinese medicine enema attenuates inflammatory response ulcerative colitis through TLR4/NF- κ B signaling pathway

Han L, Tang K, Fang XL, Xu JX, Mao XY, Li M

SYSTEMATIC REVIEWS

- 1155** Quality-adjusted life years and surgical waiting list: Systematic review of the literature

de la Plaza Llamas R, Ortega Azor L, Hernández Yuste M, Gorini L, Latorre-Fragua RA, Díaz Candelas DA, Al Shwely Abduljabar F, Gemio del Rey IA

META-ANALYSIS

- 1165** Impact of different anastomosis methods on post-recurrence after intestinal resection for Crohn's disease: A meta-analysis

Wang ZZ, Zhao CH, Shen H, Dai GP

CASE REPORT

- 1176** Suspected coexistence of perianal necrotizing sweet syndrome in chronic myelomonocytic leukemia: A case report

Yu KQ, Li HX, Wu J

- 1184** Successful splenic artery embolization in a patient with Behçet's syndrome-associated splenic rupture: A case report

Zhu GZ, Ji DH

- 1189** Stercoral perforation of the cecum: A case report

Yu HC, Pu TW, Kang JC, Chen CY, Hu JM, Su RY

- 1195** Percutaneous transhepatic stenting for acute superior mesenteric vein stenosis after pancreaticoduodenectomy with portal vein reconstruction: A case report
Lin C, Wang ZY, Dong LB, Wang ZW, Li ZH, Wang WB
- 1203** Endoscopic treatment of bleeding gastric ulcer causing gastric wall necrosis: A case report
Li WF, Gao RY, Xu JW, Yu XQ
- 1208** Intermittent melena and refractory anemia due to jejunal cavernous lymphangioma: A case report
Liu KR, Zhang S, Chen WR, Huang YX, Li XG

LETTER TO THE EDITOR

- 1215** New frontiers in ectopic pancreatic tissue management
Covantsev S

ABOUT COVER

Editorial Board Member of *World Journal of Gastrointestinal Surgery*, Arunachalam Rathnaswami, FACS, MBBS, MS, MNAMS, MCh, Ex-Professor and Head, Department of Surgical Gastroenterology, SRM Medical College Hospital and Research Institute SRMIST, Kattankulathur, Chennai, TamilNadu 603203, India. arunarathna@gmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Gastrointestinal Surgery* (WJGS, *World J Gastrointest Surg*) is to provide scholars and readers from various fields of gastrointestinal surgery with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGS mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal surgery and covering a wide range of topics including biliary tract surgical procedures, biliopancreatic diversion, colectomy, esophagectomy, esophagostomy, pancreas transplantation, and pancreatectomy, *etc.*

INDEXING/ABSTRACTING

The WJGS is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports/Science Edition, PubMed, PubMed Central, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJGS as 2.0; IF without journal self cites: 1.9; 5-year IF: 2.2; Journal Citation Indicator: 0.52; Ranking: 113 among 212 journals in surgery; Quartile category: Q3; Ranking: 81 among 93 journals in gastroenterology and hepatology; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Zi-Hang Xu, Production Department Director: Xiang Li, Cover Editor: Jia-Ru Fan.

NAME OF JOURNAL

World Journal of Gastrointestinal Surgery

ISSN

ISSN 1948-9366 (online)

LAUNCH DATE

November 30, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Peter Schemmer

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/1948-9366/editorialboard.htm>

PUBLICATION DATE

April 27, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Impact of different anastomosis methods on post-recurrence after intestinal resection for Crohn's disease: A meta-analysis

Zheng-Zuo Wang, Chun-Hua Zhao, Hui Shen, Gui-Ping Dai

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): C, C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Hawkins J, United States; Triggi M, Greece

Received: December 22, 2023

Peer-review started: December 22, 2023

First decision: January 9, 2024

Revised: January 18, 2024

Accepted: March 28, 2024

Article in press: March 28, 2024

Published online: April 27, 2024



Zheng-Zuo Wang, Department of Anorectal, Jiashan Hospital of Traditional Chinese Medicine, Jiaxing 314100, Zhejiang Province, China

Chun-Hua Zhao, Department of Oncology, Jiashan Hospital of Traditional Chinese Medicine, Jiaxing 314100, Zhejiang Province, China

Hui Shen, Department of Gastroenterology, Jiashan Hospital of Traditional Chinese Medicine, Jiaxing 314100, Zhejiang Province, China

Gui-Ping Dai, Department of Oncology, The Second Hospital of Jiaxing, Jiaxing 314000, Zhejiang Province, China

Corresponding author: Hui Shen, MBBS, Attending Doctor, Department of Gastroenterology, Jiashan Hospital of Traditional Chinese Medicine, No. 38 Gujiadai, Jiashan County, Jiaxing 314100, Zhejiang Province, China. 13758305062@163.com

Abstract

BACKGROUND

Crohn's disease (CD) is a chronic inflammatory disease of the gastrointestinal tract, often requiring intestinal resection as a common treatment. However, recurrence after surgery is common. The anastomotic configuration after bowel resection appears to be associated with the recurrence of CD. Previous studies have suggested that the Kono-S anastomosis may help to reduce the recurrence rate. However, the results remain controversial. Therefore, evidence-based evidence is needed to prove the advantages of Kono-S anastomosis.

AIM

To measure the influence of anastomosis techniques on the long-term relapse rate of CD by conducting a meta-analysis.

METHODS

PubMed, Scopus, and Cochrane Library were searched until October 8, 2023. Patients who underwent intestinal resection due to CD were included. The intervention measures included Kono-S anastomosis, whereas the control group received traditional anastomosis such as end-to-end, end-to-side, and side-to-side anastomosis. Only randomized clinical trials and observational studies were included. The primary outcome measures were hospital stay post-surgery, overall postoperative complication incidence, the proportion of Clavien-Dindo grade IIIa or higher, overall postoperative recurrence rate, and Rutgeerts score.

RESULTS

From 2011 to 2023, six articles met the inclusion and exclusion criteria. The results indicated that Kono-S anastomosis can reduce the hospital stay post-surgery of patients with CD [MD = -0.26, 95%CI: -0.42 to -0.10, $P = 0.002$] than other traditional anastomosis methods. Compared to other traditional anastomosis methods, Kono-S anastomosis can significantly reduce the total recurrence rate [MD = 0.40, 95%CI: 0.17 to 0.98, $P = 0.05$] and postoperative Rutgeerts score [MD = -0.81, 95%CI: -0.96 to -0.66, $P < 0.001$] in patients with CD. However, there is no significant disparity in the overall occurrence of postoperative complications and the proportion of Clavien-Dindo \geq IIIa.

CONCLUSION

Kono-S anastomosis has the potential to expedite the recuperation of CD and diminish relapse hazards; however, additional larger trials are necessary to authenticate its effectiveness.

Key Words: Kono-S; Crohn's disease; Traditional anastomosis; Postoperative recurrence; Meta-analysis

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: There is no agreement in academic studies regarding the impact of Kono-S and traditional anastomosis on patients' recurrence after Crohn's surgery. Here, we present a primary inclusive meta-analysis exploring the effect of Kono-S and traditional anastomosis on the recurrence of Crohn's disease following surgery. The hospital stay post-surgery, the incidence of postoperative complications, the proportion of Clavien-Dindo \geq IIIa, the postoperative recurrence rate, and the Rutgeerts score were examined in patients who underwent the two techniques.

Citation: Wang ZZ, Zhao CH, Shen H, Dai GP. Impact of different anastomosis methods on post-recurrence after intestinal resection for Crohn's disease: A meta-analysis. *World J Gastrointest Surg* 2024; 16(4): 1165-1175

URL: <https://www.wjgnet.com/1948-9366/full/v16/i4/1165.htm>

DOI: <https://dx.doi.org/10.4240/wjgs.v16.i4.1165>

INTRODUCTION

Crohn's disease (CD) is a chronic granulomatous enteritis that can affect any part of the digestive tract, with a particular predilection to the terminal ileum and cecum[1]. The disease is characterized by clinical features, such as abdominal pain, diarrhea, weight loss, and recurrent episodes, which could result in severe malnutrition in patients[2]. In Europe, the annual incidence rate of CD stands at 12.7 cases/100000 individuals, while in North America, it is nearly 20.2 cases/100000 individuals. Furthermore, the incidence rate has demonstrated a consistent upward trend in recent years[3]. Recent research has indicated that CD pathogenesis is linked to several factors, such as genetic susceptibility, imbalanced intestinal microbial communities, environmental factors, abnormal immune regulation, and excessive expression of inflammatory mediators and cytokines[4-6]. Despite this evidence, the pathogenesis remains unclear. Currently, there is no definitive cure for CD. Drug therapy involves mainly glucocorticoids, immunosuppressants, and other medications, yet long-term adherence proves challenging and severe side effects may occur. Consequently, surgical resection of the affected intestinal segment is required for patients with severe symptoms, significant complications, or inadequate drug response[7,8]. Although surgical resection can efficiently manage symptoms, high postoperative recurrence rates can significantly affect patients' quality of life[9]. Statistics show that in the absence of drug treatment, endoscopic examination following ileocolic resection reveals endoscopic recurrence in approximately 65%-90% of patients within 12 months, and this proportion can increase to 80%-100% within three years[10].

Intestinal anastomosis is an essential determinant of CD recurrence after surgery[11]. The traditional techniques of anastomosis, including end-to-end anastomosis (EEA), end-to-side anastomosis (ESA), and side-to-side anastomosis (SSA), are widely used due to their ease of operation and high repeatability[12]. Recently, there have been several reports on a new anastomosis technique, known as Kono-S anastomosis, which aims to decrease the likelihood of postoperative recurrence in patients. This anastomosis technique excludes the mesentery from the anastomosis, limits lumen deformation, and preserves innervation and angiogenesis. It has demonstrated impressive safety and disease recurrence prevention results[13]. In recent years, multiple small-scale randomized controlled trials and cohort studies have explored the effects of the two anastomosis techniques on the recurrence rate of CD. However, the findings are inconclusive, with some studies indicating that Kono-S anastomosis may be superior to the traditional method, whilst others revealing no significant difference between the two techniques[12,14,15]. Currently, evidence-based medical data from large sample sizes to assess the impact of the two anastomosis methods on the long-term recurrence rate of CD remains insufficient. Therefore, we aimed to conduct a thorough meta-analysis of the existing research to measure the influence of both anastomosis techniques on the long-term relapse rate of CD to provide reliable, evidence-based medical evidence for clinical use. Simultaneously, we sought to clarify the application value of Kono-S anastomosis.

MATERIALS AND METHODS

Literature search

To gather additional experimental evidence, we conducted independent searches on PubMed, EMBASE, and Cochrane Library databases. Our selection criteria included articles with human participants, reported in English. We imposed no restrictions on publication time and type of articles. For the intervention group, we used Kono-S anastomosis whereas traditional anastomosis was used as the control group, following the PICOS scheme. The participants consisted of patients who underwent intestinal resection due to CD. We employed a combination of subject words and free words for our retrieval strategy. The search formula of the PubMed database was: (((('Crohn Disease' [Mesh]) OR ((Crohn's Enteritis) OR (Crohn's Disease))) AND ((postoperative) OR (intestinal resection))) AND (((Kono-S anastomosis) OR (Kono-S Anastomosis)) OR (anastomosis)).

The following combination was used for the EMBASE database: (1) "crohn disease"/ exp OR "crohn disease"; (2) postoperative; (3) intestinal resection; (4) Kono-S anastomosis; (5) Kono-S Anastomosis; (6) anastomosis; (7) 4 OR 5; (8) 2 OR 3; and (9) 1 AND 7 AND 8.

The same keywords were inserted into the Cochrane Library, which used the following search strategy: (1) MeSH descriptor: (Crohn Disease) explode all trees; (2) (Kono-S anastomosis): ti, ab, kw; (3) (Kono-S Anastomosis): ti, ab, kw; (4) (Anastomosis): ti, ab, kw; (5) 2 OR 3 OR 4; (6) (postoperative): ti, ab, kw; (7) (intestinal resection): ti, ab, kw; (8) 6 OR 7; and (9) 1 AND 5 AND 8.

In the field of management, the search scope is expanded through extensive cross-checking of the reference lists of all articles that meet the inclusion criteria. The search deadline for all databases was October 8, 2023. This study was conducted based on the PRISMA 2020 statement guidelines, with complete reporting of the meta-analysis data^[16].

Study selection

The primary studies' titles and abstracts were screened independently by the same two authors. Employing Cohen's kappa statistic, the authors evaluated their agreement during the initial article screening. Non-conforming and duplicate articles were excluded after reviewing the topic, abstract, and keywords. The meta-analysis applied the following criteria: a clear, comprehensive, and objective style, simple language, consistent citation, standard structure, formal register, meticulous and precise vocabulary, and grammatical accuracy.

The inclusion criteria consisted of the following: (1) Studies evaluating patients who had undergone intestinal resection due to CD; (2) studies in which the intervention measure used was Kono-S anastomosis, while the control group received traditional anastomosis (including EEA, ESA, and SSA); (3) studies where relevant data for extraction were available in the results report; (4) randomized clinical trials (RCTs) and observational studies; and (5) studies evaluating the influence of Kono-S anastomosis on the occurrence of CD recurrence after intestinal resection. Review articles, letters, comments, case reports, as well as studies that did not evaluate postoperative recurrence indicators, were excluded from this analysis.

Data extraction

We examined all pertinent literature, tables, and charts and extracted relevant data as shown in Tables 1 and 2. The primary contents encompassed the following: (1) The name of the primary author, their country, publication date, and research type; (2) fundamental clinical data for patients with CD in the Kono-S anastomosis group and the traditional anastomosis group; and (3) postoperative observation measures, such as the hospital stay post-surgery, total incidence of postoperative complications, Clavien-Dindo scores \geq IIIa (%), total recurrence rate, and Rutgeerts score^[17]. If further information was required, we reached out to the corresponding author of the paper for assistance *via* email. Furthermore, any discrepancies among the reviewers were resolved through sound and sensible discussion.

Risk of bias

The Cochrane Risk Bias Assessment Tool was implemented to gauge bias risks in RCTs. The potential for bias was evaluated by examining factors, such as random sequence generation, allocation concealment, blinding, completeness of outcome data, selective reporting, and other possible sources of bias. Based on three aspects: Selectivity, comparability, and outcome, the Newcastle-Ottawa Scale (NOS) score was employed to assess the quality of cohort studies. The funnel plot was utilized to assess the prospect of publication bias by analyzing the scatter points' distribution.

Statistical analysis

RevMan 5.3 was utilized to perform a meta-analysis. The OR and the average difference between groups (MD) were employed as summary measures of dichotomous and continuous variables, respectively. The 95%CI was applied to both methods. In cases where studies provided median and quartile ranges, the average SD was calculated^[18]. Heterogeneity was evaluated using the I^2 test. If $I^2 < 50\%$ and $P > 0.1$, we ascertained that heterogeneity among the studies was low; thus, a fixed-effects model was applied. However, if $P < 0.1$ and $I^2 > 50\%$, we determined that heterogeneity among the studies was high, and a random-effects model was employed for analyzing the sources of heterogeneity. To assess publication bias, we generated funnel plots. We conducted sensitivity analyses using the one-by-one exclusion method. If there was a significant change in the I^2 value after removing an item, the study results were regarded as the origin of heterogeneity. All statistical analysis was significant at the $P < 0.05$ level.

Table 1 Characteristics of the included studies

Ref.	Country	Type of study	Total number of patients	Kono-S anastomosis			Traditional anastomosis		
				No. of patients	Age	M/F	No. of patients	Age	M/F
Shimada <i>et al</i> [15], 2019	Japan	Retro/cohort study	215	117	39.00 ± 3.13	84/33	98	35.06 ± 11.29	74/24
Tyrode <i>et al</i> [19], 2023	France	Retro/cohort study	85	30	36.11 ± 14.01	13/17	55	33.83 ± 12.79	24/31
Kelm <i>et al</i> [20], 2022	Germany	Retro/cohort study	51	22	38.41 ± 11.00	14/8	29	38.42 ± 13.57	14/15
Alibert <i>et al</i> [21], 2023	France	Retro/cohort study	432	155	38.20 ± 13.90	79/76	277	37.60 ± 14.20	107/170
Kono <i>et al</i> [22], 2011	Japan	Retro/cohort study	142	69	32.36 ± 9.09	57/12	73	29.38 ± 10.06	58/15
Luglio <i>et al</i> [23], 2020	Italy	RCT	79	36	34.75 ± 5.91	18/18	43	43.10 ± 7.55	22/21

Retro: Retrospective; RCT: Randomized clinical trial.

Table 2 Raw data of each included study

Ref.	Hospital stay (d)	Postoperative morbidity (%)	Recurrence rate (%)	Mean Rutgeerts score	Clavien-Dindo ≥ IIIa (%)
Shimada <i>et al</i> [15]					
Kono-S anastomosis	NR	64/117(54.70)	4/117 (3.4)	NR	NR
Traditional anastomosis	NR	68/98(69.39)	24/98 (24.5)	NR	NR
Tyrode <i>et al</i> [19]					
Kono-S anastomosis	6.71 ± 1.56	2/30(6.66)	17/30 (56.66)	NR	5/30 (16.67)
Traditional anastomosis	7.00 ± 1.52	6/55(10.90)	27/55 (49.10)	NR	9/55 (16.36)
Kelm <i>et al</i> [20]					
Kono-S anastomosis	8.1 ± 2.34	15/22(68.19)	7/22 (31.82)	1.70 ± 0.32	3/22 (13.64)
Traditional anastomosis	8.1 ± 2.56	9/29(31.03)	13/29 (44.83)	2.50 ± 0.86	4/29 (13.79)
Alibert <i>et al</i> [21]					
Kono-S anastomosis	6.20 ± 4.00	70/155(45.16)	NR	NR	7/155 (4.52)
Traditional anastomosis	7.80 ± 5.60	79/277(28.52)	NR	NR	21/277 (7.58)
Kono <i>et al</i> [22]					
Kono-S anastomosis	NR	3/69(4.34)	49/69 (71.01)	2.58 ± 0.63	NR
Traditional anastomosis	NR	7/73(9.59)	58/73 (79.45)	3.34 ± 0.42	NR
Luglio <i>et al</i> [23]					
Kono-S anastomosis	7.00 ± 3.00	16.67	8/36 (22.22)	1.05 ± 1.06	NR
Traditional anastomosis	7.60 ± 3.08	25.58	27/43 (62.79)	2.30 ± 1.32	NR

NR: Not reported.

RESULTS

Study selection

A total of 1328 articles were collected. Among them, 215 were repetitive and therefore removed. We then analyzed the titles and abstracts of the remaining 1113 articles. Among these, 76 articles were meta-analyses, 181 were review articles, and 850 did not involve the experimental comparison of Kono-S and traditional anastomosis. Finally, we chose six articles [15,19-23] for a full-text review, from which, we extracted all of the necessary literature data. A meta-analysis of six articles was conducted, all published between 2011 and 2023. The studies consisted of 1 randomized controlled trial and 5 retrospective cohort studies (Figure 1), collectively involving 1004 patients with CD. Among these patients, 429 underwent Kono-S anastomosis and 575 underwent traditional anastomosis (such as EEA, ESA, and SSA). Table 1 summarizes the features of the included studies.

Literature quality evaluation

The evaluation of RCT literature quality was conducted based on several criteria, such as random number generation, allocation blindness, researcher and subject blindness, result evaluation blindness, result integrity, reporting bias, and other factors. Based on the Cochrane risk bias assessment tool, the included randomized controlled study [23] exhibited a low risk in random sequence generation; nevertheless, the blind allocation component was unclear. Minimal risk was observed in incomplete results data, as well as in selective results reporting. Furthermore, other potential bias factors meeting the requirements of this article were assessed as low risk. The literature's quality was evaluated concerning selection bias, comparability bias, and result evaluation bias according to the NOS scale. A research quality score of 9 was achieved by one paper [15], while three papers [19,20,22] received a research quality score of 8. Additionally, one paper [21] secured a research quality score of 7. Table 3 summarizes the quality evaluation of all retrospective cohort studies. There was no disagreement between the two authors regarding the eligibility of the full-text articles (Cohen's kappa = 1).

Hospital stay post-surgery

The impact of alternative anastomosis methods (Kono-S and traditional) on the hospital stay following intestinal resection for CD was evaluated using a meta-analysis. The fixed effect model indicated moderate heterogeneity among the included literature, with $I^2 = 51\%$ and $P = 0.73$. However, the use of the random effect model revealed no heterogeneity in the included studies with $I^2 = 0\%$ and $P = 0.73$. The study findings indicated that utilization of the Kono-S anastomosis approach led to a statistically significant reduction in postoperative hospitalization duration among patients with CD, as compared with traditional anastomosis methods [MD = -0.26, 95%CI: -0.42 to -0.10, $P = 0.002$, Figure 2A].

Total incidence of postoperative complications

The impact of various anastomosis methods (Kono-S and traditional), post intestinal resection, on the overall incidence of postoperative complications in CD was evaluated through a meta-analysis. The analysis revealed a high degree of heterogeneity among the included literature in the fixed effect model ($I^2 = 79\%$, $P = 0.02$). Notably, even after using the random effect model, the included literature remained significantly heterogeneous ($I^2 = 79\%$, $P = 0.02$). After conducting sensitivity analysis by gradually excluding included literature, we found that Alibert *et al*'s study had the most significant impact on the outcomes [21], potentially serving as the primary source of heterogeneity. The findings indicated that the Kono-S anastomosis method has minimal effect on the total incidence of postoperative complications in patients with CD compared with other traditional anastomosis techniques. However, these results did not achieve statistical significance [MD = 1.00, 95%CI: 0.46 to 2.20, $P = 0.99$ Figure 2B].

Proportion of Clavien-Dindo greater than IIIa

No heterogeneity was observed among the included studies, with $I^2 = 0\%$ and $P = 0.40$ under the fixed effect model. The study findings indicate that the Kono-S anastomosis method did not significantly affect the proportion of Clavien-Dindo \geq IIIa in patients with CD, compared with other conventional anastomosis techniques [MD = 0.82, 95%CI: 0.43 to 1.54, $P = 0.53$, Figure 2C].

Total postoperative recurrence rate

The impact of Kono-S and traditional anastomosis methods on the overall recurrence rate of CD following intestinal resection was analyzed through a meta-analysis. The fixed effect model revealed a high level of heterogeneity amongst the included studies, with $I^2 = 76\%$ and $P = 0.002$. Subsequently, the random effect model also indicated a high level of heterogeneity among the included literature. Sensitivity analysis revealed that the studies conducted by Shimada *et al* [15] and Luglio *et al* [23] had the most significant influence on the results, possibly originating as the primary source of heterogeneity. The study results demonstrate that Kono-S anastomosis yields a statistically significant reduction in the total recurrence rate of patients with CD compared with other traditional anastomosis methods [MD = 0.40, 95%CI: 0.17 to 0.98, $P = 0.05$, Figure 2D].

Rutgeerts score

A meta-analysis indicated low heterogeneity ($I^2 = 34\%$, $P = 0.22$) using the fixed-effect model. The study results indicate that Kono-S anastomosis is a more effective method in reducing postoperative Rutgeerts scores for patients with CD compared to other conventional anastomosis techniques. This difference was statistically significant [MD = -0.81, 95%CI: -0.96 to -0.66, $P < 0.001$, Figure 2E].

Table 3 Quality of the included studies based on the Newcastle-Ottawa scale

Ref.	Representativeness of the exposed cohort	Selection of the non-exposed cohort	Ascertainment of exposure	Demonstration that outcome of interest was not present at the start of the study	Comparability of cohorts based on the design or analysis	Assessment of outcome	Was follow-up long enough for outcomes to occur	Adequacy of follow-up of cohorts	Total scores
Shimada <i>et al</i> [15]	1	1	1	1	2	1	1	1	9
Tyrode <i>et al</i> [19]	1	1	1	0	2	1	1	1	8
Kelm <i>et al</i> [20]	1	1	1	1	1	0	1	1	8
Alibert <i>et al</i> [21]	1	1	1	1	1	1	1	0	7
Kono <i>et al</i> [22]	1	1	1	0	2	1	1	1	8

Evaluation of literature publication bias

A funnel plot was created for each impact indicator identified within the six included articles. Continuous variables for the length of hospital stay post-surgery and Rutgeerts score were analyzed, along with binary variables for the total incidence of postoperative complications, proportion of Clavien-Dindo ≥ IIIa, and total postoperative recurrence rate. Technical abbreviations were explained upon first use. The funnel plot meta-analysis demonstrated that hospital stay post-surgery, proportion of Clavien-Dindo ≥ IIIa, and Rutgeerts score were all aligned using a vertical line. The research was evenly dispersed on both sides of the center line, creating a pyramidal dotted line range, with no evident separation phenomenon (Figure 3; no conspicuous bias). However, the meta-analysis funnel plot of the remaining impact indicators exhibited bias, possibly due to inadequate literature included in this study.

DISCUSSION

Since its introduction by Kono *et al*[22] in 2003, the efficacy of the EEA technique for anti-mesenteric function has been a subject of debate. The Kono-S anastomosis procedure involves employing a linear cutter to section the intestine, after which the intestinal ends are sutured together to form a support column. Subsequently, a functional end-to-end manual suture is performed[24,25]. Kono *et al*[22] posited that anastomosis effectively prevents postoperative recurrence in patients with CD. Historical data comparison revealed that the endoscopic recurrence rate of patients who underwent Kono-S anastomosis was lower than those who underwent traditional anastomosis. However, the persuasiveness of a single study is limited. Therefore, this study used meta-analysis to compare and analyze the impact of Kono-S anastomosis and traditional anastomosis on postoperative recurrence in patients with CD.

The meta-analysis results of this study showed that Kono-S anastomosis could shorten the hospital stay of patients with CD, compared with other traditional anastomosis methods [MD = -0.26, 95%CI: -0.42 to -0.10, *P* = 0.002]. Notably, no heterogeneity was observed in the included literature. Many studies evaluating recurrence factors tend to overlook the




Figure 1 PRISMA flow diagram.

postoperative hospitalization time, which can indicate the patient's recovery rate. Regarding surgical operations, Kono-S anastomosis preserves a portion of the mesenteric blood perfusion, potentially resulting in a reduction of ischemic injury at the anastomosis[11,22]. Adequate blood supply creates favorable biological circumstances for anastomotic healing, consequently lowering the risk of early complications like anastomotic leakage and stenosis[26]. This in turn avoids additional surgery and extended hospital stays caused by complications. Furthermore, Kono-S anastomosis optimizes the anatomical position and three-dimensional structure of the intestinal segment, promoting swift recovery of postoperative intestinal peristalsis function, early elimination, and bowel movements in patients, and faster rehabilitation. These factors may explain why Kono-S anastomosis is advantageous for hastening the postoperative recovery of patients with CD.

Regarding postoperative complications in these patients, the number of incidences, such as intestinal obstruction, anastomotic leakage, infection, and fever were tallied for the purposes of comparing the safety of the two anastomosis methods. The outcomes indicated that the Kono-S anastomosis had no considerable impact on the overall occurrence of postoperative complications in patients with CD when compared to the traditional anastomosis method [MD = 1.00, 95%CI: 0.46 to 2.20, $P = 0.99$]. However, this meta-analysis has a relatively small sample size, as only 6 studies, comprising a total of 1004 patients, were included. There is some degree of uncertainty, making it necessary to interpret the results with caution. Additionally, the incidence of complications is correlated with various factors, including the surgical procedure, perioperative treatment, and the patient's underlying health condition[27]. Therefore, due to these confounding factors, it is difficult to show the advantages and disadvantages of the two anastomosis methods just by comparing the number of complications in this study. Therefore, we further analyzed the Clavien-Dindo score, which describes the severity of complications. A higher proportion of Clavien-Dindo \geq IIIa indicates a greater number of severe complications. The study findings demonstrated that Kono-S anastomosis did not significantly impact the proportion of Clavien-Dindo \geq IIIa in patients with CD compared to other conventional methods of anastomosis [MD = 0.82, 95%CI: 0.43 to 1.54, $P = 0.53$]. While there were no notable variations observed in the incidence of complications or the percentage of severe cases between the groups, Kono-S anastomosis has been assessed for its safety and feasibility as a new surgical technique [28,29]. Further studies are required to confirm its advantages. It is imperative to implement techniques, such as increasing the sample size, rigorous control over confounding variables, and analysis of long-term outcomes.

Regarding the postoperative recurrence rate of patients with CD, the study findings demonstrate that the Kono-S anastomosis method resulted in lower total recurrence rates compared to other traditional methods [MD = 0.40, 95%CI: 0.17 to 0.98, $P = 0.05$]. Endoscopic anastomotic recurrence is typically evaluated through the Rutgeerts score (\geq IIIa) at 6 to 12 months. Therefore, we performed a meta-analysis of Rutgeerts score. Our findings indicate that Kono-S anastomosis can significantly reduce Rutgeerts score in patients with CD following surgery when compared to traditional anastomosis [MD = -0.81, 95%CI: -0.96 to -0.66, $P < 0.001$]. These findings align with those of a meta-analysis published in 2020, which showed that Kono-S anastomosis presents a reduced risk of recurrence in comparison to traditional anastomosis[30]. Although the sample size for this study is limited, the outcomes are clinically significant as postoperative recurrence directly impacts patients' life quality and prognosis. On the contrary, Kono-S anastomosis could enhance healing conditions at the anastomosis site by preserving a section of mesenteric blood flow and sustaining the physiological and anatomical structure of the intestine, thus mitigating the risk of inflammation and recurrence[31]. Conversely, Kono-S anastomosis can create a broad lumen and a supportive column at the anastomotic site, preserving a three-dimensional configuration. This is a crucial factor in preventing anastomotic stenosis[12,22] and preventing distortion while keeping




Figure 2 Meta-analysis forest plot. A: Hospital stay; B: Total incidence of postoperative complications; C: Proportion of Clavien-Dindo greater than IIIa; D: Total postoperative recurrence rate; E: Rutgeerts score.

the intestinal tract straight, which facilitates postoperative endoscopy. However, the conventional U-shaped EEA anastomosis is not suitable for endoscopic observation or balloon dilation. As a result, the Kono-S anastomosis structure surpasses the traditional EEA anastomosis structure in facilitating postoperative follow-up and treatment of anastomotic problems[22,32].

The limitations of this study stem from its small sample size and sole location of data collection at a single center. We anticipate the need for more studies of high quality to verify this outcome and establish the precise mechanism by which Kono-S anastomosis diminishes recurrence. It is important to recognize that the effectiveness of this surgical approach is contingent upon rigorous regulation of operative procedures and the selection of suitable candidates. Future research must develop a uniform training program for Kono-S anastomosis and delineate the patient cohorts who can gain the maximum benefit from this technique.




Figure 3 Literature publication bias evaluation chart. A: Hospital stay; B: Total incidence of postoperative complications; C: Proportion of Clavien-Dindo greater than IIIa; D: Total postoperative recurrence rate; E: Rutgeerts score.

CONCLUSION

To summarize, this study showed that Kono-S anastomosis is a suitable technique for patients with CD. It has proven to speed up postoperative recovery and is particularly effective in reducing long-term recurrence when compared to other conventional anastomosis methods. While this study has limitations, including a limited number of included studies and sample size, the results still provide robust support for the clinical application of Kono-S anastomosis. It is recommended that clinicians select the appropriate anastomotic technique based on the specific conditions of patients, and we anticipate further validation of the long-term efficacy of Kono-S through larger sample studies in the future.

FOOTNOTES

Co-corresponding authors: Hui Shen and Gui-Ping Dai.

Author contributions: Wang ZZ designed and performed the research; Shen H and Dai GP designed the research and guided and supervised the report; Zhao CH collected information and contributed to the analysis; all authors approved the final manuscript. Shen H and Dai GP contributed equally to this work as co-corresponding authors. To justify the designation of Shen H and Dai GP as co-corresponding authors, we offer the following reasons: Firstly, as integral leaders in our interdisciplinary project, Shen H guided research design and analysis, while Dai GP contributed to research and result interpretation, collectively driving the study's success. Secondly, their coordination bridged diverse expertise, enhancing team collaboration and fostering an innovative research environment, underscoring the value of co-authorship. Lastly, both authors made substantial, equal contributions across all research phases, warranting their joint recognition as co-corresponding authors. In brief, Shen H and Dai GP's co-authorship accurately captures their shared responsibilities and contributions, reflecting our team's collaborative ethos and interdisciplinary approach.

Conflict-of-interest statement: Dr. Shen has nothing to disclose.

PRISMA 2009 Checklist statement: The authors have read the PRISMA 2020 Checklist of items, and the manuscript was prepared and revised according to the PRISMA 2020 Checklist.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/Licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Zheng-Zuo Wang 0009-0006-4543-9801; Chun-Hua Zhao 0009-0005-6535-4242; Hui Shen 0009-0004-6249-8251; Gui-Ping Dai 0009-0007-6685-5043.

S-Editor: Lin C

L-Editor: A

REFERENCES

- Freeman HJ. Natural history and long-term clinical course of Crohn's disease. *World J Gastroenterol* 2014; **20**: 31-36 [PMID: 24415855 DOI: 10.3748/wjg.v20.i1.31]
- Torres J, Mehandru S, Colombel JF, Peyrin-Biroulet L. Crohn's disease. *Lancet* 2017; **389**: 1741-1755 [PMID: 27914655 DOI: 10.1016/S0140-6736(16)31711-1]
- Ng SC, Shi HY, Hamidi N, Underwood FE, Tang W, Benchimol EI, Panaccione R, Ghosh S, Wu JCY, Chan FKL, Sung JY, Kaplan GG. Worldwide incidence and prevalence of inflammatory bowel disease in the 21st century: a systematic review of population-based studies. *Lancet* 2017; **390**: 2769-2778 [PMID: 29050646 DOI: 10.1016/S0140-6736(17)32448-0]
- Khanna S, Raffals LE. The Microbiome in Crohn's Disease: Role in Pathogenesis and Role of Microbiome Replacement Therapies. *Gastroenterol Clin North Am* 2017; **46**: 481-492 [PMID: 28838410 DOI: 10.1016/j.gtc.2017.05.004]
- Petagna L, Antonelli A, Ganini C, Bellato V, Campanelli M, Divizia A, Efrati C, Franceschilli M, Guida AM, Ingallinella S, Montagnese F, Sensi B, Siragusa L, Sica GS. Pathophysiology of Crohn's disease inflammation and recurrence. *Biol Direct* 2020; **15**: 23 [PMID: 33160400 DOI: 10.1186/s13062-020-00280-5]
- Ballester Ferré MP, Boscá-Watts MM, Mínguez Pérez M. Crohn's disease. *Med Clin (Barc)* 2018; **151**: 26-33 [PMID: 29246562 DOI: 10.1016/j.medcli.2017.10.036]
- Seyedian SS, Nokhostin F, Malamir MD. A review of the diagnosis, prevention, and treatment methods of inflammatory bowel disease. *J Med Life* 2019; **12**: 113-122 [PMID: 31406511 DOI: 10.25122/jml-2018-0075]
- Rodríguez-Lago I, Gisbert JP. The Role of Immunomodulators and Biologics in the Medical Management of Stricturing Crohn's Disease. *J Crohns Colitis* 2020; **14**: 557-566 [PMID: 31541235 DOI: 10.1093/ecco-jcc/jjz158]
- Barnes EL. Postoperative Crohn's disease management. *Curr Opin Gastroenterol* 2020; **36**: 277-283 [PMID: 32304384 DOI: 10.1097/MOG.0000000000000638]
- Cottone M, Mocciano F, Orlando A. Infliximab prevents Crohn's disease recurrence after ileal resection. *Expert Rev Gastroenterol Hepatol* 2009; **3**: 231-234 [PMID: 19485805 DOI: 10.1586/egh.09.13]
- Meima-van Praag EM, Buskens CJ, Hompes R, Bemelman WA. Surgical management of Crohn's disease: a state of the art review. *Int J Colorectal Dis* 2021; **36**: 1133-1145 [PMID: 33528750 DOI: 10.1007/s00384-021-03857-2]
- Bislenghi G, Devriendt S, Wolthuis A, D'Hoore A. Totally stapled Kono-S anastomosis for Crohn's disease. *Tech Coloproctol* 2022; **26**: 213-216 [PMID: 35088189 DOI: 10.1007/s10151-021-02551-3]
- Dasharathy SS, Limketkai BN, Sauk JS. What's New in the Postoperative Management of Crohn's Disease? *Dig Dis Sci* 2022; **67**: 3508-3517 [PMID: 34406585 DOI: 10.1007/s10620-021-07205-w]
- Horisberger K, Birrer DL, Rickenbacher A, Turina M. Experiences with the Kono-S anastomosis in Crohn's disease of the terminal ileum-a cohort study. *Langenbecks Arch Surg* 2021; **406**: 1173-1180 [PMID: 33025079 DOI: 10.1007/s00423-020-01998-6]
- Shimada N, Ohge H, Kono T, Sugitani A, Yano R, Watanabe Y, Uemura K, Murakami Y, Sueda T. Surgical Recurrence at Anastomotic Site After Bowel Resection in Crohn's Disease: Comparison of Kono-S and End-to-end Anastomosis. *J Gastrointest Surg* 2019; **23**: 312-319 [PMID: 30353491 DOI: 10.1007/s11605-018-4012-6]
- Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L, Tetzlaff JM, Akl EA, Brennan SE, Chou R, Glanville J, Grimshaw JM, Hróbjartsson A, Lalu MM, Li T, Loder EW, Mayo-Wilson E, McDonald S, McGuinness LA, Stewart LA, Thomas J, Tricco AC, Welch VA, Whiting P, Moher D. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021; **372**: n71 [PMID: 33782057 DOI: 10.1136/bmj.n71]
- Rutgeerts P, Geboes K, Vantrappen G, Beyls J, Kerremans R, Hiele M. Predictability of the postoperative course of Crohn's disease. *Gastroenterology* 1990; **99**: 956-963 [PMID: 2394349 DOI: 10.1016/0016-5085(90)90613-6]
- Luo D, Wan X, Liu J, Tong T. Optimally estimating the sample mean from the sample size, median, mid-range, and/or mid-quartile range. *Stat Methods Med Res* 2018; **27**: 1785-1805 [PMID: 27683581 DOI: 10.1177/0962280216669183]
- Tyrode G, Lakkis Z, Vernerey D, Falcoz A, Clairet V, Alibert L, Koch S, Vuitton L. KONO-S Anastomosis Is Not Superior to Conventional Anastomosis for the Reduction of Postoperative Endoscopic Recurrence in Crohn's Disease. *Inflamm Bowel Dis* 2023 [PMID: 37776562 DOI: 10.1093/ibd/izad214]
- Kelm M, Reibetanz J, Kim M, Schoettker K, Brand M, Meining A, Germer CT, Flemming S. Kono-S Anastomosis in Crohn's Disease: A Retrospective Study on Postoperative Morbidity and Disease Recurrence in Comparison to the Conventional Side-To-Side Anastomosis. *J Clin Med* 2022; **11** [PMID: 36498490 DOI: 10.3390/jcm11236915]
- Alibert L, Betton L, Falcoz A, Manceau G, Benoist S, Zerbib P, Podevin J, Maggiori L, Brouquet A, Tyrode G, Vuitton L, Vernerey D, Lefevre JH, Lakkis Z. Does KONO-S anastomosis reduce recurrence in Crohn's disease compared to conventional ileocolonic anastomosis? A nationwide propensity score-matched study from GETAID Chirurgie Group (KoCoRICCO study). *J Crohns Colitis* 2023 [PMID: 37850342 DOI: 10.1093/ecco-jcc/jjad176]
- Kono T, Ashida T, Ebisawa Y, Chisato N, Okamoto K, Katsuno H, Maeda K, Fujiya M, Kohgo Y, Furukawa H. A new antimesenteric functional end-to-end handsewn anastomosis: surgical prevention of anastomotic recurrence in Crohn's disease. *Dis Colon Rectum* 2011; **54**: 586-592 [PMID: 21471760 DOI: 10.1007/DCR.0b013e318208b90f]
- Luglio G, Rispo A, Imperatore N, Giglio MC, Amendola A, Tropeano FP, Peltrini R, Castiglione F, De Palma GD, Bucci L. Surgical Prevention of Anastomotic Recurrence by Excluding Mesentery in Crohn's Disease: The SuPREMe-CD Study - A Randomized Clinical Trial. *Ann Surg* 2020; **272**: 210-217 [PMID: 32675483 DOI: 10.1097/SLA.0000000000003821]
- Alshantti A, Hind D, Hancock L, Brown SR. The role of Kono-S anastomosis and mesenteric resection in reducing recurrence after surgery for Crohn's disease: a systematic review. *Colorectal Dis* 2021; **23**: 7-17 [PMID: 32418300 DOI: 10.1111/codi.15136]
- Fichera A, Zoccali M, Kono T. Antimesenteric functional end-to-end handsewn (Kono-S) anastomosis. *J Gastrointest Surg* 2012; **16**: 1412-1416 [PMID: 22580840 DOI: 10.1007/s11605-012-1905-7]
- Fornaro R, Caristo G, Stratta E, Oliva A, Giovinazzo D, Mascherini M, Frascio M. Crohn's disease and postoperative recurrence. The role of

- anastomotic configurations and the kono-s anastomosis. *Ann Ital Chir* 2019; **90**: 111-120 [PMID: [31182693](#)]
- 27 **Kann BR.** Anastomotic Considerations in Crohn's Disease. *Clin Colon Rectal Surg* 2023; **36**: 63-73 [PMID: [36619281](#) DOI: [10.1055/s-0042-1758770](#)]
 - 28 **Saeed M,** Algahiny AT, Elmitwalli OSMMS, Ahmed MMMRM, Juma IM. An Analysis of the Literature and a Case Study on the Successful Surgical Treatment of a Crohn's Patient Using a Kono-S Anastomosis. *Euroasian J Hepatogastroenterol* 2023; **13**: 36-39 [PMID: [37554976](#) DOI: [10.5005/jp-journals-10018-1392](#)]
 - 29 **Katsuno H,** Maeda K, Hanai T, Masumori K, Koide Y, Kono T. Novel antimesenteric functional end-to-end handsewn (Kono-S) anastomoses for Crohn's disease: a report of surgical procedure and short-term outcomes. *Dig Surg* 2015; **32**: 39-44 [PMID: [25678306](#) DOI: [10.1159/000371857](#)]
 - 30 **Ng CH,** Chin YH, Lin SY, Koh JWH, Lieske B, Koh FH, Chong CS, Foo FJ. Kono-S anastomosis for Crohn's disease: a systemic review, meta-analysis, and meta-regression. *Surg Today* 2021; **51**: 493-501 [PMID: [32894346](#) DOI: [10.1007/s00595-020-02130-3](#)]
 - 31 **Ibrahim R,** Abounozha S, Kheder A, Alawad A. Does anastomotic technique affects the recurrence rate of Crohn's disease after ileocolic resection? *Ann Med Surg (Lond)* 2021; **62**: 164-167 [PMID: [33520216](#) DOI: [10.1016/j.amsu.2021.01.027](#)]
 - 32 **Clark DA,** Smith N, Clark I, Giddings H, Rajan R. Time for an Australian and New Zealand randomized controlled trial to study the modified Kono S anastomosis. *ANZ J Surg* 2022; **92**: 3132-3134 [PMID: [36468272](#) DOI: [10.1111/ans.17991](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: office@baishideng.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

