

World Journal of *Gastrointestinal Surgery*

World J Gastrointest Surg 2024 April 27; 16(4): 974-1217



EDITORIAL

- 974 How to identify early complications in patients undergoing distal gastrectomy?
Tropeano G, Chiarello MM, Fico V, Brisinda G
- 982 Quality assessment of surgery for colorectal cancer: Where do we stand?
Morarasu S, Livadaru C, Dimofte GM
- 988 Emerging molecules, tools, technology, and future of surgical knife in gastroenterology
Kumar A, Goyal A
- 999 Carcinoembryonic antigen in the diagnosis, treatment, and follow-up of focal liver lesions
Dilek ON, Arslan Kahraman Dİ, Kahraman G
- 1008 Relationship between *Helicobacter pylori* infection and colorectal polyp/colorectal cancer
Liu Y, Yang DQ, Jiang JN, Jiao Y

MINIREVIEWS

- 1017 Near-infrared cholangiography with intragallbladder indocyanine green injection in minimally invasive cholecystectomy
Symeonidis S, Mantzoros I, Anestiadou E, Ioannidis O, Christidis P, Bitsianis S, Bisbinas V, Zapsalis K, Karastergiou T, Athanasiou D, Apostolidis S, Angelopoulos S
- 1030 Blastomas of the digestive system in adults: A review
Liu Y, El Jabbour T, Somma J, Nakanishi Y, Ligato S, Lee H, Fu ZY

ORIGINAL ARTICLE

Retrospective Study

- 1043 Single-center retrospective study of the diagnostic value of double-balloon enteroscopy in Meckel's diverticulum with bleeding
He T, Yang C, Wang J, Zhong JS, Li AH, Yin YJ, Luo LL, Rao CM, Mao NF, Guo Q, Zuo Z, Zhang W, Wan P
- 1055 Prognostic value of a nomogram model for postoperative liver metastasis of colon cancer
Cheng DX, Xu KD, Liu HB, Liu Y
- 1066 Computer-assisted three-dimensional individualized extreme liver resection for hepatoblastoma in proximity to the major liver vasculature
Xiu WL, Liu J, Zhang JL, Wang JM, Wang XF, Wang FF, Mi J, Hao XW, Xia N, Dong Q
- 1078 Research on the prognostic value of adjusting intraperitoneal three-dimensional quality evaluation mode in laparoscopic cholecystectomy patients
Zhou Y, Chen ZQ

- 1087** Construction of a predictive model for acute liver failure after hepatectomy based on neutrophil-to-lymphocyte ratio and albumin-bilirubin score

Li XP, Bao ZT, Wang L, Zhang CY, Yang W

- 1097** Predicting short-term thromboembolic risk following Roux-en-Y gastric bypass using supervised machine learning

Ali H, Inayat F, Moond V, Chaudhry A, Afzal A, Anjum Z, Tahir H, Anwar MS, Dahiya DS, Afzal MS, Nawaz G, Sohail AH, Aziz M

- 1109** Comparative analysis of two digestive tract reconstruction methods in total laparoscopic radical total gastrectomy

Dong TX, Wang D, Zhao Q, Zhang ZD, Zhao XF, Tan BB, Liu Y, Liu QW, Yang PG, Ding PA, Zheng T, Li Y, Liu ZJ

- 1121** Incidence of surgical site infection in minimally invasive colorectal surgery

Ni LT, Zhao R, Ye YR, Ouyang YM, Chen X

Observational Study

- 1130** Burden of gallstone disease in the United States population: Prepandemic rates and trends

Unalp-Arida A, Ruhl CE

Prospective Study

- 1149** Kuicolong-yu enema decoction retains traditional Chinese medicine enema attenuates inflammatory response ulcerative colitis through TLR4/NF- κ B signaling pathway

Han L, Tang K, Fang XL, Xu JX, Mao XY, Li M

SYSTEMATIC REVIEWS

- 1155** Quality-adjusted life years and surgical waiting list: Systematic review of the literature

de la Plaza Llamas R, Ortega Azor L, Hernández Yuste M, Gorini L, Latorre-Fragua RA, Díaz Candelas DA, Al Shwely Abduljabar F, Gemio del Rey IA

META-ANALYSIS

- 1165** Impact of different anastomosis methods on post-recurrence after intestinal resection for Crohn's disease: A meta-analysis

Wang ZZ, Zhao CH, Shen H, Dai GP

CASE REPORT

- 1176** Suspected coexistence of perianal necrotizing sweet syndrome in chronic myelomonocytic leukemia: A case report

Yu KQ, Li HX, Wu J

- 1184** Successful splenic artery embolization in a patient with Behçet's syndrome-associated splenic rupture: A case report

Zhu GZ, Ji DH

- 1189** Stercoral perforation of the cecum: A case report

Yu HC, Pu TW, Kang JC, Chen CY, Hu JM, Su RY

- 1195** Percutaneous transhepatic stenting for acute superior mesenteric vein stenosis after pancreaticoduodenectomy with portal vein reconstruction: A case report
Lin C, Wang ZY, Dong LB, Wang ZW, Li ZH, Wang WB
- 1203** Endoscopic treatment of bleeding gastric ulcer causing gastric wall necrosis: A case report
Li WF, Gao RY, Xu JW, Yu XQ
- 1208** Intermittent melena and refractory anemia due to jejunal cavernous lymphangioma: A case report
Liu KR, Zhang S, Chen WR, Huang YX, Li XG

LETTER TO THE EDITOR

- 1215** New frontiers in ectopic pancreatic tissue management
Covantsev S

ABOUT COVER

Editorial Board Member of *World Journal of Gastrointestinal Surgery*, Arunachalam Rathnaswami, FACS, MBBS, MS, MNAMS, MCh, Ex-Professor and Head, Department of Surgical Gastroenterology, SRM Medical College Hospital and Research Institute SRMIST, Kattankulathur, Chennai, TamilNadu 603203, India. arunarathna@gmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Gastrointestinal Surgery* (WJGS, *World J Gastrointest Surg*) is to provide scholars and readers from various fields of gastrointestinal surgery with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGS mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal surgery and covering a wide range of topics including biliary tract surgical procedures, biliopancreatic diversion, colectomy, esophagectomy, esophagostomy, pancreas transplantation, and pancreatectomy, *etc.*

INDEXING/ABSTRACTING

The WJGS is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports/Science Edition, PubMed, PubMed Central, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJGS as 2.0; IF without journal self cites: 1.9; 5-year IF: 2.2; Journal Citation Indicator: 0.52; Ranking: 113 among 212 journals in surgery; Quartile category: Q3; Ranking: 81 among 93 journals in gastroenterology and hepatology; and Quartile category: Q4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Zi-Hang Xu, Production Department Director: Xiang Li, Cover Editor: Jia-Ru Fan.

NAME OF JOURNAL

World Journal of Gastrointestinal Surgery

ISSN

ISSN 1948-9366 (online)

LAUNCH DATE

November 30, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Peter Schemmer

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/1948-9366/editorialboard.htm>

PUBLICATION DATE

April 27, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Intermittent melena and refractory anemia due to jejunal cavernous lymphangioma: A case report

Kai-Rui Liu, Sheng Zhang, Wei-Run Chen, You-Xing Huang, Xu-Guang Li

Specialty type: Gastroenterology and hepatology

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0

Grade B (Very good): B

Grade C (Good): 0

Grade D (Fair): 0

Grade E (Poor): 0

P-Reviewer: Sanayeh EB, United States

Received: January 16, 2024

Peer-review started: January 16, 2024

First decision: February 2, 2024

Revised: February 14, 2024

Accepted: March 21, 2024

Article in press: March 21, 2024

Published online: April 27, 2024



Kai-Rui Liu, Sheng Zhang, Wei-Run Chen, You-Xing Huang, Department of Abdominal Surgery, The Second Affiliated Hospital of Guangzhou University of Chinese Medicine, Guangzhou 510000, Guangdong Province, China

Xu-Guang Li, Department of General Surgery, The Affiliated Brain Hospital of Guangzhou Medical University, Guangzhou 510000, Guangdong Province, China

Xu-Guang Li, Key Laboratory of Neurogenetics and Channelopathies of Guangdong Province and The Ministry of Education of China, Guangzhou Medical University, Guangzhou 510000, Guangdong Province, China

Corresponding author: Xu-Guang Li, PhD, Chief Doctor, Department of General Surgery, The Affiliated Brain Hospital of Guangzhou Medical University, No. 36 Mingxin Road, Liwan District, Guangzhou 510000, Guangdong Province, China. 42312801@qq.com

Abstract

BACKGROUND

Lymphangiomas in the gastrointestinal tract are extremely rare in adults. As a benign lesion, small intestine lymphangiomas often remain asymptomatic and pose challenges for definitive diagnosis. However, lymphangiomas can give rise to complications such as abdominal pain, bleeding, volvulus, and intussusception. Here, we report a case of jejunal cavernous lymphangioma that presented with intermittent melena and refractory anemia in a male adult.

CASE SUMMARY

A 66-year-old man presented with intermittent melena, fatigue and refractory anemia nine months prior. Esophagogastroduodenoscopy and colonoscopy were performed many times and revealed no apparent bleeding. Conservative management, including transfusion, hemostasis, gastric acid secretion inhibition and symptomatic treatment, was performed, but the lesions tended to recur shortly after surgery. Ultimately, the patient underwent capsule endoscopy, which revealed a more than 10 cm lesion accompanied by active bleeding. After single-balloon enteroscopy and biopsy, a diagnosis of jejunal cavernous lymphangioma was confirmed, and the patient underwent surgical resection. No complications or recurrences were observed postoperatively.

CONCLUSION

Jejunal cavernous lymphangioma should be considered a cause of obscure gastrointestinal bleeding. Capsule endoscopy and single-balloon enteroscopy can

facilitate diagnosis. Surgical resection is an effective management method.

Key Words: Intermittent melena; Refractory anemia; Capsule endoscopy; Single-balloon enteroscopy; Jejunal cavernous lymphangioma; Enterectomy; Case report

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: We report a patient with recurrent melena and refractory anemia that was misdiagnosed as obscure gastrointestinal bleeding or intestinal ischemia for 9 months. A definite diagnosis of jejunal cavernous lymphangioma was made through capsule endoscopy, double-balloon enteroscopy, and biopsy. After surgical resection of the lesion, both the melena and anemia resolved.

Citation: Liu KR, Zhang S, Chen WR, Huang YX, Li XG. Intermittent melena and refractory anemia due to jejunal cavernous lymphangioma: A case report. *World J Gastrointest Surg* 2024; 16(4): 1208-1214

URL: <https://www.wjgnet.com/1948-9366/full/v16/i4/1208.htm>

DOI: <https://dx.doi.org/10.4240/wjgs.v16.i4.1208>

INTRODUCTION

Lymphangioma is an uncommon benign neoplasm that arises from congenital dysplasia of the lymphatic system and occurs more rarely within the small intestine in adults, accounting for less than 1% of all cases[1,2]. Small bowel lymphangiomas lack characteristic manifestations and symptoms. The clinical presentations of asymptomatic chronic wasting diseases vary and include weight loss, hypoproteinemia, lymphopenia, hypogammaglobulinemia and iron deficiency anemia; acute abdominalgia, for instance, volvulus; intussusception; and acute intestinal obstruction[1,3-7]. In this report, we present a case of recurrent melena and refractory anemia caused by jejunal cavernous lymphangioma, which was initially misdiagnosed for nine months but then was successfully diagnosed through capsule endoscopy, single-balloon enteroscopy and biopsy at our hospital. Following surgical resection, there has been no recurrence of symptoms.

CASE PRESENTATION

Chief complaints

A 66-year-old man presented with recurrent melena and fatigue for 9 months.

History of present illness

The patient presented with melena approximately of unknown etiology for 3 to 4 times a day and that started nine months earlier, with approximately 100 mL of melena per occurrence, and the melena was accompanied by dull pain below the xiphoid and fatigue, without other significant discomfort. The symptoms resolved spontaneously after a few days. He underwent multiple hospital visits for treatment, during which esophagogastroduodenoscopy and colonoscopy were performed, revealing no evident bleeding lesions. He was diagnosed with moderate to severe anemia. The lowest recorded hemoglobin level was 5.5 g/dL. Following conservative management, including transfusion, hemostasis, and gastric acid secretion inhibition, his symptoms were temporarily alleviated. However, the symptoms recurred after discharge, leading to his presentation at our hospital in March 2023.

History of past illness

The patient had a history of hypertension, which was under control with medical treatment. He denied any history of trauma or surgery.

Personal and family history

No hereditary diseases or history of allergies, alcoholism or epidemiologic linkage were described.

Physical examination

The patient presented with an anemic appearance. The abdomen was soft. No tenderness, peritonitis or palpable mass was observed. No abnormalities were detected during a digital rectal examination. Cardiopulmonary and neurological examinations revealed no abnormalities.

Laboratory examinations

His hemoglobin level was 6.7 g/dL, and his fecal occult blood test was strongly positive at admission. The C-reactive

protein level was 11.7 mg/L. His leukocyte count and procalcitonin level were normal. The lymphocyte count and lymphocyte percentage were $0.47 \times 10^9/L$ and 14.6%, respectively. The thrombocyte count and coagulation function were also within the normal range. No positive laboratory results were detected for carcinoembryonic antigen, cancer antigen 19-9, tuberculosis antibody, autoimmune disease-related antibodies, immunoglobulin A (IgA), IgG, or IgM. There were no abnormalities in liver or renal function tests.

Imaging examinations

No hemorrhagic foci were detected during esophagogastroduodenoscopy or colonoscopy. Contrast-enhanced computed tomography (CT) of the entire abdomen revealed ischemic changes on the mesenteric side of the distal jejunal and adjacent mesenteric nonenhancing masses (Figure 1). During capsule endoscopy, distal jejunal mucosal erosion and active bleeding were found (Figure 2). Single-balloon enteroscopy was performed for biopsy. As revealed by hematoxylin and eosin staining, the biopsy specimens exhibited a significant abundance of buried tubular structures in the mucosal layer (Figure 3). Immunohistochemical analysis revealed negative CD34 expression and positive D2-40 expression in the biopsy specimen.

FINAL DIAGNOSIS

According to the enteroscopy findings and pathological results, a definitive diagnosis of jejunal cavernous lymphangioma was established.

TREATMENT

After positive preoperative preparation, exploratory laparotomy was performed. During intraoperative exploration, edema and thickening of the intestinal wall were observed approximately 2 meters away from the Treitz ligament. The length of the diseased jejunum exceeded 10 cm. Numerous cystic and hemorrhagic masses of varying sizes were observed on the mesentery, and these lesions were infiltrating into the intestinal lumen (Figure 4). The patient underwent resection of the diseased jejunum and mesentery, followed by side-to-side jejunal anastomosis. Following a three-day fasting period postsurgery, accompanied by prophylactic anti-infection measures and parenteral nutrition support, the patient started a full-fluid diet and gradually transitioned to a normal diet. No postoperative complications were observed, and the patient was discharged seven days after the surgical procedure.

OUTCOME AND FOLLOW-UP

The anemia resolved, and 4 months after the operation, his hemoglobin level increased to 12.9 g/dL without transfusion. No complications or recurrences were observed during the 6-month follow-up period.

DISCUSSION

Lymphangioma can arise from malformations of the lymphatic system during embryonic development or from acquired factors such as trauma, surgery, inflammation, and parasitic infections[8]. Lymphangioma can be histopathologically classified into three subtypes: Cavernous, capillary, and cystic lymphangioma, which may manifest in any part of the body[9,10]. The preferential localization of the fungus is predominantly in the head, neck, and axilla, encompassing approximately 95% of all cases, especially in young children[11,12]. However, limited reports on the small intestine, spleen, esophagus, and other organs exist[13]. Intestinal lymphangioma is extremely rare in adults. A mere total of 52 cases have been reported in the English literature since 1961[2,3,6,8,13-23].

There is no characteristic clinical presentation of lymphangioma. The manifestation may be asymptomatic at times and can occasionally encompass a range of symptoms, including abdominal pain, melena, and acute peritonitis. Hemorrhage, perforation, torsion, intussusception, and rupture are frequent causes of hospital admission[16]. Occult gastrointestinal bleeding or obscure intermittent melena may be the predominant clinical manifestations, accounting for more than half of the cases (28 out of 52, 53.8%) reported in the English literature. The mechanism of bleeding may be attributed to the obstruction of lymphatic flow, resulting in increased pressure on lymphatic-venous connections and subsequent retrograde blood flow into lymphatic vessels[16]. In our case, the patient presented with undiagnosed intermittent melena, fatigue and refractory anemia for a duration of nine months, which highlights the necessity of considering intestinal lymphangioma as a potential etiology for obscure gastrointestinal bleeding.

The laboratory tests conducted for lymphangiomias lack specificity. In this case, we observed a mild decrease in the lymphocyte count, which has been previously reported in two similar cases[3,4]. It is speculated that lymphopenia may serve as one of the clinical features associated with lymphangioma. However, the underlying mechanism is unclear. One postulation is that this condition could be due to the excessive loss of lymphatic fluid containing lymphocytes into the intestinal lumen[24].




Figure 1 Different sections of abdominal contrast-enhanced computed tomography image showing nonenhancing masses on diseased jejunal mesentery (orange arrow). A: Sagittal plane image; B: Coronal plane image; C: Transverse plane image.




Figure 2 Representative images of single-balloon enteroscopy. A: Characteristic endoscopic manifestations of edematous white-yellow mucosa accompanied by hemorrhagic red spots (white arrow); B and C: Images demonstrate multiple active bleeding lesions (white arrow).

The use of CT, magnetic resonance imaging (MRI), and ultrasound, although lacking specificity and having limited accuracy, is advantageous for facilitating the detection of lesions and differential diagnosis. On CT, intestinal lymphangiomas are nonenhanced, well-demarcated and low-density oval masses beneath the submucous membrane[8]. CT scans may yield inconclusive results, with a positive rate of 39.1%[3]. However, it plays a crucial role in the detection of masses that manifest as asymptomatic large abdominal masses or those that give rise to complications, such as perforation, volvulus, and intussusception. MRI scans often reveal a prolonged T1 and T2 signal, a high compression lipid signal and homogeneous/hypodense lesions[22]. Ultrasound can detect multiple cystic anechoic-hypoechoic masses that exhibit thickening of the mucosal and submucosal layers and that display a hypoechoic heterogeneous component[15]. Positron emission tomography/CT is helpful for identifying malignant tumors.

The utilization of capsule endoscopy may represent one of the most efficacious approaches for the diagnosis of lymphangioma. Characteristic endoscopic manifestations include edematous white-yellow mucosa accompanied by hemorrhagic red spots[14]. Once the typical features are identified, a single or double enteroscopy can be performed to conduct a biopsy and establish a definitive diagnosis. Among the 52 cases reported in the English literature, 40 included a definitive diagnosis established using capsule endoscopy or enteroscopy (76.9%, 40/52), and 42 patients (80.8%, 42/52) were diagnosed since 2010. This finding may be attributed to the widespread clinical utilization of endoscopy in the past decade and indicates the indispensability of capsule endoscopy for diagnosing lymphangioma.

Histopathologic presentation of cavernous lymphangioma revealed large dilated and congested lymphatic lumens expanding to the mucosa of the intestine[6,23]. D2-40, CD31, VEGFR3 and ERG expression is valuable for demonstrating lymphangioma endothelial cells[4,25,26]. The expression of CD34 can serve as a reliable marker for distinguishing lymphangioma tissue from hemangioma tissue, particularly in lymphangioma tissue with hemorrhage, where the dilated vessels are filled with erythrocytes[27]. Immunohistochemical analysis of this patient revealed D2-40 expression and negative CD34 expression, consistent with a diagnosis of cavernous lymphangioma.

Specific interventions are rarely necessary for asymptomatic lymphangiomas, whereas surgical resection remains the established approach for individuals experiencing symptoms.

Recently, compelling evidence has demonstrated the feasibility, efficacy, and safety of endoscopic resection for patients with lymphangiomas smaller than 2 cm[16,18,19,21,28]. Conservative therapies, such as cryotherapy, laser therapy, and local administration of sclerosing agents, are disfavored[29]. The implementation of meticulous follow-up is crucial, as recurrence rates range from approximately 10% to 27% for completely excised lesions and from 50% to 100% for partially resected lesions[13,29].




Figure 3 Microscopic features of the biopsy specimens. A: The mucosal lamina propria and submucosa exhibited variably sized cysts, which were filled with lymphatic fluid and contained red blood cells and lymphocytes within the lumen (hematoxylin–eosin staining); B: Positive immunohistochemical staining for D2-40 was observed.




Figure 4 Intraoperative findings and resection specimen. A: Numerous cystic and hemorrhagic masses of varying sizes were observed on the mesentery; B: The length of the diseased jejunum exceeded 10 cm. Longitudinal gross dissected specimens revealed that various sizes of nodular eminence were distributed on the lumen surface.

CONCLUSION

Jejunal cavernous lymphangioma, albeit rare, should be considered one of the etiologies of obscure gastrointestinal bleeding. Capsule endoscopy and single-balloon enteroscopy play pivotal roles in facilitating diagnosis. Surgical resection of the lesion represents a highly effective therapeutic intervention.

FOOTNOTES

Author contributions: Liu KR and Li XG conceived and designed the study, read and analyzed the documents, and drafted the manuscript; Zhang S collected the medical data; Chen WR participated in the clinical assessment at each visit during the follow-up; Huang YX provided suggestions on the diagnosis and operation of the patients; Li XG performed the surgical resection. All the authors read and approved the final manuscript.

Informed consent statement: Informed written consent was obtained from the patient for the publication of this report and any accompanying images.

Conflict-of-interest statement: The authors declare that they have no conflicts of interest to disclose.

CARE Checklist (2016) statement: The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the

original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: You-Xing Huang 0000-0001-9012-184X; Xu-Guang Li 0000-0002-1312-0015.

S-Editor: Fan JR

L-Editor: A

P-Editor: Xu ZH

REFERENCES

- Kohga A, Kawabe A, Hasegawa Y, Yajima K, Okumura T, Yamashita K, Isogaki J, Suzuki K, Komiyama A. Ileo-ileal intussusception caused by lymphangioma of the small bowel treated by single-incision laparoscopic-assisted ileal resection. *World J Gastroenterol* 2017; **23**: 167-172 [PMID: 28104992 DOI: 10.3748/wjg.v23.i1.167]
- Akashige T, Sato K, Odajima H, Yamazaki S. A case report of recurrent intussusception caused by small bowel lymphangioma in an adult. *Int J Surg Case Rep* 2020; **75**: 126-130 [PMID: 32950942 DOI: 10.1016/j.ijscr.2020.09.030]
- Tan B, Zhang SY, Wang YN, Li Y, Shi XH, Qian JM. Jejunal cavernous lymphangioma manifested as gastrointestinal bleeding with hypogammaglobulinemia in adult: A case report and literature review. *World J Clin Cases* 2020; **8**: 140-148 [PMID: 31970180 DOI: 10.12998/wjcc.v8.i1.140]
- Ilhan M, Oner G, Alibeyoglu A, Yeğen G, Gök AF, Akyüz F, Bicen F. Primary intestinal lymphangiomatosis of the ileum in an adult-the role of surgical approach. *J Surg Case Rep* 2016; **2016** [PMID: 27534888 DOI: 10.1093/jscr/rjw133]
- Alfadhel SF, Alghamdi AA, Alzahrani SA. Ileal volvulus secondary to cystic lymphangioma: A rare case report with a literature review. *Avicenna J Med* 2019; **9**: 82-85 [PMID: 31143702 DOI: 10.4103/ajm.AJM_203_18]
- Giuliani A, Romano L, Coletti G, Walid A, Fatayer M, Calvisi G, Maffione F, Muolo C, Vicentini V, Schietroma M, Carlei F. Lymphangiomatosis of the ileum with perforation: A case report and review of the literature. *Ann Med Surg (Lond)* 2019; **41**: 6-10 [PMID: 30992989 DOI: 10.1016/j.amsu.2019.03.010]
- Samuelson H, Giannotti G, Guralnick A. Jejunal lymphangioma causing intussusception in an adult: An unusual case with review of the literature. *Ann Med Surg (Lond)* 2018; **34**: 39-42 [PMID: 30210794 DOI: 10.1016/j.amsu.2018.08.020]
- Wang ZZ, Shen LY, Zhou JJ, Tang JL, Ye LP, Shen CB, Li SW, Zhou XB. Clinical manifestation and treatment of small intestinal lymphangioma: A single center analysis of 15 cases. *Front Med (Lausanne)* 2022; **9**: 975698 [PMID: 36213633 DOI: 10.3389/fmed.2022.975698]
- Campbell WJ, Irwin ST, Biggart JD. Benign lymphangioma of the jejunal mesentery: an unusual cause of small bowel obstruction. *Gut* 1991; **32**: 1568 [PMID: 1773970 DOI: 10.1136/gut.32.12.1568]
- Chen CW, Hsu SD, Lin CH, Cheng MF, Yu JC. Cystic lymphangioma of the jejunal mesentery in an adult: a case report. *World J Gastroenterol* 2005; **11**: 5084-5086 [PMID: 16124074 DOI: 10.3748/wjg.v11.i32.5084]
- Fluke LM, Bamberger PK. Cystic Lymphangiomas: Unusual Findings Resulting in Common General Surgery Presentations. *Am Surg* 2018; **84**: e262-e264 [PMID: 30454342]
- Wei MY, Chua J, Cheng Y, Grossberg P. Small bowel volvulus in an adult with mesenteric lymphangioma and ascariasis. *ANZ J Surg* 2018; **88**: E859-E860 [PMID: 28266164 DOI: 10.1111/ans.13953]
- Teng Y, Wang J, Xi Q. Jejunal hemolymphangioma: A case report. *Medicine (Baltimore)* 2020; **99**: e18863 [PMID: 31977886 DOI: 10.1097/MD.00000000000018863]
- Costa JM, Carvalho SD, Gonçalves R, Mascarenhas-Saraiva M, Soares JB. Small bowel lymphangioma. *Clin Res Hepatol Gastroenterol* 2019; **43**: 120-122 [PMID: 30195480 DOI: 10.1016/j.clinre.2018.07.011]
- Tanaka Y, Fujii S, Kusaka T, Kokuryu H. Effective use of EUS for diagnosing a jejunal lymphangioma accompanied with hemorrhage. *Gastrointest Endosc* 2020; **91**: 199-200 [PMID: 31442393 DOI: 10.1016/j.gie.2019.08.009]
- Wu TL, Hsu HT, Yen HH. Jejunum lymphangioma: a rare case of obscure gastrointestinal bleeding with successful endoscopic therapy. *Endoscopy* 2021; **53**: E307-E308 [PMID: 33032354 DOI: 10.1055/a-1268-6992]
- Yang J, Zhang Y, Kou G, Li Y. Jejunum Hemolymphangioma Causing Refractory Anemia in a Young Woman. *Am J Gastroenterol* 2020; **115**: 810 [PMID: 31658121 DOI: 10.14309/ajg.0000000000000438]
- Ito Y, Osuga T, Kawasaki K, Ikura Y. Jejunal hemorrhage due to hemolymphangioma successfully detected and controlled by double-balloon enteroscopy. *Clin Case Rep* 2021; **9**: e05153 [PMID: 34917370 DOI: 10.1002/ccr3.5153]
- João M, Gravito-Soares E, Lopes S, Amaro P. Jejunal cavernous lymphangioma: successful endoscopic treatment of a rare cause of small bowel bleeding. *Ann Gastroenterol* 2021; **34**: 891 [PMID: 34815657 DOI: 10.20524/aog.2021.0662]
- Hokama A, Irahia A. Jejunal lymphangioma. *Rev Esp Enferm Dig* 2023; **115**: 103-104 [PMID: 36043534 DOI: 10.17235/reed.2022.9088/2022]
- Trieu J, Dua A, Gupta N, Venu RP, Venu M. Single-Balloon-Assisted Enteroscopy With Endoscopic Mucosal Resection of a Bleeding Jejunal Lymphangioma. *ACG Case Rep J* 2022; **9**: e00860 [PMID: 36072355 DOI: 10.14309/crj.00000000000000860]
- Zhao N, Fu Y, Wang Z, An Q, Jia W. Case report: Submucosal cavernous lymphangioma causing jejuno-jejunal intussusception in an adult. *Front Surg* 2022; **9**: 953840 [PMID: 36338654 DOI: 10.3389/fsurg.2022.953840]
- Xu J, Zhang C, Tang C. A case of jejunal hemolymphangioma. *Dig Liver Dis* 2023; **55**: 291-293 [PMID: 36435714 DOI: 10.1016/j.dld.2022.11.007]
- Amadori G, Micciolo R, Poletti A. A case of intra-abdominal multiple lymphangiomas in an adult in whom the immunological evaluation supported the diagnosis. *Eur J Gastroenterol Hepatol* 1999; **11**: 347-351 [PMID: 10333211 DOI: 10.1097/00042737-199903000-00022]
- Fukunaga M. Expression of D2-40 in lymphatic endothelium of normal tissues and in vascular tumours. *Histopathology* 2005; **46**: 396-402 [PMID: 15810951 DOI: 10.1111/j.1365-2559.2005.02098.x]
- Galamboos C, Nodit L. Identification of lymphatic endothelium in pediatric vascular tumors and malformations. *Pediatr Dev Pathol* 2005; **8**:

- 181-189 [PMID: [15719202](#) DOI: [10.1007/s10024-004-8104-9](#)]
- 27 **Lee YS**, Kim GW, Cho HJ, Shim CS. Colonic lymphangiomatosis resolved after excisional biopsy. *Clin Endosc* 2015; **48**: 81-84 [PMID: [25674533](#) DOI: [10.5946/ce.2015.48.1.81](#)]
- 28 **Hizawa K**, Matsumoto T, Kouzuki T, Suekane H, Esaki M, Fujishima M. Cystic submucosal tumors in the gastrointestinal tract: endosonographic findings and endoscopic removal. *Endoscopy* 2000; **32**: 712-714 [PMID: [10989996](#) DOI: [10.1055/s-2000-9025](#)]
- 29 **Kosmidis I**, Vlachou M, Koutroufinis A, Filiopoulos K. Hemolymphangioma of the lower extremities in children: two case reports. *J Orthop Surg Res* 2010; **5**: 56 [PMID: [20704732](#) DOI: [10.1186/1749-799X-5-56](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: office@baishideng.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

