

World Journal of *Gastrointestinal Oncology*

World J Gastrointest Oncol 2022 September 15; 14(9): 1604-1890





REVIEW

- 1604** Advances in postoperative adjuvant therapy for primary liver cancer
Zeng ZM, Mo N, Zeng J, Ma FC, Jiang YF, Huang HS, Liao XW, Zhu GZ, Ma J, Peng T
- 1622** Immunotherapy for nonalcoholic fatty liver disease-related hepatocellular carcinoma: Lights and shadows
Costante F, Airola C, Santopaolo F, Gasbarrini A, Pompili M, Ponziani FR
- 1637** Emerging role of caldesmon in cancer: A potential biomarker for colorectal cancer and other cancers
Alnuaimi AR, Nair VA, Malhab LJB, Abu-Gharbieh E, Ranade AV, Pintus G, Hamad M, Busch H, Kirfel J, Hamoudi R, Abdel-Rahman WM

MINIREVIEWS

- 1654** Liquid biopsy to detect resistance mutations against anti-epidermal growth factor receptor therapy in metastatic colorectal cancer
Valenzuela G, Burotto M, Marcelain K, González-Montero J
- 1665** Implication of gut microbiome in immunotherapy for colorectal cancer
Koustantas E, Trifylli EM, Sarantis P, Papadopoulos N, Aloizos G, Tsagarakis A, Damaskos C, Garmpis N, Garmpi A, Papavassiliou AG, Karamouzis MV

ORIGINAL ARTICLE

Basic Study

- 1675** Potential of six-transmembrane epithelial antigen of the prostate 4 as a prognostic marker for colorectal cancer
Fang ZX, Li CL, Chen WJ, Wu HT, Liu J

Case Control Study

- 1689** Inverse relations between *Helicobacter pylori* infection and risk of esophageal precancerous lesions in drinkers and peanut consumption
Pan D, Sun GJ, Su M, Wang X, Yan QY, Song G, Wang YY, Xu DF, Wang NN, Wang SK

Retrospective Cohort Study

- 1699** Prognostic impact of tumor deposits on overall survival in colorectal cancer: Based on Surveillance, Epidemiology, and End Results database
Wu WX, Zhang DK, Chen SX, Hou ZY, Sun BL, Yao L, Jie JZ
- 1711** Consolidation chemotherapy with capecitabine after neoadjuvant chemoradiotherapy in high-risk patients with locally advanced rectal cancer: Propensity score study
Sheng XQ, Wang HZ, Li S, Zhang YZ, Geng JH, Zhu XG, Quan JZ, Li YH, Cai Y, Wang WH

Retrospective Study

- 1727** Efficacy and safety of computed tomography-guided microwave ablation with fine needle-assisted puncture positioning technique for hepatocellular carcinoma
Hao MZ, Hu YB, Chen QZ, Chen ZX, Lin HL
- 1739** Clinicopathological characterization of ten patients with primary malignant melanoma of the esophagus and literature review
Zhou SL, Zhang LQ, Zhao XK, Wu Y, Liu QY, Li B, Wang JJ, Zhao RJ, Wang XJ, Chen Y, Wang LD, Kong LF
- 1758** Endoscopic debulking resection with additive chemoradiotherapy: Optimal management of advanced inoperable esophageal squamous cell carcinoma
Ren LH, Zhu Y, Chen R, Shrestha Sachin M, Lu Q, Xie WH, Lu T, Wei XY, Shi RH
- 1771** Nomogram for predicting the prognosis of tumor patients with sepsis after gastrointestinal surgery
Chen RX, Wu ZQ, Li ZY, Wang HZ, Ji JF
- 1785** Efficacy and safety of laparoscopic radical resection following neoadjuvant therapy for pancreatic ductal adenocarcinoma: A retrospective study
He YG, Huang XB, Li YM, Li J, Peng XH, Huang W, Tang YC, Zheng L

Observational Study

- 1798** To scope or not - the challenges of managing patients with positive fecal occult blood test after recent colonoscopy
Rattan N, Willmann L, Aston D, George S, Bassan M, Abi-Hanna D, Anandabaskaran S, Ermerak G, Ng W, Koo JH
- 1808** Clinical implications of interleukins-31, 32, and 33 in gastric cancer
Liu QH, Zhang JW, Xia L, Wise SG, Hambly BD, Tao K, Bao SS
- 1823** Construction and analysis of an ulcer risk prediction model after endoscopic submucosal dissection for early gastric cancer
Gong SD, Li H, Xie YB, Wang XH
- 1833** Percutaneous insertion of a novel dedicated metal stent to treat malignant hilar biliary obstruction
Cortese F, Acquafredda F, Mardighian A, Zurlo MT, Ferraro V, Memeo R, Spiliopoulos S, Inchingolo R

EVIDENCE-BASED MEDICINE

- 1844** Prediction of gastric cancer risk by a polygenic risk score of *Helicobacter pylori*
Wang XY, Wang LL, Liang SZ, Yang C, Xu L, Yu MC, Wang YX, Dong QJ

META-ANALYSIS

- 1856** Dissecting novel mechanisms of hepatitis B virus related hepatocellular carcinoma using meta-analysis of public data
Aljabban J, Rohr M, Syed S, Cohen E, Hashi N, Syed S, Khorfan K, Aljabban H, Borkowski V, Segal M, Mukhtar M, Mohammed M, Boateng E, Nemer M, Panahiazar M, Hadley D, Jalil S, Mumtaz K
- 1874** Prognostic and clinicopathological value of Twist expression in esophageal cancer: A meta-analysis
Song WP, Wang SY, Zhou SC, Wu DS, Xie JY, Liu TT, Wu XZ, Che GW

LETTER TO THE EDITOR

- 1886** Nutrition deprivation affects the cytotoxic effect of CD8 T cells in hepatocellular carcinoma

Zhang CY, Liu S, Yang M

ABOUT COVER

Editorial Board Member of *World Journal of Gastrointestinal Oncology*, Luigi Marano, MD, PhD, Associate Professor, Department of Medicine, Surgery, and Neurosciences, University of Siena, Siena 53100, Italy. luigi.marano@unisi.it

AIMS AND SCOPE

The primary aim of *World Journal of Gastrointestinal Oncology* (WJGO, *World J Gastrointest Oncol*) is to provide scholars and readers from various fields of gastrointestinal oncology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGO mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal oncology and covering a wide range of topics including liver cell adenoma, gastric neoplasms, appendiceal neoplasms, biliary tract neoplasms, hepatocellular carcinoma, pancreatic carcinoma, cecal neoplasms, colonic neoplasms, colorectal neoplasms, duodenal neoplasms, esophageal neoplasms, gallbladder neoplasms, *etc.*

INDEXING/ABSTRACTING

The WJGO is now abstracted and indexed in PubMed, PubMed Central, Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJGO as 3.404; IF without journal self cites: 3.357; 5-year IF: 3.250; Journal Citation Indicator: 0.53; Ranking: 162 among 245 journals in oncology; Quartile category: Q3; Ranking: 59 among 93 journals in gastroenterology and hepatology; and Quartile category: Q3. The WJGO's CiteScore for 2021 is 3.6 and Scopus CiteScore rank 2021: Gastroenterology is 72/149; Oncology is 203/360.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Ying-Yi Yuan*; Production Department Director: *Xiang Li*; Editorial Office Director: *Jia-Ru Fan*.

NAME OF JOURNAL

World Journal of Gastrointestinal Oncology

ISSN

ISSN 1948-5204 (online)

LAUNCH DATE

February 15, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Monjur Ahmed, Florin Burada

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/1948-5204/editorialboard.htm>

PUBLICATION DATE

September 15, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Case Control Study

Inverse relations between *Helicobacter pylori* infection and risk of esophageal precancerous lesions in drinkers and peanut consumption

Da Pan, Gui-Ju Sun, Ming Su, Xin Wang, Qing-Yang Yan, Guang Song, Yuan-Yuan Wang, Deng-Feng Xu, Nian-Nian Wang, Shao-Kang Wang

Specialty type: Oncology

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): C, C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Chen LW, Taiwan;
Reshetnyak VI, Russia

Received: April 7, 2022

Peer-review started: April 7, 2022

First decision: June 2, 2022

Revised: June 16, 2022

Accepted: August 5, 2022

Article in press: August 5, 2022

Published online: September 15, 2022



Da Pan, Gui-Ju Sun, Yuan-Yuan Wang, Deng-Feng Xu, Nian-Nian Wang, Shao-Kang Wang, Key Laboratory of Environmental Medicine and Engineering of Ministry of Education, Department of Nutrition and Food Hygiene, School of Public Health, Southeast University, Nanjing 210009, Jiangsu Province, China

Ming Su, Xin Wang, Qing-Yang Yan, Guang Song, Department of Chronic Disease, Huai'an District Center for Disease Control and Prevention, Huai'an 223200, Jiangsu Province, China

Corresponding author: Shao-Kang Wang, MD, PhD, Associate Professor, Key Laboratory of Environmental Medicine and Engineering of Ministry of Education, Department of Nutrition and Food Hygiene, School of Public Health, Southeast University, No. 87 Dingjiaqiao, Nanjing 210009, Jiangsu Province, China. shaokangwang@seu.edu.cn

Abstract

BACKGROUND

Helicobacter pylori (*H. pylori*) is a Gram-negative bacterium found in the upper digestive tract. Although *H. pylori* infection is an identified risk factor for gastric cancer, its role in esophageal squamous cell carcinoma (ESCC) remains a topic of much debate.

AIM

To evaluate the association between *H. pylori* infection and the risk of precancerous lesions of ESCC, and further explore the association between dietary factors and the risk of *H. pylori* infection.

METHODS

Two hundred patients with esophageal precancerous lesions (EPL) aged 63.01 ± 6.08 years and 200 healthy controls aged 62.85 ± 6.03 years were included in this case-control study. Epidemiological data and qualitative food frequency data were investigated. Enzyme-linked immunosorbent assay measuring serum immunoglobulin G antibodies was used to determine *H. pylori* seropositivity. An unconditional logistic regression model was used to assess the association between *H. pylori* infection and EPL risk dichotomized by gender, age, and the use of tobacco and alcohol, as well as the association between dietary factors and the

risk of *H. pylori* infection.

RESULTS

A total of 47 (23.5%) EPL cases and 58 (29.0%) healthy controls had positive *H. pylori* infection. An inverse relation between *H. pylori* infection and the risk of EPL was found in the group of drinkers after adjustment for covariates [odds ratio (OR) = 0.32, 95% confidence interval (95%CI): 0.11-0.95]. Additionally, peanut intake was significantly associated with a decreased risk of *H. pylori* infection (OR = 0.39, 95%CI: 0.20-0.74).

CONCLUSION

Our study suggested that *H. pylori* infection may decrease the risk of EPL for drinkers in a rural adult Chinese population, and the consumption of peanut may reduce the risk of *H. pylori* infection. These findings should be framed as preliminary evidence, and further studies are required to address whether the mechanisms are related to the localization of lesions and alcohol consumption.

Key Words: *Helicobacter pylori*; Esophageal precancerous lesions; Peanut consumption; Esophageal squamous cell carcinoma

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: The association between *Helicobacter pylori* (*H. pylori*) infection and esophageal squamous cell carcinoma (ESCC) remains a topic of much debate. This study aimed to evaluate the association between *H. pylori* infection and the risk of precancerous lesions of ESCC, and further explore the association between dietary intake and the risk of *H. pylori* infection. Our findings suggested an inverse association between *H. pylori* infection and the risk of esophageal precancerous lesions in the group of drinkers [odds ratio (OR) = 0.32, 95% confidence interval (95%CI): 0.11-0.95]. Additionally, peanut consumption was significantly associated with a reduced risk of *H. pylori* infection (OR = 0.39, 95%CI: 0.20-0.74).

Citation: Pan D, Sun GJ, Su M, Wang X, Yan QY, Song G, Wang YY, Xu DF, Wang NN, Wang SK. Inverse relations between *Helicobacter pylori* infection and risk of esophageal precancerous lesions in drinkers and peanut consumption. *World J Gastrointest Oncol* 2022; 14(9): 1689-1698

URL: <https://www.wjgnet.com/1948-5204/full/v14/i9/1689.htm>

DOI: <https://dx.doi.org/10.4251/wjgo.v14.i9.1689>

INTRODUCTION

Esophageal cancer and gastric cancer are upper gastrointestinal cancers that share many risk factors[1-3]. However, their associations with *Helicobacter pylori* (*H. pylori*) infection can be completely different. It has been determined that *H. pylori* infection is an identified risk factor for gastric cancer[4], whereas the role of *H. pylori* in the risk of esophageal cancer remains controversial. Previous meta-analyses summarized that *H. pylori* infection is likely to be related to a reduced risk of esophageal adenocarcinoma (EAC)[5-9]. One of the reliable assumptions related to this phenomenon is that *H. pylori* infection causes gastric atrophy and parietal cell loss, thus leading to alleviated reflux and consequently, a decreased incidence of reflux esophagitis and Barrett's esophagus (precursor for EAC)[10-12]. However, the impact of *H. pylori* infection on esophageal squamous cell carcinoma (ESCC) is not well understood, and research is inconclusive as to what population may be significantly influenced[9,13-15]. Previous meta-analyses also reported that in the general population, no significant association was found between *H. pylori* infection and ESCC risk[6-8], whereas an inverse association was observed in the Middle East[9]. In the other populations, the inverse relationship was found to be highly associated with age, smoking status, and drinking status[15].

H. pylori is a Gram-negative bacterium found in the upper digestive tract. In spite of the fact that *H. pylori* infection may reduce the risk of EAC, it may also cause an adverse effect on human health. Apart from the elevated risk of gastric cancer, *H. pylori* infection is also etiologically related to peptic ulcers, atrophic and non-atrophic gastritis, and lymphoma associated with gastric mucosa, and is able to induce reduced bioavailability and malabsorption of nutrients including iron and vitamin B₁₂[16-18]. This case-control study aimed to investigate the association between *H. pylori* infection and the risk of precancerous lesions of ESCC, which is an identified early stage of carcinogenesis, and further examine the association between dietary factors and the risk of *H. pylori* infection.

MATERIALS AND METHODS

This study was carried out in a high-incidence area for ESCC located in Huai'an District, Huai'an City, Jiangsu Province, China, where the crude incidence rate from 1998 to 2016 was 91.85/100000[19]. As described in our previous studies[20-22], the Early Diagnosis and Early Treatment Project of Esophageal Cancer (EDETPEC) supported by the government and Cancer Foundation of China has been carried out in the endemic regions including Huai'an District since 2010. Local residents were required to undergo routine endoscopies. A detailed introduction to esophageal precancerous lesions (EPL) based on histological criteria for dysplasia and methods for EPL diagnosis has already been given in a previous study[21]. The localization of EPL was based on the definition of upper thoracic esophagus (from thoracic inlet to level of tracheal bifurcation; 18-23 cm from incisors), mid thoracic esophagus (from tracheal bifurcation midway to gastroesophageal junction; 24-32 cm from incisors), and lower thoracic esophagus (from midway between tracheal bifurcation and gastroesophageal junction to gastroesophageal junction, including abdominal esophagus; 32-40 cm from incisors)[23]. Figure 1 shows the flowchart of the study population and data collection process. This study included 200 EPL cases aged 62.85 ± 6.03 years and 200 healthy controls aged 63.01 ± 6.08 years matched by gender, age (± 2 years), and villages. The collection of epidemiological data and dietary intake data based on questionnaire method has been introduced in detail previously[21]. Subjects were required to provide the amount of beer/wine/liquor/any other alcoholic drinks consumed per day, which meant that the average alcohol units consumed per day could be estimated. Separated serum samples were obtained by centrifuging collected fasting blood samples at 3000 rpm for 5 min. Enzyme-linked immunosorbent assay (ELISA, KingMed Diagnostics Group Co., Ltd. Guangzhou, China) measuring serum immunoglobulin G (IgG) antibodies was used to determine *H. pylori* seropositivity. Sensitivity of the ELISA test was 97.9% [95% confidence interval (95%CI): 88.9%-99.9%] and specificity was 100% (95%CI: 86.8%-100%).

Epidata version 3.1 (EpiData Association, Odense, Denmark) was used for inputting and validating the epidemiological data and dietary intake data. Then, SPSS version 22.0 (SPSS, Chicago, IL, United States) was used to establish a database and perform statistical analyses. Two independent samples *t*-test and conditional logistic regression model were used to evaluate the differences in general characteristics and potential factors between healthy controls and EPL cases, wherever appropriate. The Fisher's exact test was used to analyze the difference in localization of EPL and *H. pylori* infection. An unconditional logistic regression model was used to assess the association between *H. pylori* infection and EPL risk dichotomized by gender, age, and tobacco and alcohol use, as well as the association between dietary factors and *H. pylori* infection. Covariates including gender, age, body mass index (BMI), education level, annual income, number of cigarettes per day, and alcohol units consumed per day were adjusted in the logistic regression model. Meanwhile, odds ratio (OR) and 95%CI were calculated accordingly. Statistical significance was defined as $P < 0.05$ (two-tailed).

The study protocol was approved by the Institutional Review Board of Southeast University Zhongda Hospital (Approval No. 2016ZDKYSB017), and the written informed consent was obtained.

RESULTS

Two hundred EPL cases aged 63.01 ± 6.08 years and 200 healthy controls aged 62.85 ± 6.03 years were enrolled. Among the pairs, 100 were males and 100 were females. Table 1 shows that 47 (23.5%) and 58 (29.0%) out of 200 cases and 200 controls, respectively, had *H. pylori* infection. Two independent samples *t*-test and conditional logistic regression analysis indicated that there were no statistically significant differences in age, BMI, education level, annual income per person, current drinking status, or *H. pylori* infection between the two groups after adjustment for covariates ($P > 0.05$). Compared with non-smokers, a smoking habit of more than 20 cigarettes a day was significantly associated with an elevated risk of EPL ($P < 0.05$).

Based on routine endoscopy examination, the study found that the number of cases whose EPL developed in upper, mid, and lower thoracic esophagus was 3, 130, and 67, respectively. Table 2 shows that the control group had the highest positive rate of *H. pylori* infection (29.0%), followed by EPL cases of upper and mid thoracic esophagus (24.8%) and EPL cases of lower thoracic esophagus (20.9%), but there was no statistically significant differences.

As shown in Table 3, when subjects were dichotomized according to gender, age, and the use of tobacco and alcohol, the protective effect of *H. pylori* infection against the risk of EPL was found in the group of drinkers after adjustment for covariates (OR = 0.32, 95%CI: 0.11-0.95). Supplementary Tables 1 and 2 shows that there may be a nonsignificant decreasing trend of *H. pylori* infection rate when alcohol consumption is increasing.

Figure 2 illustrates the association between dietary factors and the risk of *H. pylori* infection after the adjustment for covariates via the unconditional logistic regression model. The result indicated that peanut intake was significantly associated with a reduced risk of *H. pylori* infection (OR = 0.39, 95%CI 0.20-0.74). Supplementary Table 3 shows that there may be a significant positive association between peanut consumption and alcohol drinking (P for trend < 0.05).

Table 1 Characteristics and potential factors in cases with esophageal precancerous lesions and controls

Category	Cases, <i>n</i> = 200	Controls, <i>n</i> = 200	Adjusted OR (95%CI) ¹	<i>P</i> value
Age (yr), mean ± SD	63.01 ± 6.08	62.85 ± 6.03		0.792 ^a
BMI (kg/m ²), mean ± SD	24.52 ± 3.33	24.36 ± 3.37		0.631 ^a
Normal (18.5-23.9)	82 (41.0%)	84 (42.0%)	1.00 (reference)	
Underweight (< 18.5)	4 (2.0%)	5 (2.5%)	0.61 (0.12-3.02)	0.545
Overweight (24.0-28.0)	84 (42.0%)	89 (44.5%)	0.95 (0.61-1.49)	0.836
Obese (> 28.0)	30 (15.0%)	22 (11.0%)	1.61 (0.83-3.12)	0.164
Education level				
Illiterate	100 (50.0%)	96 (48.0%)	1.00 (reference)	
Primary school education	74 (37.0%)	77 (38.5%)	0.79 (0.45-1.39)	0.413
Middle school education and higher	26 (13.0%)	27 (13.5%)	0.81 (0.36-1.79)	0.599
Annual income/person (RMB)				
1-5000	53 (26.5%)	42 (21.0%)	1.00 (reference)	
5001-10000	88 (44.0%)	86 (43.0%)	0.75 (0.45-1.25)	0.267
> 10000	59 (29.5%)	72 (36.0%)	0.67 (0.37-1.21)	0.183
Current smoking status (number of cigarettes/d)				
Non-smoker	126 (63.0%)	134 (67.0%)	1.00 (reference)	
1-10	20 (10.0%)	17 (8.5%)	1.39 (0.67-2.87)	0.381
11-20	38 (19.0%)	41 (20.5%)	1.10 (0.61-1.97)	0.755
> 20	16 (8.0%)	8 (4.0%)	3.11 (1.00-9.63)	0.049
Current drinking status (alcohol units consumed/d, 1 unit is 8 g or 10 mL of pure alcohol)				
Non-drinker	147 (73.5%)	151 (75.5%)	1.00 (reference)	
< 4	10 (5.0%)	10 (5.0%)	1.03 (0.41-2.59)	0.954
4-	26 (13.0%)	23 (11.5%)	1.02 (0.52-2.02)	0.946
8-	17 (8.5%)	16 (8.0%)	1.06 (0.47-2.42)	0.885
<i>H. pylori</i> infection				
Negative	153 (76.5%)	142 (71.0%)	1.00 (reference)	
Positive	47 (23.5%)	58 (29.0%)	0.75 (0.46-1.24)	0.265

^a*P* value of two independent samples *t*-test.¹Conditional logistic regression model with adjustment for gender, age, BMI, education level, annual income, number of cigarettes per day, and alcohol units consumed per day, except the specific variable itself.*H. pylori*: *Helicobacter pylori*; BMI: Body mass index; OR: Odds ratio.

DISCUSSION

This study revealed that in drinkers, there was an association between *H. pylori* infection and a reduced risk of EPL, which is an identified early stage of esophageal carcinogenesis. However, the relationship between *H. pylori* infection and ESCC is still subject to much discussion. Some researchers believed that infection with *H. pylori* can increase the risk of ESCC by causing gastric atrophy that promotes excessive bacterial growth and causes endogenous nitrosamine production[24-26]. However, other studies which held that *H. pylori* infection probably plays a protective role in ESCC postulated that the protection is mediated *via* gastric atrophy, whereas the mechanism is related to a reduced load of esophageal acid[27, 28]. Therefore, it is likely that ESCC might be affected in a double-edged manner by *H. pylori* infection, which is dependent on population and other possible external factors. For example, previous studies have indicated that acid regurgitation may be facilitated by the reduction in lower esophageal sphincter's pressure and the retard of both esophageal motility and gastric emptying due to large consumption of alcoholic beverages[29-33]. Therefore, the current hypothesis is that *H. pylori* infection just alleviates esophageal reflux caused by alcohol to some extent, thus reducing the risk of esophageal

Table 2 Difference in *Helicobacter pylori* infection among controls and cases with esophageal precancerous lesions

Group	<i>H. pylori</i> infection			P value ^a
	Negative	Positive	Positive rate	
Controls	142	58	29.0%	0.384
EPL cases (upper and mid thoracic esophagus)	100	33	24.8%	
EPL cases (lower thoracic esophagus)	53	14	20.9%	

^aP value of Fisher's exact test.*H. pylori*: *Helicobacter pylori*.**Table 3** Association between *Helicobacter pylori* infection and esophageal precancerous lesion risk dichotomized by gender, age, cigarette smoking, and alcohol drinking

	Cases	Controls	Crude OR (95%CI)	P value	Adjusted OR (95%CI) ¹	P value
Male	n = 100	n = 100				
<i>H. pylori</i> (-)	78 (78.0%)	72 (72.0%)	1.00 (reference)	-	1.00 (reference)	-
<i>H. pylori</i> (+)	22 (22.0%)	28 (28.0%)	0.73 (0.38-2.38)	0.328	0.64 (0.32-1.27)	0.200
Female	n = 100	n = 100				
<i>H. pylori</i> (-)	75 (75.0%)	70 (70.0%)	1.00 (reference)	-	1.00 (reference)	-
<i>H. pylori</i> (+)	25 (25.0%)	30 (30.0%)	0.78 (0.42-1.45)	0.429	0.82 (0.42-1.58)	0.548
Age < 65 years	n = 107	n = 107				
<i>H. pylori</i> (-)	76 (71.0%)	73 (68.2%)	1.00 (reference)	-	1.00 (reference)	-
<i>H. pylori</i> (+)	31 (29.0%)	34 (31.8%)	0.88 (0.49-1.57)	0.656	0.89 (0.47-1.67)	0.708
Age ≥ 65 years	n = 93	n = 93				
<i>H. pylori</i> (-)	77 (82.8%)	69 (74.2%)	1.00 (reference)	-	1.00 (reference)	-
<i>H. pylori</i> (+)	16 (17.2%)	24 (25.8%)	0.60 (0.29-1.22)	0.156	0.59 (0.27-1.28)	0.183
Cigarette smoking (-)	n = 126	n = 134				
<i>H. pylori</i> (-)	97 (77.0%)	93 (69.4%)	1.00 (reference)	-	1.00 (reference)	-
<i>H. pylori</i> (+)	29 (23.0%)	41 (30.6%)	0.68 (0.39-1.18)	0.170	0.74 (0.42-1.32)	0.310
Cigarette smoking (+)	n = 74	n = 66				
<i>H. pylori</i> (-)	56 (75.7%)	49 (74.2%)	1.00 (reference)	-	1.00 (reference)	-
<i>H. pylori</i> (+)	18 (24.3%)	17 (25.8%)	0.93 (0.43-1.99)	0.845	0.80 (0.34-1.86)	0.601
Alcohol drinking (-)	n = 147	n = 151				
<i>H. pylori</i> (-)	109 (74.1%)	109 (72.2%)	1.00 (reference)	-	1.00 (reference)	-
<i>H. pylori</i> (+)	38 (25.9%)	42 (27.8%)	0.91 (0.54-1.51)	0.702	0.94 (0.55-1.61)	0.831
Alcohol drinking (+)	n = 53	n = 49				
<i>H. pylori</i> (-)	44 (83.0%)	33 (67.3%)	1.00 (reference)	-	1.00 (reference)	-
<i>H. pylori</i> (+)	9 (17.0%)	16 (32.7%)	0.42 (0.17-1.07)	0.070	0.32 (0.11-0.95)	0.040

¹Adjustment for gender, age, BMI, education level, annual income, number of cigarettes per day, and alcohol units consumed per day.*H. pylori*: *Helicobacter pylori*; 95%CI: 95% confidence interval; OR: Odds ratio.

carcinogenesis caused by acid reflux. Our results also reported that EPL cases of lower thoracic esophagus had the lowest positive rate of *H. pylori* infection, which may support the hypotheses to some extent, although the difference was not statistically significant. In addition, there is more data indicating the positive role of this bacterium for humans. For example, a recent review considered the data on *H. pylori* and suggested that *H. pylori* may be a latent or opportunistic pathogen rather than a true pathogen





Figure 1 Flowchart of study population and sample collection.

of some diseases, and is possibly part of the normal human microbiome as a commensal or even a symbiont organism[34]. However, it was reported that a regular but moderate alcohol intake could possibly facilitate elimination of *H. pylori* infection[35]. **Supplementary material** also shows a nonsignificant decreasing trend of *H. pylori* infection rate when alcohol consumption is increasing. This partly supports the hypothesis that there is a possibility that drinkers without *H. pylori* infection could have more alcohol consumption. In other words, the decreased EPL risk in drinkers with *H. pylori* infection is possibly related to a lower alcohol consumption. However, because the result was not statistically significant, and there was no significant association between alcohol consumption and EPL risk in Huai'an in both this study and the previous epidemiological investigation[21], it is hard to address whether the reduced risk of EPL in drinkers with *H. pylori* infection was related to a reduced alcohol intake.

Additionally, the present study reported that the consumption of peanuts may provide protection from *H. pylori* infection. Since peanuts are high in fat, the duodenal mucosa secretes the hormone enterogastrone when fatty food is present in the stomach or small intestine[36]. Enterogastrone inhibits gastric movements and secretion of gastric acid, possibly by blocking the production or activity of gastrin, the hormone that initially leads to these functions[37]. Therefore, the reduced amount of acid produced may influence the growth of *H. pylori*, as *H. pylori* is dependent on acidity to survive for a long time[38]. In addition, in China, people are likely to drink and eat peanuts at the same time, and **Supplementary material** shows that there was a positive association between peanut consumption and alcohol drinking. Therefore, the inverse association between the consumption of peanut and the risk of *H. pylori* infection may be mediated by alcohol drinking. However, there is still a lack of evidence to verify the above hypotheses, thus further researches are required to evaluate the relationship between peanut consumption and *H. pylori* infection.

At present, about 50% of the global population and more than 70% of the population in some developing countries are infected by *H. pylori*[39]. However, this study reported that the positive rates of *H. pylori* infection were only 23.5% and 29.0% in EPL cases and healthy controls. In an early study conducted by Gao *et al*[40], Huai'an, Jiangsu Province was selected as a high incidence area of upper digestive tract cancers, and Pizhou, Jiangsu Province was selected as a low incidence area. They used ELISA and latex agglutinate test for the detection of *H. pylori* infection, and found that the prevalence of *H. pylori* infection among the gastric cancer group/upper digestive tract cancer group in the low incidence area of Pizhou (66.67%/63.46%) was significantly higher than that in the high incidence area of Huai'an (38.64%/39.33%). However, in the high incidence area of Huai'an, the prevalence of *H. pylori* infection in non-cancer controls and the healthy family members of the cancer cases was higher than that of cases. Therefore, the previous study and our study found that the prevalence of *H. pylori* infection in Huai'an may be much lower than that in other areas, and the prevalence in upper digestive tract cancers or EPL cases can be lower than that in non-cancer population in this region.



DOI: 10.4251/wjgo.v14.i9.1689 Copyright ©The Author(s) 2022.

Figure 2 Association between dietary intake and the risk of *Helicobacter pylori* infection after adjustment for covariates via unconditional logistic regression model. BMI: Body mass index; OR: Odds ratio.

CONCLUSION

In summary, our study suggested that *H. pylori* infection is likely to decrease EPL risk in drinkers for a rural adult Chinese population, and the consumption of peanuts may be related to a reduced risk of *H. pylori* infection. However, the sample size used is a limitation of the study, which may bring difficulties to evaluate statistical significance in some statistical analyses, thus the findings should be framed as preliminary evidence. A case-control study might be difficult to determine causality, so the statement of "protective role" might be overestimated. Hence, it is necessary to design a large-scale prospective cohort study to address the impact of *H. pylori* infection on ESCC, the localization of lesions, and the association with dietary intake and the use of alcohol in the future. Additionally, the low prevalence of *H. pylori* infection in Huai'an is a peculiar finding, which implies that further investigations are recommended.

ARTICLE HIGHLIGHTS

Research background

The role of *Helicobacter pylori* (*H. pylori*) infection in esophageal squamous cell carcinoma (ESCC) remains a topic of much debate.

Research motivation

To assess the relationship between *H. pylori* infection and the risk of precancerous lesions of ESCC, which is an identified early stage of carcinogenesis.

Research objectives

This study aimed to evaluate the association between *H. pylori* infection and the risk of esophageal precancerous lesions (EPL) in a high-incidence area in Huai'an, and further explore the association between dietary factors and the risk of *H. pylori* infection.

Research methods

The study was based on a case-control design. Epidemiological data were collected and *H. pylori* seropositivity was tested. An unconditional logistic regression model was used to analyze the association between *H. pylori* infection and EPL risk with adjustment for confounders, as well as the association between dietary factors and risk of *H. pylori* infection.

Research results

The control group had the highest positive rate of *H. pylori* infection (29.0%), followed by EPL cases of upper and mid thoracic esophagus (24.8%) and EPL cases of lower thoracic esophagus (20.9%). The protective effect of *H. pylori* infection against the risk of EPL was observed in the group of drinkers after adjustment for covariates [odds ratio (OR) = 0.32, 95% confidence interval (95%CI): 0.11-0.95]. Peanut intake was significantly associated with a reduced risk of *H. pylori* infection (OR = 0.39, 95%CI: 0.20-0.74).

Research conclusions

H. pylori infection may decrease the risk of EPL in drinkers for a rural adult Chinese population, and the consumption of peanuts may be related to a reduced risk of *H. pylori* infection.

Research perspectives

A well-designed prospective cohort study is required to address the impact of *H. pylori* infection on ESCC, the localization of lesions, and the association with dietary intake and alcohol drinking. Additionally, the low prevalence of *H. pylori* infection in Huai'an is a peculiar finding, which implies that further investigations are recommended.

ACKNOWLEDGEMENTS

Thanks to all the subjects and researchers for their contributions and hard work. We really appreciate Mr Rob Unwin's assistance in polishing the manuscript.

FOOTNOTES

Author contributions: Pan D, Wang SK, and Sun GJ designed the research; Pan D, Su M, Yan QY, Song G, Wang YY, Xu DF, and Wang NN conducted the research; Wang SK, Su M, Wang X, and Sun GJ provided essential materials; Pan D analyzed the data and wrote the manuscript; Wang SK had primary responsibility for final content; all authors read and approved the final manuscript.

Supported by the Fellowship of China Postdoctoral Science Foundation, No. 2022M710675; the National Natural Science Foundation of China, No. 81673147; and the Danone Dietary Nutrition Research and Education Foundation, No. DIC2020-08.

Institutional review board statement: The study was reviewed and approved by the Institutional Review Board of Southeast University Zhongda Hospital (No. 2016ZDKYSB017).

Informed consent statement: Written informed consent was obtained from all subjects.

Conflict-of-interest statement: There are no conflicts of interest to report.

Data sharing statement: All data generated or analyzed during this study are included. The technical appendix and statistical procedure are available from the corresponding author.

STROBE statement: The authors have read the STROBE Statement—checklist of items, and the manuscript was prepared and revised according to the STROBE Statement—checklist of items.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Da Pan 0000-0001-8430-9467; Gui-Ju Sun 0000-0001-5969-2185; Shao-Kang Wang 0000-0002-7503-7655.

S-Editor: Chen YL

L-Editor: Wang TQ

P-Editor: Chen YL

REFERENCES

- 1 **Yildirim M**, Kaya V, Yildiz M, Demirpence O, Gunduz S, Dilli UD. Esophageal cancer, gastric cancer and the use of pesticides in the southwestern of Turkey. *Asian Pac J Cancer Prev* 2014; **15**: 2821-2823 [PMID: 24761907 DOI: 10.7314/APJCP.2014.15.6.2821]
- 2 **Li M**, Wan X, Wang Y, Sun Y, Yang G, Wang L. Time trends of esophageal and gastric cancer mortality in China, 1991-2009: an age-period-cohort analysis. *Sci Rep* 2017; **7**: 6797 [PMID: 28754910 DOI: 10.1038/s41598-017-07071-5]
- 3 **Castro C**, Peleteiro B, Bento MJ, Lunet N. Trends in gastric and esophageal cancer incidence in northern Portugal (1994-2009) by subsite and histology, and predictions for 2015. *Tumori* 2017; **103**: 155-163 [PMID: 27647232 DOI: 10.5301/tj.5000542]
- 4 **Alipour M**. Molecular Mechanism of Helicobacter pylori-Induced Gastric Cancer. *J Gastrointest Cancer* 2021; **52**: 23-30 [PMID: 32926335 DOI: 10.1007/s12029-020-00518-5]
- 5 **Rubenstein JH**, Taylor JB. Meta-analysis: the association of oesophageal adenocarcinoma with symptoms of gastro-oesophageal reflux. *Aliment Pharmacol Ther* 2010; **32**: 1222-1227 [PMID: 20955441 DOI: 10.1111/j.1365-2036.2010.04471.x]
- 6 **Islami F**, Kamangar F. Helicobacter pylori and esophageal cancer risk: a meta-analysis. *Cancer Prev Res (Phila)* 2008; **1**: 329-338 [PMID: 19138977 DOI: 10.1158/1940-6207.CAPR-08-0109]
- 7 **Rokkas T**, Pistiolas D, Sechopoulos P, Robotis I, Margantinis G. Relationship between Helicobacter pylori infection and esophageal neoplasia: a meta-analysis. *Clin Gastroenterol Hepatol* 2007; **5**: 1413-1417, 1417.e1 [PMID: 17997357 DOI: 10.1016/j.cgh.2007.08.010]
- 8 **Zhuo X**, Zhang Y, Wang Y, Zhuo W, Zhu Y, Zhang X. Helicobacter pylori infection and oesophageal cancer risk: association studies via evidence-based meta-analyses. *Clin Oncol (R Coll Radiol)* 2008; **20**: 757-762 [PMID: 18793831 DOI: 10.1016/j.clon.2008.07.005]
- 9 **Gao H**, Li L, Zhang C, Tu J, Geng X, Wang J, Zhou X, Jing J, Pan W. Systematic Review with Meta-analysis: Association of Helicobacter pylori Infection with Esophageal Cancer. *Gastroenterol Res Pract* 2019; **2019**: 1953497 [PMID: 31871444 DOI: 10.1155/2019/1953497]
- 10 **Wang Z**, Shaheen NJ, Whiteman DC, Anderson LA, Vaughan TL, Corley DA, El-Serag HB, Rubenstein JH, Thrift AP. Helicobacter pylori Infection Is Associated With Reduced Risk of Barrett's Esophagus: An Analysis of the Barrett's and Esophageal Adenocarcinoma Consortium. *Am J Gastroenterol* 2018; **113**: 1148-1155 [PMID: 29880962 DOI: 10.1038/s41395-018-0070-3]
- 11 **Thrift AP**. The epidemic of oesophageal carcinoma: Where are we now? *Cancer Epidemiol* 2016; **41**: 88-95 [PMID: 26851752 DOI: 10.1016/j.canep.2016.01.013]
- 12 **Vohlonen IJ**, Hakama M, Härkönen M, Malila N, Pukkala E, Koistinen V, Sipponen P. Oesophageal cancer incidence in 20-year follow-up in a population-based sample of 12 000 middle-age men with or without *Helicobacter pylori* infection in Finland. *Gut* 2018; **67**: 1201-1202 [PMID: 28860351 DOI: 10.1136/gutjnl-2017-314913]
- 13 **Poyrazoglu OB**, Dulger AC, Gultepe BS. Helicobacter Pylori infection in patients with esophageal squamous cell carcinoma. *Clinics (Sao Paulo)* 2017; **72**: 150-153 [PMID: 28355360 DOI: 10.6061/clinics/2017(03)04]
- 14 **Khoshbaten M**, Zadimani A, Bonyadi MR, Mohammadzadeh M, Gachkar L, Pourhoseingholi MA. Helicobacter pylori infection reduces the risk of esophageal squamous cell carcinoma: a case-control study in iran. *Asian Pac J Cancer Prev* 2011; **12**: 149-151 [PMID: 21517248]
- 15 **Wu DC**, Wu IC, Lee JM, Hsu HK, Kao EL, Chou SH, Wu MT. Helicobacter pylori infection: a protective factor for esophageal squamous cell carcinoma in a Taiwanese population. *Am J Gastroenterol* 2005; **100**: 588-593 [PMID: 15743356 DOI: 10.1111/j.1572-0241.2005.40623.x]
- 16 **Vitale G**, Barbaro F, Ianiro G, Cesario V, Gasbarrini G, Franceschi F, Gasbarrini A. Nutritional aspects of Helicobacter pylori infection. *Minerva Gastroenterol Dietol* 2011; **57**: 369-377 [PMID: 22105725]

- 17 **Aimasso U**, D'onofrio V, D'eusebio C, Devecchi A, Pira C, Merlo FD, De Francesco A. Helicobacter pylori and nutrition: a bidirectional communication. *Minerva Gastroenterol Dietol* 2019; **65**: 116-129 [PMID: [30759976](#) DOI: [10.23736/S1121-421X.19.02568-6](#)]
- 18 **Franceschi F**, Annalisa T, Teresa DR, Giovanna D, Ianiro G, Franco S, Viviana G, Valentina T, Riccardo LL, Antonio G. Role of Helicobacter pylori infection on nutrition and metabolism. *World J Gastroenterol* 2014; **20**: 12809-12817 [PMID: [25278679](#) DOI: [10.3748/wjg.v20.i36.12809](#)]
- 19 **Wang S**, Pan D, Chen Z, Song G, Han R, Sun G, Su M. Trends in Incidence and Mortality of Esophageal Cancer in Huai'an District, a High-Risk Area in Northern Jiangsu Province, China. *Cancer Control* 2022; **29**: 10732748221076824 [PMID: [35196897](#) DOI: [10.1177/10732748221076824](#)]
- 20 **Pan D**, Su M, Huang G, Luo P, Zhang T, Fu L, Wei J, Wang S, Sun G. MTHFR C677T genetic polymorphism in combination with serum vitamin B₂, B₁₂ and aberrant DNA methylation of P16 and P53 genes in esophageal squamous cell carcinoma and esophageal precancerous lesions: a case-control study. *Cancer Cell Int* 2019; **19**: 288 [PMID: [31754346](#) DOI: [10.1186/s12935-019-1012-x](#)]
- 21 **Pan D**, Su M, Zhang T, Miao C, Fu L, Yang L, Song G, Raine PJ, Wang S, Sun G. A Distinct Epidemiologic Pattern of Precancerous Lesions of Esophageal Squamous Cell Carcinoma in a High-risk Area of Huai'an, Jiangsu Province, China. *Cancer Prev Res (Phila)* 2019; **12**: 449-462 [PMID: [31040152](#) DOI: [10.1158/1940-6207.CAPR-18-0462](#)]
- 22 **Pan D**, Wang S, Su M, Sun G, Zhu X, Ghahvechi Chaeipeima M, Guo Z, Wang N, Zhang Z, Cui M. Vitamin B₁₂ may play a preventive role in esophageal precancerous lesions: a case-control study based on markers in blood and 3-day duplicate diet samples. *Eur J Nutr* 2021; **60**: 3375-3386 [PMID: [33619628](#) DOI: [10.1007/s00394-021-02516-0](#)]
- 23 **Ashraf HH**, Palmer J, Dalton HR, Waters C, Luff T, Strugnell M, Murray IA. Can patients determine the level of their dysphagia? *World J Gastroenterol* 2017; **23**: 1038-1043 [PMID: [28246477](#) DOI: [10.3748/wjg.v23.i6.1038](#)]
- 24 **Ye W**, Held M, Lagergren J, Engstrand L, Blot WJ, McLaughlin JK, Nyrén O. Helicobacter pylori infection and gastric atrophy: risk of adenocarcinoma and squamous-cell carcinoma of the esophagus and adenocarcinoma of the gastric cardia. *J Natl Cancer Inst* 2004; **96**: 388-396 [PMID: [14996860](#) DOI: [10.1093/jnci/djh057](#)]
- 25 **Iijima K**, Koike T, Abe Y, Yamagishi H, Ara N, Asanuma K, Uno K, Imatani A, Nakaya N, Ohara S, Shimosegawa T. Gastric hyposecretion in esophageal squamous-cell carcinomas. *Dig Dis Sci* 2010; **55**: 1349-1355 [PMID: [19513836](#) DOI: [10.1007/s10620-009-0853-x](#)]
- 26 **Houben GM**, Stockbrügger RW. Bacteria in the aetio-pathogenesis of gastric cancer: a review. *Scand J Gastroenterol Suppl* 1995; **212**: 13-18 [PMID: [8578226](#) DOI: [10.3109/00365529509090296](#)]
- 27 **Richter JE**, Falk GW, Vaezi MF. Helicobacter pylori and gastroesophageal reflux disease: the bug may not be all bad. *Am J Gastroenterol* 1998; **93**: 1800-1802 [PMID: [9772034](#) DOI: [10.1111/j.1572-0241.1998.00523.x](#)]
- 28 **Raghunath A**, Hungin AP, Wooff D, Childs S. Prevalence of Helicobacter pylori in patients with gastro-oesophageal reflux disease: systematic review. *BMJ* 2003; **326**: 737 [PMID: [12676842](#) DOI: [10.1136/bmj.326.7392.737](#)]
- 29 **Akiyama T**, Inamori M, Iida H, Mawatari H, Endo H, Hosono K, Yoneda K, Fujita K, Yoneda M, Takahashi H, Goto A, Abe Y, Kobayashi N, Kubota K, Saito S, Nakajima A. Alcohol consumption is associated with an increased risk of erosive esophagitis and Barrett's epithelium in Japanese men. *BMC Gastroenterol* 2008; **8**: 58 [PMID: [19077221](#) DOI: [10.1186/1471-230X-8-58](#)]
- 30 **Kaufman SE**, Kaye MD. Induction of gastro-oesophageal reflux by alcohol. *Gut* 1978; **19**: 336-338 [PMID: [25830](#) DOI: [10.1136/gut.19.4.336](#)]
- 31 **Keshavarzian A**, Polepalle C, Iber FL, Durkin M. Esophageal motor disorder in alcoholics: result of alcoholism or withdrawal? *Alcohol Clin Exp Res* 1990; **14**: 561-567 [PMID: [2221284](#) DOI: [10.1111/j.1530-0277.1990.tb01200.x](#)]
- 32 **Mincis M**, Chebli JM, Khouri ST, Mincis R. [Ethanol and the gastrointestinal tract]. *Arq Gastroenterol* 1995; **32**: 131-139 [PMID: [8728788](#)]
- 33 **Wang F**, Li G, Ning J, Chen L, Xu H, Kong X, Bu J, Zhao W, Li Z, Wang X, Li X, Ma J. Alcohol accumulation promotes esophagitis via pyroptosis activation. *Int J Biol Sci* 2018; **14**: 1245-1255 [PMID: [30123073](#) DOI: [10.7150/ijbs.24347](#)]
- 34 **Reshetnyak VI**, Burmistrov AI, Maev IV. Helicobacter pylori: Commensal, symbiont or pathogen? *World J Gastroenterol* 2021; **27**: 545-560 [PMID: [33642828](#) DOI: [10.3748/wjg.v27.i7.545](#)]
- 35 **Kuepper-Nybelen J**, Rothenbacher D, Brenner H. Relationship between lifetime alcohol consumption and Helicobacter pylori infection. *Ann Epidemiol* 2005; **15**: 607-613 [PMID: [16118005](#) DOI: [10.1016/j.annepidem.2004.11.001](#)]
- 36 **Gough AL**, Rai VS, Mariano EC, Greco RS, Landor JH. Effect of an intravenous fat preparation on canine gastric secretion. *Am J Surg* 1980; **139**: 829-831 [PMID: [6992614](#) DOI: [10.1016/0002-9610\(80\)90391-8](#)]
- 37 **Thulin L**, Johansson C. Gastrointestinal hormones. *Acta Chir Scand Suppl* 1978; **482**: 69-72 [PMID: [356498](#)]
- 38 **Waldum HL**, Kleaveland PM, Sørdal ØF. Helicobacter pylori and gastric acid: an intimate and reciprocal relationship. *Therap Adv Gastroenterol* 2016; **9**: 836-844 [PMID: [27803738](#) DOI: [10.1177/1756283X16663395](#)]
- 39 **Li J**, Perez-Perez GL. Helicobacter pylori the Latent Human Pathogen or an Ancestral Commensal Organism. *Front Microbiol* 2018; **9**: 609 [PMID: [29666614](#) DOI: [10.3389/fmicb.2018.00609](#)]
- 40 **Gao C**, Li Z, Ding J, Hu X, Xu T, Liu T, Takezaki T, Tajima K. The Relationship between Helicobacter pylori Infection and Gastric Cancer in High and Low Incidence Areas for Upper Digestive Tract Cancers in Jiangsu Province. *Zhongguo Aizheng Zazhi* 2000; **9**: 395-396



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

