World Journal of Gastrointestinal Oncology

World J Gastrointest Oncol 2024 January 15; 16(1): 1-250





Contents

Monthly Volume 16 Number 1 January 15, 2024

EDITORIAL

Present situation and prospect of immunotherapy for unresectable locally advanced esophageal cancer 1 during peri-radiotherapy

Wang FM, Mo P, Yan X, Lin XY, Fu ZC

8 Early gastric cancer recurrence after endoscopic submucosal dissection: Not to be ignored!

Zeng Y, Yang J, Zhang JW

REVIEW

13 New trends in diagnosis and management of gallbladder carcinoma

Pavlidis ET. Galanis IN. Pavlidis TE

ORIGINAL ARTICLE

Clinical and Translational Research

30 Mechanism of pachymic acid in the treatment of gastric cancer based on network pharmacology and experimental verification

Du YH, Zhao JJ, Li X, Huang SC, Ning N, Chen GQ, Yang Y, Nan Y, Yuan L

Retrospective Cohort Study

51 Trends in colorectal cancer incidence according to an increase in the number of colonoscopy cases in Korea Kim GH, Lee YC, Kim TJ, Hong SN, Chang DK, Kim YH, Yang DH, Moon CM, Kim K, Kim HG, Kim ER

Retrospective Study

61 C-reactive protein to albumin ratio predict responses to programmed cell death-1 inhibitors in hepatocellular carcinoma patients

Li BB, Chen LJ, Lu SL, Lei B, Yu GL, Yu SP

79 Impact of propofol and sevoflurane anesthesia on cognition and emotion in gastric cancer patients undergoing radical resection

Li AH, Bu S, Wang L, Liang AM, Luo HY

90 Development and validation of a machine learning-based early prediction model for massive intraoperative bleeding in patients with primary hepatic malignancies

Li J, Jia YM, Zhang ZL, Liu CY, Jiang ZW, Hao ZW, Peng L

102 Comparative study between Embosphere® and Marine gel® as embolic agents for chemoembolization of hepatocellular carcinoma

Kim HC, Choi JW



World Journal of Gastrointestinal Oncology

Contents

Monthly Volume 16 Number 1 January 15, 2024

110 Clinical value of oral contrast-enhanced ultrasonography in diagnosis of gastric tumors

Wang CY, Fan XJ, Wang FL, Ge YY, Cai Z, Wang W, Zhou XP, Du J, Dai DW

Observational Study

118 Prognostic significance and relationship of SMAD3 phospho-isoforms and VEGFR-1 in gastric cancer: A clinicopathological study

Lv SL, Guo P, Zou JR, Chen RS, Luo LY, Huang DQ

133 Colonoscopy plays an important role in detecting colorectal neoplasms in patients with gastric neoplasms Liu XR, Wen ZL, Liu F, Li ZW, Liu XY, Zhang W, Peng D

Basic Study

- 144 Analysis of the potential biological value of pyruvate dehydrogenase E1 subunit β in human cancer Rong Y, Liu SH, Tang MZ, Wu ZH, Ma GR, Li XF, Cai H
- 182 Colorectal cancer's burden attributable to a diet high in processed meat in the Belt and Road Initiative countries

Liu G, Li CM, Xie F, Li QL, Liao LY, Jiang WJ, Li XP, Lu GM

SYSTEMATIC REVIEWS

197 Comprehensive analysis of the role of ubiquitin-specific peptidases in colorectal cancer: A systematic

Al-Balushi E, Al Marzouqi A, Tavoosi S, Baghsheikhi AH, Sadri A, Aliabadi LS, Salarabedi MM, Rahman SA, Al-Yateem N, Jarrahi AM, Halimi A, Ahmadvand M, Abdel-Rahman WM

META-ANALYSIS

214 Application of neoadjuvant chemoradiotherapy and neoadjuvant chemotherapy in curative surgery for esophageal cancer: A meta-analysis

Yuan MX, Cai QG, Zhang ZY, Zhou JZ, Lan CY, Lin JB

CASE REPORT

- 234 Emerging role of liquid biopsy in rat sarcoma virus mutated metastatic colorectal cancer: A case report Gramaça J, Fernandes IG, Trabulo C, Gonçalves J, dos Santos RG, Baptista A, Pina I
- 244 Comprehensive evaluation of rare case: From diagnosis to treatment of a sigmoid Schwannoma: A case report

Π

Li JY, Gao XZ, Zhang J, Meng XZ, Cao YX, Zhao K

Contents

Monthly Volume 16 Number 1 January 15, 2024

ABOUT COVER

Editorial Board Member of World Journal of Gastrointestinal Oncology, Wael M Abdel-Rahman, MD, PhD, Full Professor, Chairman, Department of Medical Laboratory Sciences, College of Health Sciences and Research Institute of Medical and Health Sciences, University of Sharjah, Sharjah 27272, United Arab Emirates. whassan@sharjah.ac.ae

AIMS AND SCOPE

The primary aim of World Journal of Gastrointestinal Oncology (WJGO, World J Gastrointest Oncol) is to provide scholars and readers from various fields of gastrointestinal oncology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WIGO mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal oncology and covering a wide range of topics including liver cell adenoma, gastric neoplasms, appendiceal neoplasms, biliary tract neoplasms, hepatocellular carcinoma, pancreatic carcinoma, cecal neoplasms, colonic neoplasms, colorectal neoplasms, duodenal neoplasms, esophageal neoplasms, gallbladder neoplasms, etc.

INDEXING/ABSTRACTING

The WJGO is now abstracted and indexed in PubMed, PubMed Central, Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJGO as 3.0; IF without journal self cites: 2.9; 5-year IF: 3.0; Journal Citation Indicator: 0.49; Ranking: 157 among 241 journals in oncology; Quartile category: Q3; Ranking: 58 among 93 journals in gastroenterology and hepatology; and Quartile category: Q3. The WJGO's CiteScore for 2022 is 4.1 and Scopus CiteScore rank 2022: Gastroenterology is 71/149; Oncology is 197/366.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Xiang-Di Zhang; Production Department Director: Xiang Li; Editorial Office Director: Jia-Ru Fan.

NAME OF JOURNAL

World Journal of Gastrointestinal Oncology

ISSN 1948-5204 (online)

LAUNCH DATE

February 15, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Monjur Ahmed, Florin Burada

EDITORIAL BOARD MEMBERS

https://www.wignet.com/1948-5204/editorialboard.htm

PUBLICATION DATE

January 15, 2024

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

https://www.wjgnet.com/bpg/gerinfo/204

GUIDELINES FOR ETHICS DOCUMENTS

https://www.wignet.com/bpg/GerInfo/287

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

https://www.wjgnet.com/bpg/gerinfo/240

PUBLICATION ETHICS

https://www.wignet.com/bpg/GerInfo/288

PUBLICATION MISCONDUCT

https://www.wjgnet.com/bpg/gerinfo/208

ARTICLE PROCESSING CHARGE

https://www.wjgnet.com/bpg/gerinfo/242

STEPS FOR SUBMITTING MANUSCRIPTS

https://www.wjgnet.com/bpg/GerInfo/239

ONLINE SUBMISSION

https://www.f6publishing.com

© 2024 Baishideng Publishing Group Inc. All rights reserved. 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA E-mail: office@baishideng.com https://www.wignet.com

Ш



2aishidena® WJGO https://www.wjgnet.com

Submit a Manuscript: https://www.f6publishing.com

World J Gastrointest Oncol 2024 January 15; 16(1): 8-12

ISSN 1948-5204 (online) DOI: 10.4251/wjgo.v16.i1.8

EDITORIAL

Early gastric cancer recurrence after endoscopic submucosal dissection: Not to be ignored!

Yan Zeng, Jian Yang, Jun-Wen Zhang

Specialty type: Oncology

Provenance and peer review:

Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0 Grade B (Very good): B, B Grade C (Good): 0 Grade D (Fair): 0 Grade E (Poor): 0

P-Reviewer: Dilek ON, Turkey; Kawabata H, Japan

Received: November 3, 2023 Peer-review started: November 3,

First decision: December 11, 2023 Revised: December 11, 2023 Accepted: December 18, 2023 Article in press: December 18, 2023 Published online: January 15, 2024



Yan Zeng, Department of Psychology, The Second Affiliated Hospital of Chongqing Medical University, Chongqing 400010, China

Jian Yang, Department of Gastroenterology, Changdu People's Hospital of Xizang, Changdu 854000, Tibet Autonomous Region, China

Jian Yang, Jun-Wen Zhang, Department of Gastroenterology, The First Affiliated Hospital of Chongqing Medical University, Chongqing 400016, China

Corresponding author: Jian Yang, MD, PhD, Associate Chief Physician, Deputy Director, Department of Gastroenterology, Changdu People's Hospital of Xizang, No. 168 Macaoba Road, Changdu 854000, Tibet Autonomous Region, China. yangjian@hospital.cqmu.edu.cn

Abstract

This editorial comments on the article "Efficacy of multi-slice spiral computed tomography in evaluating gastric cancer recurrence after endoscopic submucosal dissection". We focus on the importance of paying more attention to postendoscopic submucosal dissection (ESD) gastric cancer recurrence in patients with early gastric cancer (EGC) and how to manage it effectively. ESD has been a wellknown treatment and the mainstay for EGC, with the advantages of less invasion and fewer complications when compared with traditional surgical procedures. Despite a lower local recurrence rate after ESD, the problem of postoperative recurrence in patients with EGC has become increasingly non-ignorable with the global popularization of ESD technology and the increasing number of post-ESD patients.

Key Words: Early gastric cancer; Recurrence; Endoscopic submucosal dissection; Postoperative; Prediction

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Post-endoscopic submucosal dissection (ESD) gastric cancer recurrence in patients with early gastric cancer has become increasingly non-ignorable with the global popularization of ESD technology and the increasing number of post-ESD patients. A combination approach, including endoscopic techniques, radiographic examinations, predictive biological markers, and machine learning-based prediction models, should be recommended in the individualized surveillance and management of gastric cancer recurrence after ESD.

Citation: Zeng Y, Yang J, Zhang JW. Early gastric cancer recurrence after endoscopic submucosal dissection: Not to be ignored!

World J Gastrointest Oncol 2024; 16(1): 8-12

URL: https://www.wjgnet.com/1948-5204/full/v16/i1/8.htm

DOI: https://dx.doi.org/10.4251/wjgo.v16.i1.8

INTRODUCTION

Early gastric cancer (EGC) is defined as gastric cancer (GC) confined to the mucosa or submucosa with or without nodal involvement[1]. Endoscopic submucosal dissection (ESD) is considered an effective treatment for EGC and is recommended worldwide due to its safer and more cost-effective advantages[2]. Although the recurrence rate after ESD for EGC is relatively low, the problem of post-ESD recurrence in EGC patients has grasped momentous attention in recent years due to the popularization of ESD technology and the increase of post-ESD patients worldwide. In particular, published studies have pointed out that ESD had inferior disease-free survival, recurrence-free survival, and higher recurrence when compared with surgery [3,4], which has also sounded the alarm for paying more attention to GC recurrence after ESD.

We are very interested in the original article by Yin et al[5] published in the September 2023 issue of the World Journal of Gastrointestinal Oncology. We consider this a qualified and enlightening study, as the authors explore a practical approach to monitoring post-ESD GC recurrence in patients with EGC based on the urgent need to simplify follow-up procedures and reduce operational risks. In addition, by comparing with tissue pathology, the gold standard, the authors found that the inadequacy of multi-slice spiral computed tomography (MSCT) lay in the inferior accuracy in predicting GC recurrence. We thank Yin et al[5] for their study, which has been instrumental in raising attention to GC recurrence after ESD and developing recurrence risk monitoring programs.

FACTORS ASSOCIATED WITH GC RECURRENCE AFTER ESD

The factors affecting EC recurrence after ESD in EGC patients may be split into three main sections, including several preoperative, intraoperative, and postoperative risk factors.

First of all, preoperative assessment focuses on the evaluation of the invasion depth and lymphatic metastasis, not only because multicenter studies with a large cohort found that lymphatic infiltration would significantly affect early cancer recurrence [6] but also because the recurrence rate of patients judged by preoperative endoscopic ultrasonography (EUS) to be of higher grade T stage would be significantly higher than that of patients with T1 stage [7]. Based on these results, EUS combined with MSCT has been recommended in assessing preoperative EGC patients [8]. In addition, the primary location of EGC is also considered to have a significant relationship with the local GC recurrence [9,10].

Second, intraoperative risk factors mainly concentrate on whether complete resection of primary GC has been achieved [11]. Previous research found that a cancer-positive horizontal margin (HM) more significant than 6 mm had been identified as an independent risk factor in the local GC recurrence[12]. However, subsequent studies have also shown that, without additional treatment, positive HM patients still have a comparable long-term prognosis and a low risk of lymphatic metastasis despite a non-low recurrence rate[13]. Furthermore, published research has found a higher GC recurrence risk in post-ESD patients with unevaluable or positive vertical margins than those with unevaluable or positive HM[14].

Third, postoperative risk factors may involve multiple elements. Among them, it is worth mentioning that helicobacter pylori (H.pylori) has been identified as an independent risk factor for metachronous GC recurrence after ESD[15]. Thus, H.pylori eradication after ESD in EGC patients has been recommended to reduce metachronous GC recurrence [16]. Furthermore, recent studies have found that *H.pylori* eradication can reduce GC cell migration and invasion by blocking circFNDC3B expression, which also supports H.pylori detection and treatment in EGC patients after ESD[17].

SURVEILLANCE STRATEGIES FOR POST-ESD RECURRENCE

MSCT, follow-up endoscopy, and tissue pathology are the three main methods for GC recurrence surveillance. Although MSCT is convenient and non-invasive, its most significant and non-negligible inadequacy is still its unsatisfactory accuracy. The sensitivity, specificity, and negative and positive predictive values of MSCT in Yin et all's study were all below 50% [5], while an even lower CT diagnostic accuracy was reported in another study [18]. Meanwhile, despite the increasing application of EUS and magnifying chromoendoscopy [19], follow-up endoscopy and tissue pathology applications are still restricted by scarce medical resources and potential operational risks.

Are there any practical suggestions for these seemingly unsolvable contradictions mentioned above? Combining the above three follow-up methods may be one attempt, but it may also increase total medical costs and cause an unnecessary waste of medical resources. Another promising solution may rely on the rapidly evolving field of data science. Big data and machine learning (ML) have opened up a new era for medical research and clinical practice, such as predicting risks and facilitating personalized clinical decision-making[20]. ML-based prediction models have been reported to predict non-radical resection rates and post-ESD bleeding risks in EGC patients[21,22]. Also, an ML-based predictor has been confirmed to identify post-surgery GC recurrence[23]. Therefore, we believe that the recurrence risk prediction, recurrence identification, and individualized follow-up management of EGC patients after ESD interventions through ML-based prediction models is of research significance and application prospect in the current situation of increasing post-ESD patients.

CURRENT RESEARCH FOR POST-ESD RECURRENCE

Reference Citation Analysis (RCA, https://www.referencecitationanalysis.com/) is a unique artificial intelligence system for citation evaluation of biomedical literature. RCA has been employed to analyze previous studies of EGC patients with post-ESD up to December 2023. Published studies on post-ESD GC recurrence focus mainly on risk identification and assessment model creation, imaging and endoscopy, molecular markers, and integrated treatment strategies after recurrence [24,25].

Several deficiencies exist in the current study of GC recurrence in post-ESD patients, including widely varied recurrence rates in different studies[26-28], inconsistent follow-up periods after ESD[5,29,30], the absence of standardized treatment protocols for postoperative GC recurrence, and lack of large-scale, multicenter, and prospective clinical data [31]. Research in these areas could be of interest in the future.

CONCLUSION

In conclusion, the recurrence and follow-up management of EGC patients after ESD should not be ignored, and further research is needed to assess the recurrence risk better and develop individualized management plans.

FOOTNOTES

Author contributions: Zeng Y, Yang J, and Zhang JW conceptualized and designed the research; Zeng Y and Yang J performed the literature search, analyzed the data, and wrote the original manuscript; Zhang JW edited the final manuscript; All authors have read and approved the final manuscript.

Supported by Program for Youth Innovation in Future Medicine, Chongqing Medical University, No. W0138.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: https://creativecommons.org/Licenses/by-nc/4.0/

Country/Territory of origin: China

ORCID number: Yan Zeng 0000-0003-4935-1306; Jian Yang 0000-0001-8170-0727; Jun-Wen Zhang 0000-0003-2911-598X.

Corresponding Author's Membership in Professional Societies: Digestive endoscopy Branch of Chongqing Medical Association.

S-Editor: Li L L-Editor: A P-Editor: Li L

REFERENCES

Chen D, Chen G, Jiang W, Fu M, Liu W, Sui J, Xu S, Liu Z, Zheng X, Chi L, Lin D, Li K, Chen W, Zuo N, Lu J, Chen J, Li G, Zhuo S, Yan J. Association of the Collagen Signature in the Tumor Microenvironment With Lymph Node Metastasis in Early Gastric Cancer. *JAMA Surg* 2019; **154**: e185249 [PMID: 30698615 DOI: 10.1001/jamasurg.2018.5249]



- Wang J, Li SJ, Yan Y, Yuan P, Li WF, Cao CQ, Chen WG, Chen KN, Wu Q. Feasibility of same-day discharge following endoscopic submucosal dissection for esophageal or gastric early cancer. World J Gastroenterol 2022; 28: 5957-5967 [PMID: 36405109 DOI: 10.3748/wjg.v28.i41.5957]
- 3 Jeon HK, Kim GH, Lee BE, Park DY, Song GA, Kim DH, Jeon TY. Long-term outcome of endoscopic submucosal dissection is comparable to that of surgery for early gastric cancer: a propensity-matched analysis. Gastric Cancer 2018; 21: 133-143 [PMID: 28397011 DOI: 10.1007/s10120-017-0719-4]
- Yang HJ, Kim JH, Kim NW, Choi IJ. Comparison of long-term outcomes of endoscopic submucosal dissection and surgery for undifferentiated-type early gastric cancer meeting the expanded criteria: a systematic review and meta-analysis. Surg Endosc 2022; 36: 3686-3697 [PMID: 35194664 DOI: 10.1007/s00464-022-09126-9]
- Yin JJ, Hu X, Hu S, Sheng GH. Efficacy of multi-slice spiral computed tomography in evaluating gastric cancer recurrence after endoscopic 5 submucosal dissection. World J Gastrointest Oncol 2023; 15: 1636-1643 [PMID: 37746651 DOI: 10.4251/wjgo.v15.i9.1636]
- Yamada S, Hatta W, Shimosegawa T, Takizawa K, Oyama T, Kawata N, Takahashi A, Oka S, Hoteya S, Nakagawa M, Hirano M, Esaki M, 6 Matsuda M, Nakaya N, Gotoda T. Different risk factors between early and late cancer recurrences in patients without additional surgery after noncurative endoscopic submucosal dissection for early gastric cancer. Gastrointest Endosc 2019; 89: 950-960 [PMID: 30465769 DOI: 10.1016/j.gie.2018.11.015]
- Smith JW, Brennan MF, Botet JF, Gerdes H, Lightdale CJ. Preoperative endoscopic ultrasound can predict the risk of recurrence after operation for gastric carcinoma. J Clin Oncol 1993; 11: 2380-2385 [PMID: 8246026 DOI: 10.1200/JCO.1993.11.12.2380]
- Li JH, Shen WZ, Gu XQ, Hong WK, Wang ZQ. Prognostic value of EUS combined with MSCT in predicting the recurrence and metastasis of patients with gastric cancer. Jpn J Clin Oncol 2017; 47: 487-493 [PMID: 28334806 DOI: 10.1093/jjco/hyx024]
- 9 Takenaka R, Kawahara Y, Okada H, Hori K, Inoue M, Kawano S, Tanioka D, Tsuzuki T, Yagi S, Kato J, Uemura M, Ohara N, Yoshino T, Imagawa A, Fujiki S, Takata R, Yamamoto K. Risk factors associated with local recurrence of early gastric cancers after endoscopic submucosal dissection. Gastrointest Endosc 2008; 68: 887-894 [PMID: 18565523 DOI: 10.1016/j.gie.2008.03.1089]
- 10 Lee JY, Cho KB, Kim ES, Park KS, Lee YJ, Lee YS, Jang BK, Chung WJ, Hwang JS. Risk factors for local recurrence after en bloc endoscopic submucosal dissection for early gastric cancer. World J Gastrointest Endosc 2016; 8: 330-337 [PMID: 27076871 DOI: 10.4253/wjge.v8.i7.330]
- Japanese Gastric Cancer Association. Japanese Gastric Cancer Treatment Guidelines 2021 (6th edition). Gastric Cancer 2023; 26: 1-25 11 [PMID: 36342574 DOI: 10.1007/s10120-022-01331-8]
- Sekiguchi M, Suzuki H, Oda I, Abe S, Nonaka S, Yoshinaga S, Taniguchi H, Sekine S, Kushima R, Saito Y. Risk of recurrent gastric cancer 12 after endoscopic resection with a positive lateral margin. Endoscopy 2014; 46: 273-278 [PMID: 24505020 DOI: 10.1055/s-0034-1364938]
- 13 Yang HJ, Lee WS, Lee BE, Ahn JY, Jang JY, Lim JH, Nam SY, Kim JH, Min BH, Joo MK, Park JM, Shin WG, Lee HL, Gweon TG, Park MI, Choi J, Tae CH, Kim YI, Choi IJ. Long-term Outcomes of Undifferentiated-Type Early Gastric Cancer with Positive Horizontal Margins after Endoscopic Resection. Gut Liver 2021; 15: 723-731 [PMID: 33790056 DOI: 10.5009/gnl20291]
- Figueiredo PC, Pimentel-Nunes P, Libânio D, Dinis-Ribeiro M. A systematic review and meta-analysis on outcomes after Rx or R1 14 endoscopic resection of superficial gastric cancer. Eur J Gastroenterol Hepatol 2015; 27: 1249-1258 [PMID: 26225870 DOI: 10.1097/MEG.00000000000000440]
- 15 Chung CS, Woo HS, Chung JW, Jeong SH, Kwon KA, Kim YJ, Kim KO, Park DK. Risk Factors for Metachronous Recurrence after Endoscopic Submucosal Dissection of Early Gastric Cancer. J Korean Med Sci 2017; 32: 421-426 [PMID: 28145644 DOI: 10.3346/jkms.2017.32.3.421]
- Bang CS, Baik GH, Shin IS, Kim JB, Suk KT, Yoon JH, Kim YS, Kim DJ. Helicobacter pylori Eradication for Prevention of Metachronous Recurrence after Endoscopic Resection of Early Gastric Cancer. J Korean Med Sci 2015; 30: 749-756 [PMID: 26028928 DOI: 10.3346/jkms.2015.30.6.749]
- Zhang J, Bai J, Zhu H, Li W, An Q, Wang D. The upregulation of circFNDC3B aggravates the recurrence after endoscopic submucosal 17 dissection (ESD) in early gastric cancer (EGC) patients. Sci Rep 2022; 12: 6178 [PMID: 35418175 DOI: 10.1038/s41598-022-07154-y]
- Choi KS, Kim SH, Kim SG, Han JK. Early Gastric Cancers: Is CT Surveillance Necessary after Curative Endoscopic Submucosal Resection 18 for Cancers That Meet the Expanded Criteria? Radiology 2016; 281: 444-453 [PMID: 27243549 DOI: 10.1148/radiol.2016152866]
- Kawabata H, Kawakatsu Y, Yamaguchi K, Ueda Y, Okazaki Y, Hitomi M, Miyata M, Motoi S, Enoki Y, Minamikawa S. A Rare Case of 19 Local Recurrence Following Curative Endoscopic Submucosal Dissection of Intramucosal Differentiated-Type Gastric Cancer. Gastroenterology Res 2019; **12**: 103-106 [PMID: 31019622 DOI: 10.14740/gr1159]
- 20 Radenkovic D, Keogh SB, Maruthappu M. Data science in modern evidence-based medicine. J R Soc Med 2019; 112: 493-494 [PMID: 31526210 DOI: 10.1177/0141076819871055]
- 21 Yun HR, Huh CW, Jung DH, Lee G, Son NH, Kim JH, Youn YH, Park JC, Shin SK, Lee SK, Lee YC. Machine Learning Improves the Prediction Rate of Non-Curative Resection of Endoscopic Submucosal Dissection in Patients with Early Gastric Cancer. Cancers (Basel) 2022; 14 [PMID: 35954406 DOI: 10.3390/cancers14153742]
- Na JE, Lee YC, Kim TJ, Lee H, Won HH, Min YW, Min BH, Lee JH, Rhee PL, Kim JJ. Utility of a deep learning model and a clinical model 22 for predicting bleeding after endoscopic submucosal dissection in patients with early gastric cancer. World J Gastroenterol 2022; 28: 2721-2732 [PMID: 35979158 DOI: 10.3748/wjg.v28.i24.2721]
- Zhou C, Hu J, Wang Y, Ji MH, Tong J, Yang JJ, Xia H. A machine learning-based predictor for the identification of the recurrence of patients with gastric cancer after operation. Sci Rep 2021; 11: 1571 [PMID: 33452440 DOI: 10.1038/s41598-021-81188-6]
- 24 Xie B, Xia Y, Wang X, Xiong Y, Chen SB, Zhang J, He WW. Factors associated with heterochronic gastric cancer development postendoscopic mucosal dissection in early gastric cancer patients. World J Gastrointest Oncol 2023; 15: 1644-1652 [PMID: 37746653 DOI: 10.4251/wjgo.v15.i9.1644]
- Jiang XC, Yao XB, Xia HB, Su YZ, Luo PQ, Sun JR, Song ED, Wei ZJ, Xu AM, Zhang LX, Lan YH. Nomogram established using risk factors of early gastric cancer for predicting the lymph node metastasis. World J Gastrointest Oncol 2023; 15: 665-676 [PMID: 37123061 DOI: 10.4251/wjgo.v15.i4.665]
- Sano T, Sasako M, Kinoshita T, Maruyama K. Recurrence of early gastric cancer. Follow-up of 1475 patients and review of the Japanese 26 literature. Cancer 1993; 72: 3174-3178 [PMID: 8242540 DOI: 10.1002/1097-0142(19931201)72:11<3174::AID-CNCR2820721107>3.0.CO;2-H]
- Min BH, Kim ER, Kim KM, Park CK, Lee JH, Rhee PL, Kim JJ. Surveillance strategy based on the incidence and patterns of recurrence after 27 curative endoscopic submucosal dissection for early gastric cancer. Endoscopy 2015; 47: 784-793 [PMID: 26111362 DOI:



10.1055/s-0034-1392249]

- Na HK, Choi KD, Ahn JY, Lee JH, Kim do H, Song HJ, Lee GH, Jung HY, Kim JH. Endoscopic prediction of recurrence in patients with early 28 gastric cancer after margin-negative endoscopic resection. J Gastroenterol Hepatol 2016; 31: 1284-1290 [PMID: 26820101 DOI: 10.1111/jgh.13301]
- Ryu DG, Kim SJ, Choi CW, Park SB, Nam HS, Lee SH, Hwang SH. Local Recurrence after Endoscopic Submucosal Dissection of Early 29 Gastric Cancer. J Clin Med 2023; 12 [PMID: 36902804 DOI: 10.3390/jcm12052018]
- Nishizawa T, Yahagi N. Long-Term Outcomes of Using Endoscopic Submucosal Dissection to Treat Early Gastric Cancer. Gut Liver 2018; 30 **12**: 119-124 [PMID: 28673068 DOI: 10.5009/gnl17095]
- Moon HS, Yun GY, Kim JS, Eun HS, Kang SH, Sung JK, Jeong HY, Song KS. Risk factors for metachronous gastric carcinoma development after endoscopic resection of gastric dysplasia: Retrospective, single-center study. World J Gastroenterol 2017; 23: 4407-4415 [PMID: 28706423 DOI: 10.3748/wjg.v23.i24.4407]

12



Published by Baishideng Publishing Group Inc

7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: https://www.f6publishing.com/helpdesk

https://www.wjgnet.com

