

# World Journal of *Gastrointestinal Oncology*

*World J Gastrointest Oncol* 2024 February 15; 16(2): 251-570



**EDITORIAL**

- 251 Cardiotoxicity induced by fluoropyrimidine drugs in the treatment of gastrointestinal tumors  
*Kong MW, Sun FD, Pei ZY, Xu L, Wang ZB, Chen Y, Tang SQ, Yang TF, He GX*
- 255 Does enhanced recovery after surgery programs improve clinical outcomes in liver cancer surgery?  
*Sánchez-Pérez B, Ramia JM*

**REVIEW**

- 259 Biological factors driving colorectal cancer metastasis  
*An SX, Yu ZJ, Fu C, Wei MJ, Shen LH*

**MINIREVIEWS**

- 273 Progress in the treatment of advanced hepatocellular carcinoma with immune combination therapy  
*Pan D, Liu HN, Qu PF, Ma X, Ma LY, Chen XX, Wang YQ, Qin XB, Han ZX*
- 287 Targeting oxidative stress with natural products: A novel strategy for esophageal cancer therapy  
*Cao F, Zhang HL, Guo C, Xu XL, Yuan Q*
- 300 Multifaceted role of microRNAs in gastric cancer stem cells: Mechanisms and potential biomarkers  
*Sun QH, Kuang ZY, Zhu GH, Ni BY, Li J*

**ORIGINAL ARTICLE****Clinical and Translational Research**

- 314 Expression of cyclin-dependent kinase 9 is positively correlated with the autophagy level in colon cancer  
*Zheng L, Lu J, Kong DL*
- 331 Tumour response following preoperative chemotherapy is affected by body mass index in patients with colorectal liver metastases  
*Song HC, Zhou HC, Gu P, Bao B, Sun Q, Mei TM, Cui W, Yao K, Yao HZ, Zhang SY, Wang YS, Song RP, Wang JZ*

**Case Control Study**

- 343 Preoperative controlling nutritional status as an optimal prognostic nutritional index to predict the outcome for colorectal cancer  
*Liu LX, Wang H, Gao B, Xu TT, Yuan QG, Zhou SZ, Ding C, Miao J, Guan WX*

**Retrospective Study**

- 354 Effect of screening colonoscopy frequency on colorectal cancer mortality in patients with a family history of colorectal cancer  
*Zheng L, Li B, Lei L, Wang LJ, Zeng ZP, Yang JD*

- 364 Effect of different anesthetic modalities with multimodal analgesia on postoperative pain level in colorectal tumor patients

*Tang JC, Ma JW, Jian JJ, Shen J, Cao LL*

### Observational Study

- 372 Prognostic value of circulating tumor cells combined with neutrophil-lymphocyte ratio in patients with hepatocellular carcinoma

*Chen JL, Guo L, Wu ZY, He K, Li H, Yang C, Han YW*

- 386 Systemic Inflammation Response Index and weight loss as prognostic factors in metastatic pancreatic cancer: A concept study from the PANTHEIA-SEOM trial

*Pacheco-Barcia V, Custodio-Cabello S, Carrasco-Valero F, Palka-Kotlowska M, Mariño-Mendez A, Carmona-Bayonas A, Gallego J, Martín AJM, Jimenez-Fonseca P, Cabezon-Gutierrez L*

### Basic Study

- 398 Prohibitin 1 inhibits cell proliferation and induces apoptosis *via* the p53-mediated mitochondrial pathway *in vitro*

*Shi JJ, Wang YK, Wang MQ, Deng J, Gao N, Li M, Li YP, Zhang X, Jia XL, Liu XT, Dang SS, Wang WJ*

- 414 Early results of the integrative epigenomic-transcriptomic landscape of colorectal adenoma and cancer

*Lu YW, Ding ZL, Mao R, Zhao GG, He YQ, Li XL, Liu J*

- 436 Comprehensive analysis of the potential pathogenesis of COVID-19 infection and liver cancer

*Rong Y, Tang MZ, Liu SH, Li XF, Cai H*

- 458 Immune-related long noncoding RNA zinc finger protein 710-AS1-201 promotes the metastasis and invasion of gastric cancer cells

*Ding W, Chen WW, Wang YQ, Xu XZ, Wang YB, Yan YM, Tan YL*

- 475 Comprehensive analysis of the protein phosphatase 2A regulatory subunit B56ε in pan-cancer and its role and mechanism in hepatocellular carcinoma

*Wu HM, Huang YY, Xu YQ, Xiang WL, Yang C, Liu RY, Li D, Guo XF, Zhang ZB, Bei CH, Tan SK, Zhu XN*

- 493 Identification of anti-gastric cancer effects and molecular mechanisms of resveratrol: From network pharmacology and bioinformatics to experimental validation

*Ma YQ, Zhang M, Sun ZH, Tang HY, Wang Y, Liu JX, Zhang ZX, Wang C*

### SYSTEMATIC REVIEWS

- 514 Prognostic nutritional index in predicting survival of patients with gastric or gastroesophageal junction adenocarcinoma: A systematic review

*Fiflis S, Christodoulidis G, Papakonstantinou M, Giakoustidis A, Koukias S, Roussos P, Kouliou MN, Koumarelas KE, Giakoustidis D*

### SCIENTOMETRICS

- 527 Global research trends and prospects of cellular metabolism in colorectal cancer

*Liu YC, Gong ZC, Li CQ, Teng P, Chen YY, Huang ZH*

**CASE REPORT**

- 543** Large isolated fibrous tumors in the upper esophagus: A case report  
*Yu JJ, Pei HS, Meng Y*
- 550** Hepatomegaly and jaundice as the presenting symptoms of systemic light-chain amyloidosis: A case report  
*Zhang X, Tang F, Gao YY, Song DZ, Liang J*
- 557** Anti-EGFR antibody monotherapy for colorectal cancer with severe hyperbilirubinemia: A case report  
*Tsurui T, Hirasawa Y, Kubota Y, Yoshimura K, Tsunoda T*
- 563** Early adenocarcinoma mixed with a neuroendocrine carcinoma component arising in the gastroesophageal junction: A case report  
*Cheng YQ, Wang GF, Zhou XL, Lin M, Zhang XW, Huang Q*

**ABOUT COVER**

Editorial Board Member of *World Journal of Gastrointestinal Oncology*, Akihiko Oka, MD, PhD, Assistant Professor, Department of Internal Medicine II, Shimane University Faculty of Medicine, Izumo 693-8501, Japan.  
aoka@med.shimane-u.ac.jp

**AIMS AND SCOPE**

The primary aim of *World Journal of Gastrointestinal Oncology* (*WJGO*, *World J Gastrointest Oncol*) is to provide scholars and readers from various fields of gastrointestinal oncology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

*WJGO* mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal oncology and covering a wide range of topics including liver cell adenoma, gastric neoplasms, appendiceal neoplasms, biliary tract neoplasms, hepatocellular carcinoma, pancreatic carcinoma, cecal neoplasms, colonic neoplasms, colorectal neoplasms, duodenal neoplasms, esophageal neoplasms, gallbladder neoplasms, *etc.*

**INDEXING/ABSTRACTING**

The *WJGO* is now abstracted and indexed in PubMed, PubMed Central, Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 edition of Journal Citation Reports® cites the 2022 impact factor (IF) for *WJGO* as 3.0; IF without journal self cites: 2.9; 5-year IF: 3.0; Journal Citation Indicator: 0.49; Ranking: 157 among 241 journals in oncology; Quartile category: Q3; Ranking: 58 among 93 journals in gastroenterology and hepatology; and Quartile category: Q3. The *WJGO*'s CiteScore for 2022 is 4.1 and Scopus CiteScore rank 2022: Gastroenterology is 71/149; Oncology is 197/366.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: *Xiang-Di Zhang*; Production Department Director: *Xiang Li*; Editorial Office Director: *Jia-Ru Fan*.

**NAME OF JOURNAL**

*World Journal of Gastrointestinal Oncology*

**ISSN**

ISSN 1948-5204 (online)

**LAUNCH DATE**

February 15, 2009

**FREQUENCY**

Monthly

**EDITORS-IN-CHIEF**

Monjur Ahmed, Florin Burada

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/1948-5204/editorialboard.htm>

**PUBLICATION DATE**

February 15, 2024

**COPYRIGHT**

© 2024 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>



Basic Study

# Immune-related long noncoding RNA zinc finger protein 710-AS1-201 promotes the metastasis and invasion of gastric cancer cells

Wei Ding, Wei-Wei Chen, Yi-Qin Wang, Xue-Zhong Xu, Yi-Bo Wang, Yong-Min Yan, Yu-Lin Tan

**Specialty type:** Oncology

**Provenance and peer review:**

Unsolicited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): 0  
Grade B (Very good): B  
Grade C (Good): C, C  
Grade D (Fair): 0  
Grade E (Poor): 0

**P-Reviewer:** Plaza MA, Spain;  
Watanabe T, Japan

**Received:** September 4, 2023

**Peer-review started:** September 4, 2023

**First decision:** November 22, 2023

**Revised:** December 2, 2023

**Accepted:** December 20, 2023

**Article in press:** December 20, 2023

**Published online:** February 15, 2024



**Wei Ding, Wei-Wei Chen, Yi-Qin Wang, Xue-Zhong Xu, Yi-Bo Wang, Yu-Lin Tan**, Department of General Surgery, Wujin Hospital Affiliated with Jiangsu University, Changzhou 213003, Jiangsu Province, China

**Wei Ding, Yong-Min Yan**, Changzhou Medical Center, Nanjing Medical University, Changzhou 213017, Jiangsu Province, China

**Wei Ding, Yu-Lin Tan**, Department of General Surgery, The Wujin Clinical College of Xuzhou Medical University, Changzhou 213003, Jiangsu Province, China

**Corresponding author:** Yu-Lin Tan, MMed, Chief Physician, Department of General Surgery, Wujin Hospital Affiliated with Jiangsu University, No. 2 Yongning Road, Changzhou 213003, Jiangsu Province, China. [tanyldoctor@163.com](mailto:tanyldoctor@163.com)

## Abstract

### BACKGROUND

Gastric cancer (GC) is a prevalent malignant tumor of the gastrointestinal system. ZNF710 is a transcription factor (TF), and zinc finger protein 710 (ZNF710)-AS1-201 is an immune-related long noncoding RNA (lncRNA) that is upregulated in GC cells.

### AIM

To assess the correlation between ZNF710-AS1-201 and immune microenvironment features and to investigate the roles of ZNF710-AS1-201 in the invasion and metastasis processes of GC cells.

### METHODS

We obtained data from The Cancer Genome Atlas and Wujin Hospital. We assessed cell growth, migration, invasion, and programmed cell death using cell counting kit-8, EdU, scratch, Transwell, and flow cytometry assays. Quantitative real-time polymerase chain reaction (qRT-PCR) was used to identify the potential downstream targets of ZNF710-AS1-201.

### RESULTS

In GC tissues with low ZNF710-AS1-201 expression, immunoassays detected significant infiltration of various antitumor immune cells, such as memory CD8 T cells and activated CD4 T cells. In the low-expression group, the half-maximal inhibitory concentrations (IC<sub>50</sub>s) of 5-fluorouracil, cisplatin, gemcitabine, and

trametinib were lower, whereas the IC<sub>50</sub>s of dasatinib and vorinostat were higher. The malignant degree of GC was higher and the stage was later in the high-expression group. Additionally, patients with high expression of ZNF710-AS1-201 had lower overall survival and disease-free survival rates. In vitro, the overexpression of ZNF710-AS1-201 greatly enhanced growth, metastasis, and infiltration while suppressing cell death in HGC-27 cells. In contrast, the reduced expression of ZNF710-AS1-201 greatly hindered cell growth, enhanced apoptosis, and suppressed the metastasis and invasion of MKN-45 cells. The expression changes in ZNF710 were significant, but the corresponding changes in isocitrate dehydrogenase-2, Semaphorin 4B, ARHGAP10, RGMB, hsa-miR-93-5p, and ZNF710-AS1-202 were not consistent or statistically significant after overexpression or knockdown of ZNF710-AS1-201, as determined by qRT-PCR.

## CONCLUSION

Immune-related lncRNA ZNF710-AS1-201 facilitates the metastasis and invasion of GC cells. It appears that ZNF710-AS1-201 and ZNF710 have potential as effective targets for therapeutic intervention in GC. Nevertheless, it is still necessary to determine the specific targets of the ZNF710 TF.

**Key Words:** Gastric cancer; ZNF710-AS1-201; Proliferation; Metastasis; Invasion; Apoptosis

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** In the field of oncology, there is significant interest in long noncoding RNAs (lncRNAs). They have a significant impact on the immune microenvironments of tumors and immunotherapy. In addition, lncRNAs regulate many crucial mechanisms of cancer immunity, including the presentation of antigens and the exhaustion of T cells. According to our prior investigation, the immune-associated lncRNA zinc finger protein 710 (ZNF710)-AS1-201 has the potential to function as an indicator of GC patient prognosis (recurrence, metastasis and survival). The primary focus of this study was to assess the correlation between ZNF710-AS1-201 and the immune microenvironment and to investigate the roles of ZNF710-AS1-201 in the cellular growth and metastasis processes within GC cells.

**Citation:** Ding W, Chen WW, Wang YQ, Xu XZ, Wang YB, Yan YM, Tan YL. Immune-related long noncoding RNA zinc finger protein 710-AS1-201 promotes the metastasis and invasion of gastric cancer cells. *World J Gastrointest Oncol* 2024; 16(2): 458-474

**URL:** <https://www.wjgnet.com/1948-5204/full/v16/i2/458.htm>

**DOI:** <https://dx.doi.org/10.4251/wjgo.v16.i2.458>

## INTRODUCTION

Gastric cancer (GC) is a prevalent malignant tumor affecting the gastrointestinal tract. At present, surgical excision is the favored approach for treating stomach cancer[1]. The prognosis of GC is mainly influenced by the recurrence and spread of the disease after surgery[2]. To prevent the recurrence and spread of tumors after surgery, adjuvant chemotherapy, radiotherapy, targeted therapy, and immunotherapy are employed[3]. The progression of GC is significantly influenced by recent advancements in comprehending the immune microenvironment[4]. Compared to traditional chemotherapy and radiotherapy, immunotherapy is more effective and less toxic[5]. The current methods of immunotherapy consist immune checkpoint inhibitors, adoptive cell therapy, vascular endothelial growth factor inhibitors, cancer vaccines, and chimeric antigen receptor T cell therapy[6]. Immune checkpoint therapy has yielded surprising results[7]. The related drugs include pembrolizumab, nivolumab, avelumab and so on[8]. Nevertheless, the overall prognosis for advanced GC remains poor. Currently, there is a pressing requirement to identify additional, more efficient therapeutic targets.

In the field of oncology, there is a significant interest in long noncoding RNAs (lncRNAs). It has a significant impact on the immune microenvironments of tumors and immunotherapy[9,10]. In addition, lncRNAs regulate many crucial mechanisms of cancer immunity, including the presentation of antigens and the exhaustion of T cells[11]. According to our prior investigation, the immune-associated lncRNA zinc finger protein 710 (ZNF710)-AS1-201 has the potential to function as an indicator of patient prognosis (recurrence, metastasis and survival) in cases of GC[12]. The study illustrated that patients with high ZNF710-AS1-201 expression in tumor tissues had poorer disease-free survival (DFS) than patients with low ZNF710-AS1-201 expression in both The Cancer Genome Atlas (TCGA) data and our data. ZNF710-AS1-201 is one of the transcripts of the ZNF710-AS1 gene, which acts as the opposite strand to ZNF710. The primary focus of this study was to assess the correlation between ZNF710-AS1-201 and the immune microenvironment and to investigate the roles of ZNF710-AS1-201 in the cellular growth and metastasis processes within GC cells.

## MATERIALS AND METHODS

### Data collection

The RNAseq data of all patients in the TCGA-stomach adenocarcinoma database for stomach cancer were acquired from The UCSC Xena website (<https://xenabrowser.net/datapages/>). To standardize the transcriptome sequencing data, the utilization of fragments per kilobase of transcript per million mapped reads was implemented. According to the median ZNF710-AS1-201 expression level, the patients were categorized into two groups: High expression (HExp) and low expression (LExp) groups.

### Immune correlation analysis

To compare the infiltration of immune cells among each sample, we employed single-sample gene set enrichment analysis using the R package “GSVA (Ver.1.44.5)” [13]. We acquired the feature genes of various immune cells from the TISIDB website (<http://cis.hku.hk/TISIDB/>). Additionally, we utilized the R package “estimate (Ver. 1.0.13)” to assess the stromal and immune cells within every sample, and a measurement was conducted. Wilcoxon analysis with the R package “stats (Ver. 4.2.0)” was utilized to compare the infiltration of various immune cells, including stromal and immune cells, as well as immune checkpoints, between the HExp and LExp groups.

### Analysis of antitumor drug sensitivity

The R package “oncoPredict (Ver. 1.0.13)” was utilized to compute the half-maximal inhibitory concentrations (IC<sub>50</sub>s) of chemotherapy and targeted medications for every individual [14]. We compared the IC<sub>50</sub>s of 19 antitumor drugs (5-fluorouracil, afatinib, lapatinib, axitinib, cisplatin, crizotinib, dabrafenib, dasatinib, epirubicin, gefitinib, gemcitabine, irinotecan, lapatinib, nilotinib, oxaliplatin, paclitaxel, sorafenib, trametinib and vorinostat) between the HExp and LExp groups using Wilcoxon analysis.

### Analysis of functional enrichment

The R package “limma (Ver. 1.3.54.2)” was used to screen for differentially expressed genes (DEGs) between the HExp and LExp groups, with a false discovery rate threshold (FDR) of less than 0.05 and a log<sub>2</sub> | fold change (FC) | greater than 1. Utilizing the R package “clusterProfiler (Ver. 4.4.4)” to assess these DEGs, we performed an enrichment analysis using Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) databases.

### Patients and specimens

A total of 68 patients who underwent radical surgery for GC in Changzhou Wujin People’s Hospital were enrolled from January 2014 to November 2014. Below are the criteria for inclusion. First, the pathological diagnosis was gastric adenocarcinoma, and no distant metastasis was found on imaging. Second, none of the patients received preoperative chemotherapy or targeted therapy. Finally, all follow-up data were complete. Tumor and adjacent tissues were collected immediately after tumor isolation and stored in liquid nitrogen.

### Cell culture

The HGC-27 (CL-0107) and MKN-45 (CL-0292) cell lines of GC, obtained from Procell Life Science and Technology Co. (Wuhan, China), were cultured in Roswell Park Memorial Institute (RPMI)-1640 medium (Procella, Wuhan, China) at 37 °C in a humid atmosphere consisting of 5% CO<sub>2</sub> and 95% air. RPMI-1640 contained 10% fetal bovine serum (FBS) (Procella, Wuhan, China) and 1% penicillin-streptomycin solution (Procell, Wuhan, China) as the medium.

### Cell transfection

Yuanzi Biotech Co. (Shanghai, China) synthesized the overexpression and shRNA plasmids of ZNF710-AS1-201. The most efficient shRNA of the three was selected based on the experimental findings. For all cell transfections, Lipofectamine 3000 (Invitrogen, Carlsbad, CA, United States) was administered according to the instructions provided by Thermo Fisher for Lipofectamine 3000 Transfection Reagent.

### Quantitative real-time polymerase chain reaction

TRIzol reagent (TaKaRa, Dalian, China) was used to extract RNA from HGC-27 and MKN-45 cells following the supplier's instructions, resulting in a total amount of RNA. The RNA sample was converted into complementary DNA using a kit from Moloney-murine leukemia virus (Promega Biotech, Beijing, China). To assess the RNA expression level in GC cells, SYBR Green Master Mix from Takara (Dalian, China) was utilized. Glyceraldehyde-3-phosphate dehydrogenase was employed as an endogenous control. The calculation of RNA expression was determined using the 2<sup>-ΔΔCT</sup> approach. Table 1 displays the introductory details.

### Assessment of cellular viability

The cell suspension (100 μL/well) was cultured in 96-well plates and incubated for 24, 48, and 72 h in the incubators. Next, we added 10 μL of cell counting kit-8 (CCK-8) (Beyotime, Shanghai, China) solution to every well. An enzyme label instrument (BioTek, Beijing, China) was used to measure the absorbance at 450 nm.

### Assessment of cellular growth

The Cell-Light EdU Apollo567 In Vitro kit (RiboBio, China) was utilized to conduct EdU incorporation assays. The

exponential increase was calculated for vascular smooth muscle cells plated in 96-well dishes at a density of  $4 \times 10^3$  cells per well. Following 24 h of the specified treatment, each well was exposed to 100  $\mu$ L of 50  $\mu$ M EdU medium for a duration of 2 h. Subsequently, the cells were immobilized in a PBS solution containing 4% paraformaldehyde for a period of 30 min. Next, a solution containing 2 milligrams per milliliter of glycine was introduced for a duration of 5 min. Following rinsing with phosphate buffered saline (PBS), the cells were subjected to an incubation period of 30 min with a 1  $\times$  Apollo staining solution. After discarding the staining solution, the cells were rinsed with PBS containing 0.5% Triton X-100 for 10 min. Following another round of washing with PBS, 1  $\times$  Hoechst 33342 was introduced and incubated at room temperature for 30 min. Following washing with PBS, fluorescence microscopy was used to observe the positive cells.

### Cell apoptosis assay

The evaluation of cell apoptosis was conducted through a flow cytometry test (BD, New Jersey, United States). After being rinsed twice in PBS, the cells were then suspended in 1  $\times$  binding buffer. For the control group, 400 microliters of 1  $\times$  binding buffer was utilized, while for the remaining treatment groups, 100 microliters of 1  $\times$  binding buffer was employed. Then, 100 microliters was extracted from the aforementioned solution and transferred to the flow tube. Five microliters of Fluorescein isothiocyanate-Annexin V and five microliters of propidium iodide (at a concentration of 50 micrograms per milliliter) was added. The sample was left at room temperature (25  $^{\circ}$ C) and away from light for a duration of 15 min. Next, 400  $\mu$ L of 1  $\times$  binding buffer was added to each tube and mixed well. Analysis using flow cytometry was conducted within one hour. FlowJo software (BD, New Jersey, United States) was utilized to analyze the data.

### Assays for cell metastasis

To evaluate the cell metastasis characteristics, a scratch assay was conducted in vitro by creating a linear scratch across the center of the layer of cultured cells. Pictures were captured at regular intervals (every 24 h), and the gathered data were utilized to analyze the experimental outcomes.

### Cell invasion assays

The invasion of cells was observed by employing the Transwell assay with Transwell kits from Corning, United States. The Matrigel adhesive was melted overnight at a temperature of 4  $^{\circ}$ C and then mixed with the serum-free culture after dilution. The diluted Matrigel adhesive was applied onto the upper chamber and left in the incubator for 1-1.5 h until the adhesive hardened. After terminating digestion, the cells were centrifuged while waiting for the Matrigel gum to solidify. To adjust the cell density to  $5 \times 10^5$  cells/mL, the cells were suspended in medium that included 2% FBS. The upper chamber was inoculated with 200  $\mu$ L of cell suspension, with 3 replicates per group. The lower chamber was supplemented with 800  $\mu$ L of medium containing 20% FBS and cultured at 37  $^{\circ}$ C and 5% CO<sub>2</sub> for 48 h. After removing the chamber, the medium was discarded, and the chamber was washed twice with PBS. Then, the cells were fixed with 4% paraformaldehyde at room temperature for 30 min. The paraformaldehyde was disposed of and rinsed two times with PBS. After being thoroughly dried, the chamber was treated with crystal violet stain for 15 min. The excess crystal violet dye was washed away with PBS. A microscope (Olympus, Japan) was used to count the cells.

### Assays for western blotting (WB)

WB was used to analyze the protein expression levels associated with apoptosis. Protein concentrations were measured with a bicinchoninic acid protein assay kit from Thermo, United States. Bax (Abcam, ab32503, 1:1000), Bcl2 (Abcam, ab182858, 1:2000), and  $\beta$ -actin (Affinity, AF7018, 1:3000) were left to react with the membrane overnight at 4  $^{\circ}$ C. The blots were visualized using the Millipore Enterochromaffin-like detection system.


### Statistical analysis

All analyses were conducted using GraphPad Prism (Version 9.4.1) and R (Version 4.2.0). mean  $\pm$  SD deviation values were used to present all continuous variable results. The *t* test was used if the variance was homogeneous, and the Mann-Whitney test was used if the variance was not homogeneous. Categorical variables were displayed using ratios. Differences between components were tested using chi-square tests. The grayscale values of bands on WB images were calculated using ImageJ from NIH (Bethesda, MD). The screening criteria for a significant difference were  $P < 0.05$ ,  $P < 0.01$ , or  $P < 0.001$ .

## RESULTS

### ZNF710-AS1-201-associated immune microenvironment

To assess the relationship between ZNF710-AS1-200 and the immune microenvironment of GC patients, we analyzed the infiltration of immune cells in each sample. The findings demonstrated that the level of infiltration by central memory CD8 T cells, activated CD4 T cells, gamma delta T cells, type 17 T helper cells, CD56bright natural killer cells, CD56dim natural killer cells, and natural killer T cells was elevated in GC samples exhibiting reduced expression of ZNF710-AS1-201 (Figure 1A). Conversely, GC samples exhibiting high levels of ZNF710-AS1-201 displayed an increased infiltration level of effector memory CD4 T cells and memory B cells. Therefore, there was no significant variation in immune scores observed between the two groups exhibiting high or LExp (Figure 1B). Remarkably, the stromal scores was increased the high-expression cohort. Figure 1C demonstrates that B and T lymphocyte attenuator, butyrophilin-like 9, CD40LG,



DOI: 10.4251/wjgo.v16.i2.458 Copyright ©The Author(s) 2024.

**Figure 1 Immune correlation analysis of zinc finger protein 710-AS1-201.** A: The infiltration levels of 28 immune cells in the high and low expression (LExp) groups; B: The stromal score, immune score, and ESTIMATE score in the high and LExp groups; C: The expression levels of 19 immune checkpoints in the high and LExp groups. <sup>a</sup>*P* < 0.05; <sup>b</sup>*P* < 0.01; <sup>c</sup>*P* < 0.001; ns: Not significant.


CD160, and tumor necrosis factor superfamily member 14 exhibited elevated levels of expression in GC samples with HExp of ZNF710-AS1-201. Conversely, PVR, HLA-DRB1, CD47, HLA-DMA, KIR2DL4, HAVCR2, HLA-DQB1, CD276, CD80, HLA-A, HLA-C, TNFRSF18, CD274, and CD86 displayed increased expression in GC samples with LExp of ZNF710-AS1-201.

#### Antitumor drug sensitivity analysis based on ZNF710-AS1-201 expression level

We additionally evaluated the correlation between the expression of ZNF710-AS1-201 and the responsiveness to antitumor drugs in GC. Figure 2 demonstrates that the IC<sub>50</sub>s of 5-fluorouracil, cisplatin, gemcitabine, and trametinib were reduced in the low-expression group, whereas the IC<sub>50</sub>s of dasatinib and vorinostat were elevated. The HExp and LExp groups showed no notable disparity in the IC<sub>50</sub>s of other medications.

#### Functional enrichment analysis (FEA) based on ZNF710-AS1-201

Initially, we examined the disparity between the groups with high and LExp levels and identified 647 genes that exhibited differential expression (Supplementary Table 1). Then, we conducted FEA of these DEGs. In conclusion, KEGG analysis revealed significant enrichment in 28 mechanisms, whereas GO analysis showed significant enrichment in 565 functions (Figure 3). In the GO analysis, there was a significant enrichment of collagen-containing extracellular matrix, actin binding, G protein-coupled peptide receptor activity, ion channel activity, and others (Figure 3A). Significant



**Figure 2** Sensitivity to different antitumor drugs in the high and low expression groups. A: Half-maximal inhibitory concentrations ( $IC_{50}$ s) of 5-fluorouracil; B:  $IC_{50}$ s of axitinib; C:  $IC_{50}$ s of cisplatin; D:  $IC_{50}$ s of gemcitabine; E:  $IC_{50}$ s of trametinib; F:  $IC_{50}$ s of vorinostat.  $IC_{50}$ s: Half-maximal inhibitory concentrations.

enrichment of gastric acid secretion, chemical carcinogenesis-receptor activation, and other pathways was observed in the KEGG analysis (Figure 3B).

### ZNF710-AS1-201 was closely related to GC


Clinical specimens were gathered to measure the expression level of ZNF710-AS1-201 in the samples. The findings indicated a notable increase in the expression level of ZNF710-AS1-201 in GC tissues compared to adjacent tissues (Figure 4A). Subsequently, we divided patients into high-expression and low-expression groups based on the expression level of ZNF710-AS1-201 in cancer tissues. The comparison of the clinicopathological features of patients between the low and HExp groups is shown in Table 2. The malignant degree of GC was higher and the stage was later in the high-expression group. Additionally, patients with HExp of ZNF710-AS1-201 had lower OS and DFS (Figure 4B and C). Consistent with this, the expression level of ZNF710-AS1-201 in GC cell lines (HGC-27, AGS, NCI-N87, and MKN-45) was elevated compared to that in the normal gastric mucosa cell line GES-1 (Figure 4D). To examine the biological role of ZNF710-AS1-201 in GC, HGC-27 cells were transfected with a plasmid overexpressing ZNF710-AS1-201 and an empty plasmid, while MKN-45 cells were transfected with a plasmid containing ZNF710-AS1-201 shRNA and an empty plasmid. The results demonstrated a significant increase in ZNF710-AS1-201 expression following the transfection of ZNF710-AS1-201 mimic in HGC-27 cells. Conversely, in MKN-45 cells, ZNF710-AS1-201 expression was markedly decreased upon transfection with ZNF710-AS1-201 shRNA (Figure 4E and F).

### ZNF710-AS1-201 promoted GC cell viab

Cell viability was assessed using the CCK-8 assay. After 24 h, 48 h, and 72 h of transfection with the ZNF710-AS1-201 overexpression plasmid, the viability of HGC-27 cells was significantly increased (Figure 5A). The viability of MKN-45 cells decreased when transfected with ZNF710-AS1-201 shRNA plasmid for 24, 48, and 72 h (Figure 5A). Furthermore, the EdU assay was utilized to assess cellular proliferation. The proliferation of HGC-27 cells was enhanced following transfection of the plasmid overexpressing ZNF710-AS1-201 for 72 h (Figure 5B). The growth rate of MKN-45 cells decreased after treatment with the ZNF710-AS1-201 shRNA plasmid for 72 h (Figure 5B).

### ZNF710-AS1-201 inhibited GC cell apoptosis

The quantification of proteins related to apoptosis and the assessment of cell apoptosis were conducted using qRT-PCR, western blot, and flow cytometry. qRT-PCR and Western blot analysis revealed that ZNF710-AS1-201 overexpression in HGC-27 cells significantly decreased the expression of Bax and increased the expression of Bcl-2 (Figure 6A and B). In contrast, ZNF710-AS1-201 knockdown in MKN-45 cells led to an increase in Bax levels and a decrease in Bcl-2 levels (Figure 6A and B). The flow cytometry assay results showed that the overexpression of ZNF710-AS1-200 significantly reduced HGC-27 cell apoptosis after 24 h, 48 h, and 72 h (Figure 6C). In contrast, knockdown of ZNF710-AS1-201 significantly enhanced MKN-45 cell apoptosis at 48 h and 72 h (Figure 6D).




**Figure 3 Gene ontology and kyoto encyclopedia of genes and genomes analysis of zinc finger protein 710-AS1-201.** A: Gene ontology assessment of differentially expressed genes (DEGs) between the high and low expression (LExp) groups; B: Kyoto encyclopedia of genes and genomes evaluation of DEGs between the high and LExp groups.

**ZNF710-AS1-201 facilitated GC cell metastasis and invasion.**

The study utilized the scratch assay to measure cell metastasis. The findings indicated that the overexpression of ZNF710-AS1-201 significantly enhanced the metastasis of HGC-27 cells after 48 and 72 h (Figure 7A). In contrast, the knockdown of ZNF710-AS1-201 significantly inhibited the metastasis of MKN-45 cells at 48 h and 72 h (Figure 7B). The study utilized the transwell assay to quantify cell invasion. The findings indicated that the overexpression of ZNF710-AS1-201 significantly enhanced HGC-27 cell invasion after 72 h (Figure 7C). In contrast, the invasion of MKN-45 cells was significantly inhibited by knockdown of ZNF710-AS1-201 for 72 h (Figure 7C).

**Downstream target prediction**

After conducting a thorough search in the StarBase database (<https://starbase.sysu.edu.cn/>), we identified the potential targets [ZNF710, isocitrate dehydrogenase-2 (IDH2), Semaphorin 4B (SEMA4B), ARHGAP10, RGMB, hsa-miR-93-5p, and ZNF710-AS1-202] for ZNF710-AS1-201. Additionally, we employed qRT-PCR to analyze the expression levels of these potential targets following cell transfection. The findings indicated that after overexpressing ZNF710-AS1-201 in HGC-27 cells, there were no significant changes observed in the expression levels of IDH2 and miR-93-5p. Moreover, in MKN-45 cells with decreased expression of ZNF710-AS1-201, the expression levels of SEMA4B, ARHGAP10, and ZNF710-AS1-202 also showed no significant changes. Additionally, whether ZNF710-AS1-202 was overexpressed or knocked down in GC



DOI: 10.4251/wjgo.v16.i2.458 Copyright ©The Author(s) 2024.


**Figure 4** Zinc finger protein 710-AS1-201 is upregulated in gastric cancer tissues and cell lines. A: The expression level of zinc finger protein 710 (ZNF710)-AS1-201 in 64 paired gastric cancer and nontumoral tissues was determined by quantitative real-time polymerase chain reaction (qRT-PCR) assay; B and C: Patients with high expression of ZNF710-AS1-201 had lower overall survival and disease-free survival rates; D: The expression level of ZNF710-AS1-201 in a normal human gastric mucosa cell line and gastric cancer cell lines (HGC-27, AGS, NCI-N87 and MKN-45) was determined by qRT-PCR; E: The expression level of ZNF710-AS1-201 in HGC-27 cells after transfection with the overexpression plasmid; F: The expression level of ZNF710-AS1-201 in MKN-45 cells after transfection with the shRNA plasmid. Data are shown as the mean  $\pm$  SD. ZNF710: Zinc finger protein 710. \* $P < 0.05$ ; <sup>a</sup> $P < 0.01$ ; <sup>b</sup> $P < 0.001$ ; ns: Not significant.

cells, there was a consistent decrease in the expression levels of RGMB. However, the only expression change that was statistically significant and logically coherent was that for ZNF710 (Figure 8A and B).

## DISCUSSION

This study focused on examining the contribution of ZNF710-AS1-201, an immune-related lncRNA, in facilitating the advancement and spread of GC. In line with prior research, GC tissues with low levels of ZNF710-AS1-201 exhibited significant infiltration of various antitumor immune cells[15-18], such as memory CD8 T cells, activated CD4 T cells, tumor-infiltrating gamma delta T cells, natural killer cells, and natural killer T cells. Furthermore, there was a significant presence of memory B cells in GC tissues exhibiting elevated levels of ZNF710-AS1-201 expression. These findings align with the outcomes of prior research indicating that memory B cells have the potential to facilitate the development of cancer[19]. Nevertheless, the recruitment and activation of these immune cells by ZNF710-AS1-201 are still uncertain. While the ESTIMATE scores of GC samples with elevated expression of ZNF710-AS1-201 were greater than those of samples with LExp, the observed disparity did not reach statistical significance. In contrast, the HExp group exhibited a considerably elevated stromal score. The reason for this could be an escalation in the remodeling of the extracellular matrix, which has the potential to stimulate the growth of tumors, their spread to other parts of the body, and the formation of new blood vessels[20].

Chemotherapy, targeted therapy, and immunotherapy are the present methods used for conservatively treating advanced GC[21]. By analyzing immune checkpoints and drug sensitivity, we have the ability to choose drugs that exhibit high sensitivity and immunotherapies targeting highly expressed immune checkpoints. Currently, the first-line chemotherapy treatment for GC is the combination of cisplatin and 5-fluorouracil[22]. The survival of patients with




**Figure 5** Zinc finger protein 710-AS1-201 promotes gastric cancer cell viability and proliferation. A: Cell viability was examined by cell counting kit-8 assay after cells were transfected for 24 h, 48 h, and 72 h; B: The cell proliferation rate was detected by EdU assay. <sup>a</sup>*P* < 0.05; <sup>b</sup>*P* < 0.01.

human epidermal growth factor receptor 2-positive GC is improved by adding trastuzumab to the initial chemotherapy [23]. Furthermore, pembrolizumab, an immune checkpoint inhibitor, has been authorized in various contexts based on the presence of programmed death ligand 1[24]. According to prior studies, axitinib exhibited strong antineoplastic effects on human GC, either by itself or when combined with 5-fluorouracil or cisplatin[25]. Considering these results and those of our study, the combination of axitinib and 5-fluorouracil may demonstrate enhanced antitumor impacts in individuals exhibiting elevated ZNF710-AS1-201 expression.


Based on the enrichment analysis, we identified several biological processes and mechanisms that are associated with tumors, such as collagen-containing extracellular matrix, actin binding, G protein-coupled peptide receptor activity and ion channel activity. According to the study conducted by Chen *et al*[26], the presence of collagen in the tumor microenvironment is an independent predictor of lymphatic metastasis in early GC. Jia *et al*[27] discovered that Anillin, a protein that binds to actin, facilitated the advancement of GC both in laboratory settings and in mice. Yan *et al*[28] demonstrated that the abnormal activation of G-protein-coupled receptors stimulated the growth and spread of GC cells, altered the tumor microenvironment, and enhanced immune evasion[28]. In GC, Chen *et al*[29] observed modified expression and/or malfunction of ion channels/transporters, which is believed to have a significant impact on tumor cell proliferation, metastasis, invasion, and apoptosis. In summary, ZNF710-AS1-201 might exert its influence on GC *via* these pathways, thereby facilitating the onset and progression of GC.

In a recent study, Li and colleagues found that ZNF710-AS1-202 played a role in enhancing cell growth and suppressing cell death in clear cell renal cell carcinoma through the reverse regulation of ZNF710[30]. Nevertheless, there is no existing report on the regulatory impact of ZNF710-AS1-201 on tumors. The current research examined the influence of ZNF710-AS1-201 on the promotion of tumors in GC. In HGC-27 cells, we observed that the upregulation of ZNF710-AS1-201 stimulated cell growth, hindered cell death, and enhanced cell metastasis and infiltration. Conversely, in MKN-45 cells, the downregulation of ZNF710-AS1-201 suppressed cell proliferation, promoted cell death, and impeded cell metastasis and invasion.

qRT-PCR was used to validate possible targets for ZNF710-AS1-201. IDH2, also known as isocitrate dehydrogenase 2, is an isocitrate dehydrogenase that relies on NADP and is located in the mitochondria. Through inhibition of matrix



**Figure 6 Zinc finger protein 710-AS1-201 inhibits gastric cancer cell apoptosis.** A: Quantitative real-time polymerase chain reaction was used to detect the expression levels of apoptosis-related genes (Bax and Bcl-2); B: Western blot analysis was used to detect the expression levels of apoptosis-related proteins (Bax and Bcl-2); C: Cell apoptosis in HGC-27 cells was examined by flow cytometry after transfection with the zinc finger protein 710 (ZNF710)-AS1-201 overexpression plasmid for 24 h, 48 h, and 72 h; D: Cell apoptosis in MKN-45 cells was examined by flow cytometry after transfection with the ZNF710-AS1-201 shRNA plasmid for 24 h, 48 h, and 72 h. <sup>a</sup> $P < 0.05$ ; <sup>b</sup> $P < 0.01$ ; <sup>c</sup> $P < 0.001$ ; ns: Not significant.




DOI: 10.4251/wjgo.v16.i2.458 Copyright ©The Author(s) 2024.

**Figure 7 Zinc finger protein 710-AS1-201 promotes gastric cancer cell metastasis and invasion.** A: Cell metastasis in HGC-27 cells was examined by scratch assays after transfection of zinc finger protein 710 (ZNF710)-AS1-201-overexpressing plasmids for 24 h, 48 h, and 72 h; B: Cell metastasis in MKN-45 cells was examined by scratch assay after transfection with ZNF710-AS1-201 shRNA plasmid for 24 h, 48 h, and 72 h; C: Cell invasion was examined by Transwell assay after transfection for 72 h. ZNF710: Zinc finger protein 710. <sup>a</sup>*P* < 0.05; <sup>b</sup>*P* < 0.01; ns: Not significant.

metalloproteinase 7, it has been discovered to impede the invasion of GC cells[31]. However, a significant reduction in IDH2 mRNA expression after ZNF710-AS1-201 overexpression appeared in MKN-45 cells. ARHGAP10 can promote gastric tumor metastasis[32]. Nonetheless, the upregulation of ZNF710-AS1-201 suppressed the expression of ARHGAP10 mRNA in HGC-27 cells. In their study, Jian and colleagues demonstrated that the invasion of non-small cell lung cancer was hindered by SEMA4B through the suppression of matrix metalloproteinase 9 activation, which was dependent on phosphoinositide 3-kinase[33]. According to the study conducted by Li *et al*[34], the overexpression of miR-93-5p was found to enhance the growth, movement, infiltration, and resistance to chemotherapy in SGC-7901 cells. In line with these perspectives, the mRNA expression of SEMA4B was inhibited when ZNF710-AS1-201 was overexpressed in HGC-27 cells, whereas the expression of miR-93-5p was enhanced when ZNF710-AS1-201 was knocked down in MKN-45 cells. Further investigation is needed to explore the role of these genes in GC, even though they might be controlled by ZNF710-AS1-201.

ZNF710-AS1-201 had a negative regulatory effect on the expression level of ZNF710 in both HGC-27 and MKN-45 cells. However, this result only showed that ZNF710 is negatively regulated by ZNF710-AS1-201. The function of ZNF710 in GC is currently not understood. ZNF710, as stated in the UniProt repository, is a transcription factor (TF)/cofactor of the Kruppel Cys2-His2-type zinc-finger protein[35]. ZNF710 may function as a TF to regulate the occurrence and development of GC. TFs are key factors in gene regulation[36]. TFs can regulate the migration of tumor cells through specific signaling pathways[37]. In their study, Guo and colleagues discovered that the TF RUNX2 stimulated the yes-associated protein 1 signaling pathway, contributing to the development of GC[38]. In their study, Zhao and colleagues



**Figure 8 Prediction of downstream targets of zinc finger protein 710-AS1-201.** A: The expression levels of genes in HGC-27 cells transfected with the zinc finger protein 710 (ZNF710)-AS1-201 overexpression plasmid were determined by quantitative real-time polymerase chain reaction (qRT-PCR); B: The expression levels of genes in MKN-45 cells transfected with the ZNF710-AS1-201 shRNA plasmid were determined by qRT-PCR. <sup>a</sup>*P* < 0.05; <sup>b</sup>*P* < 0.01; <sup>c</sup>*P* < 0.001; ns: Not significant.

**Table 1 Sequences of primers used in this study**

Name	Primer sequence (5' to 3')
ZNF710-AS1-201-F	ACCGCAGGCACITTTGAAGA
ZNF710-AS1-201-R	AAGGGACATCAGAGGGGAG
Bcl-2-F	GGTGGGGTCATGTGTGTGG
Bcl-2-R	CGGTTCAAGTACTCAGTCATCC
Bax-F	CCCAGAGGTCTTTTTCCGAG
Bax-R	CCAGCCCATGATGGTTCTGAT
ZNF710-F	AACAGGAGGTCTATGAGGTTTCT
ZNF710-R	CGCTGAGGTCGATCATCTTGA
IDH2-F	CGCCACTATGCCGACAAAAG
IDH2-R	ACTGCCAGATAATACGGGTCA
SEMA4B-F	GAGCGGCCATTCTCAGATTC
SEMA4B-R	CACCCACGTACAGGGTCTCT
ARHGAP10-F	CCCAGCGGAAGTTTGCTCAT
ARHGAP10-R	ACAGCTCCAAGTTGCTTTTTTC
RGMB-F	CCAGCCCAATGTGAATCCA
RGMB-R	GGTCACTGATACCCAACACGG
miR-93-5p-F	CGCAAAGTGCTGTTCCGTGC
miR-93-5p-R	AGTGCAGGGTCCGAGGTATT
ZNF710-AS1-202-F	GTCCGATCACAGTTCACTAC
ZNF710-AS1-202-R	ATACCAAACCAGGCAACATA

ZNF710: Zinc finger protein 710; IDH2: Isocitrate dehydrogenase-2; SEMA4B: Semaphorin 4B.

**Table 2 Comparison of the clinical features of patients between the low- and high-expression groups, mean  $\pm$  SD**

Features	ZNF710-AS1-201		P value
	Low expression (n = 34)	High expression (n = 34)	
Age, yr	65.62 $\pm$ 6.85	67.85 $\pm$ 8.25	0.228
Sex, male (%)	19 (55.9)	22 (64.7)	0.457
CEA, U/mL	18.32 $\pm$ 69.96	11.09 $\pm$ 24.58	0.415 <sup>1</sup>
Neutrophil count, $\times 10^9$ /L	3.56 $\pm$ 0.97	4.67 $\pm$ 2.08	0.006 <sup>1,a</sup>
Lymphocyte count, $\times 10^9$ /L	1.89 $\pm$ 0.62	1.42 $\pm$ 0.47	< 0.001 <sup>a</sup>
Monocyte count, $\times 10^9$ /L	0.38 $\pm$ 0.18	0.44 $\pm$ 0.24	0.240
Differentiation degree, III-IV (%)	18 (52.9)	26 (76.5)	0.042 <sup>a</sup>
T stage, III-IV (%)	15 (44.1)	28 (82.4)	0.001 <sup>a</sup>
Lymphatic invasion, positive (%)	15 (44.1)	24 (70.6)	0.044 <sup>a</sup>

<sup>a</sup>P < 0.05.

<sup>1</sup>Mann-Whitney test.

CEA: Carcinoembryonic antigen; ZNF710: Zinc finger protein 710.

found that the growth and metastasis of GC cells were suppressed by the TF kruppel-like factor 15 through the regulation of the TFAP2A-AS1/Neonatal Ichthyosis Sclerosing Cholangitis axis[39]. This demonstrated that transcription regulators had a controlling function in the metastasis and invasion of stomach cancer.

## CONCLUSION

The findings from this study indicate that ZNF710-AS1-201 facilitated growth, metastasis and invasion while suppressing cell death in GC cells. These findings suggest that ZNF710-AS1-201 and ZNF710 have potential as effective targets for therapeutic intervention in GC. Nevertheless, these findings have not been observed in living organisms. Furthermore, rescue experiments are needed to demonstrate the interaction mechanism between ZNF710-AS1-201 and ZNF710. Moreover, it is still necessary to determine the specific targets of the ZNF710 TF.

## ARTICLE HIGHLIGHTS

### Research background

Gastric cancer (GC) is a prevalent malignant tumor of the gastrointestinal system. ZNF710 is a transcription factor, and ZNF710-AS1-201 is an immune-related long noncoding RNA (lncRNA) that is upregulated in GC cells.

### Research motivation

The research is motivated by the pressing challenges posed by GC and the need for a deeper understanding of its underlying molecular mechanisms. Specifically, investigating the functional role of lncRNA ZNF710-AS1-201 and exploring the dynamics of the immune microenvironment are crucial for unraveling the complexities of GC progression. The study aims to contribute valuable insights that may have implications for precision therapy approaches in addressing this formidable disease.

### Research objectives

The primary objectives of this research are to elucidate the functional significance of lncRNA ZNF710-AS1-201 in GC, analyze its impact on cellular processes and pathways, investigate its potential as a biomarker, and comprehensively characterize the immune microenvironment in GC tissues. By achieving these objectives, the study aims to provide a foundation for developing targeted therapeutic strategies, advancing our understanding of GC pathogenesis, and contributing to the broader field of cancer research.

### Research methods

This research employed a multidisciplinary approach, integrating molecular biology techniques, bioinformatics analyses, and *in vitro* experiments. The study involved the profiling of lncRNA ZNF710-AS1-201 expression in GC tissues, functional assays to assess its impact on cell behavior, and bioinformatics tools to unravel potential interacting pathways. Additionally, immune microenvironment analysis was conducted to explore the relationship between lncRNA expression and immune responses in GC. These methods collectively provided a comprehensive platform for investigating the role of ZNF710-AS1-201 in GC progression.

### Research results

The research revealed a significant upregulation of lncRNA ZNF710-AS1-201 in GC tissues compared to adjacent normal tissues. Functional assays demonstrated that ZNF710-AS1-201 overexpression promoted cell proliferation, migration, and invasion, suggesting its potential oncogenic role. Bioinformatics analyses unveiled the involvement of ZNF710-AS1-201 in key cancer-related pathways. Moreover, immune microenvironment analysis indicated a correlation between ZNF710-AS1-201 expression and immune cell infiltration, providing insights into its potential influence on the tumor immune response in GC.

### Research conclusions

In conclusion, this study elucidates the crucial role of lncRNA ZNF710-AS1-201 in GC progression, highlighting its potential as a diagnostic and therapeutic target. The findings underscore the significance of ZNF710-AS1-201 in modulating cellular processes and its association with cancer-related pathways. Furthermore, insights into the immune microenvironment suggest its impact on the tumor immune response. Overall, these results contribute to our understanding of the molecular mechanisms underlying GC and open avenues for future research and targeted therapeutic interventions.

### Research perspectives

The study opens new perspectives for further investigation into the intricate mechanisms of lncRNA ZNF710-AS1-201 in GC. Future research should explore its potential as a biomarker for early diagnosis and prognosis. Additionally, understanding its interactions within the tumor microenvironment and elucidating its role in therapeutic resistance could pave the way for innovative treatment strategies. Collaborative efforts integrating multi-omics approaches may provide a

comprehensive view of ZNF710-AS1-201's involvement in GC, offering valuable insights for precision medicine and targeted therapies.

---

## ACKNOWLEDGEMENTS

---

The authors would like to acknowledge Professor Li Yang for guidance and support.

---

## FOOTNOTES

---

**Author contributions:** Ding W contributed to conceptualization; Chen WW and Wang YB contributed to methodology; Wang YQ contributed to validation; Wang YB contributed to formal analysis; Xu XZ contributed to resources; Ding W contributed to writing original draft preparation; Ding W and Tan YL contributed to writing-review and editing; Yan YM contributed to supervision; Tan YL contributed to project administration; Ding W and Tan YL contributed to funding acquisition; all authors contributed to the study and approved the submitted version of the manuscript.

**Supported by** Changzhou Sci and Tech Program, No. CJ20220008; Young Talent Development Plan of Changzhou Health Commission, No. CZQM2020118; Changzhou High-Level Medical Talents Training Project, No. 2022CZBJ105; Cultivation Project of Changzhou Medical Center, Nanjing Medical University, No. CMCB202211; and Development Foundation of Affiliated Hospital of Xuzhou Medical University, No. XYFC202304, and No. XYFM202307.

**Institutional review board statement:** The study was reviewed and approved by the Institutional Review Board at Wujin Hospital affiliated with Jiangsu University approved this study.

**Institutional animal care and use committee statement:** All procedures involving animals were reviewed and approved by the Institutional Animal Care and Use Committee of the Wujin Hospital affiliated with Jiangsu University.

**Conflict-of-interest statement:** All other authors have nothing to disclose.

**Data sharing statement:** No additional data are available.

**ARRIVE guidelines statement:** The authors have read the ARRIVE guidelines, and the manuscript was prepared and revised according to the ARRIVE guidelines.

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** China

**ORCID number:** Wei Ding [0000-0001-6990-8621](https://orcid.org/0000-0001-6990-8621); Yu-Lin Tan [0000-0002-3278-7624](https://orcid.org/0000-0002-3278-7624).

**S-Editor:** Qu XL

**L-Editor:** A

**P-Editor:** Cai YX

---

## REFERENCES

---

- 1 Sun H, Wang H, Pan H, Zuo Y, Zhao R, Huang R, Xue Y, Song H. CD19 (+) B Cell Combined with Prognostic Nutritional Index Predicts the Clinical Outcomes of Patients with Gastric Cancer Who Underwent Surgery. *Cancers (Basel)* 2023; **15** [PMID: [37173997](https://pubmed.ncbi.nlm.nih.gov/37173997/) DOI: [10.3390/cancers15092531](https://doi.org/10.3390/cancers15092531)]
- 2 Zhang T, Wang B, Su F, Gu B, Xiang L, Gao L, Zheng P, Li XM, Chen H. TCF7L2 promotes anoikis resistance and metastasis of gastric cancer by transcriptionally activating PLAUR. *Int J Biol Sci* 2022; **18**: 4560-4577 [PMID: [35864968](https://pubmed.ncbi.nlm.nih.gov/35864968/) DOI: [10.7150/ijbs.69933](https://doi.org/10.7150/ijbs.69933)]
- 3 Joshi SS, Badgwell BD. Current treatment and recent progress in gastric cancer. *CA Cancer J Clin* 2021; **71**: 264-279 [PMID: [33592120](https://pubmed.ncbi.nlm.nih.gov/33592120/) DOI: [10.3322/caac.21657](https://doi.org/10.3322/caac.21657)]
- 4 Chen Y, Jia K, Sun Y, Zhang C, Li Y, Zhang L, Chen Z, Zhang J, Hu Y, Yuan J, Zhao X, Gong J, Dong B, Zhang X, Li J, Shen L. Predicting response to immunotherapy in gastric cancer via multi-dimensional analyses of the tumour immune microenvironment. *Nat Commun* 2022; **13**: 4851 [PMID: [35982052](https://pubmed.ncbi.nlm.nih.gov/35982052/) DOI: [10.1038/s41467-022-32570-z](https://doi.org/10.1038/s41467-022-32570-z)]
- 5 Xiang Y, Gong M, Deng Y, Wang H, Ye D. T cell effects and mechanisms in immunotherapy of head and neck tumors. *Cell Commun Signal* 2023; **21**: 49 [PMID: [36872320](https://pubmed.ncbi.nlm.nih.gov/36872320/) DOI: [10.1186/s12964-023-01070-y](https://doi.org/10.1186/s12964-023-01070-y)]
- 6 Jin X, Liu Z, Yang D, Yin K, Chang X. Recent Progress and Future Perspectives of Immunotherapy in Advanced Gastric Cancer. *Front Immunol* 2022; **13**: 948647 [PMID: [35844558](https://pubmed.ncbi.nlm.nih.gov/35844558/) DOI: [10.3389/fimmu.2022.948647](https://doi.org/10.3389/fimmu.2022.948647)]
- 7 Li K, Zhang A, Li X, Zhang H, Zhao L. Advances in clinical immunotherapy for gastric cancer. *Biochim Biophys Acta Rev Cancer* 2021; **1876**:

- 188615 [PMID: 34403771 DOI: 10.1016/j.bbcan.2021.188615]
- 8 **Bang YJ**, Ruiz EY, Van Cutsem E, Lee KW, Wyrwicz L, Schenker M, Alsina M, Ryu MH, Chung HC, Evesque L, Al-Batran SE, Park SH, Lichinitser M, Boku N, Moehler MH, Hong J, Xiong H, Hallwachs R, Conti I, Taieb J. Phase III, randomised trial of avelumab vs physician's choice of chemotherapy as third-line treatment of patients with advanced gastric or gastro-oesophageal junction cancer: primary analysis of JAVELIN Gastric 300. *Ann Oncol* 2018; **29**: 2052-2060 [PMID: 30052729 DOI: 10.1093/annonc/mdy264]
- 9 **Chen L**, Deng J. Role of non-coding RNA in immune microenvironment and anticancer therapy of gastric cancer. *J Mol Med (Berl)* 2022; **100**: 1703-1719 [PMID: 36329206 DOI: 10.1007/s00109-022-02264-6]
- 10 **Park EG**, Pyo SJ, Cui Y, Yoon SH, Nam JW. Tumor immune microenvironment lncRNAs. *Brief Bioinform* 2022; **23** [PMID: 34891154 DOI: 10.1093/bib/bbab504]
- 11 **Wu M**, Fu P, Qu L, Liu J, Lin A. Long Noncoding RNAs, New Critical Regulators in Cancer Immunity. *Front Oncol* 2020; **10**: 550987 [PMID: 33194608 DOI: 10.3389/fonc.2020.550987]
- 12 **Ding W**, Sun P, Tan Y, Jiang H, Xi C, Zhuang L, Xu Y, Xu X. Construction of a Prognostic Immune-Related lncRNA Risk Model for Gastric Cancer. *J Oncol* 2022; **2022**: 5137627 [PMID: 35794986 DOI: 10.1155/2022/5137627]
- 13 **Hänzelmann S**, Castelo R, Guinney J. GSVA: gene set variation analysis for microarray and RNA-seq data. *BMC Bioinformatics* 2013; **14**: 7 [PMID: 23323831 DOI: 10.1186/1471-2105-14-7]
- 14 **Maeser D**, Gruener RF, Huang RS. oncoPredict: an R package for predicting *in vivo* or cancer patient drug response and biomarkers from cell line screening data. *Brief Bioinform* 2021; **22** [PMID: 34260682 DOI: 10.1093/bib/bbab260]
- 15 **Han J**, Khatwani N, Searles TG, Turk MJ, Angeles CV. Memory CD8(+) T cell responses to cancer. *Semin Immunol* 2020; **49**: 101435 [PMID: 33272898 DOI: 10.1016/j.smim.2020.101435]
- 16 **Borst J**, Ahrends T, Bąbala N, Melief CJM, Kastenmüller W. CD4(+) T cell help in cancer immunology and immunotherapy. *Nat Rev Immunol* 2018; **18**: 635-647 [PMID: 30057419 DOI: 10.1038/s41577-018-0044-0]
- 17 **Saura-Esteller J**, de Jong M, King LA, Ensing E, Winograd B, de Gruijl TD, Parren PWHI, van der Vliet HJ. Gamma Delta T-Cell Based Cancer Immunotherapy: Past-Present-Future. *Front Immunol* 2022; **13**: 915837 [PMID: 35784326 DOI: 10.3389/fimmu.2022.915837]
- 18 **Kucuksezer UC**, Aktas Cetin E, Esen F, Tahrali I, Akdeniz N, Gelmez MY, Deniz G. The Role of Natural Killer Cells in Autoimmune Diseases. *Front Immunol* 2021; **12**: 622306 [PMID: 33717125 DOI: 10.3389/fimmu.2021.622306]
- 19 **Downs-Canner SM**, Meier J, Vincent BG, Serody JS. B Cell Function in the Tumor Microenvironment. *Annu Rev Immunol* 2022; **40**: 169-193 [PMID: 35044794 DOI: 10.1146/annurev-immunol-101220-015603]
- 20 **Oya Y**, Hayakawa Y, Koike K. Tumor microenvironment in gastric cancers. *Cancer Sci* 2020; **111**: 2696-2707 [PMID: 32519436 DOI: 10.1111/cas.14521]
- 21 **Nakamura Y**, Kawazoe A, Lordick F, Janjigian YY, Shitara K. Biomarker-targeted therapies for advanced-stage gastric and gastro-oesophageal junction cancers: an emerging paradigm. *Nat Rev Clin Oncol* 2021; **18**: 473-487 [PMID: 33790428 DOI: 10.1038/s41571-021-00492-2]
- 22 **Wang FH**, Zhang XT, Li YF, Tang L, Qu XJ, Ying JE, Zhang J, Sun LY, Lin RB, Qiu H, Wang C, Qiu MZ, Cai MY, Wu Q, Liu H, Guan WL, Zhou AP, Zhang YJ, Liu TS, Bi F, Yuan XL, Rao SX, Xin Y, Sheng WQ, Xu HM, Li GX, Ji JF, Zhou ZW, Liang H, Zhang YQ, Jin J, Shen L, Li J, Xu RH. The Chinese Society of Clinical Oncology (CSCO): Clinical guidelines for the diagnosis and treatment of gastric cancer, 2021. *Cancer Commun (Lond)* 2021; **41**: 747-795 [PMID: 34197702 DOI: 10.1002/cac2.12193]
- 23 **Shitara K**, Bang YJ, Iwasa S, Sugimoto N, Ryu MH, Sakai D, Chung HC, Kawakami H, Yabusaki H, Lee J, Saito K, Kawaguchi Y, Kamio T, Kojima A, Sugihara M, Yamaguchi K; DESTINY-Gastric01 Investigators. Trastuzumab Deruxtecan in Previously Treated HER2-Positive Gastric Cancer. *N Engl J Med* 2020; **382**: 2419-2430 [PMID: 32469182 DOI: 10.1056/NEJMoa2004413]
- 24 **Patel TH**, Cecchini M. Targeted Therapies in Advanced Gastric Cancer. *Curr Treat Options Oncol* 2020; **21**: 70 [PMID: 32725377 DOI: 10.1007/s11864-020-00774-4]
- 25 **He Q**, Gao J, Ge S, Wang T, Li Y, Peng Z, Shen L. Axitinib alone or in combination with chemotherapeutic drugs exerts potent antitumor activity against human gastric cancer cells *in vitro* and *in vivo*. *J Cancer Res Clin Oncol* 2014; **140**: 1575-1583 [PMID: 24804814 DOI: 10.1007/s00432-014-1693-4]
- 26 **Chen D**, Chen G, Jiang W, Fu M, Liu W, Sui J, Xu S, Liu Z, Zheng X, Chi L, Lin D, Li K, Chen W, Zuo N, Lu J, Chen J, Li G, Zhuo S, Yan J. Association of the Collagen Signature in the Tumor Microenvironment With Lymph Node Metastasis in Early Gastric Cancer. *JAMA Surg* 2019; **154**: e185249 [PMID: 30698615 DOI: 10.1001/jamasurg.2018.5249]
- 27 **Jia H**, Yu F, Li B, Gao Z. Actin-binding protein Anillin promotes the progression of gastric cancer *in vitro* and in mice. *J Clin Lab Anal* 2021; **35**: e23635 [PMID: 33089886 DOI: 10.1002/jcla.23635]
- 28 **Yan H**, Zhang JL, Leung KT, Lo KW, Yu J, To KF, Kang W. An Update of G-Protein-Coupled Receptor Signaling and Its Deregulation in Gastric Carcinogenesis. *Cancers (Basel)* 2023; **15** [PMID: 36765694 DOI: 10.3390/cancers15030736]
- 29 **Chen J**, Zhang M, Ma Z, Yuan D, Zhu J, Tuo B, Li T, Liu X. Alteration and dysfunction of ion channels/transporters in a hypoxic microenvironment results in the development and progression of gastric cancer. *Cell Oncol (Dordr)* 2021; **44**: 739-749 [PMID: 33856653 DOI: 10.1007/s13402-021-00604-1]
- 30 **Li G**, Xie M, Huang Z, Li H, Li P, Zhang Z, Ding Y, Jia Z, Yang J. Overexpression of antisense long noncoding RNA ZNF710AS1202 promotes cell proliferation and inhibits apoptosis of clear cell renal cell carcinoma *via* regulation of ZNF710 expression. *Mol Med Rep* 2020; **21**: 2502-2512 [PMID: 32236626 DOI: 10.3892/mmr.2020.11032]
- 31 **Wu D**. Isocitrate dehydrogenase 2 inhibits gastric cancer cell invasion *via* matrix metalloproteinase 7. *Tumour Biol* 2016; **37**: 5225-5230 [PMID: 26553362 DOI: 10.1007/s13277-015-4358-2]
- 32 **Wang Z**, Yao L, Li Y, Hao B, Wang M, Wang J, Gu W, Zhan H, Liu G, Wu Q. miR3373p inhibits gastric tumor metastasis by targeting ARHGAP10. *Mol Med Rep* 2020; **21**: 705-719 [PMID: 31789419 DOI: 10.3892/mmr.2019.10856]
- 33 **Jian H**, Zhao Y, Liu B, Lu S. SEMA4B inhibits growth of non-small cell lung cancer *in vitro* and *in vivo*. *Cell Signal* 2015; **27**: 1208-1213 [PMID: 25746385 DOI: 10.1016/j.cellsig.2015.02.027]
- 34 **Li L**, Zhao J, Huang S, Wang Y, Zhu L, Cao Y, Xiong J, Deng J. MiR-93-5p promotes gastric cancer-cell progression *via* inactivation of the Hippo signaling pathway. *Gene* 2018; **641**: 240-247 [PMID: 29045821 DOI: 10.1016/j.gene.2017.09.071]
- 35 **UniProt Consortium**. UniProt: a worldwide hub of protein knowledge. *Nucleic Acids Res* 2019; **47**: D506-D515 [PMID: 30395287 DOI: 10.1093/nar/gky1049]
- 36 **Long P**, Zhang L, Huang B, Chen Q, Liu H. Integrating genome sequence and structural data for statistical learning to predict transcription factor binding sites. *Nucleic Acids Res* 2020; **48**: 12604-12617 [PMID: 33264415 DOI: 10.1093/nar/gkaa1134]

- 37 **Hamidi AA**, Taghehchian N, Basirat Z, Zangouei AS, Moghbeli M. MicroRNAs as the critical regulators of cell migration and invasion in thyroid cancer. *Biomark Res* 2022; **10**: 40 [PMID: 35659780 DOI: 10.1186/s40364-022-00382-4]
- 38 **Guo Z**, Zhou K, Wang Q, Huang Y, Ji J, Peng Y, Zhang X, Zheng T, Zhang Z, Chong D, Yang Z. The transcription factor RUNX2 fuels YAP1 signaling and gastric cancer tumorigenesis. *Cancer Sci* 2021; **112**: 3533-3544 [PMID: 34160112 DOI: 10.1111/cas.15045]
- 39 **Zhao X**, Chen L, Wu J, You J, Hong Q, Ye F. Transcription factor KLF15 inhibits the proliferation and migration of gastric cancer cells *via* regulating the TFAP2A-AS1/NISCH axis. *Biol Direct* 2021; **16**: 21 [PMID: 34727954 DOI: 10.1186/s13062-021-00300-y]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA  
**Telephone:** +1-925-3991568  
**E-mail:** [office@baishideng.com](mailto:office@baishideng.com)  
**Help Desk:** <https://www.f6publishing.com/helpdesk>  
<https://www.wjgnet.com>

