

# World Journal of *Gastrointestinal Oncology*

*World J Gastrointest Oncol* 2024 April 15; 16(4): 1091-1675



### EDITORIAL

- 1091** Parallel pathways: A chronicle of evolution in rectal and breast cancer surgery  
*Pesce A, Fabbri N, Iovino D, Feo CV*
- 1097** Hepatitis B virus genotypes in precision medicine of hepatitis B-related hepatocellular carcinoma: Where we are now  
*Sukowati CHC, Jayanti S, Turyadi T, Muljono DH, Tiribelli C*

### REVIEW

- 1104** Novel milestones for early esophageal carcinoma: From bench to bed  
*Qi JH, Huang SL, Jin SZ*
- 1119** Colorectal cancer screening: A review of current knowledge and progress in research  
*Lopes SR, Martins C, Santos IC, Teixeira M, Gamito É, Alves AL*
- 1134** New avenues for the treatment of immunotherapy-resistant pancreatic cancer  
*Silva LGO, Lemos FFB, Luz MS, Rocha Pinheiro SL, Calmon MDS, Correa Santos GL, Rocha GR, de Melo FF*

### MINIREVIEWS

- 1154** Present situation of minimally invasive surgical treatment for early gastric cancer  
*Li CY, Wang YF, Luo LK, Yang XJ*
- 1166** Mixed neuroendocrine non-neuroendocrine neoplasms in gastroenteropancreatic tract  
*Díaz-López S, Jiménez-Castro J, Robles-Barraza CE, Ayala-de Miguel C, Chaves-Conde M*
- 1180** Esophageal cancer screening, early detection and treatment: Current insights and future directions  
*Qu HT, Li Q, Hao L, Ni YJ, Luan WY, Yang Z, Chen XD, Zhang TT, Miao YD, Zhang F*

### ORIGINAL ARTICLE

#### Retrospective Cohort Study

- 1192** Pre-operative enhanced magnetic resonance imaging combined with clinical features predict early recurrence of hepatocellular carcinoma after radical resection  
*Chen JP, Yang RH, Zhang TH, Liao LA, Guan YT, Dai HY*
- 1204** Clinical analysis of multiple primary gastrointestinal malignant tumors: A 10-year case review of a single-center  
*Zhu CL, Peng LZ*

### Retrospective Study

- 1213** Predictive model for non-malignant portal vein thrombosis associated with cirrhosis based on inflammatory biomarkers  
*Nie GL, Yan J, Li Y, Zhang HL, Xie DN, Zhu XW, Li X*
- 1227** Predictive modeling for postoperative delirium in elderly patients with abdominal malignancies using synthetic minority oversampling technique  
*Hu WJ, Bai G, Wang Y, Hong DM, Jiang JH, Li JX, Hua Y, Wang XY, Chen Y*
- 1236** Efficacy and predictive factors of transarterial chemoembolization combined with lenvatinib plus programmed cell death protein-1 inhibition for unresectable hepatocellular carcinoma  
*Ma KP, Fu JX, Duan F, Wang MQ*
- 1248** Should we perform sigmoidoscopy for colorectal cancer screening in people under 45 years?  
*Leong W, Guo JQ, Ning C, Luo FF, Jiao R, Yang DY*
- 1256** Computed tomography-based radiomics diagnostic approach for differential diagnosis between early- and late-stage pancreatic ductal adenocarcinoma  
*Ren S, Qian LC, Cao YY, Daniels MJ, Song LN, Tian Y, Wang ZQ*
- 1268** Prognostic analysis of related factors of adverse reactions to immunotherapy in advanced gastric cancer and establishment of a nomogram model  
*He XX, Du B, Wu T, Shen H*

### Clinical Trials Study

- 1281** Safety and efficacy of a programmed cell death 1 inhibitor combined with oxaliplatin plus S-1 in patients with Borrmann large type III and IV gastric cancers  
*Bao ZH, Hu C, Zhang YQ, Yu PC, Wang Y, Xu ZY, Fu HY, Cheng XD*

### Observational Study

- 1296** Computed tomography radiogenomics: A potential tool for prediction of molecular subtypes in gastric stromal tumor  
*Yin XN, Wang ZH, Zou L, Yang CW, Shen CY, Liu BK, Yin Y, Liu XJ, Zhang B*
- 1309** Application of texture signatures based on multiparameter-magnetic resonance imaging for predicting microvascular invasion in hepatocellular carcinoma: Retrospective study  
*Nong HY, Cen YY, Qin M, Qin WQ, Xie YX, Li L, Liu MR, Ding K*
- 1319** Causal roles of gut microbiota in cholangiocarcinoma etiology suggested by genetic study  
*Chen ZT, Ding CC, Chen KL, Gu YJ, Lu CC, Li QY*
- 1334** Is recovery enhancement after gastric cancer surgery really a safe approach for elderly patients?  
*Li ZW, Luo XJ, Liu F, Liu XR, Shu XP, Tong Y, Lv Q, Liu XY, Zhang W, Peng D*
- 1344** Establishment of a cholangiocarcinoma risk evaluation model based on mucin expression levels  
*Yang CY, Guo LM, Li Y, Wang GX, Tang XW, Zhang QL, Zhang LF, Luo JY*

- 1361** Effectiveness of fecal DNA syndecan-2 methylation testing for detection of colorectal cancer in a high-risk Chinese population

*Luo WF, Jiao YT, Lin XL, Zhao Y, Wang SB, Shen J, Deng J, Ye YF, Han ZP, Xie FM, He JH, Wan Y*

#### Clinical and Translational Research

- 1374** Clinical and socioeconomic determinants of survival in biliary tract adenocarcinomas

*Sahyoun L, Chen K, Tsay C, Chen G, Protiva P*

- 1384** Risk factors, prognostic factors, and nomograms for distant metastasis in patients with diagnosed duodenal cancer: A population-based study

*Shang JR, Xu CY, Zhai XX, Xu Z, Qian J*

- 1421** NOX4 promotes tumor progression through the MAPK-MEK1/2-ERK1/2 axis in colorectal cancer

*Xu YJ, Huo YC, Zhao QT, Liu JY, Tian YJ, Yang LL, Zhang Y*

#### Basic Study

- 1437** Curcumin inhibits the growth and invasion of gastric cancer by regulating long noncoding RNA AC022424.2

*Wang BS, Zhang CL, Cui X, Li Q, Yang L, He ZY, Yang Z, Zeng MM, Cao N*

- 1453** MicroRNA-298 determines the radio-resistance of colorectal cancer cells by directly targeting human dual-specificity tyrosine(Y)-regulated kinase 1A

*Shen MZ, Zhang Y, Wu F, Shen MZ, Liang JL, Zhang XL, Liu XJ, Li XS, Wang RS*

- 1465** Human  $\beta$ -defensin-1 affects the mammalian target of rapamycin pathway and autophagy in colon cancer cells through long non-coding RNA TCONS\_00014506

*Zhao YX, Cui Y, Li XH, Yang WH, An SX, Cui JX, Zhang MY, Lu JK, Zhang X, Wang XM, Bao LL, Zhao PW*

- 1479** FAM53B promotes pancreatic ductal adenocarcinoma metastasis by regulating macrophage M2 polarization

*Pei XZ, Cai M, Jiang DW, Chen SH, Wang QQ, Lu HM, Lu YF*

- 1500** Transcriptome sequencing reveals novel biomarkers and immune cell infiltration in esophageal tumorigenesis

*Sun JR, Chen DM, Huang R, Wang RT, Jia LQ*

- 1514** Construction of CDKN2A-related competitive endogenous RNA network and identification of GAS5 as a prognostic indicator for hepatocellular carcinoma

*Pan Y, Zhang YR, Wang LY, Wu LN, Ma YQ, Fang Z, Li SB*

- 1532** Two missense STK11 gene variations impaired LKB1/adenosine monophosphate-activated protein kinase signaling in Peutz-Jeghers syndrome

*Liu J, Zeng SC, Wang A, Cheng HY, Zhang QJ, Lu GX*

- 1547** Long noncoding RNAs HAND2-AS1 ultrasound microbubbles suppress hepatocellular carcinoma progression by regulating the miR-873-5p/tissue inhibitor of matrix metalloproteinase-2 axis

*Zou Q, Wang HW, Di XL, Li Y, Gao H*

- 1564** Upregulated lncRNA PRNT promotes progression and oxaliplatin resistance of colorectal cancer cells by regulating HIPK2 transcription

*Li SN, Yang S, Wang HQ, Hui TL, Cheng M, Zhang X, Li BK, Wang GY*

### SYSTEMATIC REVIEWS

- 1578** Prognosis value of heat-shock proteins in esophageal and esophagogastric cancer: A systematic review and meta-analysis

*Nakamura ET, Park A, Pereira MA, Kikawa D, Tustumi F*

- 1596** Risk factors for hepatocellular carcinoma associated with hepatitis C genotype 3 infection: A systematic review

*Farooq HZ, James M, Abbott J, Oyibo P, Divall P, Choudhry N, Foster GR*

### META-ANALYSIS

- 1613** Effectiveness and tolerability of programmed cell death protein-1 inhibitor + chemotherapy compared to chemotherapy for upper gastrointestinal tract cancers

*Zhang XM, Yang T, Xu YY, Li BZ, Shen W, Hu WQ, Yan CW, Zong L*

- 1626** Success rate of current human-derived gastric cancer organoids establishment and influencing factors: A systematic review and meta-analysis

*Jiang KL, Wang XX, Liu XJ, Guo LK, Chen YQ, Jia QL, Yang KM, Ling JH*

### CASE REPORT

- 1647** Pathologically successful conversion hepatectomy for advanced giant hepatocellular carcinoma after multidisciplinary therapy: A case report and review of literature

*Chu JH, Huang LY, Wang YR, Li J, Han SL, Xi H, Gao WX, Cui YY, Qian MP*

- 1660** Clinical pathological characteristics of "crawling-type" gastric adenocarcinoma cancer: A case report

*Xu YW, Song Y, Tian J, Zhang BC, Yang YS, Wang J*

- 1668** Primary pancreatic peripheral T-cell lymphoma: A case report

*Bai YL, Wang LJ, Luo H, Cui YB, Xu JH, Nan HJ, Yang PY, Niu JW, Shi MY*

**ABOUT COVER**

Peer Reviewer of *World Journal of Gastrointestinal Oncology*, Lie Zheng, Director, Professor, Department of Gastroenterology, Shaanxi Provincial Hospital of Traditional Chinese Medicine, Xi'an 730000, Shaanxi Province, China. xinliwen696@126.com

**AIMS AND SCOPE**

The primary aim of *World Journal of Gastrointestinal Oncology* (WJGO, *World J Gastrointest Oncol*) is to provide scholars and readers from various fields of gastrointestinal oncology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGO mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal oncology and covering a wide range of topics including liver cell adenoma, gastric neoplasms, appendiceal neoplasms, biliary tract neoplasms, hepatocellular carcinoma, pancreatic carcinoma, cecal neoplasms, colonic neoplasms, colorectal neoplasms, duodenal neoplasms, esophageal neoplasms, gallbladder neoplasms, *etc.*

**INDEXING/ABSTRACTING**

The WJGO is now abstracted and indexed in PubMed, PubMed Central, Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJGO as 3.0; IF without journal self cites: 2.9; 5-year IF: 3.0; Journal Citation Indicator: 0.49; Ranking: 157 among 241 journals in oncology; Quartile category: Q3; Ranking: 58 among 93 journals in gastroenterology and hepatology; and Quartile category: Q3. The WJGO's CiteScore for 2022 is 4.1 and Scopus CiteScore rank 2022: Gastroenterology is 71/149; Oncology is 197/366.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: Xiang-Di Zhang; Production Department Director: Xiang Li; Cover Editor: Jia-Ru Fan.

**NAME OF JOURNAL**

*World Journal of Gastrointestinal Oncology*

**ISSN**

ISSN 1948-5204 (online)

**LAUNCH DATE**

February 15, 2009

**FREQUENCY**

Monthly

**EDITORS-IN-CHIEF**

Monjur Ahmed, Florin Burada

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/1948-5204/editorialboard.htm>

**PUBLICATION DATE**

April 15, 2024

**COPYRIGHT**

© 2024 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>



Observational Study

# Causal roles of gut microbiota in cholangiocarcinoma etiology suggested by genetic study

Zhi-Tao Chen, Chen-Chen Ding, Kai-Lei Chen, Yang-Jun Gu, Chi-Cheng Lu, Qi-Yong Li

**Specialty type:** Oncology

**Provenance and peer review:**

Invited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): 0

Grade B (Very good): B

Grade C (Good): 0

Grade D (Fair): 0

Grade E (Poor): 0

**P-Reviewer:** Lee H, South Korea

**Received:** October 23, 2023

**Peer-review started:** October 23, 2023

**First decision:** December 12, 2023

**Revised:** December 20, 2023

**Accepted:** January 15, 2024

**Article in press:** January 15, 2024

**Published online:** April 15, 2024



**Zhi-Tao Chen, Yang-Jun Gu, Qi-Yong Li**, Division of Hepatobiliary and Pancreatic Surgery, Shulan (Hangzhou) Hospital Affiliated to Zhejiang Shuren University Shulan International Medical College, Hangzhou 310000, Zhejiang Province, China

**Chen-Chen Ding**, Pediatric Psychology, The Affiliated Mental Health Centre & Hangzhou Seventh People's Hospital, Zhejiang University School of Medicine, Hangzhou 310000, Zhejiang Province, China

**Kai-Lei Chen**, School of Medicine, Zhejiang Shuren University, Hangzhou 310000, Zhejiang Province, China

**Chi-Cheng Lu**, School of Medicine, Zhejiang Chinese Medical University Zhejiang Shuren College, Hangzhou 310000, Zhejiang Province, China

**Corresponding author:** Qi-Yong Li, PhD, Deputy Director, Researcher, Surgeon, Division of Hepatobiliary and Pancreatic Surgery, Shulan (Hangzhou) Hospital Affiliated to Zhejiang Shuren University Shulan International Medical College, No. 848 Dongxin Road, Hangzhou 310000, Zhejiang Province, China. [liqiyong@zju.edu.cn](mailto:liqiyong@zju.edu.cn)

## Abstract

### BACKGROUND

Cholangiocarcinoma (CCA) is a highly malignant biliary tract cancer with poor prognosis. Previous studies have implicated the gut microbiota in CCA, but evidence for causal mechanisms is lacking.

### AIM

To investigate the causal relationship between gut microbiota and CCA risk.

### METHODS

We performed a two-sample mendelian randomization study to evaluate potential causal associations between gut microbiota and CCA risk using genome-wide association study summary statistics for 196 gut microbial taxa and CCA. Genetic variants were used as instrumental variables. Multiple sensitivity analyses assessed result robustness.

### RESULTS

Fifteen gut microbial taxa showed significant causal associations with CCA risk. Higher genetically predicted abundance of genus *Eubacteriumnodatum* group, genus

*Ruminococcustorques* group, genus *Coproccoccus*, genus *Dorea*, and phylum *Actinobacteria* were associated with reduced risk of gallbladder cancer and extrahepatic CCA. Increased intrahepatic CCA risk was associated with higher abundance of family *Veillonellaceae*, genus *Alistipes*, order *Enterobacteriales*, and phylum *Firmicutes*. Protective effects against CCA were suggested for genus *Collinsella*, genus *Eisenbergiella*, genus *Anaerostipes*, genus *Paraprevotella*, genus *Parasutterella*, and phylum *Verrucomicrobia*. Sensitivity analyses indicated these findings were reliable without pleiotropy.

## CONCLUSION

This pioneering study provides novel evidence that specific gut microbiota may play causal roles in CCA risk. Further experimental validation of these candidate microbes is warranted to consolidate causality and mechanisms.

**Key Words:** Cholangiocarcinoma; Mendelian randomization; Gut microbiota; Instrumental variables; Sensitivity analyses

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** Cholangiocarcinoma (CCA) is a highly malignant biliary tract cancer with poor prognosis. Emerging evidence suggests the gut microbiota may play a causal role in CCA pathogenesis, but robust genetic evidence is still lacking. Using genome-wide association study summary statistics, our study provides novel evidence that 15 gut microbial taxa may confer either protective or detrimental causal effects on CCA risk.

**Citation:** Chen ZT, Ding CC, Chen KL, Gu YJ, Lu CC, Li QY. Causal roles of gut microbiota in cholangiocarcinoma etiology suggested by genetic study. *World J Gastrointest Oncol* 2024; 16(4): 1319-1333

**URL:** <https://www.wjgnet.com/1948-5204/full/v16/i4/1319.htm>

**DOI:** <https://dx.doi.org/10.4251/wjgo.v16.i4.1319>

## INTRODUCTION

Cholangiocarcinoma (CCA) originates from the biliary epithelium and is among the most prevalent malignancies due to its significant malignancy potential[1,2]. Based on anatomical site of origin, CCA manifests as three distinct subtypes: Intrahepatic CCA (iCCA), extrahepatic CCA (eCCA), and gallbladder cancer (GC)[1]. Established risk factors for CCA include fluke infections, inflammatory bowel disease, intrahepatic bile duct stones, choledochal cysts, and primary sclerosing cholangitis (PSC)[3,4]. Despite recent progress in diagnosis and therapy, CCA prognosis remains poor with 5-year survival below 5% for advanced disease[5-7]. Further elucidation of CCA pathogenesis at the molecular, epigenetic and genomic levels is therefore critical to enable novel treatment approaches.

In recent years, the gut microbiota has emerged as a key factor governing health[8,9]. Microbiota dysbiosis can impact immune function, metabolism and physiology, contributing to diseases like obesity, diabetes, non-alcoholic fatty liver disease and cancer[10-12]. Anatomically and physiologically, the hepatobiliary duct and gastrointestinal tract comprise a “gut-liver axis” that regulates liver pathology and intrahepatic/systemic immunity[13]. The microbiota likely contributes to diverse hepatobiliary conditions including cancer, PSC, choledocholithiasis and cholelithiasis[14-17]. Previous research has revealed that the gut microbiota plays a pivotal role in the diagnosis and treatment of CCA[18,19]. In-depth investigations into the role of the gut microbiota in CCA have significantly improved the prognostic outlook for individuals affected by this disease. However, the causal relationship between the gut microbiota and CCA remains unclear. Elucidating such mechanisms would enable microbiome modulation as an early preventative approach aligning with precancer interception paradigms.


Establishing causality is challenged by limited clinical trial follow-up and potential confounding in observational studies. Mendelian randomization (MR) helps address this by using genetic variants as instrumental variables (IVs)[20]. The present study represents a pioneering effort in employing a two-sample MR approach to discern a potential causal link between particular gut microbiota taxa and CCA, thereby offering valuable insights for subsequent mechanistic inquiries.

## MATERIALS AND METHODS

### Ethical statement and study design

This research adheres to the STROBE-MR Guidelines, and all data employed in this study are openly available and appropriately cited[21]. Consequently, our study did not require additional ethics committee approval.

Figure 1 presents the directed acyclic graph guiding the design of the current MR study. In this framework, the gut microbiota represents the exposure variables, while CCA constitutes the outcome variable. Genetic variants associated with gut microbiota taxa were leveraged as IVs to evaluate potential causal associations of gut microbiota composition



**Figure 1** The process of present mendelian randomization analyses is shown in flow chart. A: Principle diagram of mendelian randomization study; B: Diagrammatic illustration of the complete mendelian randomization analysis process. Other risk factors influencing the occurrence of cholangiocarcinoma include exposure to fluke infections, inflammatory bowel disease, intrahepatic bile duct stones, choledochal cysts, and primary sclerosing cholangitis. Assumption 1: The instrumental variables (IVs) selected for this study should demonstrate a significant association with gut microbiota; Assumption 2: The IVs chosen for present study are required to have no significant associations with other potential confounding factors; Assumption 3: The IVs utilized in present study do not have any independent causal pathways leading to the outcome (CCA) other than through gut microbiota. MR: Mendelian randomization; IV: Instrumental variable; CCA: Cholangiocarcinoma; GC: Gallbladder cancer; eCCA: Extrahepatic cholangiocarcinoma; iCCA: Intrahepatic cholangiocarcinoma; LD: Linkage disequilibrium; SNPs: Single nucleotide polymorphisms.

with CCA risk, thereby minimizing issues of confounding.

### Date sources


A large-scale genome-wide association study (GWAS) encompassed 18340 participants drawn from 24 diverse cohorts spanning multiple countries, examining 122110 Loci of genetic variation. This study provided summary statistics for gut microbiota based on 16S rRNA gene sequencing data obtained from the MiBioGen (<https://mibiogen.gcc.rug.nl/>) database[22]. Among the participants, a significant majority, 13266 individuals or 72.3%, were of European ancestry. The study encompassed a broad spectrum of 211 traits, which included members from 131 *genera*, 35 *families*, 20 *orders*, 16 *classes*, and 9 *phyla*. In the current MR study, 15 unidentified taxa were notably excluded, resulting in the analysis incorporating 196 taxonomic units, spanning 9 *phyla*, 16 *classes*, 20 *orders*, 32 *families*, and 119 *genera*. Jiang *et al*[23] conducted analyses of summary statistics for malignant neoplasms of the gallbladder and extrahepatic bile ducts (195 European ancestry cases, 456153 European ancestry controls) as well as intrahepatic CCA (104 European ancestry cases, 456244 European ancestry controls), which were provided by the GWAS Catalog (<https://www.ebi.ac.uk/gwas/>).

### The choice of IVs

MR analyses utilized IVs, primarily single nucleotide polymorphisms (SNPs), as mediators to explore causality between exposures and outcomes. The foundational assumption in MR necessitates that all SNPs robustly and independently predict the exposure variable at the genome-wide significance level. In present research, we utilized robust SNPs associated with gut microbiota as IVs for the exposure variable. However, applying a stringent threshold of  $5 \times 10^{-8}$  would have excluded the majority of these SNPs. Consequently, we opted for a relatively lenient yet still statistically significant threshold of  $1 \times 10^{-5}$ , as supported by prior studies[24,25]. This threshold was set to encompass most gut microbiota-associated SNPs, ensuring that those with an  $R^2 < 0.001$  and a physical distance (kb) of 10000 were included, thus mitigating linkage disequilibrium (LD). The  $F$  statistic was employed to assess the strength of the correlation between IVs and exposures, with an  $F$  statistic exceeding 10 typically indicating a substantial correlation. These screening criteria serve to establish the reliability of the findings in present MR study.

### Statistical analysis

The analyses were conducted in RStudio (Version: 2023.06.1 + 524) using the TwoSampleMR package (version 0.5.7) and MRPRESSO package. In the context of a global-level test, a two-sided  $P$  value of 0.05 was considered statistically significant. In present study, we utilized a comprehensive approach, incorporating MR-Egger, weighted median, inverse



**Figure 2** The forest plot illustrates the relationships between gut microbiota and cholangiocarcinoma. A: An increased abundance of gut microbial taxa was observed to be linked with a reduced risk of cholangiocarcinoma (CCA); B: An increased abundance of gut microbial taxa was observed to be linked with an elevated risk of CCA. OR: Odds ratio.

variance weighted (IVW), simple mode, and weighted mode methodologies for MR analysis, enabling a thorough assessment of the causal relationship between gut microbiota and CCA. In cases where pleiotropy among IVs is absent, IVW is selected as the primary analytical method due to its superior statistical power[26]. To evaluate the reliability of our findings, we conducted a set of sensitivity analyses, including Cochran's *Q* test, MR-Egger intercept test, and MR-PRESSO global test. Both the Cochran's *Q* test and MR-Egger intercept test were employed to assess the presence of SNP-associated heterogeneity and horizontal pleiotropy for each gut microbiota trait. The outcomes revealed *P* values exceeding 0.05, indicating the absence of heterogeneity and horizontal pleiotropy. Outliers were identified through the application of MR-PRESSO analysis. Additionally, we also employed funnel plots and conducted leave-one-out sensitivity tests to assess heterogeneity. The leave-one-out analysis was utilized to identify potential pleiotropic effects originating from individual SNPs. Scatter plots, forest plots, funnel plots, and leave-one-out sensitivity tests serve as valuable tools for visualizing MR results in a comprehensive manner.

## RESULTS

### Selection of IVs







Initially, we identified 122100 SNPs associated with gut microbiota traits through the MiBioGen Consortium dataset. Following a rigorous sequence of quality control procedures based on locus-wide statistical significance ( $P < 1 \times 10^{-5}$ ) and the LD threshold ( $R^2 < 0.001$ , with a clumping distance of 10000 kb), 2236 SNPs associated with 196 gut microbiota traits were selected as IVs. Notably, all IVs exhibited *F*-statistics exceeding 10, thereby indicating the absence of evidence for weak instrument bias (Supplementary Table 1). Based on these SNPs, we have extracted corresponding pieces of information from the outcome variable dataset (Supplementary Tables 2 and 3).


### Causal inference of the relationship of the gut microbiota with GC and eCCA risk

According to the results of the IVW method, the higher genetically predicted abundance of genus *Eubacteriumnodatum* group [odds ratio (OR) = 0.489, standard error (SE) = 0.267, 95% confidence interval (CI): 0.290–0.826,  $P = 0.007$ ], genus *Ruminococcustorques* group (OR = 0.291, SE = 0.535, 95% CI: 0.102–0.830,  $P = 0.021$ ), genus *Coprococcus* (OR = 0.296, SE = 0.593, 95% CI: 0.093–0.946,  $P = 0.040$ ), genus *Dorea* (OR = 0.250, SE = 0.622, 95% CI: 0.074–0.847,  $P = 0.026$ ), phylum *Actinobacteria* (OR = 0.348, SE = 0.483, 95% CI: 0.135–0.898,  $P = 0.029$ ) were associated with a reduced risk of GC and eCCA (Figure 2A). In contrast, genetically predicted abundance of genus *Collinsella* (OR = 6.977, SE = 0.621, 95% CI: 2.006–23.560,  $P = 0.002$ ), genus *Eisenbergiella* (OR = 2.018, SE = 0.342, 95% CI: 1.033–3.941,  $P = 0.040$ ) was positively related to GC and eCCA risk (Figure 2B). The weighted median, simple mode, and weighted mode exhibited the same directional impact as IVW, although the *P* values were not consistently statistically significant (Supplementary Table 4, Figure 3).

### Causal inference of the relationship of the gut microbiota with iCCA risk

Employing the IVW method, our study found suggestive evidence of a potential causal link between genetically predicted increases in the family *Veillonellaceae* ( $P = 0.014$ , 95% CI: 1.292–9.929, OR = 3.582, SE = 0.520), order *Enterobacteriales*/family *Enterobacteriaceae* ( $P = 0.032$ , 95% CI: 1.156–27.429, OR = 5.632, SE = 0.808), genus *Alistipes* ( $P = 0.020$ , 95% CI: 1.316–24.245, OR = 5.648, SE = 0.743), and phylum *Firmicutes* ( $P = 0.046$ , 95% CI: 1.025–12.258, OR = 3.545, SE = 0.633) with an increased risk of iCCA (Figure 2B). From the earlier mentioned traits, it was noted that both the order *Enterobacteriales* and the family *Enterobacteriaceae* fall under the same bacterial category and share identical IVs. Furthermore, our findings





**Figure 3** Scatter plots of causal estimates of specific gut microbiota taxa on gallbladder cancer and extrahepatic cholangiocarcinoma.

The slope of each line corresponding to the estimated mendelian randomization effect in different models, including the conventional inverse variance weighted, weighted median, simple mode, and weighted mode. A: *Genus Eubacteriumnodatum* group; B: *Genus Ruminococcustorques* group; C: *Genus Collinsella*; D: *Genus Coprococcus*; E: *Genus Dorea*; F: *Genus Eisenbergiella*; G: *Phylum Actinobacteria*. MR: Mendelian randomization; SNP: Single nucleotide polymorphisms.

suggest that genetically predicted increases in the *genus Anaerostipes* ( $P = 0.006$ , 95%CI: 0.033-0.564, OR = 0.135, SE = 0.728), the *genus Parasutterella* ( $P = 0.032$ , 95%CI: 0.115-0.907, OR = 0.323, SE = 0.527), the *genus Paraprevotella* ( $P = 0.005$ , 95%CI: 0.107-0.672, OR = 0.268, SE = 0.470), and the *phylum Verrucomicrobia* ( $P = 0.005$ , 95%CI: 0.048-0.588, OR = 0.168, SE = 0.640) are associated with protective effects against iCCA (Figure 3A). Additionally, the causal effect estimates derived from the weighted median, simple mode, and weighted mode methods showed similar magnitudes and directions as those obtained with the previously mentioned IVW method (Supplementary Table 4, Figure 4).

### Sensitivity analysis

Subsequently, a comprehensive sensitivity analysis was carried out to assess the stability and reliability of the inferred causal relationship between gut microbiota and CCA. The detailed description is summarized in Table 1. Cochran's  $Q$  statistics revealed the absence of significant heterogeneity among the selected IVs, with  $P$  values exceeding 0.05 in both IVW and MR-Egger methods. Furthermore, our analysis, employing the MR-Egger intercept method, did not reveal any indications of a horizontal pleiotropic effect ( $P > 0.05$ ). The outcomes from the MR-PRESSO trial indicated the absence of any horizontal pleiotropic outliers. The leave-one-out analysis results demonstrated that none of the SNPs were influential outliers (Figures 5 and 6). Additionally, the use of funnel and forest plots to depict a symmetrical pattern serves to visually affirm the reliability of the study's results (Supplementary Figures 1-4).

## DISCUSSION

CCA is a highly diverse form of cancer, with its global incidence steadily on the rise[27]. With surgical resection being the exclusive curative treatment modality, the prognosis for individuals afflicted with CCA remains bleak[28]. In recent years, the prevention of tumor initiation and the inhibition of tumor progression have emerged as pivotal milestones in cancer management within the field of oncology. Changes in the composition of gut microbiota are closely associated with the initiation and progression of cancer[8,29]. This study represents the inaugural attempt to evaluate the causal link between gut microbiota and CCA while also endeavoring to identify particular causative microbial taxa through two-sample MR analyses based on GWAS summary statistics.

The gut microbiota constitutes a complex and dynamically evolving assembly of ecological microbial communities that reside within the human gastrointestinal tract, often referred to as a "neglected organ"[30-32]. These microorganisms assume a pivotal role in maintaining the homeostasis of the digestive system, exerting multifaceted metabolic, immunological, and protective functions that contribute to the overall health of the host[32]. Although gut microbiota plays a crucial role in facilitating various essential and advantageous physiological processes, such as the digestion of macronutrients and the synthesis of certain vitamins, a wealth of empirical data underscores their potential involvement in the emergence of detrimental phenotypes[33,34]. Notably, discernible alterations in both the structure and function of the microbial community have been linked to numerous disease states, including cancer[31]. Due to the bidirectional communication between the gastrointestinal tract and the biliary system, the liver excretes bile acids and other biologically active components *via* the bile duct to interface with the intestine. Simultaneously, the gut microbiota and its metabolites are transported to the liver through the bile duct. Therefore, the gut microbiota plays a pivotal role in the


**Table 1 Heterogeneity and pleiotropy analysis of the mendelian randomization study on gut microbiota and cholangiocarcinoma**


Exposure	Outcome	Method	Heterogeneity		Horizontal pleiotropy		MR-PRESSO
			Q	Q, P value	Egger intercept	P value	P value
<i>Verrucomicrobia</i>	iCCA	MRE	4.795	0.904	-0.048	0.762	0.951
		IVW	4.892	0.936			
<i>Firmicutes</i>	iCCA	MRE	16.151	0.513	-0.117	0.342	0.527
		IVW	17.108	0.516			
<i>Enterobacteriales/Enterobacteriaceae</i>	iCCA	MRE	5.600	0.692	0.231	0.433	0.725
		IVW	6.282	0.711			
<i>Parasutterella</i>	iCCA	MRE	9.901	0.769	-0.062	0.638	0.805
		IVW	10.133	0.811			
<i>Paraprevotella</i>	iCCA	MRE	10.402	0.495	-0.057	0.740	0.619
		IVW	10.518	0.571			
<i>Anaerostipes</i>	iCCA	MRE	9.312	0.593	-0.095	0.526	0.679
		IVW	9.740	0.639			
<i>Alistipes</i>	iCCA	MRE	8.587	0.803	0.203	0.373	0.808
		IVW	9.437	0.802			
<i>Veillonellaceae</i>	iCCA	MRE	13.399	0.818	0.037	0.672	0.850
		IVW	13.584	0.851			
<i>Eubacteriumnodatum group</i>	GC and eCCA	MRE	6.622	0.676	0.106	0.555	0.755
		IVW	6.997	0.726			
<i>Ruminococcustorques group</i>	GC and eCCA	MRE	6.527	0.836	-0.113	0.309	0.812
		IVW	7.665	0.811			
<i>Collinsella</i>	GC and eCCA	MRE	14.609	0.147	0.024	0.894	0.229
		IVW	14.636	0.200			
<i>Coproccoccus</i>	GC and eCCA	MRE	4.421	0.817	-0.311	0.068	0.442
		IVW	8.877	0.449			
<i>Dorea</i>	GC and eCCA	MRE	13.383	0.203	-0.031	0.820	0.304
		IVW	13.456	0.265			
<i>Eisenbergiella</i>	GC and eCCA	MRE	10.563	0.393	-0.052	0.857	0.493
		IVW	10.599	0.477			
<i>Actinobacteria</i>	GC and eCCA	MRE	14.593	0.481	0.182	0.153	0.406
		IVW	16.855	0.395			

MR: Mendelian randomization; CCA: Cholangiocarcinoma; iCCA: Intrahepatic cholangiocarcinoma; GC: Gallbladder cancer; eCCA: Extrahepatic cholangiocarcinoma; Q: Cochran's Q test; MRE: MR egger; IVW: Inverse variance weighted.

pathogenesis and progression CCA[18,35]. Hence, there is a compelling need for additional investigations to elucidate the causal connection between gut microbiota and CCA, thereby establishing a novel theoretical foundation for the prevention and treatment of CCA.

The impact of the gut microbiota in the field of oncology is a double-edged sword, and our research has equally substantiated this perspective from a genetic standpoint. The gut microbiota actively fosters the development of extraintestinal cancers by facilitating bacterial translocation and the generation of bioactive molecules within the biliary tract. Numerous research investigations have demonstrated notable distinctions in gut microbiota composition between individuals with extraintestinal cancers and those without the disease[32,35]. *Bacteroides* and *Ruminococcaceae* have been shown to potentially contribute to the pathogenesis of hepatocellular carcinoma by exacerbating hepatic inflammation, accumulating toxic compounds, and inducing liver steatosis[36]. Zhang *et al*[35] observed a depletion of *Saccharomyces cerevisiae* (*S. cerevisiae*) in iCCA. Significantly, past studies have shown that *S. cerevisiae* has the capacity to impede the





**Figure 4** Scatter plots of causal estimates of specific gut microbiota taxa on intrahepatic cholangiocarcinoma. The slope of each line corresponding to the estimated Mendelian randomization effect in different models, including the conventional inverse variance weighted, weighted median, simple mode, and weighted mode. A: *Family Veillonellaceae*; B: *Genus Alistipes*; C: *Order Enterobacteriales*; D: *Phylum Firmicutes*; E: *Genus Anaerostipes*; F: *Genus Paraprevotella*; G: *Genus Parasutterella*; H: *Phylum Verrucomicrobia*. MR: Mendelian randomization; SNP: Single nucleotide polymorphisms.


growth of colorectal tumors. It achieves this by inducing apoptosis in epithelial cells, modulating intestinal immunity, and altering the composition of the gut microbiota[37]. The microbiota can indirectly influence tumor progression by generating and metabolizing bioactive molecules, which, when carried through systemic circulation, such as bacterial LPS entering the bloodstream, can impact tumor formation in distant tissues from the gastrointestinal tract[38]. *Lactic acid bacteria* and *Bifidobacterium* play a role in the regulation of pH and bile acid processes[39]. Furthermore, their enzymatic capacity to degrade potential carcinogens and their metabolites, including heterocyclic amines, nitrosamines, and aflatoxins, contributes to the inhibition of the development of various cancers, such as gastric and liver cancers[40].


In this MR study, we determined that 15 microbial taxa are causally associated with CCA. Elevated genetic predisposition towards higher abundance of the *genus Eubacteriumnodatum* group, *genus Ruminococcustorques* group, *genus Coprococcus*, *genus Dorea*, *phylum Actinobacteria*, *family Veillonellaceae*, *genus Alistipes*, *order Enterobacteriales/family Enterobacteriaceae*, and *phylum Firmicutes* were found to be associated with a decreased risk of CCA. Conversely, a genetically predicted increase in the abundance of the *genus Collinsella*, *genus Eisenbergiella*, *genus Anaerostipes*, *genus Paraprevotella*, *genus Parasutterella*, and *phylum Verrucomicrobia* exhibited a positive correlation with the risk of CCA. The abundance of the *genus Eubacteriumnodatum* group was found significantly reduced in colorectal cancer patients, and its functionality appeared to be associated with processes related to protein digestion and absorption, as well as the renin-angiotensin system pathway[41]. The *genus Ruminococcustorques* group and *phylum Actinobacteria* also demonstrated an association with an elevated risk of bladder cancer[42]. *Genus Coprococcus* is a bacterium known for producing butyrate, and its presence may be associated with a reduction in the effectiveness of neoadjuvant chemoradiation therapy for rectal cancer [43]. Elevated abundance of the gut *genus Dorea* has the potential to serve as a predictive factor for farnesoid X receptor deactivation, which is recognized as a risk factor for metabolic dysfunction-associated steatotic liver disease[44]. The aforementioned findings indicate that certain protective or risky gut microbiota identified in this MR analysis are consistent with previous research and are likely to play significant roles as reference points in future clinical studies. Furthermore, the causal relationship between gut microbiota and CCA warrant further investigation through clinical and *in vivo* experiments to enhance our understanding of the “gut-liver axis” theories.

One major strength of this study is the utilization of the MR method, which assists in mitigating the impact of confounding variables, thereby enhancing the persuasiveness of the findings compared to observational research. Nevertheless, our analysis has several limitations that warrant consideration. Firstly, it is important to note that MR analyses were conducted at the bacterial genus level, as opposed to a more specific species level, due to the limited resolution provided by 16S rRNA sequencing. Secondly, the significance threshold for exposure IVs was set at  $1 \times 10^{-05}$  due to the inadequacy of IVs reaching genome-wide significance. However, IVs with F-statistics below 10 were excluded to mitigate the potential bias associated with weak instruments.





## CONCLUSION


In summary, this two-sample MR study offers new insights, suggesting a potential causal link between certain gut microbiota taxa and CCA. By utilizing genetic variants as IVs, we identified 15 microbial taxa that may confer either protective or detrimental effects on CCA risk. This study sheds new light on the intricate gut-liver axis interactions and microbiota-mediated mechanisms underlying CCA. Further experimental validations are warranted to consolidate the causality, delineate the molecular events, and exploit the clinical values of these candidate microbes.





**Figure 5** Leave-one-out stability tests causal estimates of specific gut microbiota taxa on gallbladder cancer and extrahepatic cholangiocarcinoma. Calculate the Mendelian randomization results of the remaining single nucleotide polymorphisms (SNPs) after removing the SNP one by one. A: *Genus Eubacteriumnodatum* group; B: *Genus Ruminococcustorques* group; C: *Genus Collinsella*; D: *Genus Coprococcus*; E: *Genus Dorea*; F: *Genus Eisenbergiella*; G: *Phylum Actinobacteria*. MR: Mendelian randomization; SNP: Single nucleotide polymorphisms.





**Figure 6** Leave-one-out stability tests causal estimates of specific gut microbiota taxa on intrahepatic cholangiocarcinoma. Calculate the mendelian randomization results of the remaining single nucleotide polymorphisms (SNPs) after removing the SNP one by one. A: Family *Veillonellaceae*; B: Genus *Alistipes*; C: Order *Enterobacteriales*; D: Phylum *Firmicutes*; E: Genus *Anaerostipes*; F: Genus *Paraprevotella*; G: Genus *Parasutterella*; H: Phylum *Verrucomicrobia*. MR: Mendelian randomization; SNP: Single nucleotide polymorphisms.

## ARTICLE HIGHLIGHTS

### Research background

Cholangiocarcinoma (CCA) is a highly malignant biliary tract cancer with poor prognosis. Previous studies have implicated the gut microbiota in CCA, but evidence for causal mechanisms is lacking.

### Research motivation

To investigate the causal relationship between gut microbiota and CCA risk.

### Research objectives

To investigate the causal relationship between gut microbiota and CCA risk.

### Research methods

We performed a two-sample mendelian randomization study to evaluate potential causal associations between gut microbiota and CCA risk using genome-wide association study summary statistics for 196 gut microbial taxa and CCA. Genetic variants were used as instrumental variables. Multiple sensitivity analyses assessed result robustness.

### Research results

Fifteen gut microbial taxa showed significant causal associations with CCA risk. Higher genetically predicted abundance of genus *Eubacteriumnodatum* group, Genus *Ruminococcustorques* group, *Coprococcus*, *Dorea*, and *Actinobacteria* were

associated with reduced risk of gallbladder cancer and extrahepatic CCA. Increased intrahepatic CCA risk was associated with higher abundance of *Veillonellaceae*, *Alistipes*, *Enterobacteriales*, and *Firmicutes*. Protective effects against CCA were suggested for *Collinsella*, *Eisenbergiella*, *Anaerostipes*, *Paraprevotella*, *Parasutterella*, and *Verrucomicrobia*. Sensitivity analyses indicated these findings were reliable without pleiotropy.

### Research conclusions

This pioneering study provides novel evidence that specific gut microbiota may play causal roles in CCA risk. Further experimental validation of these candidate microbes is warranted to consolidate causality and mechanisms.

### Research perspectives

Experimental validation of the candidate microbes identified to be causally associated with CCA risk. Further *in vitro* and *in vivo* studies could be conducted to consolidate the causal effects and explore the underlying molecular mechanisms. Analysis of species-level resolution of gut microbiota through metagenomic shotgun sequencing or other techniques. The current study was limited to genus-level associations due to 16S rRNA gene sequencing. A more detailed characterization at the species level could provide further insights.

## FOOTNOTES

**Author contributions:** Li QY interpreted the study design; Chen ZT and Ding CC downloaded data, performed statistical analysis, and drafted the manuscript; Chen KL, Gu YJ, and Lu CC performed data analysis and revised manuscript; Chen ZT, Ding CC, and Li QY helped revised our manuscript; all authors agreed to submit to the current journal, gave final approval of the version to be published, and agree to be accountable for all aspects of the work.

**Institutional review board statement:** The research does not involve any animal experiments and clinical data or human subjects, which does not require any special ethical clearance as per the guidelines of our institution.

**Informed consent statement:** Informed consent statement is not required since our manuscript solely utilizes publicly available data for analysis.

**Conflict-of-interest statement:** The authors declare that they have no competing interests.

**Data sharing statement:** The datasets used and analyzed in the present study are available from the corresponding authors on reasonable request. The datasets generated and/or analyzed during the current study are available in GWAS Catalog (<https://www.ebi.ac.uk/gwas/>) and MiBioGen (<https://mibiogen.gcc.rug.nl>) database.

**STROBE statement:** The authors have read the STROBE Statement—checklist of items, and the manuscript was prepared and revised according to the STROBE Statement—checklist of items.

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** China

**ORCID number:** Zhi-Tao Chen 0000-0001-9469-7176; Yang-Jun Gu 0000-0001-7666-8697; Qi-Yong Li 0000-0001-8421-5612.

**S-Editor:** Chen YL

**L-Editor:** A

**P-Editor:** Yu HG

## REFERENCES

- Brindley PJ, Bachini M, Ilyas SI, Khan SA, Loukas A, Sirica AE, Teh BT, Wongkham S, Gores GJ. Cholangiocarcinoma. *Nat Rev Dis Primers* 2021; 7: 65 [PMID: 34504109 DOI: 10.1038/s41572-021-00300-2]
- Rodrigues PM, Olaizola P, Paiva NA, Olaizola I, Aguirre-Lizaso A, Landa A, Bujanda L, Perugorria MJ, Banales JM. Pathogenesis of Cholangiocarcinoma. *Annu Rev Pathol* 2021; 16: 433-463 [PMID: 33264573 DOI: 10.1146/annurev-pathol-030220-020455]
- Labib PL, Goodchild G, Pereira SP. Molecular Pathogenesis of Cholangiocarcinoma. *BMC Cancer* 2019; 19: 185 [PMID: 30819129 DOI: 10.1186/s12885-019-5391-0]
- Khan SA, Tavolari S, Brandi G. Cholangiocarcinoma: Epidemiology and risk factors. *Liver Int* 2019; 39 Suppl 1: 19-31 [PMID: 30851228 DOI: 10.1111/liv.14095]
- Rizvi S, Khan SA, Hallemeier CL, Kelley RK, Gores GJ. Cholangiocarcinoma - evolving concepts and therapeutic strategies. *Nat Rev Clin Oncol* 2018; 15: 95-111 [PMID: 28994423 DOI: 10.1038/nrclinonc.2017.157]
- Banales JM, Marin JJG, Lamarca A, Rodrigues PM, Khan SA, Roberts LR, Cardinale V, Carpino G, Andersen JB, Braconi C, Calvisi DF,

- Perugorria MJ, Fabris L, Boulter L, Macias RIR, Gaudio E, Alvaro D, Gradilone SA, Strazzabosco M, Marzioni M, Coulouarn C, Fouassier L, Raggi C, Invernizzi P, Mertens JC, Moncsek A, Rizvi S, Heimbach J, Koerkamp BG, Bruix J, Forner A, Bridgewater J, Valle JW, Gores GJ. Cholangiocarcinoma 2020: the next horizon in mechanisms and management. *Nat Rev Gastroenterol Hepatol* 2020; **17**: 557-588 [PMID: 32606456 DOI: 10.1038/s41575-020-0310-z]
- 7 Elvevi A, Laffusa A, Scaravaglio M, Rossi RE, Longarini R, Stagno AM, Cristoferi L, Ciaccio A, Cortinovis DL, Invernizzi P, Massironi S. Clinical treatment of cholangiocarcinoma: an updated comprehensive review. *Ann Hepatol* 2022; **27**: 100737 [PMID: 35809836 DOI: 10.1016/j.aohp.2022.100737]
- 8 Wong SH, Yu J. Gut microbiota in colorectal cancer: mechanisms of action and clinical applications. *Nat Rev Gastroenterol Hepatol* 2019; **16**: 690-704 [PMID: 31554963 DOI: 10.1038/s41575-019-0209-8]
- 9 Fernandes MR, Aggarwal P, Costa RGF, Cole AM, Trinchieri G. Targeting the gut microbiota for cancer therapy. *Nat Rev Cancer* 2022; **22**: 703-722 [PMID: 36253536 DOI: 10.1038/s41568-022-00513-x]
- 10 Cai J, Sun L, Gonzalez FJ. Gut microbiota-derived bile acids in intestinal immunity, inflammation, and tumorigenesis. *Cell Host Microbe* 2022; **30**: 289-300 [PMID: 35271802 DOI: 10.1016/j.chom.2022.02.004]
- 11 Zhou CB, Zhou YL, Fang JY. Gut Microbiota in Cancer Immune Response and Immunotherapy. *Trends Cancer* 2021; **7**: 647-660 [PMID: 33674230 DOI: 10.1016/j.trecan.2021.01.010]
- 12 Lu Y, Yuan X, Wang M, He Z, Li H, Wang J, Li Q. Gut microbiota influence immunotherapy responses: mechanisms and therapeutic strategies. *J Hematol Oncol* 2022; **15**: 47 [PMID: 35488243 DOI: 10.1186/s13045-022-01273-9]
- 13 Tilg H, Adolph TE, Trauner M. Gut-liver axis: Pathophysiological concepts and clinical implications. *Cell Metab* 2022; **34**: 1700-1718 [PMID: 36208625 DOI: 10.1016/j.cmet.2022.09.017]
- 14 Zhang X, Coker OO, Chu ES, Fu K, Lau HCH, Wang YX, Chan AWH, Wei H, Yang X, Sung JJY, Yu J. Dietary cholesterol drives fatty liver-associated liver cancer by modulating gut microbiota and metabolites. *Gut* 2021; **70**: 761-774 [PMID: 32694178 DOI: 10.1136/gutjnl-2019-319664]
- 15 Jia X, Lu S, Zeng Z, Liu Q, Dong Z, Chen Y, Zhu Z, Hong Z, Zhang T, Du G, Xiang J, Wu D, Bai W, Yang B, Li Y, Huang J, Li H, Safadi R, Lu Y. Characterization of Gut Microbiota, Bile Acid Metabolism, and Cytokines in Intrahepatic Cholangiocarcinoma. *Hepatology* 2020; **71**: 893-906 [PMID: 31298745 DOI: 10.1002/hep.30852]
- 16 Lemoine S, Kemgang A, Ben Belkacem K, Straube M, Jegou S, Corpechot C, Saint-Antoine IBD Network, Chazouillères O, Housset C, Sokol H. Fungi participate in the dysbiosis of gut microbiota in patients with primary sclerosing cholangitis. *Gut* 2020; **69**: 92-102 [PMID: 31003979 DOI: 10.1136/gutjnl-2018-317791]
- 17 Philips CA, Phadke N, Ganesan K, Rajesh S, Padsalgi G, Ahamed R, John SK, Valiathan GC, Augustine P. Gut Microbiota in Alcoholic Hepatitis is Disparate from Those in Acute Alcoholic Pancreatitis and Biliary Disease. *J Clin Exp Hepatol* 2019; **9**: 690-698 [PMID: 31889749 DOI: 10.1016/j.jceh.2019.04.001]
- 18 Zhang T, Zhang S, Jin C, Lin Z, Deng T, Xie X, Deng L, Li X, Ma J, Ding X, Liu Y, Shan Y, Yu Z, Wang Y, Chen G, Li J. A Predictive Model Based on the Gut Microbiota Improves the Diagnostic Effect in Patients With Cholangiocarcinoma. *Front Cell Infect Microbiol* 2021; **11**: 751795 [PMID: 34888258 DOI: 10.3389/fcimb.2021.751795]
- 19 Herraiz E, Romero MR, Macias RIR, Monte MJ, Marin JGG. Clinical relevance of the relationship between changes in gut microbiota and bile acid metabolism in patients with intrahepatic cholangiocarcinoma. *Hepatobiliary Surg Nutr* 2020; **9**: 211-214 [PMID: 32355682 DOI: 10.21037/hbsn.2019.10.11]
- 20 Smith GD, Ebrahim S. 'Mendelian randomization': can genetic epidemiology contribute to understanding environmental determinants of disease? *Int J Epidemiol* 2003; **32**: 1-22 [PMID: 12689998 DOI: 10.1093/ije/dyg070]
- 21 Skrivankova VW, Richmond RC, Woolf BAR, Yarmolinsky J, Davies NM, Swanson SA, VanderWeele TJ, Higgins JPT, Timpson NJ, Dimou N, Langenberg C, Golub RM, Loder EW, Gallo V, Tybjaerg-Hansen A, Davey Smith G, Egger M, Richards JB. Strengthening the Reporting of Observational Studies in Epidemiology Using Mendelian Randomization: The STROBE-MR Statement. *JAMA* 2021; **326**: 1614-1621 [PMID: 34698778 DOI: 10.1001/jama.2021.18236]
- 22 Kurilshikov A, Medina-Gomez C, Bacigalupe R, Radjabzadeh D, Wang J, Demirkan A, Le Roy CI, Raygoza Garay JA, Finnicum CT, Liu X, Zhernakova DV, Bonder MJ, Hansen TH, Frost F, Rühlemann MC, Turpin W, Moon JY, Kim HN, Lüll K, Barkan E, Shah SA, Fornage M, Szopinska-Tokov J, Wallen ZD, Borisevich D, Agreus L, Andreasson A, Bang C, Bedrani L, Bell JT, Bisgaard H, Boehnke M, Boomsma DI, Burk RD, Claringbould A, Croitoru K, Davies GE, van Duijn CM, Duijts L, Falony G, Fu J, van der Graaf A, Hansen T, Homuth G, Hughes DA, Ijzerman RG, Jackson MA, Jaddoe VWV, Joossens M, Jørgensen T, Keszthelyi D, Knight R, Laakso M, Laudes M, Launer LJ, Lieb W, Lusis AJ, Masclee AAM, Moll HA, Mujagic Z, Qibin Q, Rothschild D, Shin H, Sørensen SJ, Steves CJ, Thorsen J, Timpson NJ, Tito RY, Vieira-Silva S, Völker U, Völzke H, Vösa U, Wade KH, Walter S, Watanabe K, Weiss S, Weiss FU, Weissbrod O, Westra HJ, Willemsen G, Payami H, Jonkers DMAE, Arias Vasquez A, de Geus EJC, Meyer KA, Stokholm J, Segal E, Org E, Wijmenga C, Kim HL, Kaplan RC, Spector TD, Uitterlinden AG, Rivadeneira F, Franke A, Lerch MM, Franke L, Sanna S, D'Amato M, Pedersen O, Paterson AD, Kraaij R, Raes J, Zhernakova A. Large-scale association analyses identify host factors influencing human gut microbiome composition. *Nat Genet* 2021; **53**: 156-165 [PMID: 33462485 DOI: 10.1038/s41588-020-00763-1]
- 23 Jiang L, Zheng Z, Fang H, Yang J. A generalized linear mixed model association tool for biobank-scale data. *Nat Genet* 2021; **53**: 1616-1621 [PMID: 34737426 DOI: 10.1038/s41588-021-00954-4]
- 24 Chen S, Zhou G, Han H, Jin J, Li Z. Causal effects of specific gut microbiota on bone mineral density: a two-sample Mendelian randomization study. *Front Endocrinol (Lausanne)* 2023; **14**: 1178831 [PMID: 37645419 DOI: 10.3389/fendo.2023.1178831]
- 25 Li Z, Chen Y, Ke H. Investigating the Causal Relationship Between Gut Microbiota and Crohn's Disease: A Mendelian Randomization Study. *Gastroenterology* 2023 [PMID: 37678500 DOI: 10.1053/j.gastro.2023.08.047]
- 26 Burgess S, Davey Smith G, Davies NM, Dudbridge F, Gill D, Glymour MM, Hartwig FP, Kutalik Z, Holmes MV, Minelli C, Morrison JV, Pan W, Relton CL, Theodoratou E. Guidelines for performing Mendelian randomization investigations: update for summer 2023. *Wellcome Open Res* 2019; **4**: 186 [PMID: 32760811 DOI: 10.12688/wellcomeopenres.15555.3]
- 27 Razumilava N, Gores GJ. Cholangiocarcinoma. *Lancet* 2014; **383**: 2168-2179 [PMID: 24581682 DOI: 10.1016/S0140-6736(13)61903-0]
- 28 Cillo U, Fondevila C, Donadon M, Gringeri E, Mocchegiani F, Schlitt HJ, Ijzermans JNM, Vivarelli M, Zieniewicz K, Olde Damink SWM, Groot Koerkamp B. Surgery for cholangiocarcinoma. *Liver Int* 2019; **39** Suppl 1: 143-155 [PMID: 30843343 DOI: 10.1111/liv.14089]
- 29 Yang J, Wei H, Zhou Y, Szeto CH, Li C, Lin Y, Coker OO, Lau HCH, Chan AWH, Sung JJY, Yu J. High-Fat Diet Promotes Colorectal Tumorigenesis Through Modulating Gut Microbiota and Metabolites. *Gastroenterology* 2022; **162**: 135-149.e2 [PMID: 34461052 DOI: 10.1053/j.gastro.2021.08.041]

- 30 **Mohr AE**, Jäger R, Carpenter KC, Kerksick CM, Purpura M, Townsend JR, West NP, Black K, Gleeson M, Pyne DB, Wells SD, Arent SM, Kreider RB, Campbell BI, Bannock L, Scheiman J, Wissent CJ, Pane M, Kalman DS, Pugh JN, Ortega-Santos CP, Ter Haar JA, Arciero PJ, Antonio J. The athletic gut microbiota. *J Int Soc Sports Nutr* 2020; **17**: 24 [PMID: 32398103 DOI: 10.1186/s12970-020-00353-w]
- 31 **Muscolino P**, Granata B, Omero F, De Pasquale C, Campana S, Calabrò A, D'Anna F, Drommi F, Pezzino G, Cavaliere R, Ferlazzo G, Silvestris N, Speranza D. Potential predictive role of gut microbiota to immunotherapy in HCC patients: a brief review. *Front Oncol* 2023; **13**: 1247614 [PMID: 37692859 DOI: 10.3389/fonc.2023.1247614]
- 32 **Xue X**, Li R, Chen Z, Li G, Liu B, Guo S, Yue Q, Yang S, Xie L, Zhang Y, Zhao J, Tan R. The role of the symbiotic microecosystem in cancer: gut microbiota, metabolome, and host immunome. *Front Immunol* 2023; **14**: 1235827 [PMID: 37691931 DOI: 10.3389/fimmu.2023.1235827]
- 33 **Popa AD**, Niță O, Gherasim A, Enache AI, Caba L, Mihalache L, Arhire LI. A Scoping Review of the Relationship between Intermittent Fasting and the Human Gut Microbiota: Current Knowledge and Future Directions. *Nutrients* 2023; **15** [PMID: 37432222 DOI: 10.3390/nu15092095]
- 34 **Barber TM**, Kabisch S, Pfeiffer AFH, Weickert MO. The Effects of the Mediterranean Diet on Health and Gut Microbiota. *Nutrients* 2023; **15** [PMID: 37432307 DOI: 10.3390/nu15092150]
- 35 **Zhang L**, Chen C, Chai D, Kuang T, Deng W, Wang W. Alterations of gut mycobiota profiles in intrahepatic cholangiocarcinoma. *Front Microbiol* 2022; **13**: 1090392 [PMID: 36687597 DOI: 10.3389/fmicb.2022.1090392]
- 36 **Ponziani FR**, Nicoletti A, Gasbarrini A, Pompili M. Diagnostic and therapeutic potential of the gut microbiota in patients with early hepatocellular carcinoma. *Ther Adv Med Oncol* 2019; **11**: 1758835919848184 [PMID: 31205505 DOI: 10.1177/1758835919848184]
- 37 **Li JQ**, Li JL, Xie YH, Wang Y, Shen XN, Qian Y, Han JX, Chen YX, Fang JY. *Saccharomyces cerevisiae* may serve as a probiotic in colorectal cancer by promoting cancer cell apoptosis. *J Dig Dis* 2020; **21**: 571-582 [PMID: 33245627 DOI: 10.1111/1751-2980.12930]
- 38 **Shock T**, Badang L, Ferguson B, Martinez-Guryn K. The interplay between diet, gut microbes, and host epigenetics in health and disease. *J Nutr Biochem* 2021; **95**: 108631 [PMID: 33789148 DOI: 10.1016/j.jnutbio.2021.108631]
- 39 **Jia W**, Xie G, Jia W. Bile acid-microbiota crosstalk in gastrointestinal inflammation and carcinogenesis. *Nat Rev Gastroenterol Hepatol* 2018; **15**: 111-128 [PMID: 29018272 DOI: 10.1038/nrgastro.2017.119]
- 40 **Lili Z**, Junyan W, Hongfei Z, Baoqing Z, Bolin Z. Detoxification of cancerogenic compounds by lactic acid bacteria strains. *Crit Rev Food Sci Nutr* 2018; **58**: 2727-2742 [PMID: 29053003 DOI: 10.1080/10408398.2017.1339665]
- 41 **Huang R**, He K, Duan X, Xiao J, Wang H, Xiang G. Changes of Intestinal Microflora in Colorectal Cancer Patients after Surgical Resection and Chemotherapy. *Comput Math Methods Med* 2022; **2022**: 1940846 [PMID: 35251295 DOI: 10.1155/2022/1940846]
- 42 **Mingdong W**, Xiang G, Yongjun Q, Mingshuai W, Hao P. Causal associations between gut microbiota and urological tumors: a two-sample mendelian randomization study. *BMC Cancer* 2023; **23**: 854 [PMID: 37697271 DOI: 10.1186/s12885-023-11383-3]
- 43 **Huang X**, Chen C, Xie W, Zhou C, Tian X, Zhang Z, Wang Q, Chang H, Xiao W, Zhang R, Gao Y. Metagenomic Analysis of Intratumoral Microbiome Linking to Response to Neoadjuvant Chemoradiotherapy in Rectal Cancer. *Int J Radiat Oncol Biol Phys* 2023; **117**: 1255-1269 [PMID: 37433373 DOI: 10.1016/j.ijrobp.2023.06.2515]
- 44 **Yang G**, Liu R, Rezaei S, Liu X, Wan YY. Uncovering the Gut-Liver Axis Biomarkers for Predicting Metabolic Burden in Mice. *Nutrients* 2023; **15** [PMID: 37571345 DOI: 10.3390/nu15153406]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

**Telephone:** +1-925-3991568

**E-mail:** [office@baishideng.com](mailto:office@baishideng.com)

**Help Desk:** <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

