

# World Journal of *Gastrointestinal Oncology*

*World J Gastrointest Oncol* 2024 April 15; 16(4): 1091-1675



## EDITORIAL

- 1091 Parallel pathways: A chronicle of evolution in rectal and breast cancer surgery  
*Pesce A, Fabbri N, Iovino D, Feo CV*
- 1097 Hepatitis B virus genotypes in precision medicine of hepatitis B-related hepatocellular carcinoma: Where we are now  
*Sukowati CHC, Jayanti S, Turyadi T, Muljono DH, Tiribelli C*

## REVIEW

- 1104 Novel milestones for early esophageal carcinoma: From bench to bed  
*Qi JH, Huang SL, Jin SZ*
- 1119 Colorectal cancer screening: A review of current knowledge and progress in research  
*Lopes SR, Martins C, Santos IC, Teixeira M, Gamito É, Alves AL*
- 1134 New avenues for the treatment of immunotherapy-resistant pancreatic cancer  
*Silva LGO, Lemos FFB, Luz MS, Rocha Pinheiro SL, Calmon MDS, Correa Santos GL, Rocha GR, de Melo FF*

## MINIREVIEWS

- 1154 Present situation of minimally invasive surgical treatment for early gastric cancer  
*Li CY, Wang YF, Luo LK, Yang XJ*
- 1166 Mixed neuroendocrine non-neuroendocrine neoplasms in gastroenteropancreatic tract  
*Díaz-López S, Jiménez-Castro J, Robles-Barraza CE, Ayala-de Miguel C, Chaves-Conde M*
- 1180 Esophageal cancer screening, early detection and treatment: Current insights and future directions  
*Qu HT, Li Q, Hao L, Ni YJ, Luan WY, Yang Z, Chen XD, Zhang TT, Miao YD, Zhang F*

## ORIGINAL ARTICLE

## Retrospective Cohort Study

- 1192 Pre-operative enhanced magnetic resonance imaging combined with clinical features predict early recurrence of hepatocellular carcinoma after radical resection  
*Chen JP, Yang RH, Zhang TH, Liao LA, Guan YT, Dai HY*
- 1204 Clinical analysis of multiple primary gastrointestinal malignant tumors: A 10-year case review of a single-center  
*Zhu CL, Peng LZ*

**Retrospective Study**

- 1213** Predictive model for non-malignant portal vein thrombosis associated with cirrhosis based on inflammatory biomarkers  
*Nie GL, Yan J, Li Y, Zhang HL, Xie DN, Zhu XW, Li X*
- 1227** Predictive modeling for postoperative delirium in elderly patients with abdominal malignancies using synthetic minority oversampling technique  
*Hu WJ, Bai G, Wang Y, Hong DM, Jiang JH, Li JX, Hua Y, Wang XY, Chen Y*
- 1236** Efficacy and predictive factors of transarterial chemoembolization combined with lenvatinib plus programmed cell death protein-1 inhibition for unresectable hepatocellular carcinoma  
*Ma KP, Fu JX, Duan F, Wang MQ*
- 1248** Should we perform sigmoidoscopy for colorectal cancer screening in people under 45 years?  
*Leong W, Guo JQ, Ning C, Luo FF, Jiao R, Yang DY*
- 1256** Computed tomography-based radiomics diagnostic approach for differential diagnosis between early- and late-stage pancreatic ductal adenocarcinoma  
*Ren S, Qian LC, Cao YY, Daniels MJ, Song LN, Tian Y, Wang ZQ*
- 1268** Prognostic analysis of related factors of adverse reactions to immunotherapy in advanced gastric cancer and establishment of a nomogram model  
*He XX, Du B, Wu T, Shen H*

**Clinical Trials Study**

- 1281** Safety and efficacy of a programmed cell death 1 inhibitor combined with oxaliplatin plus S-1 in patients with Borrmann large type III and IV gastric cancers  
*Bao ZH, Hu C, Zhang YQ, Yu PC, Wang Y, Xu ZY, Fu HY, Cheng XD*

**Observational Study**

- 1296** Computed tomography radiogenomics: A potential tool for prediction of molecular subtypes in gastric stromal tumor  
*Yin XN, Wang ZH, Zou L, Yang CW, Shen CY, Liu BK, Yin Y, Liu XJ, Zhang B*
- 1309** Application of texture signatures based on multiparameter-magnetic resonance imaging for predicting microvascular invasion in hepatocellular carcinoma: Retrospective study  
*Nong HY, Cen YY, Qin M, Qin WQ, Xie YX, Li L, Liu MR, Ding K*
- 1319** Causal roles of gut microbiota in cholangiocarcinoma etiology suggested by genetic study  
*Chen ZT, Ding CC, Chen KL, Gu YJ, Lu CC, Li QY*
- 1334** Is recovery enhancement after gastric cancer surgery really a safe approach for elderly patients?  
*Li ZW, Luo XJ, Liu F, Liu XR, Shu XP, Tong Y, Lv Q, Liu XY, Zhang W, Peng D*
- 1344** Establishment of a cholangiocarcinoma risk evaluation model based on mucin expression levels  
*Yang CY, Guo LM, Li Y, Wang GX, Tang XW, Zhang QL, Zhang LF, Luo JY*

- 1361** Effectiveness of fecal DNA syndecan-2 methylation testing for detection of colorectal cancer in a high-risk Chinese population

*Luo WF, Jiao YT, Lin XL, Zhao Y, Wang SB, Shen J, Deng J, Ye YF, Han ZP, Xie FM, He JH, Wan Y*

#### Clinical and Translational Research

- 1374** Clinical and socioeconomic determinants of survival in biliary tract adenocarcinomas

*Sahyoun L, Chen K, Tsay C, Chen G, Protiva P*

- 1384** Risk factors, prognostic factors, and nomograms for distant metastasis in patients with diagnosed duodenal cancer: A population-based study

*Shang JR, Xu CY, Zhai XX, Xu Z, Qian J*

- 1421** NOX4 promotes tumor progression through the MAPK-MEK1/2-ERK1/2 axis in colorectal cancer

*Xu YJ, Huo YC, Zhao QT, Liu JY, Tian YJ, Yang LL, Zhang Y*

#### Basic Study

- 1437** Curcumin inhibits the growth and invasion of gastric cancer by regulating long noncoding RNA AC022424.2

*Wang BS, Zhang CL, Cui X, Li Q, Yang L, He ZY, Yang Z, Zeng MM, Cao N*

- 1453** MicroRNA-298 determines the radio-resistance of colorectal cancer cells by directly targeting human dual-specificity tyrosine(Y)-regulated kinase 1A

*Shen MZ, Zhang Y, Wu F, Shen MZ, Liang JL, Zhang XL, Liu XJ, Li XS, Wang RS*

- 1465** Human  $\beta$ -defensin-1 affects the mammalian target of rapamycin pathway and autophagy in colon cancer cells through long non-coding RNA TCONS\_00014506

*Zhao YX, Cui Y, Li XH, Yang WH, An SX, Cui JX, Zhang MY, Lu JK, Zhang X, Wang XM, Bao LL, Zhao PW*

- 1479** FAM53B promotes pancreatic ductal adenocarcinoma metastasis by regulating macrophage M2 polarization

*Pei XZ, Cai M, Jiang DW, Chen SH, Wang QQ, Lu HM, Lu YF*

- 1500** Transcriptome sequencing reveals novel biomarkers and immune cell infiltration in esophageal tumorigenesis

*Sun JR, Chen DM, Huang R, Wang RT, Jia LQ*

- 1514** Construction of CDKN2A-related competitive endogenous RNA network and identification of GAS5 as a prognostic indicator for hepatocellular carcinoma

*Pan Y, Zhang YR, Wang LY, Wu LN, Ma YQ, Fang Z, Li SB*

- 1532** Two missense STK11 gene variations impaired LKB1/adenosine monophosphate-activated protein kinase signaling in Peutz-Jeghers syndrome

*Liu J, Zeng SC, Wang A, Cheng HY, Zhang QJ, Lu GX*

- 1547** Long noncoding RNAs HAND2-AS1 ultrasound microbubbles suppress hepatocellular carcinoma progression by regulating the miR-873-5p/tissue inhibitor of matrix metalloproteinase-2 axis

*Zou Q, Wang HW, Di XL, Li Y, Gao H*

- 1564** Upregulated lncRNA PRNT promotes progression and oxaliplatin resistance of colorectal cancer cells by regulating HIPK2 transcription

*Li SN, Yang S, Wang HQ, Hui TL, Cheng M, Zhang X, Li BK, Wang GY*

### SYSTEMATIC REVIEWS

- 1578** Prognosis value of heat-shock proteins in esophageal and esophagogastric cancer: A systematic review and meta-analysis

*Nakamura ET, Park A, Pereira MA, Kikawa D, Tustumi F*

- 1596** Risk factors for hepatocellular carcinoma associated with hepatitis C genotype 3 infection: A systematic review

*Farooq HZ, James M, Abbott J, Oyibo P, Divall P, Choudhry N, Foster GR*

### META-ANALYSIS

- 1613** Effectiveness and tolerability of programmed cell death protein-1 inhibitor + chemotherapy compared to chemotherapy for upper gastrointestinal tract cancers

*Zhang XM, Yang T, Xu YY, Li BZ, Shen W, Hu WQ, Yan CW, Zong L*

- 1626** Success rate of current human-derived gastric cancer organoids establishment and influencing factors: A systematic review and meta-analysis

*Jiang KL, Wang XX, Liu XJ, Guo LK, Chen YQ, Jia QL, Yang KM, Ling JH*

### CASE REPORT

- 1647** Pathologically successful conversion hepatectomy for advanced giant hepatocellular carcinoma after multidisciplinary therapy: A case report and review of literature

*Chu JH, Huang LY, Wang YR, Li J, Han SL, Xi H, Gao WX, Cui YY, Qian MP*

- 1660** Clinical pathological characteristics of “crawling-type” gastric adenocarcinoma cancer: A case report

*Xu YW, Song Y, Tian J, Zhang BC, Yang YS, Wang J*

- 1668** Primary pancreatic peripheral T-cell lymphoma: A case report

*Bai YL, Wang LJ, Luo H, Cui YB, Xu JH, Nan HJ, Yang PY, Niu JW, Shi MY*

**ABOUT COVER**

Peer Reviewer of *World Journal of Gastrointestinal Oncology*, Lie Zheng, Director, Professor, Department of Gastroenterology, Shaanxi Provincial Hospital of Traditional Chinese Medicine, Xi'an 730000, Shaanxi Province, China. xinliwen696@126.com

**AIMS AND SCOPE**

The primary aim of *World Journal of Gastrointestinal Oncology* (WJGO, *World J Gastrointest Oncol*) is to provide scholars and readers from various fields of gastrointestinal oncology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGO mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal oncology and covering a wide range of topics including liver cell adenoma, gastric neoplasms, appendiceal neoplasms, biliary tract neoplasms, hepatocellular carcinoma, pancreatic carcinoma, cecal neoplasms, colonic neoplasms, colorectal neoplasms, duodenal neoplasms, esophageal neoplasms, gallbladder neoplasms, *etc.*

**INDEXING/ABSTRACTING**

The WJGO is now abstracted and indexed in PubMed, PubMed Central, Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJGO as 3.0; IF without journal self cites: 2.9; 5-year IF: 3.0; Journal Citation Indicator: 0.49; Ranking: 157 among 241 journals in oncology; Quartile category: Q3; Ranking: 58 among 93 journals in gastroenterology and hepatology; and Quartile category: Q3. The WJGO's CiteScore for 2022 is 4.1 and Scopus CiteScore rank 2022: Gastroenterology is 71/149; Oncology is 197/366.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: Xiang-Di Zhang; Production Department Director: Xiang Li; Cover Editor: Jia-Ru Fan.

**NAME OF JOURNAL**

*World Journal of Gastrointestinal Oncology*

**ISSN**

ISSN 1948-5204 (online)

**LAUNCH DATE**

February 15, 2009

**FREQUENCY**

Monthly

**EDITORS-IN-CHIEF**

Monjur Ahmed, Florin Burada

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/1948-5204/editorialboard.htm>

**PUBLICATION DATE**

April 15, 2024

**COPYRIGHT**

© 2024 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>



Observational Study

## Effectiveness of fecal DNA syndecan-2 methylation testing for detection of colorectal cancer in a high-risk Chinese population

Wen-Feng Luo, Yu-Ting Jiao, Xiao-Ling Lin, Ying Zhao, Sheng-Bo Wang, Jian Shen, Jie Deng, Yu-Feng Ye, Ze-Ping Han, Fang-Mei Xie, Jin-Hua He, Yu Wan

**Specialty type:** Oncology

**Provenance and peer review:**

Unsolicited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): 0

Grade B (Very good): B

Grade C (Good): 0

Grade D (Fair): 0

Grade E (Poor): 0

**P-Reviewer:** Miyoshi E, Japan

**Received:** December 21, 2023

**Peer-review started:** December 21, 2023

**First decision:** December 28, 2023

**Revised:** January 12, 2024

**Accepted:** February 7, 2024

**Article in press:** February 7, 2024

**Published online:** April 15, 2024



**Wen-Feng Luo, Xiao-Ling Lin, Ying Zhao, Jian Shen, Ze-Ping Han, Fang-Mei Xie, Jin-Hua He,** Central Laboratory of Panyu Central Hospital, Guangzhou Panyu Central Hospital, Guangzhou 511400, Guangdong Province, China

**Yu-Ting Jiao,** South China Normal University-Panyu Central Hospital Joint Laboratory of Translational Medical Research, Guangzhou Panyu Central Hospital, Guangzhou 511400, Guangdong Province, China

**Sheng-Bo Wang, Yu Wan,** Digestive Disease Center, Guangzhou Panyu Central Hospital, Guangzhou 511400, Guangdong Province, China

**Jie Deng,** Shunde Vocational and Technical College, Foshan 528300, Guangdong Province, China

**Yu-Feng Ye,** Medical Imaging Institute of Panyu, Guangzhou Panyu Central Hospital, Guangzhou 511400, Guangdong Province, China

**Corresponding author:** Wen-Feng Luo, MS, PhD, Assistant Professor, Doctor, Central Laboratory of Panyu Central Hospital, Guangzhou Panyu Central Hospital, No. 8 East Fuyou Road, Qiaonan Street, Panyu District, Guangzhou 511400, Guangdong Province, China.

[luowenfeng163@163.com](mailto:luowenfeng163@163.com)

### Abstract

#### BACKGROUND

Colorectal cancer (CRC) is among the most prevalent and life-threatening malignancies worldwide. Syndecan-2 methylation (mSDC2) testing has emerged as a widely used biomarker for early detection of CRC in stool and serum samples. Cancer (CRC) is among the most prevalent and life-threatening malignancies worldwide. mSDC2 testing has emerged as a widely used biomarker for early detection of CRC in stool and serum samples.

#### AIM

To validate the effectiveness of fecal DNA mSDC2 testing in the detection of CRC among a high-risk Chinese population to provide evidence-based data for the development of diagnostic and/or screening guidelines for CRC in China.

#### METHODS



A high-risk Chinese cohort consisting of 1130 individuals aged 40-79 years was selected for evaluation *via* fecal mSDC2 testing. Sensitivity and specificity for CRC, advanced adenoma (AA) and advanced colorectal neoplasia (ACN) were determined. High-risk factors for the incidence of colorectal lesions were determined and a logistic regression model was constructed to reflect the efficacy of the test.

## RESULTS

A total of 1035 high-risk individuals were included in this study according to established criteria. Among them, 16 suffered from CRC (1.55%), 65 from AA (6.28%) and 189 from non-AAAs (18.26%); 150 patients were diagnosed with polyps (14.49%). Diagnoses were established based upon colonoscopic and pathological examinations. Sensitivities of the mSDC2 test for CRC and AA were 87.50% and 40.00%, respectively; specificities were 95.61% for other groups. Positive predictive values of the mSDC2 test for CRC, AA and ACN were 16.09%, 29.89% and 45.98%, respectively; the negative predictive value for CRC was 99.79%. After adjusting for other high-risk covariates, mSDC2 test positivity was found to be a significant risk factor for the occurrence of ACN ( $P < 0.001$ ).

## CONCLUSION

Our findings confirmed that offering fecal mSDC2 testing and colonoscopy in combination for CRC screening is effective for earlier detection of malignant colorectal lesions in a high-risk Chinese population.

**Key Words:** Colorectal cancer; Syndecan-2; DNA methylation; Chinese population

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** A high-risk Chinese cohort composed of 1130 individuals 40-79 years of age was selected for evaluation using the fecal syndecan-2 methylation (mSDC2) test. Sensitivity and specificity to colorectal cancer (CRC), advanced adenoma and advanced colorectal neoplasia were quantified. High-risk factors for the incidence of colorectal lesions were analyzed; a logistic regression model was subsequently constructed to better reflect the efficacy of fecal mSDC2 testing. The results of this CRC screening study revealed that offering patients a combination of fecal mSDC2 testing and colonoscopy is ideal for facilitating early detection of CRC among a high-risk Chinese population.

**Citation:** Luo WF, Jiao YT, Lin XL, Zhao Y, Wang SB, Shen J, Deng J, Ye YF, Han ZP, Xie FM, He JH, Wan Y. Effectiveness of fecal DNA syndecan-2 methylation testing for detection of colorectal cancer in a high-risk Chinese population. *World J Gastrointest Oncol* 2024; 16(4): 1361-1373

**URL:** <https://www.wjgnet.com/1948-5204/full/v16/i4/1361.htm>

**DOI:** <https://dx.doi.org/10.4251/wjgo.v16.i4.1361>

## INTRODUCTION

Colorectal cancer (CRC) is a life-threatening malignancy that is highly prevalent in China. Indeed, its incidence and mortality rates have steadily increased. According to data from the International Agency for Research on Cancer, in 2020 alone, China witnessed 555.5 thousand new cases and 283.8 thousand deaths from CRC, ranking second and fifth, respectively, among malignancies[1]. Although CRC incidence is relatively low among individuals aged less than 50 years, incidence markedly increases with age. Importantly, recent data has suggested a rise in early-onset CRC cases[2-4]. Progression from colorectal adenoma to CRC takes approximately a decade on average[5]. Consequently, implementation of screening programs to detect precancerous lesions among average-risk populations is a highly effective strategy for reducing the incidence of CRC.

Currently, several screening methods for CRC include colonoscopy, fecal immunochemical testing (FIT) and serum biomarker evaluation. However, each approach has its limitations. Although colonoscopy remains the gold standard for CRC diagnosis, its invasiveness, complicated intestinal preparation and poor compliance rates have hindered widespread adoption[6]. Importantly, colonoscopy may miss certain lesions[7]. While FIT offers an efficient, non-invasive screening option for the average-risk population, it has modest sensitivity in detecting CRC[7,8]. Levels of certain serum proteins such as carbohydrate antigen 19-9 (CA19-9) and carcinoembryonic antigen (CEA) have been utilized as pan-cancer biomarkers for various malignancies such as those of the pancreas, breast and colon[9-12]. However, their specificity for detecting CRC is limited. Such limitations highlight the need for superior alternative screening methods to improve rates of CRC detection and diagnosis.

Due to advancements in biotechnology, molecular diagnostics have attracted rapidly increasing interest among *in vitro* techniques. Importantly, such methods outperform biochemical and immunological diagnostics in terms of accuracy and prognostic worth. Stool-based DNA testing is an emerging CRC screening method that detects tumors by identifying relevant biomarkers in stool samples. Aberrant genetic and epigenetic alterations in the setting of CRC may be tracked by extracting DNA from exfoliated cells[13-15]. In the setting of carcinogenesis, DNA methylation is also highly relevant and considered to be a good biomarker of malignancy[16-18]. Genes such as *Vimentin*, *NDRG4*, *BMP3* and *TFP12* have been



employed as molecular markers to develop sensitive and simple screening approaches for evaluation of DNA methylation[19,20]. Due to its non-invasive, convenient and sensitive characteristics, stool-based DNA testing is predicted to become extensively employed in large-scale screening among average-risk populations. Cologuard® (Exact Sciences, Madison WI) was the first such commercial product authorized for clinical use by the United States Food and Drug Administration. It incorporates fecal hemoglobin immunoassays and quantitative molecular detection of methylated *BMP3*, methylated *NDRG4* and mutant *KRAS*[21]. Furthermore, the United States Preventive Services Task Force initially recommended stool DNA testing as a screening approach for CRC in 2016[22]. In particular, syndecan-2 methylation (mSDC2) testing has emerged as a tool for early detection of CRC in stool and serum samples[23,24]. The *SDC2* gene is a transmembrane heparan sulfate proteoglycan that is a member of the *SDC* proteoglycan family. It is relevant in cell proliferation, angiogenesis and cell migration, and is widely expressed in colonic mesenchymal cells. Zhang *et al*[25] reported that levels of *SDC2* regulatory region methylation are significantly higher in colorectal tumor specimens as compared to paired adjacent non-malignant tissues. Zhao *et al*[26] showed that the methylation level of *SDC2* in tumor tissues surpasses that in corresponding non-cancerous tissues and exceeds levels observed in polyps. As mSDC2 can be detected in both adenomas as well as early carcinogenesis, its methylation is considered to be a gradual process. Relative to normal colorectal tissues and polyps, *SDC2* was found to be significantly more methylated in different stages of CRC pathogenesis and advanced intestinal adenoma formation[27,28]. On meta analyses, pooled sensitivity and specificity of mSDC2 testing for CRC was 0.81 [95% confidence interval (CI): 0.74-0.86] and 0.95 (95% CI: 0.93-0.96), respectively[20]. No significant differences in sensitivity (0.82, 95% CI: 0.58-0.94; and 0.83, 95% CI: 0.77-0.88, respectively) or specificity (0.95, 95% CI: 0.91-0.97; and 0.94, 95% CI: 0.90-0.96, respectively) between blood and stool samples were noted.

Here, we aimed to evaluate the efficacy of mSDC2 testing as a screening tool for CRC in a high-risk population. Our primary objective was to assess both sensitivity and specificity of mSDC2 testing for detection of CRC and advanced adenoma (AA). Furthermore, we study aimed to evaluate clinical usefulness of mSDC2 testing as an adjunctive tool for CRC and precancerous lesion identification.

## MATERIALS AND METHODS

### Study design

From April 2020 to May 2022, CRC screening with stool-based mSDC2 testing was performed at the Panyu Center Hospital of Guangzhou (Guangdong, China) as a primary screening method. Over 1000 high-risk participants were enrolled in this study. Fresh stool was collected from every participant for mSDC2 testing prior to initiating bowel preparation for colonoscopy; subjects subsequently underwent colonoscopy for further diagnostic evaluation. This study was approved by the institutional review board, IRB approval number: No. [2019]62, and written, informed consent was obtained from all study participants.

### Inclusion and exclusion criteria

Individual outpatients 40-79 years old completed a high-risk factor questionnaire (HRFQ). A positive HRFQ was defined as: (1) A family history of CRC in first-degree relatives; (2) A history of polyps; and/or (3) A history of two or more of the following: (1) Chronic diarrhea; (2) Chronic constipation; (3) Mucoid bloody feces; (4) Chronic appendicitis or appendectomy; (5) Chronic cholecystitis or cholecystectomy; and (6) Psychiatric trauma (*e.g.*, divorce, death of first-degree relatives)[29-31].

Individuals unsuitable for colonoscopic examination such as those suffering severe cardiac or chronic renal conditions or pregnant patients were excluded from analyses. Participants previously diagnosed with any malignancies, those who had undergone chemo- or immuno-therapy treatments or subjects who were unable or unwilling to provide written, informed consent were also excluded from analyses.

### SDC2 methylation test

**Specimen collection and processing:** Each participant was required to provide about 4.5 g of fresh stool for mSDC2 testing using a stool collection device (Creative Biosciences Co. Ltd., Guangzhou, China) at home. Samples stored in preservation buffer were sent to the designated clinical laboratory within two days of collection. Stool specimens were immediately homogenized and centrifuged upon receipt. Supernatants were then aliquoted and frozen at -80 °C until further use. Frozen aliquots were subsequently tested in batches by experienced laboratory technicians using Colosafe® testing reagent kits (Creative Biosciences).

**mSDC2 testing:** For blind testing, each stool collection device was labeled with a unique code and no other identifying information. Target and control genes [*SDC2* and  $\beta$ -actin (*ACTB*)] were extracted from stool supernatant after centrifugation and subsequently enriched and purified *via* sequence-specific capture technology. Positive and negative controls *SDC2* gene controls were also tested in parallel. Real-time quantitative methylation-specific polymerase chain reaction was employed to quantitatively detect *SDC2* and *ACTB* methylation status in stool samples using a LightCycler 480 II machine (Roche, Basel, Switzerland). *ACTB* was amplified as a reference for DNA input. Primers, probes and amplification conditions in this study were all as previously described[32].

**Positive criteria:** Stool samples with CT values of *SDC2*  $\leq$  38 were classified as positive while those with CT values of *SDC2*  $>$  38 were classified as negative under the precondition that CT values of *ACTB*  $\leq$  36. Stool samples with CT values of *ACTB*  $>$  36 were considered invalid.

## Colonoscopy and pathological examination

After mSDC2 testing, all participants were required to undergo colonoscopy. Colonoscopic examinations were performed by gastroenterology specialist at the endoscopy center of Panyu Center Hospital. After standard bowel preparation, endoscopy with a minimum withdrawal time of 6 min was performed. All lesions noted were measured with opened biopsy forceps and recorded based on size, morphology and localization. Neoplastic lesions observed on colonoscopy were immediately removed and/or biopsied for histologic diagnosis. Individuals suspected of having CRC or polyps that could not be removed endoscopically were referred for surgery. If more than one polyp was noted, the most advanced pathological lesion or largest lesion was considered in analyses.

CRC was staged according to eight edition guidelines of the American Joint Committee on Cancer[33]. AA was defined as an adenomatous lesion with either a diameter of  $\geq 10$  mm, of  $\geq 25\%$  villous character or characterized by high-grade dysplasia[29,30]. Advanced colorectal neoplasia (ACN) was defined as evident CRC and AA lesions. Non-adenomatous polyps (NAP) included inflammatory, hyperplastic and juvenile polyps[29]. Other findings of diagnostic evaluation included completely normal appearance as well as the presence of colitis and colonic diverticulum.

## Statistical analyses

All data were represented as numbers and relevant detection rates (DR). Pearson chi-squared or Fisher's exact tests were used, as deemed appropriate, to compare detection data and patient clinicopathological characteristics. A logistic regression model was constructed to compare high-risk factors for colorectal lesions as well as odds ratios and relevant 95%CI to determine exposure risk. Additionally, *P* values of logistic model data were adjusted *via* false discover rate methods to ensure reliability of significance testing. Data analyses were performed using SPSS software version 23.0 (IBM, Armonk, United States). Receiver operating characteristic (ROC) curve and predictive value [sensitivity, specificity, positive predictive value (PPV), negative predictive value (NPV), area under the curve and relevant 95%CI] analyses were performed using GraphPad Prism version 8.0. *P* < 0.05 was considered statistically significant at  $\alpha = 0.05$ .

# RESULTS

## Baseline data

This study enrolled a total of 1130 high-risk subjects who were either outpatients at our hospital's department of gastroenterology or attending well-adult examination at the health examination center. After excluding 49 persons with inadequate colonoscopy findings and 29 persons who did not complete mSDC2 testing, a total of 1052 subjects were effectively screened and the pass rate was 93.10% (1052/1130). After excluding three patients who lacked pathological data, 10 subjects without qualifying stool samples and four persons who did not meet the age requirement, 1035 cases were finally evaluated in this study (Figure 1). Our studied sample included 502 males and 533 females 40-79 years of age (median age: 52). Most subjects were 50-59 years of age (39.22%); few were 70-79 years of age (4.35%). Statistical data are shown in Table 1. A total of 87 subjects (87/1035, 8.41%) were positive on mSDC2 testing, including 52 males (52/502, 10.36%) and 35 females (35/533, 6.57%); data significantly different among the two groups ( $\chi^2 = 4.828$ , *P* = 0.028). Furthermore, rates of positivity for different age groups (*i.e.*, 40-49, 50-59, 60-69, 70-79) were 4.02% (15/373), 9.61% (39/406), 11.85% (25/211), and 17.78% (8/45), respectively. Rates of positivity on mSDC2 testing increased with age; rates of positivity significantly differed among age groups ( $\chi^2 = 18.454$ , *P* < 0.001).

## General outcomes of screening

As shown in Figure 1 and Table 1, a total of 420 patients (420/1035, 40.58%) with intestinal lesions of different severity and 615 others (*e.g.*, normal findings on colonoscopy, colitis, colonic diverticulum, *etc.*; 615/1035, 59.42%) were evaluated on colonoscopy. Among all intestinal lesions, 16 cases of CRC diverticulum (16/1035, 1.55%), 65 cases of AA (65/1035, 6.28%), 189 cases of non-AA (189/1035, 18.26%), and 150 cases of polyps (150/1035, 14.49%) were diagnosed. All lesions were verified on histopathology. The DR of intestinal lesions in males was significantly higher than that in females (47.41% *vs* 34.15%,  $\chi^2 = 18.862$ , *P* < 0.001; Table 2). DR markedly increased with age ( $\chi^2 = 55.920$ , *P* < 0.001) with the highest rate of positivity among individuals 70-79 years of age (28/45, 62.22%). Pathological CRC characteristics of lesions detected in this study are detailed in Table 3.

Importantly, mSDC2 test positivity, sex, age, a family history of CRC and a history of polyps were all significant risk factors for the development of ACN (*i.e.*, CRC and AA). Otherwise, chronic diarrhea, constipation, appendicitis and cholecystitis, as well as hematochezia and a history of psychiatric trauma, were not significant risk factors for CRC (Table 4).

## Efficacy of the mSDC2 test for the screening of colorectal neoplasms

To assess the performance of the mSDC2 test in identifying intestinal malignancies, an ROC curve was constructed (Figure 2). The value of the area under the ROC curve (AUC) in regards to detecting SDC2 gene methylation in CRC cases was 0.914 (95%CI: 0.807-1.000), highlighting the capacity of this test to accurately distinguish CRC lesions from non-advanced or benign tissue (Figure 2A). For AAs, the AUC value decreased to 0.707 (95%CI: 0.631-0.782), indicating relatively standard test efficacy in detection of precancerous lesions (Figure 2B). For ACN, the AUC value was 0.756 (95%CI: 0.689-0.823; Figure 2C). Furthermore, compared with fecal occult blood testing, mSDC2 testing exhibited significantly improved diagnostic sensitivity, specificity and AUC values for CRC, AA and ACN, suggesting that mSDC2 is effective for use in the early diagnosis of malignancy. Although the price of mSDC2 testing far exceeds that of fecal

Table 1 Baseline data of subjects evaluated

Characteristics	Cases (N%)	mSDC2 (N%)		P value
		(+) n = 87	(-) n = 948	
Gender				0.028 <sup>a</sup>
Male	502 (48.50)	52 (5.02)	450 (43.48)	
Female	533 (51.50)	35 (3.38)	498 (48.12)	
Age				< 0.001 <sup>b</sup>
40-49	373 (36.04)	15 (1.45)	358 (34.59)	
50-59	406 (39.22)	39 (3.77)	367 (35.46)	
60-69	211 (20.39)	25 (2.42)	186 (17.97)	
70-79	45 (4.35)	8 (0.77)	37 (3.57)	
Family history of CRC				< 0.001 <sup>b</sup>
Yes	99 (9.57)	18 (1.74)	81 (7.83)	
No	936 (90.43)	69 (6.67)	867 (83.77)	
History of colorectal polyps				0.406
Yes	252 (24.35)	18 (1.74)	234 (22.61)	
No	783 (75.65)	69 (6.67)	714 (68.99)	
History of chronic diarrhea				0.108
Yes	342 (33.04)	22 (2.13)	320 (30.92)	
No	693 (66.96)	65 (6.28)	628 (60.68)	
History of chronic constipation				0.747
Yes	493 (47.63)	40 (3.86)	453 (43.77)	
No	542 (52.37)	47 (4.54)	495 (47.83)	
History of hematochezia				0.692
Yes	580 (56.04)	47 (4.54)	533 (51.50)	
No	455 (43.96)	40 (3.86)	415 (40.10)	
History of appendicitis				0.243
Yes	66 (6.38)	3 (0.29)	63 (6.09)	
No	969 (93.62)	84 (8.12)	885 (85.51)	
History of cholecystitis				0.836
Yes	66 (6.38)	6 (0.58)	60 (5.80)	
No	969 (93.62)	81 (7.83)	888 (85.80)	
History of psychiatric trauma				0.890
Yes	63 (6.09)	5 (0.48)	58 (5.60)	
No	972 (93.91)	82 (7.92)	890 (85.99)	
Pathological classification				< 0.001 <sup>b</sup>
CRC	16 (1.55)	14 (1.35)	2 (0.19)	
AA	65 (6.28)	26 (2.51)	39 (3.77)	
NAA	189 (18.26)	8 (0.77)	181 (17.49)	
Polyps	150 (14.49)	12 (1.16)	138 (13.33)	
Others	615 (59.42)	27 (2.61)	588 (56.81)	

<sup>a</sup>P < 0.05.

<sup>b</sup>*P* < 0.001.


mSDC: Syndecan-2 methylation; CRC: Colorectal cancer; AA: Advanced adenoma; NAA: Non-advanced adenoma; Polyps: Non-adenomatous polyps; Others: Totally normal colonoscopy, colitis and colonic diverticulum, *etc.*

**Table 2 Primary outcomes of pathological findings on screening**

Characteristics	Intestinal lesions (N%)				Others (N%)	P value
	CRC	AA	NAA	Polyps		
<b>Gender</b>						< 0.001 <sup>a</sup>
Male	9 (0.87)	43 (4.15)	109 (10.53)	77 (7.44)	264 (25.51)	
Female	7 (0.68)	22 (2.13)	80 (7.73)	73 (7.05)	351 (33.91)	
<b>Age (yr)</b>						< 0.001 <sup>a</sup>
40-49	3 (0.29)	8 (0.77)	48 (4.64)	46 (4.44)	268 (25.89)	
50-59	5 (0.48)	30 (2.90)	73 (7.05)	59 (5.70)	239 (23.09)	
60-69	7 (0.68)	18 (1.74)	57 (5.51)	38 (3.67)	91 (8.79)	
70-79	1 (0.10)	9 (0.87)	11 (1.06)	7 (0.68)	17 (1.64)	
<b>mSDC2</b>						< 0.001 <sup>a</sup>
Positive	14 (1.35)	26 (2.51)	8 (0.77)	12 (1.16)	27 (2.61)	
Negative	2 (0.19)	39 (3.77)	181 (17.49)	138 (13.33)	588 (56.81)	

<sup>a</sup>*P* < 0.001.

mSDC: Syndecan-2 methylation; CRC: Colorectal cancer; AA: Advanced adenoma; NAA: Non-advanced adenoma; Polyps: Non-adenomatous polyps; Others: Totally normal colonoscopy, colitis and colonic diverticulum, *etc.*



**Figure 1 Screening process flow chart.** mSDC: Syndecan-2 methylation; CRC: Colorectal cancer; AA: Advanced adenoma; NAA: Non-advanced adenoma; Polyps: Non-adenomatous polyps; Others: Totally normal colonoscopy, colitis and colonic diverticulum, *etc.*

occult blood testing, early detection of CRC has the potential to save lives and direct more effective clinical intervention. As such, the long-term benefits of mSDC2 testing warrant its use in clinic.

In this study, as shown in Tables 1, 2 and 5, the mSDC2 test was able to identify 14 out of 16 CRC cases with a sensitivity of 87.50% (95%CI: 60.41-97.80). For AA cases, sensitivity was 40.00% (95%CI: 28.28-52.90). The specificity of the mSDC2 test for detecting other conditions was 95.61% (95%CI: 93.59-97.03) among 615 individuals. For 954 subjects who were diagnosed with conditions other than AA or CRC, mSDC2 test specificity was 95.07% (95%CI: 93.45-96.32).

**Table 3 Characteristics of colorectal cancer detected on screening**

Characteristics	Colonoscopy (N = 16)	mSDC2 (N%)	
		Positive (n = 14)	Negative (n = 2)
Gender			
Male	9 (56.25)	8 (50.00)	1 (6.25)
Female	7 (43.75)	6 (37.50)	1 (6.25)
Age (yr)			
40-49	3 (18.75)	2 (12.50)	1 (6.25)
50-59	5 (31.25)	5 (31.25)	0 (0)
60-69	7 (43.75)	7 (43.75)	0 (0)
70-79	1 (6.25)	0 (0)	1 (6.25)
TNM stage			
0/I/II	8 (50.00)	6 (43.7)	2 (12.50)
III/IV	5 (31.25)	5 (31.25)	0 (0)
Unknown	3 (18.75)	3 (18.75)	0 (0)
Tumor location			
Proximal	3 (18.75)	3 (18.75)	0 (0)
Distal	13 (81.25)	11 (68.75)	2 (12.50)
Tumor size (mm)			
≤ 30	9 (56.25)	7 (43.75)	2 (12.50)
> 30	7 (43.75)	7 (43.75)	0 (0)
Dysplasia			
Low	0 (0)	0 (0)	0 (0)
Median	12 (75.00)	12 (75.00)	0 (0)
High	4 (25.00)	2 (12.50)	2 (12.50)

mSDC: Syndecan-2 methylation.

The PPVs of the mSDC2 test for CRC, AA and ACN were 16.09% (14/87) (95%CI: 9.38-25.87), 29.89% (26/87) (95%CI: 20.78-40.79) and 45.98% (40/87) (95%CI: 35.36-56.96), respectively. The NPVs of this test for CRC, AA, and ACN were 99.79% (946/948) (95%CI: 99.15-99.96), 95.89% (909/948) (95%CI: 94.37-97.02) and 95.68% (907/948) (95%CI: 94.13-96.84), respectively.

## DISCUSSION

The risk factors questionnaire is a cost-effective and easily administered tool, but its high rate of false positivity and subjectivity have resulted in low compliance with colonoscopy among patients it identifies as indicated to undergo the procedure. Consequently, the effectiveness of large-scale population screening remains effectively compromised. To address such limitations, this study evaluated the efficacy, specificity and sensitivity of mSDC2 testing for detection of intestinal malignancies. Our findings indicate that mSDC2 testing is a valuable complementary approach for large-scale population screening. Furthermore, mSDC2 testing can enhance patient compliance with colonoscopy, more precisely identify patients indicated to undergo colonoscopy, conserve medical resources, and improve screening efficiency.

Importantly, mSDC2 testing can also miss diagnoses. Based on our study as well as previous literature, the rate of missed diagnosis for CRC was estimated to be 10%-15%. Missed diagnoses can occur for several reasons. Firstly, as this is a self-sampling product, test accuracy may be influenced by factors such as adherence to proper sampling technique by subjects and timely sample transportation to the laboratory. Secondly, while mSDC2 is a reliable marker for CRC detection, there may be a small number of patients who do not exhibit abnormal mSDC2. To minimize missed diagnoses, it is beneficial to combine mSDC2 testing with other methods such as questionnaires and fecal occult blood tests to enhance overall disease detection. Additionally, regular and repeated testing can also effectively reduce missed diagnoses.

**Table 4 Risk factors for detection of advanced colorectal neoplasia**

Characteristics	B	SE	Wald	df	P value	OR	95%CI	
							Lower limit	Upper limit
Gender								
Male	-0.768	0.294	6.816	1	0.009 <sup>b</sup>	0.464	0.261	0.826
Age								
40-49			22.792	3	< 0.001 <sup>c</sup>			
50-59	-2.627	0.567	21.459	1	< 0.001 <sup>c</sup>	0.072	0.024	0.220
60-69	-1.612	0.499	10.422	1	0.001 <sup>c</sup>	0.200	0.075	0.531
70-79	-1.266	0.516	6.010	1	0.014 <sup>a</sup>	0.282	0.102	0.776
mSDC2								
Positive	-2.798	0.320	76.308	1	< 0.001 <sup>c</sup>	0.061	0.033	0.114
Family history of CRC								
Yes	-1.674	0.530	9.958	1	0.002 <sup>b</sup>	0.188	0.066	0.530
History of colorectal polyps								
Yes	2.055	0.580	12.560	1	< 0.001 <sup>c</sup>	7.809	2.506	24.335
History of chronic diarrhea								
Yes	0.077	0.405	0.036	1	0.850	1.080	0.488	2.388
History of chronic constipation								
Yes	0.417	0.399	1.090	1	0.296	1.518	0.694	3.320
History of hematochezia								
Yes	0.322	0.442	0.529	1	0.467	1.380	0.580	3.284
History of appendicitis								
Yes	0.860	0.818	1.105	1	0.293	2.362	0.476	11.729
History of cholecystitis								
Yes	-0.306	0.555	0.305	1	0.581	0.736	0.248	2.183
History of psychiatric trauma								
Yes	1.287	0.912	1.991	1	0.158	3.621	0.606	21.633

<sup>a</sup>*P* < 0.05.<sup>b</sup>*P* < 0.01.<sup>c</sup>*P* < 0.001.

CI: Confidence interval; B: Beta; SE: Standard error; df: Degree of freedom; OR: Odds ratio; CRC: Colorectal cancer; mSDC: Syndecan-2 methylation.

Despite the excellent efficacy of mSDC2 testing, its price currently remains high, thus limiting its accessibility and popularity among the general public. Many individuals are unaware of this product or even lack knowledge concerning the importance of CRC screening in general. As such, there is an urgent need to raise awareness among the general patient population. Continuous product optimization is also essential, focusing on enhancing sampling success and simplifying laboratory testing. Such improvements are bound to contribute towards expanding mSDC2 utilization worldwide.

Usually, CRC develops from adenomas and certain genes relevant for CRC pathogenesis undergo alterations in methylation levels during the adenomatous stage. Most gene promoters associated with CRC contain CpG islands; abnormal methylation of cytosine within these sequences can result in gene inactivation. As disease progresses, gene methylation increases and the number of malignant cells with methylation mutations increases. One such relevant gene is *SDC2*, which encodes a transmembrane proteoglycan that influences CRC cell proliferation, migration and invasive capabilities. Methylation of *SDC2* leads to the silencing of DNA transcription, disrupting cell growth and differentiation, promoting tumor cell proliferation and enhancing invasive and metastatic characteristics. Interestingly, levels of mSDC2 in CRC are higher as compared to AA tissue.


Recently, enteroscopy was reported to not be a suitable method for large-scale population screening[34]. Considering China's large population as well as an uneven distribution of medical technology across different geographic regions, comprehensive colorectal screening poses significant challenges. As such, optimization of CRC screening methods and



**Table 5 Performance characteristics of the syndecan-2 methylation test regarding detection of colorectal neoplasia**

Category	Colonoscopy (N = 1035)	mSDC2 test (N = 1035)		PPV % (95%CI)	NPV % (95%CI)
		Positive (n = 87)	Sensitivity % (95%CI)		
Colorectal cancer	16	14	87.50 (60.41-97.80)	16.09 (9.38-25.87)	99.79 (99.15-99.96)
Advanced adenoma	65	26	40.00 (28.28-52.90)	29.89 (20.78-40.79)	95.89 (94.37-97.02)
Advanced colorectal neoplasia	81	40	49.38 (38.19-60.64)	45.98 (35.36-56.96)	95.68 (94.13-96.84)
		Negative (n = 948)	Specificity % (95%CI)		
Others	615	588	95.61 (93.59-97.03)		
Others, polyps and NAA	954	907	95.07 (93.45-96.32)		
Normal	552	527	95.47 (93.29-96.99)		

mSDC: Syndecan-2 methylation; CI: Confidence interval; PPV: Positive predictive value; NPV: Negative predictive value; Others: Totally normal colonoscopy, colitis and colonic diverticulum, *etc.*; Polyps: Non-adenomatous polyps; NAA: Non-advanced adenoma; Normal: Totally normal colonoscopy.



**Figure 2 Receiver operating characteristic curve of fecal occult-blood and syndecan-2 methylation tests for detection of colorectal cancer, advanced adenoma and advanced colorectal neoplasia.** A: Colorectal cancer; B: Advanced adenoma; C: Advanced colorectal neoplasia. SCD2: Syndecan-2; FOBT: Fecal occult-blood; AUC: Area under the receiver operating characteristic curve.

efficiency is urgently warranted.

Here, we found that men and older individuals were more likely to possess intestinal lesions as compared to women and younger individuals, respectively. Based on known data, most specialists recommend that individuals over the age of 40 undergo screening for CRC. If no precancerous lesions such as polyps are detected during initial screening, subsequent screening should be performed five years later.



It is important to note that the sample used in this study was primarily native to Guangzhou, China, which naturally imposes geographic and ethnic limitations. In the future, large-scale studies that clinically validate mSDC2 while considering possible ethnic and geographic differences within a studied population will be necessary. Finally, combinations of stool mSDC2 methylation analysis with other tumor markers, such as combined mSDC2 and serum CEA detection[35], stool mSDC2 and *TFPI2* methylation detection[36], or stool mSDC2 and *NDRG4* methylation detection[37], significantly enhance both sensitivity and diagnostic accuracy of CRC. Further investigation relevant to this topic is urgently warranted to develop novel simple, accurate and efficacious CRC screening methods.

## CONCLUSION

In summary, stool mSDC2 testing carries a sensitivity of 87.50% for the detection of CRC among high-risk populations, underscoring its clinical value in early cancer screening. Importantly, mSDC2 serves as a reliable biomarker for CRC diagnosis. Use of stool samples for analyses offers a simpler and less invasive approach, minimizing patient discomfort and promoting patient compliance.

Here, we solely focused on evaluating SDC2 from the perspective of a diagnostic biomarker; as such, our study was not without its limitations. Multi-target fecal DNA testing, which includes markers such as *WIF*, *NAP*, *PENK*, *SETP9*, is described in literature as relevant to the diagnosis of CRC. Use of multiple fecal DNA biomarkers in combination was reported to enhance CRC DR, better identify early intestinal lesions and enable patients to receive treatment earlier, thereby reducing the case fatality rate. Thus, future research should consider incorporating multi-target fecal DNA testing in analyses to improve CRC screening accuracy and efficiency.

## ARTICLE HIGHLIGHTS

### Research background

Colorectal cancer (CRC) is a prevalent and life-threatening malignant tumor affecting the digestive system globally. Testing for syndecan-2 methylation (mSDC2) has emerged as a widely used screening tool for early detection of CRC in stool and serum samples.

### Research motivation

Our findings provide evidence-based data concerning diagnostic and screening methods relevant to a Chinese population at high-risk population for CRC.

### Research objectives

To validate the effectiveness of fecal mSDC2 testing in the detection of CRC among a high-risk Chinese population.

### Research methods

A high-risk Chinese cohort composed of 1130 individuals 40-79 years of age was selected for evaluation using the fecal mSDC2 test. Sensitivity and specificity to CRC, advanced adenoma (AA) and advanced colorectal neoplasia (ACN) were quantified. High-risk factors for the incidence of colorectal lesions were analyzed; a logistic regression model was subsequently constructed to better reflect the efficacy of fecal mSDC2 testing.

### Research results

According to criteria previously established, 1035 high-risk individuals were included in analyses. Among them, 16 CRC cases (1.55%), 65 AA cases (6.28%), 189 non-advanced adenoma cases (18.26%), and 150 cases of polyps (14.49%) were successfully identified on colonoscopy and pathological examination. The sensitivities of mSDC2 testing for CRC and AA were 87.50% and 40.00%, respectively; the specificity for subjects in the “others” group was 95.61%. The positive predictive values of mSDC2 testing for CRC, AA, and ACN were 16.09%, 29.89% and 45.98%, respectively. In addition, negative predictive value of mSDC2 testing for CRC was 99.79%. Positivity on mSDC2 testing is a significant risk factor for the development of ACN ( $P < 0.001$ ) after adjusting for other high-risk covariates.

### Research conclusions

The results of this CRC screening study revealed that offering patients a combination of fecal mSDC2 testing and colonoscopy is ideal for facilitating early detection of CRC among a high-risk Chinese population. CRC screening study revealed that offering patients a combination of fecal mSDC2 testing and colonoscopy is ideal for facilitating early detection of CRC among a high-risk Chinese population.

### Research perspectives

Detection of stool mSDC2 offers great promise for early and effective CRC screening.

## FOOTNOTES

**Co-first authors:** Wen-Feng Luo and Yu-Ting Jiao.

**Co-corresponding authors:** Wen-Feng Luo and Yu Wan.

**Author contributions:** Luo WF and Jiao YT contributed equally to this work; Luo WF and Wan Y were co-corresponding authors; Luo WF, Jiao YT, and Wan Y contributed to conceptualization, visualization, initial draft writing, funding acquisition, review and editing; Luo WF, Jiao YT, Wang SB, and Deng J proposed and designed the study; Lin XL, Shen J, and Xie FM performed analyses; Zhao Y, Han ZP, and Han ZP wrote the initial draft; Jiao YT, Ye YF, He JH, and Wan Y revised the manuscript; and all authors read and approved the final manuscript.

**Supported by** the Science and Technology Program of Panyu Central Hospital, No. PY-2023-003; the Science and Technology Program of Panyu, No. 2020-Z04-054; the Science and Technology Project of the Guangzhou Health Commission, No. 20211A011114; the Science and Technology Program of Guangzhou, No. 202002020023; the General University Youth Innovative Talent Project of Guangdong Province, No. 2022KQNCX281; the Guangdong Provincial Key Field Special Project for Ordinary Colleges and Universities, No. 2023ZDZX2097; and the Foshan Engineering Technology Research Center for Prepared Food Processing and Quality Evaluation, No. 2022-KJZX113.

**Institutional review board statement:** The study was reviewed and approved by the Ethics Committee of Panyu Central Hospital in China (Approval No. [2019]62).

**Informed consent statement:** The informed consent was obtained from all study participants.

**Conflict-of-interest statement:** All the authors report no relevant conflicts of interest for this article.

**Data sharing statement:** No additional data are available.

**STROBE statement:** The authors have read the STROBE Statement-checklist of items, and the manuscript was prepared and revised according to the STROBE Statement-checklist of items.

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** China

**ORCID number:** Wen-Feng Luo 0000-0003-0922-1005; Yu-Feng Ye 0000-0002-9843-475X; Jin-Hua He 0000-0001-8506-2419; Yu Wan 0009-0001-1813-773X.

**S-Editor:** Wang JJ

**L-Editor:** A

**P-Editor:** Zhang XD

## REFERENCES

- 1 Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, Bray F. Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA Cancer J Clin* 2021; **71**: 209-249 [PMID: 33538338 DOI: 10.3322/caac.21660]
- 2 Wu CW, Lui RN. Early-onset colorectal cancer: Current insights and future directions. *World J Gastrointest Oncol* 2022; **14**: 230-241 [PMID: 35116113 DOI: 10.4251/wjgo.v14.i1.230]
- 3 Saad El Din K, Loree JM, Sayre EC, Gill S, Brown CJ, Dau H, De Vera MA. Trends in the epidemiology of young-onset colorectal cancer: a worldwide systematic review. *BMC Cancer* 2020; **20**: 288 [PMID: 32252672 DOI: 10.1186/s12885-020-06766-9]
- 4 Siegel RL, Medhanie GA, Fedewa SA, Jemal A. State Variation in Early-Onset Colorectal Cancer in the United States, 1995-2015. *J Natl Cancer Inst* 2019; **111**: 1104-1106 [PMID: 31141602 DOI: 10.1093/jnci/djz098]
- 5 Hossain MS, Karuniawati H, Jairoun AA, Urbi Z, Ooi J, John A, Lim YC, Kibria KMK, Mohiuddin AKM, Ming LC, Goh KW, Hadi MA. Colorectal Cancer: A Review of Carcinogenesis, Global Epidemiology, Current Challenges, Risk Factors, Preventive and Treatment Strategies. *Cancers (Basel)* 2022; **14** [PMID: 35406504 DOI: 10.3390/cancers14071732]
- 6 Liao C, Chen X, Fu Y, Salivary. analysis: An emerging paradigm for non-invasive healthcare diagnosis and monitoring. *Interdiscip Med* 2023 [DOI: 10.1002/INMD.20230009]
- 7 Ladabaum U, Dominitz JA, Kahi C, Schoen RE. Strategies for Colorectal Cancer Screening. *Gastroenterology* 2020; **158**: 418-432 [PMID: 31394083 DOI: 10.1053/j.gastro.2019.06.043]
- 8 Imperiale TF, Gruber RN, Stump TE, Emmett TW, Monahan PO. Performance Characteristics of Fecal Immunochemical Tests for Colorectal Cancer and Advanced Adenomatous Polyps: A Systematic Review and Meta-analysis. *Ann Intern Med* 2019; **170**: 319-329 [PMID: 30802902 DOI: 10.7326/M18-2390]
- 9 Hata T, Chiba K, Mizuda M, Masuda K, Ohtsuka H, Nakagawa K, Morikawa T, Hayashi H, Motoi F, Unno M. Levels of tumor markers CEA/CA 19-9 in serum and peritoneal lavage predict postoperative recurrence in patients with pancreatic cancer. *Ann Gastroenterol Surg*

- 2022; **6**: 862-872 [PMID: [36338582](#) DOI: [10.1002/ags3.12597](#)]
- 10 **Kato H**, Kishiwada M, Hayasaki A, Chipaila J, Maeda K, Noguchi D, Gyoten K, Fujii T, Iizawa Y, Tanemura A, Murata Y, Kuriyama N, Usui M, Sakurai H, Isaji S, Mizuno S. Role of Serum Carcinoembryonic Antigen (CEA) Level in Localized Pancreatic Adenocarcinoma: CEA Level Before Operation is a Significant Prognostic Indicator in Patients With Locally Advanced Pancreatic Cancer Treated With Neoadjuvant Therapy Followed by Surgical Resection: A Retrospective Analysis. *Ann Surg* 2022; **275**: e698-e707 [PMID: [32744820](#) DOI: [10.1097/SLA.0000000000004148](#)]
- 11 **Chen R**, Jiang C, Zhu Q, You S, Li Y, Li S, Ding L, Meng H, Yang Y, Zha X, Wang J. Combining the tumor abnormal protein test with tests for carcinoembryonic antigens, cancer antigen 15-3, and/or cancer antigen 125 significantly increased their diagnostic sensitivity for breast cancer. *Medicine (Baltimore)* 2020; **99**: e21231 [PMID: [32702897](#) DOI: [10.1097/MD.00000000000021231](#)]
- 12 **Lakemeyer L**, Sander S, Wittau M, Henne-Bruns D, Kornmann M, Lemke J. Diagnostic and Prognostic Value of CEA and CA19-9 in Colorectal Cancer. *Diseases* 2021; **9** [PMID: [33802962](#) DOI: [10.3390/diseases9010021](#)]
- 13 **Ryan L**, Wong Y, Dwyer KM, Clarke D, Kyprian L, Craig JM. Coprocology: A Technical Review of Cytological Colorectal Cancer Screening in Fecal Samples. *SLAS Technol* 2021; **26**: 591-604 [PMID: [34219541](#) DOI: [10.1177/24726303211024562](#)]
- 14 **Zhang W**, Yang C, Wang S, Xiang Z, Dou R, Lin Z, Zheng J, Xiong B. SDC2 and TFPI2 Methylation in Stool Samples as an Integrated Biomarker for Early Detection of Colorectal Cancer. *Cancer Manag Res* 2021; **13**: 3601-3617 [PMID: [33958894](#) DOI: [10.2147/CMAR.S300861](#)]
- 15 **Ansa BE**, Lewis N, Hoffman Z, Datta B, Johnson JA. Evaluation of Blood Stool Test Utilization for Colorectal Cancer Screening in Georgia, USA. *Healthcare (Basel)* 2021; **9** [PMID: [34065816](#) DOI: [10.3390/healthcare9050569](#)]
- 16 **Grady WM**, Yu M, Markowitz SD. Epigenetic Alterations in the Gastrointestinal Tract: Current and Emerging Use for Biomarkers of Cancer. *Gastroenterology* 2021; **160**: 690-709 [PMID: [33279516](#) DOI: [10.1053/j.gastro.2020.09.058](#)]
- 17 **Joo JE**, Clendenning M, Wong EM, Rosty C, Mahmood K, Georgeson P, Winship IM, Preston SG, Win AK, Dugué PA, Jayasekara H, English D, Macrae FA, Hopper JL, Jenkins MA, Milne RL, Giles GG, Southey MC, Buchanan DD. DNA Methylation Signatures and the Contribution of Age-Associated Methylomic Drift to Carcinogenesis in Early-Onset Colorectal Cancer. *Cancers (Basel)* 2021; **13** [PMID: [34070516](#) DOI: [10.3390/cancers13112589](#)]
- 18 **Hong J**, Rhee JK. Genomic Effect of DNA Methylation on Gene Expression in Colorectal Cancer. *Biology (Basel)* 2022; **11** [PMID: [36290295](#) DOI: [10.3390/biology11101388](#)]
- 19 **Koch A**, Joosten SC, Feng Z, de Ruijter TC, Draht MX, Melotte V, Smits KM, Veeck J, Herman JG, Van Neste L, Van Criekinge W, De Meyer T, van Engeland M. Analysis of DNA methylation in cancer: location revisited. *Nat Rev Clin Oncol* 2018; **15**: 459-466 [PMID: [29666440](#) DOI: [10.1038/s41571-018-0004-4](#)]
- 20 **Chen J**, Sun H, Tang W, Zhou L, Xie X, Qu Z, Chen M, Wang S, Yang T, Dai Y, Wang Y, Gao T, Zhou Q, Song Z, Liao M, Liu W. DNA methylation biomarkers in stool for early screening of colorectal cancer. *J Cancer* 2019; **10**: 5264-5271 [PMID: [31602277](#) DOI: [10.7150/jca.34944](#)]
- 21 **Imperiale TF**, Ransohoff DF, Itzkowitz SH, Levin TR, Lavin P, Lidgard GP, Ahlquist DA, Berger BM. Multitarget stool DNA testing for colorectal-cancer screening. *N Engl J Med* 2014; **370**: 1287-1297 [PMID: [24645800](#) DOI: [10.1056/NEJMoa1311194](#)]
- 22 **US Preventive Services Task Force**, Bibbins-Domingo K, Grossman DC, Curry SJ, Davidson KW, Epling JW Jr, García FAR, Gillman MW, Harper DM, Kemper AR, Krist AH, Kurth AE, Landefeld CS, Mangione CM, Owens DK, Phillips WR, Phipps MG, Pignone MP, Siu AL. Screening for Colorectal Cancer: US Preventive Services Task Force Recommendation Statement. *JAMA* 2016; **315**: 2564-2575 [PMID: [27304597](#) DOI: [10.1001/jama.2016.5989](#)]
- 23 **Yue C**, Zhang Y, Wang Y, Zhang Z, Zhang M, Wang H, Chen W, Shang Z, Xin Y, Zhang X. The Application Value of Syndecan-2 Gene Methylation for Colorectal Cancer Diagnosis: A Clinical Study and Meta-Analyses. *Front Med (Lausanne)* 2022; **9**: 753545 [PMID: [35372441](#) DOI: [10.3389/fmed.2022.753545](#)]
- 24 **Wang L**, Liu Y, Zhang D, Xiong X, Hao T, Zhong L, Zhao Y. Diagnostic accuracy of DNA-based SDC2 methylation test in colorectal cancer screening: a meta-analysis. *BMC Gastroenterol* 2022; **22**: 314 [PMID: [35754025](#) DOI: [10.1186/s12876-022-02395-7](#)]
- 25 **Zhang L**, Dong L, Lu C, Huang W, Yang C, Wang Q, Lei R, Sun R, Wan K, Li T, Sun F, Gan T, Lin J, Yin L. Methylation of SDC2/TFPI2 and Its Diagnostic Value in Colorectal Tumorous Lesions. *Front Mol Biosci* 2021; **8**: 706754 [PMID: [35004840](#) DOI: [10.3389/fmolb.2021.706754](#)]
- 26 **Zhao G**, Li H, Yang Z, Wang Z, Xu M, Xiong S, Li S, Wu X, Liu X, Zhu Y, Ma Y, Fei S, Zheng M. Multiplex methylated DNA testing in plasma with high sensitivity and specificity for colorectal cancer screening. *Cancer Med* 2019; **8**: 5619-5628 [PMID: [31407497](#) DOI: [10.1002/cam4.2475](#)]
- 27 **Kim JH**, Park SC. Syndecan-2 Methylation as a New Biomarker for Early Detection of Colorectal Neoplasm. *Gut Liver* 2018; **12**: 479-480 [PMID: [30205668](#) DOI: [10.5009/gnl18286](#)]
- 28 **Ryu HY**, Lee J, Yang S, Park H, Choi S, Jung KC, Lee ST, Seong JK, Han IO, Oh ES. Syndecan-2 functions as a docking receptor for pro-matrix metalloproteinase-7 in human colon cancer cells. *J Biol Chem* 2009; **284**: 35692-35701 [PMID: [19858218](#) DOI: [10.1074/jbc.M109.054254](#)]
- 29 **Meng W**, Cai SR, Zhou L, Dong Q, Zheng S, Zhang SZ. Performance value of high risk factors in colorectal cancer screening in China. *World J Gastroenterol* 2009; **15**: 6111-6116 [PMID: [20027686](#) DOI: [10.3748/wjg.15.6111](#)]
- 30 **Lin G**, Feng Z, Liu H, Li Y, Nie Y, Liang Y, Li K. Mass screening for colorectal cancer in a population of two million older adults in Guangzhou, China. *Sci Rep* 2019; **9**: 10424 [PMID: [31320661](#) DOI: [10.1038/s41598-019-46670-2](#)]
- 31 **The Chinese Society of Digestive Endoscopy**, Tumor Endoscopy Committee of China Anti-Cancer Association. [China Guideline for the Early Colorectal Cancer Screening and Treatment by Endoscopy (2014, Beijing)]. *Chinese Journal of Digestive Endoscopy* 2015; **32**: 341-360 [DOI: [10.3760/cma.j.issn.1007-5232.2015.06.001](#)]
- 32 **Wang J**, Liu S, Wang H, Zheng L, Zhou C, Li G, Huang R, Li C, Fan X, Fu X, Wang X, Guo H, Guan J, Sun Y, Song X, Li Z, Mu D, Sun J, Liu X, Qi Y, Niu F, Chen C, Wu X, Zou H. Robust performance of a novel stool DNA test of methylated SDC2 for colorectal cancer detection: a multicenter clinical study. *Clin Epigenetics* 2020; **12**: 162 [PMID: [33126908](#) DOI: [10.1186/s13148-020-00954-x](#)]
- 33 **Amin MB**, Gress DM, Meyer Vega LR, Edge SB, Greene FL, Byrd DR, Brookland RK, Washington MK, Compton CC. *AJCC Cancer Staging Manual* 8th edition. New York: Springer, 2017: 252-274
- 34 **Bretthauer M**, Löberg M, Wieszczny P, Kalager M, Emilsson L, Garborg K, Rupinski M, Dekker E, Spaander M, Bugajski M, Holme Ø, Zauber AG, Pilonis ND, Mroz A, Kuipers EJ, Shi J, Hernán MA, Adami HO, Regula J, Hoff G, Kaminski MF; NordICC Study Group. Effect of Colonoscopy Screening on Risks of Colorectal Cancer and Related Death. *N Engl J Med* 2022; **387**: 1547-1556 [PMID: [36214590](#) DOI: [10.1056/NEJMoa2110000](#)]

10.1056/NEJMoa2208375]

- 35 **Li J**, Sun Z. [Application of noninvasive feces SDC2 methylation combined with serum CEA in the early diagnosis of colorectal cancer and the value assessment]. *Jilin Med J* 2021; **42**: 1582-1584 [DOI: [10.3969/j.issn.1004-0412.2021.07.014](https://doi.org/10.3969/j.issn.1004-0412.2021.07.014)]
- 36 **Wu H**, Fang H, Deng Y, He B. [Analysis on diagnostic efficacy of combined detection of fecal genes SDC2 andTFPI2 methylation in diagnosing colorectal cancer and precancerous lesions]. *Modern Med J China* 2023; **25**: 7-12
- 37 **Jin W**, Zhang X, Liu L, Zhou Q, Miao Y, Tan X, Tan L. [Application of SDC2 and NDRG4 gene methylation detection in colorectal cancer screening]. *Journal of Hunan Normal University (Medical Science)* 2021; **18**: 88-92 [DOI: [10.3969/j.issn.1673-016X.2021.06.026](https://doi.org/10.3969/j.issn.1673-016X.2021.06.026)]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

**Telephone:** +1-925-3991568

**E-mail:** [office@baishideng.com](mailto:office@baishideng.com)

**Help Desk:** <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

