World Journal of *Gastrointestinal Oncology*

World J Gastrointest Oncol 2024 April 15; 16(4): 1091-1675





Published by Baishideng Publishing Group Inc

World Journal of Gastrointestinal Oncology

Contents

Monthly Volume 16 Number 4 April 15, 2024

EDITORIAL

1091	Parallel pathways: A chronicle of evolution in rectal and breast cancer surgery
1097	Hepatitis B virus genotypes in precision medicine of hepatitis B-related hepatocellular carcinoma: Where we are now Sukowati CHC, Jayanti S, Turyadi T, Muljono DH, Tiribelli C
	REVIEW
1104	Novel milestones for early esophageal carcinoma: From bench to bed
	Qi JH, Huang SL, Jin SZ
1119	Colorectal cancer screening: A review of current knowledge and progress in research
	Lopes SR, Martins C, Santos IC, Teixeira M, Gamito É, Alves AL
1134	New avenues for the treatment of immunotherapy-resistant pancreatic cancer
-	Silva LGO, Lemos FFB, Luz MS, Rocha Pinheiro SL, Calmon MDS, Correa Santos GL, Rocha GR, de Melo FF
	MINIREVIEWS
1154	Present situation of minimally invasive surgical treatment for early gastric cancer
	Li CY, Wang YF, Luo LK, Yang XJ
1166	Mixed neuroendocrine non-neuroendocrine neoplasms in gastroenteropancreatic tract
	Díaz-López S, Jiménez-Castro J, Robles-Barraza CE, Ayala-de Miguel C, Chaves-Conde M
1180	Esophageal cancer screening, early detection and treatment: Current insights and future directions
1100	Qu HT, Li Q, Hao L, Ni YJ, Luan WY, Yang Z, Chen XD, Zhang TT, Miao YD, Zhang F
	ORIGINAL ARTICLE
	Retrospective Cohort Study
1192	Pre-operative enhanced magnetic resonance imaging combined with clinical features predict early recurrence of hepatocellular carcinoma after radical resection
	Chen JP, Yang RH, Zhang TH, Liao LA, Guan YT, Dai HY

Zhu CL, Peng LZ

center



Contents

World Journal of Gastrointestinal Oncology

Monthly Volume 16 Number 4 April 15, 2024

Retrospective Study

1213 Predictive model for non-malignant portal vein thrombosis associated with cirrhosis based on inflammatory biomarkers

Nie GL, Yan J, Li Y, Zhang HL, Xie DN, Zhu XW, Li X

1227 Predictive modeling for postoperative delirium in elderly patients with abdominal malignancies using synthetic minority oversampling technique

Hu WJ, Bai G, Wang Y, Hong DM, Jiang JH, Li JX, Hua Y, Wang XY, Chen Y

Efficacy and predictive factors of transarterial chemoembolization combined with lenvatinib plus 1236 programmed cell death protein-1 inhibition for unresectable hepatocellular carcinoma

Ma KP, Fu JX, Duan F, Wang MQ

1248 Should we perform sigmoidoscopy for colorectal cancer screening in people under 45 years? Leong W, Guo JQ, Ning C, Luo FF, Jiao R, Yang DY

1256 Computed tomography-based radiomics diagnostic approach for differential diagnosis between early- and late-stage pancreatic ductal adenocarcinoma

Ren S, Qian LC, Cao YY, Daniels MJ, Song LN, Tian Y, Wang ZQ

1268 Prognostic analysis of related factors of adverse reactions to immunotherapy in advanced gastric cancer and establishment of a nomogram model

He XX, Du B, Wu T, Shen H

Clinical Trials Study

1281 Safety and efficacy of a programmed cell death 1 inhibitor combined with oxaliplatin plus S-1 in patients with Borrmann large type III and IV gastric cancers

Bao ZH, Hu C, Zhang YQ, Yu PC, Wang Y, Xu ZY, Fu HY, Cheng XD

Observational Study

1296 Computed tomography radiogenomics: A potential tool for prediction of molecular subtypes in gastric stromal tumor

Yin XN, Wang ZH, Zou L, Yang CW, Shen CY, Liu BK, Yin Y, Liu XJ, Zhang B

1309 Application of texture signatures based on multiparameter-magnetic resonance imaging for predicting microvascular invasion in hepatocellular carcinoma: Retrospective study

Nong HY, Cen YY, Qin M, Qin WQ, Xie YX, Li L, Liu MR, Ding K

- 1319 Causal roles of gut microbiota in cholangiocarcinoma etiology suggested by genetic study Chen ZT, Ding CC, Chen KL, Gu YJ, Lu CC, Li QY
- 1334 Is recovery enhancement after gastric cancer surgery really a safe approach for elderly patients? Li ZW, Luo XJ, Liu F, Liu XR, Shu XP, Tong Y, Lv Q, Liu XY, Zhang W, Peng D
- 1344 Establishment of a cholangiocarcinoma risk evaluation model based on mucin expression levels Yang CY, Guo LM, Li Y, Wang GX, Tang XW, Zhang QL, Zhang LF, Luo JY



Contor	World Journal of Gastrointestinal Oncology
Conten	Monthly Volume 16 Number 4 April 15, 2024
1361	Effectiveness of fecal DNA syndecan-2 methylation testing for detection of colorectal cancer in a high-risk Chinese population
	Luo WF, Jiao YT, Lin XL, Zhao Y, Wang SB, Shen J, Deng J, Ye YF, Han ZP, Xie FM, He JH, Wan Y
	Clinical and Translational Research
1374	Clinical and socioeconomic determinants of survival in biliary tract adenocarcinomas
	Sahyoun L, Chen K, Tsay C, Chen G, Protiva P
1384	Risk factors, prognostic factors, and nomograms for distant metastasis in patients with diagnosed duodenal cancer: A population-based study
	Shang JR, Xu CY, Zhai XX, Xu Z, Qian J
1421	NOX4 promotes tumor progression through the MAPK-MEK1/2-ERK1/2 axis in colorectal cancer
	Xu YJ, Huo YC, Zhao QT, Liu JY, Tian YJ, Yang LL, Zhang Y
	Basic Study
1437	Curcumin inhibits the growth and invasion of gastric cancer by regulating long noncoding RNA AC022424.2
	Wang BS, Zhang CL, Cui X, Li Q, Yang L, He ZY, Yang Z, Zeng MM, Cao N
1453	MicroRNA-298 determines the radio-resistance of colorectal cancer cells by directly targeting human dual- specificity tyrosine(Y)-regulated kinase 1A
	Shen MZ, Zhang Y, Wu F, Shen MZ, Liang JL, Zhang XL, Liu XJ, Li XS, Wang RS
1465	Human β -defensin-1 affects the mammalian target of rapamycin pathway and autophagy in colon cancer cells through long non-coding RNA TCONS_00014506
	Zhao YX, Cui Y, Li XH, Yang WH, An SX, Cui JX, Zhang MY, Lu JK, Zhang X, Wang XM, Bao LL, Zhao PW
1479	FAM53B promotes pancreatic ductal adenocarcinoma metastasis by regulating macrophage M2 polarization
	Pei XZ, Cai M, Jiang DW, Chen SH, Wang QQ, Lu HM, Lu YF
1500	Transcriptome sequencing reveals novel biomarkers and immune cell infiltration in esophageal tumori- genesis
	Sun JR, Chen DM, Huang R, Wang RT, Jia LQ
1514	Construction of CDKN2A-related competitive endogenous RNA network and identification of GAS5 as a prognostic indicator for hepatocellular carcinoma
	Pan Y, Zhang YR, Wang LY, Wu LN, Ma YQ, Fang Z, Li SB
1532	Two missense <i>STK11</i> gene variations impaired LKB1/adenosine monophosphate-activated protein kinase signaling in Peutz-Jeghers syndrome
	Liu J, Zeng SC, Wang A, Cheng HY, Zhang QJ, Lu GX
1547	Long noncoding RNAs HAND2-AS1 ultrasound microbubbles suppress hepatocellular carcinoma progression by regulating the miR-873-5p/tissue inhibitor of matrix metalloproteinase-2 axis
	Zou Q, Wang HW, Di XL, Li Y, Gao H

Conton	World Journal of Gastrointestinal Oncology
Conten	Monthly Volume 16 Number 4 April 15, 2024
1564	Upregulated IncRNA PRNT promotes progression and oxaliplatin resistance of colorectal cancer cells by regulating HIPK2 transcription
	Li SN, Yang S, Wang HQ, Hui TL, Cheng M, Zhang X, Li BK, Wang GY
	SYSTEMATIC REVIEWS
1578	Prognosis value of heat-shock proteins in esophageal and esophagogastric cancer: A systematic review and meta-analysis
	Nakamura ET, Park A, Pereira MA, Kikawa D, Tustumi F
1596	Risk factors for hepatocellular carcinoma associated with hepatitis C genotype 3 infection: A systematic review
	Farooq HZ, James M, Abbott J, Oyibo P, Divall P, Choudhry N, Foster GR
	META-ANALYSIS
1613	Effectiveness and tolerability of programmed cell death protein-1 inhibitor + chemotherapy compared to chemotherapy for upper gastrointestinal tract cancers
	Zhang XM, Yang T, Xu YY, Li BZ, Shen W, Hu WQ, Yan CW, Zong L
1626	Success rate of current human-derived gastric cancer organoids establishment and influencing factors: A systematic review and meta-analysis
	Jiang KL, Wang XX, Liu XJ, Guo LK, Chen YQ, Jia QL, Yang KM, Ling JH
	CASE REPORT
1647	Pathologically successful conversion hepatectomy for advanced giant hepatocellular carcinoma after multidisciplinary therapy: A case report and review of literature
	Chu JH, Huang LY, Wang YR, Li J, Han SL, Xi H, Gao WX, Cui YY, Qian MP
1660	Clinical pathological characteristics of "crawling-type" gastric adenocarcinoma cancer: A case report
	Xu YW, Song Y, Tian J, Zhang BC, Yang YS, Wang J
1668	Primary pancreatic peripheral T-cell lymphoma: A case report
	Bai YL, Wang LJ, Luo H, Cui YB, Xu JH, Nan HJ, Yang PY, Niu JW, Shi MY



Contents

World Journal of Gastrointestinal Oncology

Monthly Volume 16 Number 4 April 15, 2024

ABOUT COVER

Peer Reviewer of World Journal of Gastrointestinal Oncology, Lie Zheng, Director, Professor, Department of Gastroenterology, Shaanxi Provincial Hospital of Traditional Chinese Medicine, Xi'an 730000, Shaanxi Province, China. xinliwen696@126.com

AIMS AND SCOPE

The primary aim of World Journal of Gastrointestinal Oncology (WJGO, World J Gastrointest Oncol) is to provide scholars and readers from various fields of gastrointestinal oncology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGO mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal oncology and covering a wide range of topics including liver cell adenoma, gastric neoplasms, appendiceal neoplasms, biliary tract neoplasms, hepatocellular carcinoma, pancreatic carcinoma, cecal neoplasms, colonic neoplasms, colorectal neoplasms, duodenal neoplasms, esophageal neoplasms, gallbladder neoplasms, etc.

INDEXING/ABSTRACTING

The WJGO is now abstracted and indexed in PubMed, PubMed Central, Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJGO as 3.0; IF without journal self cites: 2.9; 5-vear IF: 3.0; Journal Citation Indicator: 0.49; Ranking: 157 among 241 journals in oncology; Quartile category: Q3; Ranking: 58 among 93 journals in gastroenterology and hepatology; and Quartile category: Q3. The WJGO's CiteScore for 2022 is 4.1 and Scopus CiteScore rank 2022: Gastroenterology is 71/149; Oncology is 197/366.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Xiang-Di Zhang; Production Department Director: Xiang Li; Cover Editor: Jia-Ru Fan.

NAME OF JOURNAL	INSTRUCTIONS TO AUTHORS
World Journal of Gastrointestinal Oncology	https://www.wjgnet.com/bpg/gerinfo/204
ISSN	GUIDELINES FOR ETHICS DOCUMENTS
ISSN 1948-5204 (online)	https://www.wjgnet.com/bpg/GerInfo/287
LAUNCH DATE	GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH
February 15, 2009	https://www.wjgnet.com/bpg/gerinfo/240
FREQUENCY	PUBLICATION ETHICS
Monthly	https://www.wjgnet.com/bpg/GerInfo/288
EDITORS-IN-CHIEF	PUBLICATION MISCONDUCT
Monjur Ahmed, Florin Burada	https://www.wjgnet.com/bpg/gerinfo/208
EDITORIAL BOARD MEMBERS	ARTICLE PROCESSING CHARGE
https://www.wjgnet.com/1948-5204/editorialboard.htm	https://www.wjgnet.com/bpg/gerinfo/242
PUBLICATION DATE	STEPS FOR SUBMITTING MANUSCRIPTS
April 15, 2024	https://www.wjgnet.com/bpg/GerInfo/239
COPYRIGHT	ONLINE SUBMISSION
© 2024 Baishideng Publishing Group Inc	https://www.f6publishing.com

© 2024 Baishideng Publishing Group Inc. All rights reserved. 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA E-mail: office@baishideng.com https://www.wjgnet.com



0 WJ

World Journal of *Gastrointestinal* Oncology

Submit a Manuscript: https://www.f6publishing.com

World J Gastrointest Oncol 2024 April 15; 16(4): 1532-1546

DOI: 10.4251/wjgo.v16.i4.1532

Basic Study

ISSN 1948-5204 (online)

ORIGINAL ARTICLE

Two missense STK11 gene variations impaired LKB1/adenosine monophosphate-activated protein kinase signaling in Peutz-Jeghers syndrome

Jin Liu, Si-Cong Zeng, An Wang, Hai-Ying Cheng, Qian-Jun Zhang, Guang-Xiu Lu

Specialty type: Genetics & heredity

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0 Grade B (Very good): B Grade C (Good): 0 Grade D (Fair): 0 Grade E (Poor): 0

P-Reviewer: Gragnaniello V, Italy

Received: December 5, 2023 Peer-review started: December 5, 2023 First decision: December 14, 2023 Revised: December 29, 2023 Accepted: February 3, 2024

Article in press: February 3, 2024 Published online: April 15, 2024



Jin Liu, Si-Cong Zeng, An Wang, Hai-Ying Cheng, Qian-Jun Zhang, Guang-Xiu Lu, Hunan Guangxiu Hospital, Hunan Normal University, Changsha 410000, Hunan Province, China

Jin Liu, Si-Cong Zeng, Scientific Research Department, Reproductive and Genetic Hospital of Citic-Xiangya, Changsha 410000, Hunan Province, China

Corresponding author: Guang-Xiu Lu, PhD, Doctor, Professor, Hunan Guangxiu Hospital, Hunan Normal University, No. 8 Luyun Road, Yuelu District, Changsha 410000, Hunan Province, China. lugxdirector@aliyun.com

Abstract

BACKGROUND

Peutz-Jeghers syndrome (PJS) is a rare hereditary neoplastic disorder mainly associated with serine/threonine kinase 11 (STK11/LKB1) gene mutations. Preimplantation genetic testing can protect a patient's offspring from mutated genes; however, some variations in this gene have been interpreted as variants of uncertain significance (VUS), which complicate reproductive decision-making in genetic counseling.

AIM

To identify the pathogenicity of two missense variants and provide clinical guidance.

METHODS

Whole exome gene sequencing and Sanger sequencing were performed on the peripheral blood of patients with PJS treated at the Reproductive and Genetic Hospital of Citic-Xiangya. Software was employed to predict the protein structure, conservation, and pathogenicity of the two missense variation sites in patients with PJS. Additionally, plasmids were constructed and transfected into HeLa cells to observe cell growth. The differences in signal pathway expression between the variant group and the wild-type group were compared using western blot and immunohistochemistry. Statistical analysis was performed using one-way analysis of variance. P < 0.05 was considered statistically significant.

RESULTS

We identified two missense STK11 gene VUS [c.889A>G (p.Arg297Gly) and



c.733C>T (p.Leu245Phe)] in 9 unrelated PJS families who were seeking reproductive assistance. The two missense VUS were located in the catalytic domain of serine/threonine kinase, which is a key structure of the liver kinase B1 (LKB1) protein. *In vitro* experiments showed that the phosphorylation levels of adenosine monophosphate-activated protein kinase (AMPK) at Thr172 and LKB1 at Ser428 were significantly higher in transfected variation-type cells than in wild-type cells. In addition, the two missense *STK11* variants promoted the proliferation of HeLa cells. Subsequent immunohistochemical analysis showed that phosphorylated-AMPK (Thr172) expression was significantly lower in gastric, colonic, and uterine polyps from PJS patients with missense variations than in non-PJS patients. Our findings indicate that these two missense *STK11* variants are likely pathogenic and inactivate the *STK11* gene, causing it to lose its function of regulating downstream phosphorylated-AMPK (Thr172), which may lead to the development of PJS. The identification of the pathogenic mutations in these two clinically characterized PJS patients has been helpful in guiding them toward the most appropriate mode of pregnancy assistance.

CONCLUSION

These two missense variants can be interpreted as likely pathogenic variants that mediated the onset of PJS in the two patients. These findings not only offer insights for clinical decision-making, but also serve as a foundation for further research and reanalysis of missense VUS in rare diseases.

Key Words: Missense; STK11; Peutz-Jeghers syndrome; Rare disease; Genetic counseling; Assisted reproductive technique

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: These two missense variants, *STK11* c.889A>G (p.Arg297Gly) and *STK11* c.733C>T (p.Leu245Phe), have been found to contribute to the development of Peutz-Jeghers syndrome (PJS) by impairing the *STK11*/adenosine monophosphate-activated protein kinase signaling pathway. It clarifies that these two germline variants, *STK11* c.889A>G (p.Arg297Gly) and *STK11* c.733C>T (p.Leu245Phe), are likely pathogenic mutations, providing valuable information for fertility selection in patients with PJS.

Citation: Liu J, Zeng SC, Wang A, Cheng HY, Zhang QJ, Lu GX. Two missense *STK11* gene variations impaired LKB1/adenosine monophosphate-activated protein kinase signaling in Peutz-Jeghers syndrome. *World J Gastrointest Oncol* 2024; 16(4): 1532-1546 **URL:** https://www.wjgnet.com/1948-5204/full/v16/i4/1532.htm **DOI:** https://dx.doi.org/10.4251/wjgo.v16.i4.1532

INTRODUCTION

Peutz-Jeghers syndrome [PJS; Mendelian Inheritance in Man (MIM), #175200, www.ncbi.nlm.gov/OMIM] is a rare hereditary neoplastic disorder characterized by mucocutaneous pigmentation and the development of multiple gastrointestinal (GI) polyps[1]. The annual incidence of PJS ranges from 1: 8300 to 1: 29000 and does not differ by sex, race, or ethnicity[2-4]. PJS has extremely high penetrance and clinical heterogeneity[5] and can lead to the formation of hamartomatous polyps in many organs, such as the jejunum, ileum, duodenum, stomach, colon, and gallbladder[6,7]. Even at a very young age, PJS patients with pathological hamartomatous polyps can experience numerous GI complications, including bleeding, anemia, abdominal pain, intussusception, obstruction, and infarction[8]. Patients with PJS are at high risk for GI and extra-GI cancer[9,10]. The disease is inherited in an autosomal dominant manner and has a 50% chance of being passed on to offspring.

The serine/threonine kinase 11 (*STK11/LKB1*; MIM, *602216) gene has been identified as a pathogenic factor of PJS, with germline mutations in this gene listed in the medical records of most PJS patients as a risk factor[3,7]. *STK11* is a tumor suppressor gene located on chromosome 19p13.3 and includes ten exons[11]. *STK11* functions mainly depend on the phosphorylation and activation of adenosine monophosphate-activated protein kinase (AMPK). As an important metabolic regulator, AMPK regulates cell generation by regulating the activity of metabolic enzymes and activating adaptive transcriptional responses[12]. *STK11* is a master stress response regulator[13] and plays an important role in tumorigenesis, implicated in numerous key biological processes, including cell metabolism, cell cycle regulation, cell polarity and motility, and angiogenesis[1].

A search for *STK11* gene variants in the ClinVar database revealed a total of 2481 entries with clinical significance, of which 974 (39.26%) were variants of uncertain significance (VUS), predominantly (79.98%, 779/974) missense variants until December 18, 2023. Many VUS affect diagnosis, clinical treatment, and decision-making processes because their pathogenicity cannot be defined. Some variants have been shown to be frameshift or nonsense mutations that result in abnormal protein truncation and subsequent loss of kinase activity and are therefore classified as pathogenic or potentially pathogenic. However, missense variants usually only lead to the formation of incorrect amino acid residues at the corresponding position in the translated product, leading to an abnormal protein. Whether this protein alteration affects the normal function of the gene needs to be tested experimentally. Understanding how the function of *STK11* is

Raishideng® WJGO https://www.wjgnet.com

disrupted may contribute to a better understanding of the molecular mechanisms involved in the pathogenesis and carcinogenesis of PJS. Clinical fertility counseling and reproductive assistance need to target specific variant sites, so determining the pathogenicity of VUS is urgent and important for patients.

MATERIALS AND METHODS

Patient

Peripheral blood samples were collected from 9 PJS probands and 19 family members, all of whom were Chinese Han. These patients with PJS were diagnosed according to the PJS diagnostic criteria published by the European consensus in 2007[14]. When ≥ 2 individuals are affected within the same family, they are considered family cases. Patients without a family history of PIS are defined as sporadic cases. Patient information such as family history and clinical characteristics were collected from the patient's medical records. All patients provided informed consent for this study. Clinical data collection included demographics, medical history, family history, symptoms, and surgical interventions related to PJS. Some patients underwent endoscopic or surgical polypectomy. Tissues were fixed with formalin and embedded in paraffin. The wax blocks were successively sliced and prepared for subsequent hematoxylin-eosin (HE) staining and immunohistochemical (IHC) experiments.

Whole-exome sequencing and Sanger sequencing

The peripheral blood of all probands was sent to Beijing Genomics Research Institute for Whole-exome sequencing (WES), and the tumor-related gene exons and adjacent ± 10 bp intron variations were detected. The results were verified by polymerase chain reaction (PCR)-Sanger sequencing. The identified variations were further tested in all available family members to confirm co-segregation of these variations with the disease. We used the online site primer 3 to design primers. Variation analysis was mainly based on the American Society for Medical Genetics and Genomics (ACMG) guidelines for interpreting sequence variation^[15].

Predicting the impact of identified VUS

Amino acid conservation analysis was performed using UGENE software. Five online software programs, AlphaMissense (https://alphamissense.hegelab.org/search), PROVEAN (http://provean.jcvi.org/seq_submit.php), mutation Taster (http://www.mutationtaster.org/), PolyPhen-2 (http://genetics.bwh.harvard.edu/pph2/), and Swiss-model (http:// swissmodel.expasy.org/), were used to predict the functional and structural significance of these variations.

Tissue staining

Tissue sections were incubated with liver kinase B1 (LKB1), phosphorylated (p)-LKB1 (Ser428), AMPK, p-AMPK (Thr172) antibodies. The IHC kit used was from Leica Biosystems (catalog No.RE7280-k, Germany) and was operated according to the reagent instructions. Polyp tissue sections were stained by HE staining. An experienced pathologist reviewed these specimens and determined the histological types of these polyps. The staining of the sliced tissue with a brownish yellow color indicates positivity. Statistical analysis of IHC results was based on a direct count of the number of positive cells in five random visual fields at 70 × magnification or a visual assessment of the proportion of positive cells. Each group was repeated three times.

Cell experiments

Five plasmids including enhanced green fluorescent protein (EGFP)-N1-vector (EGFP-Empty), PEGFP-N1-STK11 (EGFP-STK11), PEGFP-N1-STK11 c.889A>G (p.Arg297Gly), PEGFP-N1-STK11 c.733C>T (p.Leu245Phe) and PEGFP-N1-STK11 c.910C>T(p.Arg304Trp) were transfected into HeLa cells with lipofectamine 3000. Cells were collected 48 h after transfection, subsequent PCR product gel electrophoresis, Western blotting (WB), Reverse transcription quantitative PCR (RT-qPCR), and Cell Counting Kit 8 (CCK 8) experiments were performed. Transfected cells were counted and seeded in 6-well plates (1000 cells/well). Cell proliferation was detected at 12 h, 24 h, 36 h, 48 h, 60 h, and 72 h after transfection. Each group was repeated three times.

Total RNA was extracted using Trizol, and cDNA was synthesized by reverse transcription from 1 µg of total RNA using oligo (dT) primers. RT-qPCR was performed using the All-in-One™ qPCR mix on a LightCycler 480 system (Roche, Switzerland). Primers were performed according to the primer website (https://www.ncbi.nlm.nih.gov/tools/primer). Glyceraldehyde 3-phosphate dehydrogenase (GAPDH) was used as an internal control for normalization. The level of expression endogenous, exogenous, and total mRNA of STK11 was calculated using the 2-^{ΔΔCt} method.

Total protein was extracted using radioimmunoprecipitation assay lysis buffer and detected using bicinchoninic acid kits. Total protein was separated on 10% SDS-PAGE. After separation, the protein was elactrotransferred to a polyvinylidene difluoride membrane, and then blocked in 5% skimmed milk for 1 h at room temperature. The membranes were incubated with primary antibodies against STK11, p-STK11, AMPK, p-AMPK and GAPDH overnight at 4°C. GAPDH served as the loading control. Membranes were subsequently incubated with secondary antibodies for 2 h at room temperature. Protein expression was detected using an enhanced chemiluminescence kit (Thermo Fisher Scientific, United States).

Statistical analysis

All statistical analyzes were performed with SPSS (version 26.0, United States) and GraphPad Prism (version 9.0, United



States). The data are expressed as mean ± SD unless otherwise noted. Comparisons among multiple groups were determined using one-way analysis of variance, followed by least significant difference or Dunnett's T3 test. P < 0.05 was considered statistically significant.

RESULTS

Clinical features of patients

Some patients with PJS and a clinical phenotype seek genetic and reproductive assistance when giving birth. The demographics and clinical characteristics of 9 patients with PJS are shown in Table 1. At the time of the last follow-up, 8 patients were married, and 1 was unmarried. All patients with PJS had skin and mucosa pigmentation. The median age of the 9 PJS patients when they visited our hospital for the first time was 30 years (ranging from 23 to 45 years), of whom 5 were male (55.6%). The age at diagnosis varied widely, with some patients being diagnosed in the first few years of life and others demonstrating a later onset and milder symptoms. Eight of the 9 patients (89.0%) with PIS developed colon polyps and underwent polypectomy, and the mean age at the onset of the first GI polyp was 19.38 years (19.38 \pm 4.39). The burden of polyps (cumulative number of polyps) varied from approximately 10 to several hundred. PJS polyps were distributed in the stomach in 5 patients (55.6%), in the small intestine in 7 (77.8%), and in the colorectum in 4 (44.4%). Two patients (22.2%) had uterine polyps, 1 (11.1%) had gallbladder polyps, and 1 (11.1%) had nasal polyps. Six (66.67%) were initially treated before the age of 20 years, and all patients received initial treatment before the age of 30 years. Patients with a family history of PJS were treated at a later age than those without; however, the difference was not significant (*P* = 0.4059). The age at first treatment differed by sex, with the first treatment occurring later in men than in women; however, the difference was not significant (P = 0.4521).

WES identified heterozygous STK11 variations

WES or STK11 variant detection was performed for the 9 unrelated PJS patients. Ten (111%) different germline variations in the STK11 gene were identified. Clinical and sociodemographic data were collected from 26 family members, 19 of whom underwent pedigree analysis for the STK11 gene (Supplementary Table 1). A total of 4 STK11 variations were identified in familial cases and 6 in sporadic cases. Of the 10 variants identified, three were nonsense variants, three were splicing variants, three were missense variants, and one was a frameshift variant. According the ACMG guidelines, the variant pathogenicity assessment indicated that 5 were pathogenic, 2 were likely pathogenic, 1 was likely benign, and 2 were VUS. Two patients with missense VUS underwent GI endoscopy at our hospital, and one had polyps removed from the stomach, colon, and uterus (Figure 1A-F). Two of the variations [c.889A>G (p.Arg297Gly) and c.733C>T (p.Leu245Phe)] were identified by Sanger sequencing (Figure 1G-J). A schematic representation of the localization of 10 STK11 variations is shown in Figure 1K; most of the variations were concentrated in the exon 1 region.

Three-dimensional structure and function prediction of proteins resulting from missense STK11 VUS

Swiss-model software was used to construct a three-dimensional (3D) model of the protein structure of STK11 (Figure 2A-D). The STK11 c.889A>G (p.Arg297Gly) variant resulted in the substitution of arginine for glycine. Arginine has a large molecular weight, electrically charged side chain, hydrophilic properties, and an alkaline R group. It plays an important role in the structure and function of proteins. Glycine has a small molecular weight, nonpolar side chain, hydrophobic properties, and a neutral R group containing only one hydrogen atom, as well as a large degree of freedom. The STK11 c.733C>T (p.Leu245Phe) variant substituted leucine with phenylalanine. Both amino acids have nonpolar side chains and are hydrophobic, but leucine is a side branched-chain amino acid, while phenylalanine has an aromatic ring in the side chain. Therefore, the variation between these two amino acids may lead to large structural changes. The two amino acids were evolutionarily conserved by UGENE (Figure 2E), while Mutation Taster, AlphaMissese, PolyPhen-2, and SIFT predicted the two missense variations (p.Arg297Gly and p.Leu245Phe) to be deleterious (Figure 2F).

Functional analysis of the two missense STK11 variations in HeLa cells

To verify whether these two missense variants affect STK11 gene transcription and translation, STK11 plasmids were established [an empty vector, wild-type vector, STK11 c.889A>G (p.Arg297Gly) variant vector, and c.733C>T (p.Leu245Phe) variant vector]. One missense variant c.910C>T (p.Arg304Trp) was interpreted as pathogenic in ClinVar; thus, a plasmid with this mutation was used as the positive control. PCR and WB were used to detect endogenous STK11 expression levels in HeLa and HEK-293T cell lines. STK11 expression levels were nearly undetectable in HeLa cells vs HEK-293T cell lines. Thus, HeLa cells were used for further studies. All plasmids were verified by gel electrophoresis and PCR product sequencing; since all carried GFP constructs, successful transfection was detected by fluorescence microscopy (Figure 3A-E). STK11 expression was examined by WB and RT-qPCR analysis 48 h after transfection. The protein and mRNA expression levels of STK11 in HeLa cells transfected with the STK11 variation and wild-type vectors were significantly increased in comparison with those in cells transfected with the empty vector. However, there was no significant difference in STK11 expression in cells transfected with the wildtype vector and the STK11 variant vectors (Figure 3F and G). These results suggest that the two missense STK11 variants have negligible effects on STK11 transcription and translation.

We next tested LKB1, p-LKB1, AMPK, and p-AMPK protein levels to detect the effects of these two missense VUS on LKB1 protein function and whether they play a role through the downstream AMPK signaling pathway. p-LKB1 and p-AMPK levels were significantly lower in all cells transfected with variant plasmids than in wild-type cells (Figure 3H).

WJGO | https://www.wjgnet.com







 Jaisbideng®
 WJGO
 https://www.wjgnet.com

April 15, 2024 Volume 16 Issue 4



Figure 1 Clinical features and sequencing data of patients with Peutz-Jeghers syndrome. A-F: Polyps and hematoxylin-eosin (HE) staining of a patient with Peutz-Jeghers syndrome (PJS). Gastroenteroscopy and hysteroscopy revealed and resected multiple polyps in the stomach, intestines, and uterus. Gastric polyp and HE (A and D), colon polyp and HE (B and E), endometrial polyp and HE (C and F), magnification, 5 × (D-F); G-J: Family pedigree and sequencing maps of two missense *STK11* variants of uncertain significance (VUS) in two PJS patients; Family pedigree of a female patient with PJS who had *STK11* c.889A>G (VUS) and c.1062C>G (likely benign) variations (G); The patient and her older brother had *STK11* c.889A>G variation, but her father and two aunts were undetected (H); Family pedigree of a male patient with PJS who had *STK11* c.733C>T (VUS) variations (I); The patient had *STK11* c.733C>T variations, but his father and mother were undetected (J); K: Schematic representation of the *STK11* gene structure and variations in 9 PJS families. Arg: Arginine; Gly: Glycine; Phe: Phenylalanine; Leu: Leucine; y: Years; Tyr: Tyrosine; Term: Termination; Lys: Lysine; del: Deletion; Glu: Glutamic acid.

Baisbidena® WJGO | https://www.wjgnet.com



Figure 2 Multiple software predict the pathogenicity of these variations. A-D: Three-dimensional model of the protein structure of liver kinase B1 and amino acid molecular structure formula. These variations result in significant changes in the properties of amino acids, which may be harmful; E: Evolutionary conservation shows that the variation sites (p.R297G and p.L245F) are highly conserved across different species; F: Four software assessed the pathogenicity of these variations, all of which were likely pathogenic or pathogenic. "R" represents arginine; "G" represents glycine; "L" represents leucine; and "F" represents phenylalanine.

Raishideng® WJGO | https://www.wjgnet.com

April 15, 2024 Volume 16 Issue 4



Baishideng® WJGO | https://www.wjgnet.com



Figure 3 Expression and functional analysis of STK11. A: Plasmids were successfully transfected into HeLa cells. HeLa cell-Lipofectamine 3000; and HeLa cell transfected with enhanced green fluorescent protein (EGFP)-Empty, EGFP-N1-STK11, EGFP-N1-STK11 c.889A>G (p.Arg297Gly), EGFP-N1-STK11 c.733C>T (p.Leu245Phe) and EGFP-N1-STK11 c.910C>T (p.Arg304Trp) plasmid vectors, Magnification: 10 ×; B: Plasmid transfection efficiency was high; C: PCR gel electrophoresis confirmed that the transfected plasmid was correct; D: The endogenous expression of STK11 gene was low in HeLa cells; E: The exogenous expression of STK11 gene in HeLa cells after transfection; F: These missense variants do not affect gene transcription; G: These variations do not affect the function of gene translation into proteins; H: These missense variants disrupted the phosphorylation of liver kinase B1 and adenosine monophosphate-activated protein kinase; I: These missense variants promote cell proliferation, °P < 0.0001. "R" represents arginine; "G" represents glycine; "L" represents leucine; "F" represents phenylalanine; and "S" represents serine. Independent sample results were obtained after at least three experiments and then obtained statistically. NS: Not significant; GAPDH: Glyceraldehyde 3-phosphate dehydrogenase; LKB1: Serine/threonine kinase 11.

We performed CCK-8 assays to further determine whether these two STK11 VUS could effectively inhibit cell growth and proliferation. Compared with the wild-type group, the growth of cells transfected with the STK11 variant vectors was significantly accelerated (Figure 31). These results indicate that both missense VUS disrupt the endogenous protein kinase activity of STK11 and inhibit the activation of the AMPK pathway, disrupting the inhibitory function of the STK11 gene on cell growth, leading to cell proliferation that may contribute to the development of polyps and tumors.

In summary, based on the above results, functional evidence is provided that these two missense STK11 variants are likely pathogenic according to the ACMG's guidelines.

STK11 gene variation may influence the occurrence of polyps in PJS patients via the LKB1/AMPK pathway

Polyp tissue from non-PJS patients with the same social factors as the PJS patients was used as the control, and the LKB1, p-LKB1, AMPK, and p-AMPK protein expression levels were compared between the two groups. The expression levels of p-STK11 and p-AMPK were decreased in the stomach, colon, and uterus in patients with PJS, and the differences were statistically significant (P < 0.05). Therefore, it is inferred that the STK11 gene may cause polyps or PJS through the LKB1/ AMPK pathway (Figures 4 and 5).

DISCUSSION

STK11 plays a crucial role in regulating cell damage repair, energy metabolism, and tumor immune responses[16]. The LKB1 protein comprises three major domains: The N-terminal non-catalytic domain (encoded by amino acids 1-49), the catalytic kinase domain (encoded by amino acids 49-309), and the C-terminal non-catalytic regulatory domain (encoded by amino acids 309-433)[1,17]. Variations in patients with PJS are predominantly located in the catalytic domain region, leading to kinase activity dysfunction and the disruption of STK11 function[18]. Germline STK11 mutation has been confirmed to be a significant cause of PJS[19].

We conducted a review of PJS patients with STK11 gene variations at our hospital. A total of 10 variations were found, primarily in exon 1, consistent with Li et al^[20]. While all PJS patients exhibited STK11 gene variations, some were VUS [21,22], and numerous *STK11* gene VUS exist in the ClinVar database, with the majority being missense variations[23]. Understanding the relationship between these variations and PJS is challenging, complicating clinical decision-making, reproduction decisions, and genetic counseling for the patient[24,25], which may lead to increased psychological stress [26]. Therefore, it is crucial to determine the pathogenicity of these VUS.

Most patients with PJS have STK11 germline mutations, and their offspring face an elevated risk of inheriting the same mutation and experiencing PJS due to the autosomal dominant inheritance pattern^[27]. PJS patients affected by the disease naturally do not want their offspring to experience the same effects [28]. Therefore, some patients opt for



WJGO | https://www.wjgnet.com



Figure 4 Expression of the serine/threonine kinase 11 in polyps of Peutz-Jeghers syndrome and non-Peutz-Jeghers syndrome patients. A-D: The phosphorylation levels of liver kinase B1 in Peutz-Jeghers syndrome (PJS) patients polyps were significantly lower than those in non-PJS patients polyps. PJS: Peutz-Jeghers syndrome; LKB1: Liver kinase B1. ^aP < 0.05; ^bP < 0.001; ^cP < 0.0001.

preimplantation genetic testing (PGT)[29] to prevent the transmission of disease-causing mutations to future generations [5,30]. Some PJS patients choose to pursue a natural pregnancy[31]. In cases where the female patient or the male patient's partner becomes pregnant, prenatal diagnosis is performed to predict whether the fetus carries STK11 or other genetic mutations[32,33]. Parents can terminate the pregnancy if the fetus is a carrier of the STK11 mutation[34]; therefore, this method also allows healthy children to be born. In our study, one PJS patient chose in vitro fertilization-embryo transfer. Unfortunately, prenatal diagnosis at 18 wk of gestation revealed that the fetus carried the STK11 gene mutation; thus, the pregnancy was terminated. Seven patients (78%) chose PGT to block the transmission of the mutated genes to their offspring. It is worth mentioning that after the two missense VUS were indicated to likely be pathogenic through experiments, both patients chose PGT to assist their pregnancies, and one patient then gave birth to a healthy girl. The other suffered a spontaneous miscarriage in the fifth week of pregnancy. Therefore, identifying the pathogenicity of genetic variations may affect the patient's reproduction decisions.

The two missense variants found in this study are VUS, complicating clinical decision-making. Clinical geneticists and reproductive specialists cannot provide help and guidance to patients with VUS since it is unclear whether the variant is related to the disease. Here, the two missense VUS were predicted by software to be harmful and affect protein structure; however, because the variants were VUS, the patients faced great difficulties in clinical fertility selection. Now that experiments have supported the idea that these variants can be interpreted as being likely pathogenic, reproductive, and genetic physicians will be able to provide better fertility counseling to patients. The findings can also be used for evidence-based genetic and reproductive counseling for PJS families. Finally, we provided valuable insights into the molecular mechanisms of PJS pathogenesis. Two patients with PJS had a child by PGT, and no pathogenic gene mutation was found in the fetus by prenatal diagnosis, suggesting that the combination of molecular and prenatal diagnosis is an effective way to prevent the recurrence or transmission of inherited diseases in these families. However, determining the genetic cause before conception is crucial in this process.

Zaishideng® WJGO | https://www.wjgnet.com





Several limitations should be noted in our study. First, because of the rarity of PJS, the number of patients included is relatively small, which may reduce the statistical analysis power. Second, the follow-up period was short; the longest follow-up was 8 years. Because the patients with PJS were generally young when they visited the doctor, no malignant tumors were found till the date of the final follow-up; thus, the correlation between PJS and cancer development could not be discussed. Third, because our hospital mainly caters to patients who need assisted reproduction and genetic counseling, there may be some bias in the selection of patients included; that is, patients visit the doctor for assisted reproduction, leading to a high proportion of patients choosing PGT.

CONCLUSION

We identified two missense variants in the *STK11* gene that were experimentally evaluated for their functional significance. Our results identify the two missense variations as being likely pathogenic, providing new guidance for genetic counseling, fertility counseling, and prenatal diagnosis in patients with *STK11* gene variants.

Zaishideng® WJGO | https://www.wjgnet.com

Proband family No.	Proband gender, age (yr)	Clinical features	Age at first discovery of polyps	Previous medicine history	Nucleotide alteration, amino acid alteration	Exon/Intron, mutation type	ACMG	PJS family history	Reproductive choice	Current Clinical outcome	Pedigree analysis
1	Man, 34	Pigmentation of skin and mucosa; multiple polyps	19	Intestinal polypectomy	c.180C>A (p.Tyr60Ter)	Exon1, nonsense	Р	De novo	IVF-ET	The fetus had the same mutation and the pregnancy was terminated	Reject
2	Woman, 30	Pigmentation of skin and mucosa; multiple polyps	25	Polypectomy was performed in the stomach, colon and uterus	c.889A>G (p.Arg297Gly)	Exon7, missense	VUS	Yes, maternal. Mother, brother	PGT	The fetus was 5 wk+ and had a spontaneous abortion	Brother carries STK11 variation but father doesn't
					c.1062C>G (p.Phe354Leu)	Exon 8, missense	LB				
3	Man, 27	Pigmentation of skin and mucosa; multiple polyps	16	Multiple gastrointestinal polypectomy	c.733C>T (p.Leu245Phe)	Exon5, missense	VUS	De novo	PGT	The fetus was prenatally diagnosed as having no mutation and gave birth to a child	Parents had no <i>STK11</i> variation
4	Man, 25	Pigmentation of skin and mucosa; multiple polyps	15	Multiple gastrointestinal polypectomy	c.250A>T (p.Lys84 [*])	Exon1, nonsense	Р	De novo	PGT	Preoperative preparation for PGT	Parents had no <i>STK11</i> variation
5	Woman, 35	Pigmentation on the face and lips; multiple polyps	19	Twice minimally invasive surgery for intestinal polyps and laparotomy operation	c.114_121delGCGCCGCA (p.Arg39Alafs [*] 121)	Exon1, frame shift	Р	Yes, paternal. Father, uncle, brother	PGT	The fetus was prenatally diagnosed as having no mutation and gave birth to a child	Mother and son had no <i>STK11</i> variation
6	Woman, 33	Pigmentation on the lower lip, fingers, soles of the feet, and oral mucosa; multiple polyps	25	Polypectomy was performed in the small intestine and uterus	c.193G>T(p.Glu65 [*])	Exon1, nonsense	Р	De novo	PGT	Preoperative preparation for PGT	Parents had no <i>STK11</i> variation
7	Man, 45	Pigmentation on the lower lip, fingers, and oral mucosa; multiple polyps	22	Multiple gastrointestinal polypectomy	c.464+1G>A	Intron 3-4, splicing	LP	De novo	PGT	Preoperative preparation for PGT	Parents had no STK11 variation
8	Man, 23	Pigmentation on the lips	/	Laser treatment for hyperpig- mentation at 19 yr, no polyps found on gastrointestinal endoscopy	c.734+3A>T	Intron 5-6, splicing	LP	Yes, paternal. Father, grandfather, sister, and niece	/	Unmarried, under consideration	Father had STK11 variation
9	Woman, 24	Pigmentation on the limbs and trunk;	13	Multiple gastrointestinal polypectomy; termination	c.734+2T>C	Intron 5-6, splicing	Р	De novo	PGT	Preoperative preparation for PGT	Parents had no STK11

Table 1 Spectrum of STK11 mutations and clinical characteristics of patients with Peutz-Jeghers syndrome

multiple polyps

because fetal had STK11 mutation variation

ACMG: American Society for Medical Genetics and Genomics; P: Pathogenic; IVF-ET: In vitro fertilization-embryo transfer; VUS: Variant of uncertain significance; LB: Likely benign; PGT: Preimplantation genetic testing; LP: Likely pathogenic.

ARTICLE HIGHLIGHTS

Research background

Peutz-Jeghers syndrome (PJS) is a rare hereditary tumor disease with autosomal dominant inheritance that primarily results from mutations in the *STK11* gene. While certain missense variations of the *STK11* gene have been identified and classified as pathogenic or likely pathogenic, numerous missense variations still remain as variants of uncertain significance (VUS). This ambiguity makes it challenging to establish the association between these variations and PJS, which poses significant challenges for clinical treatment management and fertility selection.

Research motivation

Among the 9 PJS families evaluated at our hospital, 2 patients with PJS were identified as having missense variations through whole-exome sequencing and were classified as VUS. Understanding how the function of *STK11* is disrupted may contribute to a better understanding of the pathogenesis of PJS and the molecular mechanisms involved in the carcinogenesis process. Clarifying the pathogenicity of variations can offer patients the most appropriate reproductive guidance.

Research objectives

To clarify the pathogenicity of these two missense variations and to better provide patients with health management and fertility guidance.

Research methods

This study used whole-exome gene sequencing and Sanger sequencing methods to identify gene variants, combining bioinformatics analysis, quantitative polymerase chain reaction, immunoblotting, immunohistochemistry, and a variety of *in vitro* and *in vivo* functional assays to investigate pathogenicity.

Research results

Bioinformatics software analysis indicated that these two missense variants are deleterious. The phosphorylation levels of adenosine monophosphate-activated protein kinase (AMPK) and liver kinase B1 in the variant group were significantly lower than those in the wild-type group. Both missense *STK11* variants promoted the proliferation of HeLa cells. The expression of phosphorylated-AMPK was significantly lower in PJS patients with a missense variant in gastric, colon, and uterine polyps compared to non-PJS patients. These missense variants inactivate the *STK11* gene, disrupting the regulation of downstream AMPK proteins, thereby mediating the occurrence of PJS. As a result, they are interpreted as likely pathogenic, providing valuable information for patients in reproductive decision-making.

Research conclusions

These findings provide a basis for further research on and reanalysis of clinical decision making related to rare disease missense VUS.

Research perspectives

Further investigation into the role of STK11 gene in mediating PJS is required. Identifying the pathogenicity of more missense VUS is crucial to offer more accurate medical services for patients.

ACKNOWLEDGEMENTS

We thank the patients, their families, and the referring clinicians for their valuable contributions to this study.

FOOTNOTES

Author contributions: Liu J contributed to drafting of the manuscript; Wang A and Cheng HY contributed to clinical polyp treatment; Zeng SC and Zhang QJ contributed to experiments and analysis; Lu GX contributed to critical revision of the manuscript for important intellectual content; all authors read and approved the final manuscript.

Supported by the Natural Science Foundation of Hunan Province, China, No. 2023JJ30422.

Institutional review board statement: The study was reviewed and approved by the Ethics Committee of the Reproduction and Genetics Hospital of CITIC-Xiangya [Approval No. LL-SC-2016(008)].

Conflict-of-interest statement: All authors declared no conflicts of interest to disclose.

Data sharing statement: The data that support the findings of this study are available from the corresponding author upon reasonable request.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: https://creativecommons.org/Licenses/by-nc/4.0/

Country/Territory of origin: China

ORCID number: Jin Liu 0000-0002-4576-582X; Guang-Xiu Lu 0009-0002-4732-6756.

S-Editor: Chen YL L-Editor: A P-Editor: Zhang XD

REFERENCES

- Forte G, Cariola F, De Marco K, Manghisi A, Guglielmi FA, Armentano R, Lippolis G, Giorgio P, Simone C, Disciglio V. A novel STK11 1 gene mutation (c.388dupG, p.Glu130Glyfs*33) in a Peutz-Jeghers family and evidence of higher gastric cancer susceptibility associated with alterations in STK11 region aa 107-170. Genes Dis 2022; 9: 288-291 [PMID: 35224145 DOI: 10.1016/j.gendis.2021.11.002]
- Kirakosyan E, Lokhmatov M. High-Tech Diagnostic Methods and Enteroscopic Treatment of Children with Peutz-Jeghers Syndrome. Eur J 2 Pediatr Surg 2020; 30: 529-535 [PMID: 31770782 DOI: 10.1055/s-0039-3400286]
- Klimkowski S, Ibrahim M, Ibarra Rovira JJ, Elshikh M, Javadi S, Klekers AR, Abusaif AA, Moawad AW, Ali K, Elsayes KM. Peutz-Jeghers 3 Syndrome and the Role of Imaging: Pathophysiology, Diagnosis, and Associated Cancers. Cancers (Basel) 2021; 13 [PMID: 34680270 DOI: 10.3390/cancers132051211
- Monahan KJ, Bradshaw N, Dolwani S, Desouza B, Dunlop MG, East JE, Ilyas M, Kaur A, Lalloo F, Latchford A, Rutter MD, Tomlinson I, 4 Thomas HJW, Hill J; Hereditary CRC guidelines eDelphi consensus group. Guidelines for the management of hereditary colorectal cancer from the British Society of Gastroenterology (BSG)/Association of Coloproctology of Great Britain and Ireland (ACPGBI)/United Kingdom Cancer Genetics Group (UKCGG). Gut 2020; 69: 411-444 [PMID: 31780574 DOI: 10.1136/gutjnl-2019-319915]
- Kariv R, Dahary D, Yaron Y, Petel-Galil Y, Malcov M, Rosner G. Whole Genome Sequencing Applied in Familial Hamartomatous Polyposis 5 Identifies Novel Structural Variations. Genes (Basel) 2022; 13 [PMID: 36011318 DOI: 10.3390/genes13081408]
- McGarrity TJ, Amos CI, Baker MJ. Peutz-Jeghers Syndrome. 2001 Feb 23. In: GeneReviews® [Internet]. Seattle (WA): University of 6 Washington, Seattle; 1993- [PMID: 20301443]
- Wang Z, Liang L, Liu L, Wang Z, Wang Y, Yu Z, Wu B, Chen Y. Changes in the Gut Microbiome Associated with Intussusception in Patients 7 with Peutz-Jeghers Syndrome. Microbiol Spectr 2023; 11: e0281922 [PMID: 36719190 DOI: 10.1128/spectrum.02819-22]
- Wagner A, Aretz S, Auranen A, Bruno MJ, Cavestro GM, Crosbie EJ, Goverde A, Jelsig AM, Latchford A, Leerdam MEV, Lepisto A, 8



WJGO | https://www.wjgnet.com

Puzzono M, Winship I, Zuber V, Möslein G. The Management of Peutz-Jeghers Syndrome: European Hereditary Tumour Group (EHTG) Guideline. J Clin Med 2021; 10 [PMID: 33513864 DOI: 10.3390/jcm10030473]

- 9 Liu S, Ma Y, You W, Li J, Li JN, Qian JM. Hamartomatous polyposis syndrome associated malignancies: Risk, pathogenesis and endoscopic surveillance. J Dig Dis 2021; 22: 444-451 [PMID: 34145757 DOI: 10.1111/1751-2980.13029]
- 10 Tian LL, Guo JZ, Yin YQ, Dang XH, Huo LJ. Analysis of a pedigree of Peutz-Jeghers syndrome and RET proto-oncogene mutation: one case report and literature review. Transl Cancer Res 2020; 9: 3007-3011 [PMID: 35117658 DOI: 10.21037/tcr.2020.02.64]
- Altamish M, Dahiya R, Singh AK, Mishra A, Aljabali AAA, Satija S, Mehta M, Dureja H, Prasher P, Negi P, Kapoor DN, Goyal R, 11 Tambuwala MM, Chellappan DK, Dua K, Gupta G. Role of the Serine/Threonine Kinase 11 (STK11) or Liver Kinase B1 (LKB1) Gene in Peutz-Jeghers Syndrome. Crit Rev Eukaryot Gene Expr 2020; 30: 245-252 [PMID: 32749111 DOI: 10.1615/CritRevEukaryotGeneExpr.2020033451]
- 12 Li L, Yao Y, Zhao J, Cao J, Ma H. Dehydroepiandrosterone protects against hepatic glycolipid metabolic disorder and insulin resistance induced by high fat via activation of AMPK-PGC-1α-NRF-1 and IRS1-AKT-GLUT2 signaling pathways. Int J Obes (Lond) 2020; 44: 1075-1086 [PMID: 31911660 DOI: 10.1038/s41366-019-0508-8]
- Zyla RE, Hahn E, Hodgson A. Gene of the month: STK11. J Clin Pathol 2021; 74: 681-685 [PMID: 34479873 DOI: 13 10.1136/jclinpath-2021-207906]
- Beggs AD, Latchford AR, Vasen HF, Moslein G, Alonso A, Aretz S, Bertario L, Blanco I, Bülow S, Burn J, Capella G, Colas C, Friedl W, 14 Møller P, Hes FJ, Järvinen H, Mecklin JP, Nagengast FM, Parc Y, Phillips RK, Hyer W, Ponz de Leon M, Renkonen-Sinisalo L, Sampson JR, Stormorken A, Tejpar S, Thomas HJ, Wijnen JT, Clark SK, Hodgson SV. Peutz-Jeghers syndrome: a systematic review and recommendations for management. Gut 2010; 59: 975-986 [PMID: 20581245 DOI: 10.1136/gut.2009.198499]
- 15 Richards S, Aziz N, Bale S, Bick D, Das S, Gastier-Foster J, Grody WW, Hegde M, Lyon E, Spector E, Voelkerding K, Rehm HL; ACMG Laboratory Quality Assurance Committee. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. Genet Med 2015; 17: 405-424 [PMID: 25741868 DOI: 10.1038/gim.2015.30]
- Long LL, Ma SC, Guo ZQ, Zhang YP, Fan Z, Liu LJ, Liu L, Han DD, Leng MX, Wang J, Guo XJ, Tan JL, Cai XT, Lin Y, Pan X, Wu DH, 16 Bai X, Dong ZY. PARP Inhibition Induces Synthetic Lethality and Adaptive Immunity in LKB1-Mutant Lung Cancer. Cancer Res 2023; 83: 568-581 [PMID: 36512628 DOI: 10.1158/0008-5472.CAN-22-1740]
- Hanks SK, Quinn AM, Hunter T. The protein kinase family: conserved features and deduced phylogeny of the catalytic domains. Science 17 1988; 241: 42-52 [PMID: 3291115 DOI: 10.1126/science.3291115]
- 18 Islam MJ, Khan AM, Parves MR, Hossain MN, Halim MA. Prediction of Deleterious Non-synonymous SNPs of Human STK11 Gene by Combining Algorithms, Molecular Docking, and Molecular Dynamics Simulation. Sci Rep 2019; 9: 16426 [PMID: 31712642 DOI: 10.1038/s41598-019-52308-01
- 19 Huber-Keener KJ. Cancer genetics and breast cancer. Best Pract Res Clin Obstet Gynaecol 2022; 82: 3-11 [PMID: 35272929 DOI: 10.1016/j.bpobgyn.2022.01.007]
- Li YX, LV XS, Xia JH. Mutation characteristic of STK11 gene in Chinese with Peutz-Jeghers syndrome. Zhonghua Yixue Yichuanxue Zazhi 20 2001; **18**: 4-7 [DOI: 10.1186/1471-2350-12-161]
- Frebourg T. The challenge for the next generation of medical geneticists. Hum Mutat 2014; 35: 909-911 [PMID: 24838402 DOI: 21 10.1002/humu.22592
- Brito S, Póvoas M, Dupont J, Lopes AI. Peutz-Jeghers syndrome: early clinical expression of a new STK11 gene variant. BMJ Case Rep 2015; 22 2015 [PMID: 26430231 DOI: 10.1136/bcr-2015-211345]
- Landrum MJ, Chitipiralla S, Brown GR, Chen C, Gu B, Hart J, Hoffman D, Jang W, Kaur K, Liu C, Lyoshin V, Maddipatla Z, Maiti R, 23 Mitchell J, O'Leary N, Riley GR, Shi W, Zhou G, Schneider V, Maglott D, Holmes JB, Kattman BL. ClinVar: improvements to accessing data. Nucleic Acids Res 2020; 48: D835-D844 [PMID: 31777943 DOI: 10.1093/nar/gkz972]
- 24 Ataei-Kachouei M, Nadaf J, Akbari MT, Atri M, Majewski J, Riazalhosseini Y, Garshasbi M. Double Heterozygosity of BRCA2 and STK11 in Familial Breast Cancer Detected by Exome Sequencing. Iran J Public Health 2015; 44: 1348-1352 [PMID: 26576347]
- Gunawardena K, Sirisena ND, Anandagoda G, Neththikumara N, Dissanayake VHW. Germline variants of uncertain significance, their 25 frequency, and clinico-pathological features in a cohort of Sri Lankan patients with hereditary breast cancer. BMC Res Notes 2023; 16:95 [PMID: 37277882 DOI: 10.1186/s13104-023-06365-4]
- van Lier MG, Mathus-Vliegen EM, van Leerdam ME, Kuipers EJ, Looman CW, Wagner A, Vanheusden K. Quality of life and psychological 26 distress in patients with Peutz-Jeghers syndrome. Clin Genet 2010; 78: 219-226 [PMID: 20695872 DOI: 10.1111/j.1399-0004.2010.01469.x]
- 27 Jelsig AM, van Overeem Hansen T, Gede LB, Qvist N, Christensen LL, Lautrup CK, Frederiksen JH, Sunde L, Ousager LB, Ljungmann K, Bertelsen B, Karstensen JG. Survival, surveillance, and genetics in patients with Peutz-Jeghers syndrome: A nationwide study. Clin Genet 2023; **104**: 81-89 [PMID: 37017260 DOI: 10.1111/cge.14337]
- Wang S, Huang G, Wang JX, Tian L, Zuo XL, Li YQ, Yu YB. Altered Gut Microbiota in Patients With Peutz-Jeghers Syndrome. Front 28 Microbiol 2022; 13: 881508 [PMID: 35910641 DOI: 10.3389/fmicb.2022.881508]
- Villy MC, Frydman N, Moutou C, Thierry G, Raad J, Colas C, Steffann J, Metras J, Chabbert-Buffet N, Parc Y, Richard S, Benusiglio PR. 29 Preimplantation genetic testing in patients with genetic susceptibility to cancer. Fam Cancer 2023; 22: 119-125 [PMID: 35415820 DOI: 10.1007/s10689-022-00293-11
- Xu X, Song R, Hu K, Li Y, Jin H, Chen B, Song W, Zhang Y, Xu J, Sun Y. Multidisciplinary management for Peutz-Jeghers syndrome and 30 prevention of vertical transmission to offspring using preimplantation genetic testing. Orphanet J Rare Dis 2022; 17: 64 [PMID: 35189935 DOI: 10.1186/s13023-022-02221-z]
- 31 Wang Z, Liu S, Wang Y, Chen J, Wu B. Prenatal diagnosis in a hereditary Peutz-Jeghers syndrome family with high cancer risk. BMC Med Genet 2018; 19: 66 [PMID: 29720104 DOI: 10.1186/s12881-018-0594-9]
- Leaver M, Wells D. Non-invasive preimplantation genetic testing (niPGT): the next revolution in reproductive genetics? Hum Reprod Update 32 2020; 26: 16-42 [PMID: 31774124 DOI: 10.1093/humupd/dmz033]
- Sullivan-Pyke C, Dokras A. Preimplantation Genetic Screening and Preimplantation Genetic Diagnosis. Obstet Gynecol Clin North Am 2018; 33 45: 113-125 [PMID: 29428279 DOI: 10.1016/j.ogc.2017.10.009]
- 34 Byrjalsen A, Roos L, Diemer T, Karstensen JG, Løssl K, Jelsig AM. Preimplantation genetic testing in two Danish couples affected by Peutz-Jeghers syndrome. Scand J Gastroenterol 2023; 58: 314-318 [PMID: 36200740 DOI: 10.1080/00365521.2022.2129031]





Published by Baishideng Publishing Group Inc 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA Telephone: +1-925-3991568 E-mail: office@baishideng.com Help Desk: https://www.f6publishing.com/helpdesk https://www.wjgnet.com

