

World Journal of *Stem Cells*

World J Stem Cells 2024 February 26; 16(2): 54-227



Contents

Monthly Volume 16 Number 2 February 26, 2024

EDITORIAL

- 54 Human dental pulp stem/stromal cells in clinical practice
Grawish ME
- 58 Multiple pretreatments can effectively improve the functionality of mesenchymal stem cells
Wan XX, Hu XM, Xiong K
- 64 Cellular preconditioning and mesenchymal stem cell ferroptosis
Zineldeen DH, Mushtaq M, Haider KH

REVIEW

- 70 Therapeutic utility of human umbilical cord-derived mesenchymal stem cells-based approaches in pulmonary diseases: Recent advancements and prospects
Meng M, Zhang WW, Chen SF, Wang DR, Zhou CH
- 89 Unlocking the versatile potential: Adipose-derived mesenchymal stem cells in ocular surface reconstruction and oculoplastics
Surico PL, Scarabosio A, Miotti G, Grando M, Salati C, Parodi PC, Spadea L, Zeppieri M

MINIREVIEWS

- 102 Crosstalk between Wnt and bone morphogenetic protein signaling during osteogenic differentiation
Arya PN, Saranya I, Selvamurugan N
- 114 Human pluripotent stem cell-derived kidney organoids: Current progress and challenges
Long HY, Qian ZP, Lan Q, Xu YJ, Da JJ, Yu FX, Zha Y
- 126 Recent progress in hair follicle stem cell markers and their regulatory roles
Xing YZ, Guo HY, Xiang F, Li YH
- 137 Advances in the differentiation of pluripotent stem cells into vascular cells
Jiao YC, Wang YX, Liu WZ, Xu JW, Zhao YY, Yan CZ, Liu FC

ORIGINAL ARTICLE

Basic Study

- 151 Silencing of Jumonji domain-containing 1C inhibits the osteogenic differentiation of bone marrow mesenchymal stem cells *via* nuclear factor- κ B signaling
Li JY, Wang TT, Ma L, Zhang Y, Zhu D
- 163 Effects of different concentrations of nicotinamide on hematopoietic stem cells cultured *in vitro*
Ren Y, Cui YN, Wang HW

- 176** High quality repair of osteochondral defects in rats using the extracellular matrix of antler stem cells
Wang YS, Chu WH, Zhai JJ, Wang WY, He ZM, Zhao QM, Li CY
- 191** Extracellular vesicles derived from mesenchymal stem cells mediate extracellular matrix remodeling in osteoarthritis through the transport of microRNA-29a
Yang F, Xiong WQ, Li CZ, Wu MJ, Zhang XZ, Ran CX, Li ZH, Cui Y, Liu BY, Zhao DW
- 207** VX-509 attenuates the stemness characteristics of colorectal cancer stem-like cells by regulating the epithelial-mesenchymal transition through Nodal/Smad2/3 signaling
Yuan Y, Zhang XF, Li YC, Chen HQ, Wen T, Zheng JL, Zhao ZY, Hu QY

ABOUT COVER

Editorial Board Member of *World Journal of Stem Cells*, Khawaja Husnain Haider, PhD, Professor, Department of Basic Sciences, Sulaiman Al-Rajhi University, AlQaseem 52726, Saudi Arabia. kh.haider@gmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Stem Cells (WJSC, World J Stem Cells)* is to provide scholars and readers from various fields of stem cells with a platform to publish high-quality basic and clinical research articles and communicate their research findings online. *WJSC* publishes articles reporting research results obtained in the field of stem cell biology and regenerative medicine, related to the wide range of stem cells including embryonic stem cells, germline stem cells, tissue-specific stem cells, adult stem cells, mesenchymal stromal cells, induced pluripotent stem cells, embryonal carcinoma stem cells, hemangioblasts, lymphoid progenitor cells, *etc.*

INDEXING/ABSTRACTING

The *WJSC* is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Journal Citation Reports/Science Edition, PubMed, PubMed Central, Scopus, Biological Abstracts, BIOSIS Previews, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for *WJSC* as 4.1; IF without journal self cites: 3.9; 5-year IF: 4.5; Journal Citation Indicator: 0.53; Ranking: 15 among 29 journals in cell and tissue engineering; Quartile category: Q3; Ranking: 99 among 191 journals in cell biology; and Quartile category: Q3. The *WJSC*'s CiteScore for 2022 is 8.0 and Scopus CiteScore rank 2022: Histology is 9/57; Genetics is 68/325; Genetics (clinical) is 19/90; Molecular Biology is 119/380; Cell Biology is 95/274.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Xiang-Di Zhang; Production Department Director: Xu Guo; Editorial Office Director: Jia-Ru Fan.

NAME OF JOURNAL

World Journal of Stem Cells

ISSN

ISSN 1948-0210 (online)

LAUNCH DATE

December 31, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Shengwen Calvin Li, Carlo Ventura

EDITORIAL BOARD MEMBERS

<https://www.wjnet.com/1948-0210/editorialboard.htm>

PUBLICATION DATE

February 26, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Recent progress in hair follicle stem cell markers and their regulatory roles

Yi-Zhan Xing, Hai-Ying Guo, Fei Xiang, Yu-Hong Li

Specialty type: Cell and tissue engineering

Provenance and peer review: Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Wang X, China

Received: October 27, 2023

Peer-review started: October 27, 2023

First decision: December 5, 2023

Revised: December 19, 2023

Accepted: January 16, 2024

Article in press: January 16, 2024

Published online: February 26, 2024



Yi-Zhan Xing, Hai-Ying Guo, Yu-Hong Li, Department of Cell Biology, Army Medical University, Chongqing 400038, China

Fei Xiang, Institute of Burn Research, Southwest Hospital, Army Medical University, Chongqing 400038, China

Corresponding author: Yu-Hong Li, PhD, Associate Professor, Department of Cell Biology, Army Medical University, No. 30 Gaotanyan Street, Shapingba District, Chongqing 400038, China. liyuhongtmmu@hotmail.com

Abstract

Hair follicle stem cells (HFSCs) in the bulge are a multipotent adult stem cell population. They can periodically give rise to new HFs and even regenerate the epidermis and sebaceous glands during wound healing. An increasing number of biomarkers have been used to isolate, label, and trace HFSCs in recent years. Considering more detailed data from single-cell transcriptomics technology, we mainly focus on the important HFSC molecular markers and their regulatory roles in this review.

Key Words: Hair follicle stem cells; Bulge; Secondary hair germ; Marker; Single-cell RNA-sequencing

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Hair follicle stem cells (HFSCs) in the bulge are a multipotent adult stem cell population. They can periodically give rise to new HFs and even regenerate the epidermis and sebaceous glands during wound healing. An increasing number of biomarkers have been used to isolate, label, and trace HFSCs in recent years. Considering more detailed data from single-cell transcriptomics technology, we mainly focus on the important HFSC molecular markers and their regulatory roles in this review.

Citation: Xing YZ, Guo HY, Xiang F, Li YH. Recent progress in hair follicle stem cell markers and their regulatory roles. *World J Stem Cells* 2024; 16(2): 126-136

URL: <https://www.wjgnet.com/1948-0210/full/v16/i2/126.htm>

DOI: <https://dx.doi.org/10.4252/wjsc.v16.i2.126>

INTRODUCTION

Hair follicle (HF) is an appendage of mammalian skin that develops from the interaction of dermal mesenchymal cells and epithelial cells during embryonic development[1,2]. Postnatal HFs undergo cycles of growth (anagen), regression (catagen), and rest (telogen), namely, the hair cycle[3]. The dysfunction of hair regeneration may lead to several diseases. For example, HFs become miniaturization in androgenic alopecia[4]. HF stem cells (HFSCs) reside in the bulge region near the attachment site of the arrector pili muscle and drive periodical hair regeneration. Usually, the new HFs are oriented from HFSCs. Thus, HFSCs may be useful in treating hair loss-related diseases. The regulation of the activation and quiescence of HFSCs is a critical dimension in this research area[5]. The concentric layers of HFs are composed of the outer root sheath (ORS), inner root sheath (IRS), and hair shaft (HS)[6]. Dermal papilla (DP) is a condensed mesenchymal cell compartment at the base of the hair bulb and is important for the activation of HFSCs[7]. There is a secondary group of cells named the secondary hair germ (HG) that resides between the bulge and the DP and becomes activated as the telogen progresses. Although derived from bulge SCs, HG cells respond more quickly to stimulus signals. During early anagen, there is a two-step activation process. First, HG cells become activated before bulge cells[8]. Second, these cells undergo several self-renewals and proliferate downward to form the ORS. In mature HFs, the ORS extends from the bulge to the matrix (Mx) bottom. Mx cells undergo rapid cell growth and differentiation to form new hairs. During the degenerative phase, the lower keratinocytes of HFs begin to undergo apoptosis, and the residual epithelial strands shrink, which pulls the DP upward. At the transition of catagen to telogen, some mid-zone ORS cells survive and generate HG, while the surviving upper slow-cycling ORS cells eventually form a new bulge[8,9]. The stages of HF cycle and the roles played by critical cell types are summarized in Figure 1.

Since the discovery of slow cycling, label-retaining cells in the outermost layer of the bulge region[10], the characteristics and activity of multipotent HFSCs have been intensively studied in recent decades[11-14]. Various markers have been successfully used for the purification and enrichment of HFSCs. Multiple markers have been used to trace HFSCs, and to some extent, there was some overlap and differences among these marked cell populations. Here, we focus on some important biomarkers of HFSCs and their regulatory roles in HFSCs (Table 1).

LEUKOCYTE DIFFERENTIATION ANTIGEN 34

Leukocyte differentiation antigen 34 (CD34) is a transmembrane glycoprotein initially identified on hematopoietic stem and progenitor cells[15]. The expression of CD34 is substantially localized in the outer layer bulge region and HG in mouse HFs[14,16]. In anagen, CD34 expression levels decrease in HFs[17] and at the most peripheral layer of the ORS between the isthmus and the bulb[18]. At present, CD34 is the most commonly used indicator in tracing or sorting bulge SCs and is often combined with $\alpha 6$ -integrin (or CD49f)[16,18]. These isolated cells were characterized as having the ability for long-term self-renewal and pluripotent differentiation[19]. However, CD34 is not found in the bulge of human HFs during catagen or telogen[20-22].

CD200

CD200, formerly known as OX-2, is a type-1 transmembrane glycoprotein that mediates an immunoregulatory signal through binding to the CD200 receptor. CD200+ cells are observed in the bulge region of murine[23] and human HFs[20, 22,24,25] and colocalize with keratin 14 (K14). In murine HFs, CD200 expression is relatively uniform throughout the whole HF, except for the Mx, DP, and HS[23]. Deficiency of CD200 in C57BL/6 mice led to significant perifollicular and intrafollicular inflammation in skin grafts and an increase in graft-infiltrating T cells after 14 d[23]. Alopecia areata is a common hair loss disease characterized by cellular autoimmune reactions. In older women with alopecia areata, CD200 expression was downregulated in the affected bulge lesions[26]. These results suggest that CD200 may play a role in attenuating inflammatory reactions and promoting immune tolerance.

K15

K15, an intermediate filament protein, is expressed in the epidermal basal layer and HF bulge in neonatal mice[27]. However, the expression level of K15 significantly decreases at approximately 2 wk of age, while strong expression of K15 is still observed in the bulge cells of mature mouse skin, similar to the human scalp[28]. In addition, all four types of HFs (guard, awl, auchene, and zigzag) show strong staining of K15 protein in the bulge cells of the dorsal skin in mice.

Table 1 Expression profile in postnatal hair cycle and related functions of some hair follicle stem cell markers

	Expression location in telogen	Expression location in anagen	Related functions	Ref.
CD34	Outer layer of bulge and HG (but not found in human HF)	Most peripheral layer of the ORS	Sustain the stemness of HFSCs	[14-22]
CD200	Bulge and HG	Relatively uniform throughout the whole HF, except for the Mx, DP and HS	Sustain the stemness of HFSCs	[20,22-26]
K15	Epidermal basal layer and HF bulge in neonatal mice; still strong in bulge in mature mice skin	ORS, the bulge is the strongest area	Not found	[22,27,28]
K19	Bulge	Mainly the U-ORS and the pb-ORS of HFs, stronger within the pb-ORS versus the bulge	Not found	[27,29-31]
Lgr5	Bulge, stronger in HG	The lower portion of the ORS, not overlapped with CD34	In the telogen-to-anagen transition	[32-38]
Lhx2	Bulge and HG	Through the ORS	Epidermal differentiation, HFSC quiescence, hair anchoring	[31,39-41]
Sox9	Bulge and HG	ORS, stronger in the sub-bulge region	Maintainance and differentiation of HFSCs	[12,31,42,43]
Runx1	Bulge, stronger in HG	The infundibulum, bulge, ORS, Mx and cortex	Regulating anagen initiation and HFSC proliferation	[12,42,44-50]
Axin2	Outer bulge cells	ORS	Maintainance of hair-forming ability	[51-60]
Nfatc1	Bulge	Bulge, overlapped with CD34 and Lhx2	Maintainance of HFSC quiescence	[46,61-65]
Foxp1	Bulge	Bulge, ORS, Mx	Maintainance of HFSC quiescence	[66-70]
Foxc1	Bulge, isthmus, sebaceous glands	Bulge, K6+ inner bulge layer, isthmus, IRS and sebaceous glands	Maintainance of the old bulge	[16,40,71-73]
Msi2	Basal layer of bulge and HG	Basal and suprabasal bulge, ORS and IRS	In the telogen-to-anagen transition	[74-77]

CD34: Leukocyte differentiation antigen 34; CD200: Leukocyte differentiation antigen 200; K15: Keratin 15; K19: Keratin 19; Lgr5: Leucine-rich repeat-containing G protein-coupled receptor 5; Lhx2: Lim-homeodomain transcription factor 2; Sox9: Sex-determining region Y-Box 9; Runx1: Runx family transcription factor 1; Axin2: Axin-related protein 2; Nfatc1: Nuclear factor of activated T cells c1; Foxp1: Forkhead Box P1; Foxc1: Forkhead Box C1; Msi2: Musashi 2; Mx: Matrix; HFSC: Hair follicle stem cell; HS: Hair shaft; HG: Hair germ; DP: Dermal papilla; ORS: Outer root sheath; IRS: Inner root sheath; pb-ORS: Proximal bulb outer root sheath; U-ORS: Upper outer root sheath.

Previously, K5 or K14 promoters were usually used to target epithelial SCs in the skin, which traces not only epidermal basal SCs but also HFSCs and even transient amplifying (TA) cells. Thus, the K15 promoter is presently used to selectively target HFSCs in the bulge[22].

K19

K19, another Keratin protein, is expressed in different stages of the hair cycle[27,29,30]. In anagen, the expression of K19 is mainly restricted to two areas, one in the upper ORS (U-ORS) and the other in the proximal bulb ORS (pb-ORS) of HFs. The K19 staining intensity is stronger in the pb-ORS than in the basal layer of the bulge, and the opposite is true for K15 [31]. K15+ and K19+ cells represent two distinct progenitor cell populations located in the bulge and pb-ORS. These K19+ cells in the pb-ORS are K15- and Ki67+[27,29,31], which indicates that this cell population contains more differentiated TA cells. In catagen, the K19+ region of pb-ORS moves upward with the club hair; eventually, it fuses with the K19+ portion of U-ORS during telogen. These two regions are separated again at the beginning of the next growth period.

LEUCINE-RICH REPEAT-CONTAINING G PROTEIN-COUPLED RECEPTOR 5

Leucine-rich repeat-containing G protein-coupled receptor 5 (Lgr5) is an orphan seven-pass transmembrane protein located on the cell membrane. Lgr5 is a marker of SCs at the bottom of small intestinal crypts[32,33]. In the E18.5 Lgr5^{LacZ} mouse model, Lgr5+ cells are observed in the ORS of larger HFs. During anagen and catagen, Lgr5-LacZ expression is located in the lower ORS up to Mx around the HFs, while in telogen, the expression of Lgr5 is concentrated in the basal




Figure 1 Hair follicle stem cells and the hair cycle. Telogen: The length of the hair follicle (HF) is the shortest. HF stem cells (HFSCs) remain quiescent until dermal papilla (DP) releases a new round of stimulations, which activate the hair germ cells at first, then HFSCs. Anagen: The activated HFSCs proliferate and differentiate, forming the outer root sheath and matrix via down growth. After that, the inner root sheath and hair shaft are formed from the matrix. Catagen: The cells in the matrix undergo apoptosis and form an epithelial strand, which retracts the DP. The HF shrinks gradually until entering the next telogen. HS: Hair shaft; SG: Sebaceous gland; HG: Hair germ; DP: Dermal papilla; IRS: Inner root sheath; ORS: Outer root sheath.

cells of the bulge and HG, especially in the HG[34].

Through fluorescence-activated cell sorting (FACS) and flow cytometry, Jaks *et al*[34] analyzed Lgr5-LacZ expression in mouse keratinocytes during telogen and compared it with other known markers. The results showed that approximately 80% of keratinocytes had medium to high levels of $\alpha 6$ -integrin expression, whereas Lgr5^{high} cells composed 0.6% of the $\alpha 6$ -integrin population. Moreover, 80% of Lgr5⁺ cells overlapped with the CD34⁺ cell population in telogen HFs, which constituted only 10% of CD34⁺ cells. However, CD34⁺ and Lgr5⁺ cells, found in the lower ORS, had no overlap in anagen HFs[34,35]. Similarly, K15⁺ cells showed a partial overlap with Lgr5⁺ cells in the bulge in telogen but not in anagen[34]. Thus, Lgr5⁺ cells constitute a particular cell compartment in mouse HFs.

Lgr5⁺ cells are essential for the telogen-to-anagen transition. Hoeck *et al*[35] utilized diphtheria toxin-mediated Lgr5⁺ SC ablation in mouse HFs and showed that loss of Lgr5⁺ cells abolished hair regeneration, but this effect was reversible. During recovery, CD34⁺ SCs can activate inflammatory response programs and begin to divide. When the SC population was restored, new Lgr5⁺ cells were generated from CD34⁺ SCs. This program is dependent on the Wnt signal.

Lgr5 has been identified as a target of the Wnt signaling pathway. R-spondin, a secreted agonist of the canonical Wnt/ β -catenin signal, binds to Lgr receptors and stabilizes them[36]. Through intradermal injection of recombinant R-spondin2 protein, the combination of R-spondin and Lgr5 prevents the degradation of β -catenin and improves Wnt signaling[37]. This results in persistent anagen and longer HSs.

After injury, the Lgr5⁺ bulge cells were sorted and sequenced through lineage tracing. Although the progeny of Lgr5 cells responded and contributed to wound healing, they gradually lost their bulge signature and expressed basal interfollicular epidermis-related genes[38].

LIM-HOMEODOMAIN TRANSCRIPTION FACTOR 2

Lim-homeodomain transcription factor 2 (Lhx2) is a key regulator of bulge SCs[39,40]. During mouse embryogenesis, Lhx2 is first expressed in the early hair placode. As morphogenesis progresses, Lhx2 gradually localizes in the HG and peg precursor cells[40]. At P0, Lhx2 expression is mainly concentrated in the bulge area and is downregulated in the ORS and Mx. At P44, when HFs are in telogen, Lhx2 expression persists in the bulge and secondary HG[41].

Purba *et al*[31] demonstrated that Lhx2⁺ cells were throughout the ORS, including the basal and adjacent suprabasal layers of occipital scalp HFs in humans, but not in the innermost ORS layers. Lhx2⁺ bulge cells simultaneously express K15 and CD200. However, the pb-ORS and the sub-bulge cells have brighter fluorescence and more Lhx2⁺ cell numbers than bulge cells. The Lhx2⁺ subpopulation is distinct from the sex-determining region Y-Box 9 (Sox9⁺) cells. Most of the

cells in the sub-bulge and the pb-ORS are either Lhx2+ or Sox9+. Only 17% of these cells are Lhx2 and Sox9 double positive, which reveals that these cells display distinct progenitor cell populations in the ORS. When HF enter catagen, Lhx2 expression is upregulated in the proximal hair bulb epithelium.

Rhee *et al*[40] reported that Lhx2 remarkably suppressed epidermal differentiation in K14-Lhx2 transgenic mice. However, E16 Lhx2 null embryos displayed a marked reduction in HF density. Lhx2 KO HF exhibited fewer label-retaining cells, enhanced proliferative activity within the bulge, and shortened telogen. In addition, Lhx2 heterozygous mutations lead to slower wound healing[39]. K14 promoter-mediated conditional knockout of Lhx2 causes the inability to maintain HFSC quiescence and hair anchoring. Therefore, when Lhx2 is absent, the functioning of HFSCs is seriously impaired.

SOX 9

Sox9, a member of the Sox gene family, is characterized by the presence of high mobility group boxes. Sox9 is also an important transcription factor in HFSC biology and has a similar expression pattern to Lhx2 in murine and human HFSCs [31]. Sox9 expression was first found in the epithelial component of the hair placode. However, the nuclear Sox9 protein is located in the suprabasal cells of the hair placode rather than in the basal layer, which is P-cadherin+ and Lhx2+[42]. At the peg stage, Sox9+ cells are concentrated in an U-ORS region, which exhibits moderate expression of P-cadherin.

In the postnatal hair cycle, Sox9+ cells extend down the ORS and diminish toward the Mx. Cells exhibiting nuclear Sox9 immunoreactivity are mainly distributed in the innermost, differentiated ORS layer and partially distributed in the basal layer. The most apparent Sox9+ region is the sub-bulge region, just below the K15+ bulge compartment, while the Sox9+ population partially overlaps with K15+ cells in the pb-ORS. In catagen, Sox9+ cells are evident in the pb-ORS but not in the bulge area. At P21, Sox9+ cells are concentrated in the CD34+ bulge and HG[12].

Vidal *et al*[42] constructed conditionally targeted Y10:Cre/Sox9(fl/fl) mice, which showed retained atrophic hair coats. Nowak *et al*[12] targeted Sox9 ablation using K14-Cre and further investigated the role of Sox9 in regulating HF morphogenesis and the postnatal hair cycle. Before HF morphogenesis, Sox9 loss leads to the absence of bulge SCs. Although it has no impact on the differentiation of epithelial cells at first, due to the lack of SCs, Mx cells could not be replenished. Eventually, it still affected hair production. To understand whether Sox9 functions in HFSC biology, Kadaja *et al*[43] conditionally targeted Sox9 in the HFSCs of adult mice. Once activated, Sox9-deficient HFSCs differentiate along the epidermal lineage. This differentiation leads to premature termination of the downward growth of the HF, which is related to the bulge and U-ORS. Genome-wide RNA sequencing profiling and immunofluorescence results further show that Sox9-deficient bulge cells lose SC characteristics. Overall, Sox9 is indispensable for maintaining the identity of HFSCs.

RUNX FAMILY TRANSCRIPTION FACTOR 1

Runx family transcription factor 1 (Runx1), also known as Aml1, is a master regulator of hematopoietic SCs[44]. It regulates the development and homeostasis of multiple tissues. During HF development, Runx1 expression is mainly located in the mesenchyme. At E16.5, X-Gal-positive cells can be detected in the hair placode and germ cells of the Runx1-LacZ mouse embryonic skin, with stronger staining in the upper HF cells[45]. The Runx1 expression pattern is similar to Sox9 and nuclear factor of activated T cells c1 (Nfatc1) in the embryonic epithelium[12,42,46]. The results of lineage tracing experiments showed that cells expressing Runx1 in the embryonic stage contribute to postnatal mesenchymal skin and unique epithelial HF populations.

In the anagen phase, Runx1 is expressed in the infundibulum, bulge, ORS, Mx, and cortex, in contrast to the lower ORS in the catagen phase. There is no expression in the interfollicular epidermis. In the telogen phase, Runx1 is localized to the bulge and HG, and HG cells have the highest expression levels[45]. The Runx1 protein is expressed in the bulge, HS, and IRS of human skin, which is similar to the Runx1 protein expression in mice[47].

To interpret the effect of Runx1 on HFSCs, Osorio *et al*[48] used K14-driven Cre mice to conditionally ablate Runx1 in epithelial cells during morphogenesis. Runx1 ablation mice exhibited blocked hair regeneration due to prolonged telogen. Runx1 disruption affected the activation of HFSCs, including reductions in colony formation and BrdU+ cell numbers during the telogen-anagen transition *in vitro*. If deletion of Runx1 is performed after the end of morphogenesis, it also delays anagen onset[49]. Runx1 loss affects the cell cycle progression of cultured CD34+/α6+-sorted bulge cells by upregulating the cyclin-dependent kinase (CDK) inhibitor Cdkn1a (P21). In addition, it has been reported that Runx1 inhibits the transcription of P21, P27, P57, and P15 in HFSCs *in vivo* and synergistically regulates the resting degree of HFSCs with P21[50]. Therefore, Runx1 may act as a direct regulator of anagen initiation and HFSC proliferation.

AXIN-RELATED PROTEIN 2

Axin-related protein 2 (Axin2), a scaffold protein, can participate in the formation of β-catenin degradation complexes[51-53]. Wnt/β-catenin signaling plays an important role in the development, maintenance, proliferation, and lineage determination of HFSCs[54-58]. As a Wnt target gene, Axin2 is a sensitive Wnt reporter of Wnt/β-catenin signaling in HFSCs[59]. During embryonic development of the skin epithelium, Axin2-expressing cells are located in the placode and represent

hair cell progenitors. Then, Axin2 expression is gradually lost until the adult HFSC compartment is established, in which Axin2 is expressed again.

By using RNA *in situ* hybridization and Axin2-LacZ mice, it was found that Axin2 is specifically expressed in outer bulge (OB) cells in telogen. Flow cytometry analysis confirmed that these cells exhibited a G0/G1 state. Tracking the fate of these cells and their offspring also showed that Axin2+ cells were labeled with long-lived external bulge HFSCs. During the anagen phase, Axin2 in the bulge zone is also continuously expressed[60].

The β -catenin gene was conditionally deleted in Axin2+ cells in Axin2-CreERT2/ β -cat ^{Δ ex2-6-fl/fl} mutant mice[60]. This deletion led to a decrease in Axin2 mRNA expression levels in the mutant bulge cells. Moreover, Axin2-CreERT2/ β -cat ^{Δ ex2-6-fl/fl} mutant HFs failed to grow and showed abnormal telogen-like morphology. These results suggest that bulge HFSCs require β -catenin for the expression of Axin2 and to maintain their hair-forming ability.

NFATC1

Nfatc1, a transcription factor, is a member of the NFAT family, which plays key roles in many cellular processes[61,62]. Calcineurin controls the translocation of NFAT proteins from the cytoplasm to the nucleus of activated cells. Using microarray profiling, Nfatc1 was identified as an upregulated gene that can distinguish embryonic hair buds in the epidermis[63,64].

Nuclear Nfatc1 is first detected during late HF morphogenesis. With the maturation of HFs, Nfatc1 expression is also positive in the intermediate segment. In the postnatal hair cycle, cells expressing Nfatc1 persist in the bulge not only in growing HFs but also in telogen HFs. Nuclear Nfatc1 is specifically expressed by HFSCs. Compared to other bulge markers, Nfatc1 colocalizes with CD34 and Lhx2 and partially overlaps with Tcf3 and Sox9, with no Nfatc1-positive cells in the lower ORS. Nfatc1 protein and mRNA appear to be specific to bulge cells in both embryonic and adult skin[46].

After transplantation, HFs in the skin of Nfatc1 null mice can form and reenter telogen, but the telogen stage is shortened and the cells subsequently enter anagen early. Therefore, Nfatc1 has an important regulatory effect on the resting state of HFSCs in the bulge. In K14 promoter-driven Nfatc1 knockout mice, HFs could also develop normally, but most of the HFs entered the growth phase at P56, and 75% of the HFs had BrdU-labeled HFs. However, the HFs of wild-type mice are still in telogen until P75[46]. These results suggest that in the absence of Nfatc1, the slow-cycle nature of HFSCs is selectively lost. Horsley *et al*[46] further investigated the effect of Nfatc1 on the cell cycle and found that it downregulated the expression of CDK4 in HFSCs, a key gene in regulating cell cycle progression.

In addition, Keyes *et al*[65] applied Nfatc1 chromatin immunoprecipitation-sequencing analysis and compared age-related signatures. The results showed enrichment of Nfatc1 target genes in FACS-isolated quiescent HFSCs from aged mice. Moreover, when BMP and/or Nfatc1 were inhibited, aged HFSCs showed a lower level of hair regeneration. These results suggest that upregulated Nfatc1 significantly promotes the maintenance of quiescent bulge markers in aged HFSCs.

FORKHEAD BOX P1

The forkhead box (Fox) family contains a group of evolutionarily conserved transcription factors[66], characterized by a forkhead DNA-binding domain, that is involved in numerous functions during development and beyond. As a member of the Fox family, Foxp1 regulates the development of many tissues and is both a transcriptional activator and a transcriptional inhibitor[67-69].

During embryonic development at E13.5, when the epithelium is still a monolayer of ectodermal cells, Foxp1 is uniformly expressed in the epithelium. As the epithelium becomes a multilayer structure, Foxp1 is enriched in the downward-growing HG, which develops into HFs. In postnatal HFs, Foxp1 is continuously expressed in HFSCs of the bulge zone[70]. Foxp1 transcription levels were found to be nearly 3-fold higher in the bulge during telogen than during anagen by real-time polymerase chain reaction analysis. In addition, Foxp1 is also distributed in the ORS and the Mx.

Foxp1 induction and overexpression in keratinocytes using the tetracycline derivative doxycycline system led to proliferation inhibition[70]. Cell cycle analysis showed that Foxp1-induced cells had a higher proportion of cells in the G1 phase and a decrease in the number of cells in the G2 and S phases, which means that the cell cycle was arrested. Foxp1fl/fl; K14-cre (Foxp1 cKO) mice contained 4 types of HFs on dorsal skin, but the number of awl hairs increased, the number of auchene hairs decreased, and the HS length of various hair types became shorter[70]. At P20 and P53, the HFs in the control group were in the quiescent phase, while the HFs in the Foxp1 cKO group had entered the growth phase, indicating that Foxp1 ablation shortened the quiescent phase of HFs. Immunostaining of skin sections showed that HFSCs in the skin of Foxp1 cKO mice were substantially activated and that the number of Ki67-positive cells increased in the bulge region. Furthermore, through gain- and loss-of-function studies, Leishman *et al*[70] proposed that Foxp1 deletion may be caused by its regulation by fibroblast growth factor 18 and CDK inhibitor p57KIP2 expression. Foxp4 is another family member that is similar to Foxp1 in terms of skin expression patterns and cKO phenotypes[69]. Co-immunoprecipitation results showed that there was an interaction between Foxp1 and Foxp4 in HaCaT keratinocytes and primary epidermal cells, suggesting their dimerization in epithelial cells.

FORKHEAD BOX C1

Forkhead Box C1 (Foxc1) belongs to another subtype of the Fox family, which is an early transcription factor expressed in HF and bulge HFSCs during development and at maturity[16,40]. In addition to the bulge HFSCs, Foxc1 is also expressed in the K6+ inner bulge layer, isthmus, IRS, and sebaceous glands. Conditionally ablated Foxc1 in HF has an impact on the maintenance of HFSC quiescence[71], which may be related to the reduced expression level of E-cadherin in SCs. Although a new bulge can be produced, the HF loses its ability to maintain the old bulge. Moreover, the telogen phase of Foxc1 cKO HF is shorter than that of wild-type HF. Applying ATAC-seq and ChIP-polymerase chain reaction analysis, Wang *et al*[72] confirmed the direct regulatory effect of Foxc1 on Nfatc1, Bmp6, and Hspb8. They indicated the existence of a collaborative regulatory network between Foxc1, Nfatc1, and BMP signals.

Moreover, in aged mice, the escape of HFSCs from the bulge to the dermis was observed, leading to HF miniaturization and even hair loss. This was related to the reduced expression levels of Foxc1 and Nfatc1. These two transcription factors collaborate to regulate some HFSC-specific cell adhesion and extracellular Mx genes[73].

MUSASHI 2

Musashi (Msi), first identified in *Drosophila*, is an evolutionarily conserved RNA-binding protein family[74]. Msi has two orthologs in mammals, namely, Msi1 and Msi2[75,76]. Ma *et al*[77] demonstrated that Msi2 is an important posttranscriptional regulator of HFSC quiescence by directly targeting the Shh/Gli1 signaling pathway. At P21, Msi2 is strongly expressed in the basal layer of the bulge and HG. After entering the growth phase, the expression of Msi2 is broader, including the basal and suprabasal bulge, ORS, and IRS. Thereafter, Msi2 expression gradually decreases and is concentrated again in the bulge and HG at the next telogen.

By using gain- and loss-of-function mouse models, it was demonstrated that overexpression of Msi2 significantly prolonged the telogen-to-anagen transition, hindered hair cell entry into anagen, and impaired hair cell regeneration after depilation[77]. In contrast, the lack of Msi2 accelerated hair growth. In combination with several other HFSC markers, including K15, Sox9, and Nfatc1, Msi2 induced more Nfatc1+ cells, fewer K15+ cells and Ki67+ proliferating cells, while the opposite results were observed with Msi2 knockdown. Altogether, Msi2 plays a key role in the resting state of HFSCs.

CONCLUSION

Due to their relatively superficial location, easy isolation and observation, and periodic activation in adults, epithelial HFSCs are not only considered the preferred source for adult SC research but also the ideal seed cells for tissue-engineered skin. SC markers can not only provide a basis for the labeling and identification of HFSCs and serve as a precondition for the isolation and purification of HFSCs but also provide a foundation for the molecular research on the biological characteristics and related regulatory networks of HFSCs. The HFSC markers described here have their own characteristics. Some are classic and commonly used markers, such as CD34 and K15. Others, such as Nfatc1 and Lhx2, are molecules closely related to the important life processes of HFSCs, including but not limited to resting, proliferation, and aging. There was some colocalization between these markers. The combination of multiple markers can greatly improve the labeling rate of HFSCs.

The heterogeneity of SCs in the bulge region of HF has attracted increasing attention. Some analyses of single-cell RNA-sequencing (scRNA-seq) data also support this view in recent years[2,17,25,78,79]. Through quantitative scRNA-seq to sequence cells from the murine telogen and anagen epidermis, some studies identified 13 distinct main groups of epidermal cells[17,79]. The bulge had one inner and one outer compartment, characterized by K6a and CD34, respectively. Upon taking a second round of unsupervised clustering, the OB keratinocytes, characterized by high expression of CD34, K15, and Lgr5, were further divided into five subpopulations (OB I-V). Most OB cells were proximal OBI (CD34^{high}, Postn^{high}, Lgr5^{high}, K24^{high}) and central OBII (CD34^{high}, Postn^{high}, Lgr5^{dim}, K24^{dim}) populations. However, the study was not able to distinguish previously reported cell populations containing certain specific markers, such as Gli1+ or Lgr5+ cells in the lower bulge. The cell populations distinguished by specific markers often contain more than one subpopulation in the study. The features and functions of these new cell subpopulations require further in-depth research. In addition, the study also compared SCs and progenitor cells, but did not find any significant distinguishable stemness characteristics. Considering the different spatial locations of these cells in the tissue, space transcriptomics may become an important complement to scRNA-seq.

Combined with fluorescence-activated cell sorting, another interesting scRNA-seq study was on K14-H2BGFP+CD34+ cells from the early/mid-anagen stage of murine skin[78]. During this stage, HFSCs undergo self-renewal. Four bulge subpopulations were identified as migration-, low-, mid-, and up-bulge parts. All four subpopulations were positive for several known HFSC markers, such as K15, Sox9, Lhx2, and Col17a1. However, some other used markers had different expression patterns, including CD34, Lgr5, Nfatc1, Foxc1, Runx1, and Postn. The mid-bulge subpopulation was mainly dormant and speculated to be a reserve pool. According to the differential gene expression analysis, upregulated genes included Zfp36, Atf3, Sfn, Btg2, Cdkn1a and Hspb1. Consistent with the murine study, scRNA-seq of human skin has been performed and revealed 23 primary cell clusters[25]. The bulge subpopulation was immunostained with CD200 and Cxcl14, and a Gene Ontology analysis revealed that the bulge subpopulation was enriched in cell adhesion and extracellular Mx molecules.

This review mainly summarized the recent progresses and functions of some classic and commonly used molecular markers in HFSCs. These molecules can be used as candidate molecules for the identification, labeling, and screening of HFSCs to a certain extent. Along with the development of sequencing technology, our understanding of HFSCs will increase gradually. There is still a long way to go to clarify the molecular heterogeneity of HFSCs in more detail. With the assistance of various new technologies, more in-depth characteristics of HFSCs will be revealed, such as transcriptional heterogeneity, spatial location information, and chromatin accessibility. These new researches will help in developing reasonable methods for treating hair loss in clinic.

FOOTNOTES

Author contributions: Xing YZ drafted the manuscript; Guo HY and Xiang F participated in the literature sorting; Guo HY prepared the table; Xiang F was involved in preparation of the figure; Li YH conceived and revised the manuscript; and all authors have read and approve the final manuscript.

Supported by National Natural Science Foundation of China, No. 82173446; and the Youth Training Program of the Army Medical University, No. 2018XQN01.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Yu-Hong Li 0000-0003-1684-207X.

S-Editor: Wang JJ

L-Editor: A

P-Editor: Zhang XD

REFERENCES

- Gonzales KAU, Fuchs E. Skin and Its Regenerative Powers: An Alliance between Stem Cells and Their Niche. *Dev Cell* 2017; **43**: 387-401 [PMID: 29161590 DOI: 10.1016/j.devcel.2017.10.001]
- Yang H, Adam RC, Ge Y, Hua ZL, Fuchs E. Epithelial-Mesenchymal Micro-niches Govern Stem Cell Lineage Choices. *Cell* 2017; **169**: 483-496.e13 [PMID: 28413068 DOI: 10.1016/j.cell.2017.03.038]
- Alonso L, Fuchs E. The hair cycle. *J Cell Sci* 2006; **119**: 391-393 [PMID: 16443746 DOI: 10.1242/jcs.02793]
- Yu N, Hu T, Yang H, Zhang L, Zhu L, Zhou X, Xiang F, Yang X, Li Y. Androgen receptor inhibits the hair follicle induction potential of dermal papilla cells by binding with Tcf4 at the A574 binding site. *Genes Dis* 2023; **10**: 51-54 [PMID: 37013037 DOI: 10.1016/j.gendis.2022.04.015]
- Wu P, Zhang Y, Xing Y, Xu W, Guo H, Deng F, Ma X, Li Y. The balance of Bmp6 and Wnt10b regulates the telogen-anagen transition of hair follicles. *Cell Commun Signal* 2019; **17**: 16 [PMID: 30791955 DOI: 10.1186/s12964-019-0330-x]
- Legué E, Sequeira I, Nicolas JF. Hair follicle renewal: authentic morphogenesis that depends on a complex progression of stem cell lineages. *Development* 2010; **137**: 569-577 [PMID: 20110322 DOI: 10.1242/dev.044123]
- Yu N, Hu T, Yang H, Zhang L, Song Q, Xiang F, Yang X, Li Y. Twist1 Contributes to the Maintenance of Some Biological Properties of Dermal Papilla Cells in vitro by Forming a Complex With Tcf4 and β -Catenin. *Front Cell Dev Biol* 2020; **8**: 824 [PMID: 32974352 DOI: 10.3389/fcell.2020.00824]
- Greco V, Chen T, Rendl M, Schober M, Pasolli HA, Stokes N, Dela Cruz-Racelis J, Fuchs E. A two-step mechanism for stem cell activation during hair regeneration. *Cell Stem Cell* 2009; **4**: 155-169 [PMID: 19200804 DOI: 10.1016/j.stem.2008.12.009]
- Ito M, Kizawa K, Hamada K, Cotsarelis G. Hair follicle stem cells in the lower bulge form the secondary germ, a biochemically distinct but functionally equivalent progenitor cell population, at the termination of catagen. *Differentiation* 2004; **72**: 548-557 [PMID: 15617565 DOI: 10.1111/j.1432-0436.2004.07209008.x]
- Cotsarelis G, Sun TT, Lavker RM. Label-retaining cells reside in the bulge area of pilosebaceous unit: implications for follicular stem cells, hair cycle, and skin carcinogenesis. *Cell* 1990; **61**: 1329-1337 [PMID: 2364430 DOI: 10.1016/0092-8674(90)90696-C]
- Blanpain C, Fuchs E. Epidermal homeostasis: a balancing act of stem cells in the skin. *Nat Rev Mol Cell Biol* 2009; **10**: 207-217 [PMID: 19209183 DOI: 10.1038/nrm2636]
- Nowak JA, Polak L, Pasolli HA, Fuchs E. Hair follicle stem cells are specified and function in early skin morphogenesis. *Cell Stem Cell* 2008; **3**: 33-43 [PMID: 18593557 DOI: 10.1016/j.stem.2008.05.009]
- Brownell I, Guevara E, Bai CB, Loomis CA, Joyner AL. Nerve-derived sonic hedgehog defines a niche for hair follicle stem cells capable of becoming epidermal stem cells. *Cell Stem Cell* 2011; **8**: 552-565 [PMID: 21549329 DOI: 10.1016/j.stem.2011.02.021]
- Hsu YC, Li L, Fuchs E. Emerging interactions between skin stem cells and their niches. *Nat Med* 2014; **20**: 847-856 [PMID: 25100530 DOI: 10.1038/nm.3643]

- 15 **Brown J**, Greaves MF, Molgaard HV. The gene encoding the stem cell antigen, CD34, is conserved in mouse and expressed in haemopoietic progenitor cell lines, brain, and embryonic fibroblasts. *Int Immunol* 1991; **3**: 175-184 [PMID: [1709048](#) DOI: [10.1093/intimm/3.2.175](#)]
- 16 **Blanpain C**, Lowry WE, Geoghegan A, Polak L, Fuchs E. Self-renewal, multipotency, and the existence of two cell populations within an epithelial stem cell niche. *Cell* 2004; **118**: 635-648 [PMID: [15339667](#) DOI: [10.1016/j.cell.2004.08.012](#)]
- 17 **Joost S**, Annusver K, Jacob T, Sun X, Dalessandri T, Sivan U, Sequeira I, Sandberg R, Kasper M. The Molecular Anatomy of Mouse Skin during Hair Growth and Rest. *Cell Stem Cell* 2020; **26**: 441-457.e7 [PMID: [32109378](#) DOI: [10.1016/j.stem.2020.01.012](#)]
- 18 **Trempus CS**, Morris RJ, Bortner CD, Cotsarelis G, Faircloth RS, Reece JM, Tennant RW. Enrichment for living murine keratinocytes from the hair follicle bulge with the cell surface marker CD34. *J Invest Dermatol* 2003; **120**: 501-511 [PMID: [12648211](#) DOI: [10.1046/j.1523-1747.2003.12088.x](#)]
- 19 **Najafzadeh N**, Sagha M, Heydari Tajaddod S, Golmohammadi MG, Massahi Oskoui N, Deldadeh Moghaddam M. In vitro neural differentiation of CD34 (+) stem cell populations in hair follicles by three different neural induction protocols. *In Vitro Cell Dev Biol Anim* 2015; **51**: 192-203 [PMID: [25294494](#) DOI: [10.1007/s11626-014-9818-2](#)]
- 20 **Inoue K**, Aoi N, Sato T, Yamauchi Y, Suga H, Eto H, Kato H, Araki J, Yoshimura K. Differential expression of stem-cell-associated markers in human hair follicle epithelial cells. *Lab Invest* 2009; **89**: 844-856 [PMID: [19506554](#) DOI: [10.1038/labinvest.2009.48](#)]
- 21 **Poblet E**, Jiménez F, Godínez JM, Pascual-Martín A, Izeta A. The immunohistochemical expression of CD34 in human hair follicles: a comparative study with the bulge marker CK15. *Clin Exp Dermatol* 2006; **31**: 807-812 [PMID: [16981909](#) DOI: [10.1111/j.1365-2230.2006.02255.x](#)]
- 22 **Kloepfer JE**, Tiede S, Brinckmann J, Reinhardt DP, Meyer W, Faessler R, Paus R. Immunophenotyping of the human bulge region: the quest to define useful in situ markers for human epithelial hair follicle stem cells and their niche. *Exp Dermatol* 2008; **17**: 592-609 [PMID: [18558994](#) DOI: [10.1111/j.1600-0625.2008.00720.x](#)]
- 23 **Rosenblum MD**, Olasz EB, Yancey KB, Woodliff JE, Lazarova Z, Gerber KA, Truitt RL. Expression of CD200 on epithelial cells of the murine hair follicle: a role in tissue-specific immune tolerance? *J Invest Dermatol* 2004; **123**: 880-887 [PMID: [15482475](#) DOI: [10.1111/j.0022-202X.2004.23461.x](#)]
- 24 **Ohyama M**, Terunuma A, Tock CL, Radonovich MF, Pise-Masison CA, Hopping SB, Brady JN, Udey MC, Vogel JC. Characterization and isolation of stem cell-enriched human hair follicle bulge cells. *J Clin Invest* 2006; **116**: 249-260 [PMID: [16395407](#) DOI: [10.1172/JCI26043](#)]
- 25 **Takahashi R**, Grzenda A, Allison TF, Rawnsley J, Balin SJ, Sabri S, Plath K, Lowry WE. Defining Transcriptional Signatures of Human Hair Follicle Cell States. *J Invest Dermatol* 2020; **140**: 764-773.e4 [PMID: [31676413](#) DOI: [10.1016/j.jid.2019.07.726](#)]
- 26 **Yoshida R**, Tanaka K, Amagai M, Ohyama M. Involvement of the bulge region with decreased expression of hair follicle stem cell markers in senile female cases of alopecia areata. *J Eur Acad Dermatol Venereol* 2011; **25**: 1346-1350 [PMID: [21981340](#) DOI: [10.1111/j.1468-3083.2010.03956.x](#)]
- 27 **Lyle S**, Christofidou-Solomidou M, Liu Y, Elder DE, Albelda S, Cotsarelis G. Human hair follicle bulge cells are biochemically distinct and possess an epithelial stem cell phenotype. *J Invest Dermatol Symp Proc* 1999; **4**: 296-301 [PMID: [10674385](#) DOI: [10.1038/sj.jidsp.5640233](#)]
- 28 **Liu Y**, Lyle S, Yang Z, Cotsarelis G. Keratin 15 promoter targets putative epithelial stem cells in the hair follicle bulge. *J Invest Dermatol* 2003; **121**: 963-968 [PMID: [14708593](#) DOI: [10.1046/j.1523-1747.2003.12600.x](#)]
- 29 **Commo S**, Gaillard O, Bernard BA. The human hair follicle contains two distinct K19 positive compartments in the outer root sheath: a unifying hypothesis for stem cell reservoir? *Differentiation* 2000; **66**: 157-164 [PMID: [11269941](#) DOI: [10.1046/j.1432-0436.2000.660401.x](#)]
- 30 **Gho CG**, Braun JE, Tilli CM, Neumann HA, Ramaekers FC. Human follicular stem cells: their presence in plucked hair and follicular cell culture. *Br J Dermatol* 2004; **150**: 860-868 [PMID: [15149497](#) DOI: [10.1111/j.1365-2133.2004.05862.x](#)]
- 31 **Purba TS**, Haslam IS, Shahmalak A, Bhogal RK, Paus R. Mapping the expression of epithelial hair follicle stem cell-related transcription factors LHX2 and SOX9 in the human hair follicle. *Exp Dermatol* 2015; **24**: 462-467 [PMID: [25808706](#) DOI: [10.1111/exd.12700](#)]
- 32 **Barker N**, van Es JH, Kuipers J, Kujala P, van den Born M, Cozijnsen M, Haegebarth A, Korving J, Begthel H, Peters PJ, Clevers H. Identification of stem cells in small intestine and colon by marker gene Lgr5. *Nature* 2007; **449**: 1003-1007 [PMID: [17934449](#) DOI: [10.1038/nature06196](#)]
- 33 **Ayyaz A**, Kumar S, Sangiorgi B, Ghoshal B, Gosio J, Ouladan S, Fink M, Barutcu S, Trcka D, Shen J, Chan K, Wrana JL, Gregorieff A. Single-cell transcriptomes of the regenerating intestine reveal a revival stem cell. *Nature* 2019; **569**: 121-125 [PMID: [31019301](#) DOI: [10.1038/s41586-019-1154-y](#)]
- 34 **Jaks V**, Barker N, Kasper M, van Es JH, Snippert HJ, Clevers H, Toftgård R. Lgr5 marks cycling, yet long-lived, hair follicle stem cells. *Nat Genet* 2008; **40**: 1291-1299 [PMID: [18849992](#) DOI: [10.1038/ng.239](#)]
- 35 **Hoeck JD**, Biehs B, Kurtova AV, Kljavin NM, de Sousa E Melo F, Aliche B, Koeppen H, Modrusan Z, Piskol R, de Sauvage FJ. Stem cell plasticity enables hair regeneration following Lgr5(+) cell loss. *Nat Cell Biol* 2017; **19**: 666-676 [PMID: [28553937](#) DOI: [10.1038/ncb3535](#)]
- 36 **de Lau WB**, Snel B, Clevers HC. The R-spondin protein family. *Genome Biol* 2012; **13**: 242 [PMID: [22439850](#) DOI: [10.1186/gb-2012-13-3-242](#)]
- 37 **de Lau W**, Barker N, Low TY, Koo BK, Li VS, Teunissen H, Kujala P, Haegebarth A, Peters PJ, van de Wetering M, Stange DE, van Es JE, Guardavaccaro D, Schasfoort RB, Mohri Y, Nishimori K, Mohammed S, Heck AJ, Clevers H. Lgr5 homologues associate with Wnt receptors and mediate R-spondin signalling. *Nature* 2011; **476**: 293-297 [PMID: [21727895](#) DOI: [10.1038/nature10337](#)]
- 38 **Joost S**, Jacob T, Sun X, Annusver K, La Manno G, Sur I, Kasper M. Single-Cell Transcriptomics of Traced Epidermal and Hair Follicle Stem Cells Reveals Rapid Adaptations during Wound Healing. *Cell Rep* 2018; **25**: 585-597.e7 [PMID: [30332640](#) DOI: [10.1016/j.celrep.2018.09.059](#)]
- 39 **Mardaryev AN**, Meier N, Poterlowicz K, Sharov AA, Sharova TY, Ahmed MI, Rapisarda V, Lewis C, Fessing MY, Ruenger TM, Bhawan J, Werner S, Paus R, Botchkarev VA. Lhx2 differentially regulates Sox9, Tcf4 and Lgr5 in hair follicle stem cells to promote epidermal regeneration after injury. *Development* 2011; **138**: 4843-4852 [PMID: [22028024](#) DOI: [10.1242/dev.070284](#)]
- 40 **Rhee H**, Polak L, Fuchs E. Lhx2 maintains stem cell character in hair follicles. *Science* 2006; **312**: 1946-1949 [PMID: [16809539](#) DOI: [10.1126/science.1128004](#)]
- 41 **Folgueras AR**, Guo X, Pasolli HA, Stokes N, Polak L, Zheng D, Fuchs E. Architectural niche organization by LHX2 is linked to hair follicle stem cell function. *Cell Stem Cell* 2013; **13**: 314-327 [PMID: [24012369](#) DOI: [10.1016/j.stem.2013.06.018](#)]
- 42 **Vidal VP**, Chaboissier MC, Lützkendorf S, Cotsarelis G, Mill P, Hui CC, Ortonne N, Ortonne JP, Schedl A. Sox9 is essential for outer root sheath differentiation and the formation of the hair stem cell compartment. *Curr Biol* 2005; **15**: 1340-1351 [PMID: [16085486](#) DOI: [10.1016/j.cub.2005.06.064](#)]
- 43 **Kadaja M**, Keyes BE, Lin M, Pasolli HA, Genander M, Polak L, Stokes N, Zheng D, Fuchs E. SOX9: a stem cell transcriptional regulator of

- secreted niche signaling factors. *Genes Dev* 2014; **28**: 328-341 [PMID: 24532713 DOI: 10.1101/gad.233247.113]
- 44 **Lacaud G**, Gore L, Kennedy M, Kouskoff V, Kingsley P, Hogan C, Carlsson L, Speck N, Palis J, Keller G. Runx1 is essential for hematopoietic commitment at the hemangioblast stage of development in vitro. *Blood* 2002; **100**: 458-466 [PMID: 12091336 DOI: 10.1182/blood-2001-12-0321]
- 45 **Osorio KM**, Lilja KC, Tumber T. Runx1 modulates adult hair follicle stem cell emergence and maintenance from distinct embryonic skin compartments. *J Cell Biol* 2011; **193**: 235-250 [PMID: 21464233 DOI: 10.1083/jcb.201006068]
- 46 **Horsley V**, Aliprantis AO, Polak L, Glimcher LH, Fuchs E. NFATc1 balances quiescence and proliferation of skin stem cells. *Cell* 2008; **132**: 299-310 [PMID: 18243104 DOI: 10.1016/j.cell.2007.11.047]
- 47 **Raveh E**, Cohen S, Levanon D, Negreanu V, Groner Y, Gat U. Dynamic expression of Runx1 in skin affects hair structure. *Mech Dev* 2006; **123**: 842-850 [PMID: 17011173 DOI: 10.1016/j.mod.2006.08.002]
- 48 **Osorio KM**, Lee SE, McDermitt DJ, Waghmare SK, Zhang YV, Woo HN, Tumber T. Runx1 modulates developmental, but not injury-driven, hair follicle stem cell activation. *Development* 2008; **135**: 1059-1068 [PMID: 18256199 DOI: 10.1242/dev.012799]
- 49 **Hoi CS**, Lee SE, Lu SY, McDermitt DJ, Osorio KM, Piskun CM, Peters RM, Paus R, Tumber T. Runx1 directly promotes proliferation of hair follicle stem cells and epithelial tumor formation in mouse skin. *Mol Cell Biol* 2010; **30**: 2518-2536 [PMID: 20308320 DOI: 10.1128/MCB.01308-09]
- 50 **Lee J**, Hoi CS, Lilja KC, White BS, Lee SE, Shalloway D, Tumber T. Runx1 and p21 synergistically limit the extent of hair follicle stem cell quiescence in vivo. *Proc Natl Acad Sci U S A* 2013; **110**: 4634-4639 [PMID: 23487742 DOI: 10.1073/pnas.1213015110]
- 51 **Jho EH**, Zhang T, Domon C, Joo CK, Freund JN, Costantini F. Wnt/beta-catenin/Tcf signaling induces the transcription of Axin2, a negative regulator of the signaling pathway. *Mol Cell Biol* 2002; **22**: 1172-1183 [PMID: 11809808 DOI: 10.1128/MCB.22.4.1172-1183.2002]
- 52 **Bernkopf DB**, Hadjihannas MV, Behrens J. Negative-feedback regulation of the Wnt pathway by conductin/axin2 involves insensitivity to upstream signalling. *J Cell Sci* 2015; **128**: 33-39 [PMID: 25380820 DOI: 10.1242/jcs.159145]
- 53 **Moshkovsky AR**, Kirschner MW. The nonredundant nature of the Axin2 regulatory network in the canonical Wnt signaling pathway. *Proc Natl Acad Sci U S A* 2022; **119** [PMID: 35197279 DOI: 10.1073/pnas.2108408119]
- 54 **Xu Z**, Wang W, Jiang K, Yu Z, Huang H, Wang F, Zhou B, Chen T. Embryonic attenuated Wnt/ β -catenin signaling defines niche location and long-term stem cell fate in hair follicle. *Elife* 2015; **4**: e10567 [PMID: 26653852 DOI: 10.7554/eLife.10567]
- 55 **Qiu W**, Lei M, Zhou L, Bai X, Lai X, Yu Y, Yang T, Lian X. Hair follicle stem cell proliferation, Akt and Wnt signaling activation in TPA-induced hair regeneration. *Histochem Cell Biol* 2017; **147**: 749-758 [PMID: 28185006 DOI: 10.1007/s00418-017-1540-1]
- 56 **Myung PS**, Takeo M, Ito M, Atit RP. Epithelial Wnt ligand secretion is required for adult hair follicle growth and regeneration. *J Invest Dermatol* 2013; **133**: 31-41 [PMID: 22810306 DOI: 10.1038/jid.2012.230]
- 57 **Lien WH**, Polak L, Lin M, Lay K, Zheng D, Fuchs E. In vivo transcriptional governance of hair follicle stem cells by canonical Wnt regulators. *Nat Cell Biol* 2014; **16**: 179-190 [PMID: 24463605 DOI: 10.1038/ncb2903]
- 58 **Lien WH**, Fuchs E. Wnt some lose some: transcriptional governance of stem cells by Wnt/ β -catenin signaling. *Genes Dev* 2014; **28**: 1517-1532 [PMID: 25030692 DOI: 10.1101/gad.244772.114]
- 59 **van Amerongen R**, Bowman AN, Nusse R. Developmental stage and time dictate the fate of Wnt/ β -catenin-responsive stem cells in the mammary gland. *Cell Stem Cell* 2012; **11**: 387-400 [PMID: 22863533 DOI: 10.1016/j.stem.2012.05.023]
- 60 **Lim X**, Tan SH, Yu KL, Lim SB, Nusse R. Axin2 marks quiescent hair follicle bulge stem cells that are maintained by autocrine Wnt/ β -catenin signaling. *Proc Natl Acad Sci U S A* 2016; **113**: E1498-E1505 [PMID: 26903625 DOI: 10.1073/pnas.1601599113]
- 61 **Rao A**, Luo C, Hogan PG. Transcription factors of the NFAT family: regulation and function. *Annu Rev Immunol* 1997; **15**: 707-747 [PMID: 9143705 DOI: 10.1146/annurev.immunol.15.1.707]
- 62 **Crabtree GR**, Olson EN. NFAT signaling: choreographing the social lives of cells. *Cell* 2002; **109** Suppl: S67-S79 [PMID: 11983154 DOI: 10.1016/S0092-8674(02)00699-2]
- 63 **Tumber T**, Guasch G, Greco V, Blanpain C, Lowry WE, Rendl M, Fuchs E. Defining the epithelial stem cell niche in skin. *Science* 2004; **303**: 359-363 [PMID: 14671312 DOI: 10.1126/science.1092436]
- 64 **Morris RJ**, Liu Y, Marles L, Yang Z, Trempus C, Li S, Lin JS, Sawicki JA, Cotsarelis G. Capturing and profiling adult hair follicle stem cells. *Nat Biotechnol* 2004; **22**: 411-417 [PMID: 15024388 DOI: 10.1038/nbt950]
- 65 **Keyes BE**, Segal JP, Heller E, Lien WH, Chang CY, Guo X, Oristian DS, Zheng D, Fuchs E. Nfatc1 orchestrates aging in hair follicle stem cells. *Proc Natl Acad Sci U S A* 2013; **110**: E4950-E4959 [PMID: 24282298 DOI: 10.1073/pnas.1320301110]
- 66 **Wang B**, Lin D, Li C, Tucker P. Multiple domains define the expression and regulatory properties of Foxp1 forkhead transcriptional repressors. *J Biol Chem* 2003; **278**: 24259-24268 [PMID: 12692134 DOI: 10.1074/jbc.M207174200]
- 67 **Patzelt T**, Keppler SJ, Gorka O, Thoenes S, Wartewig T, Reth M, Förster I, Lang R, Buchner M, Ruland J. Foxp1 controls mature B cell survival and the development of follicular and B-1 B cells. *Proc Natl Acad Sci U S A* 2018; **115**: 3120-3125 [PMID: 29507226 DOI: 10.1073/pnas.1711335115]
- 68 **Jepsen K**, Gleiberman AS, Shi C, Simon DI, Rosenfeld MG. Cooperative regulation in development by SMRT and FOXP1. *Genes Dev* 2008; **22**: 740-745 [PMID: 18347093 DOI: 10.1101/gad.1637108]
- 69 **Li S**, Wang Y, Zhang Y, Lu MM, DeMayo FJ, Dekker JD, Tucker PW, Morrissey EE. Foxp1/4 control epithelial cell fate during lung development and regeneration through regulation of anterior gradient 2. *Development* 2012; **139**: 2500-2509 [PMID: 22675208 DOI: 10.1242/dev.079699]
- 70 **Leishman E**, Howard JM, Garcia GE, Miao Q, Ku AT, Dekker JD, Tucker H, Nguyen H. Foxp1 maintains hair follicle stem cell quiescence through regulation of Fgf18. *Development* 2013; **140**: 3809-3818 [PMID: 23946441 DOI: 10.1242/dev.097477]
- 71 **Lay K**, Kume T, Fuchs E. FOXC1 maintains the hair follicle stem cell niche and governs stem cell quiescence to preserve long-term tissue-regenerating potential. *Proc Natl Acad Sci U S A* 2016; **113**: E1506-E1515 [PMID: 26912458 DOI: 10.1073/pnas.1601569113]
- 72 **Wang L**, Siegenthaler JA, Dowell RD, Yi R. Foxc1 reinforces quiescence in self-renewing hair follicle stem cells. *Science* 2016; **351**: 613-617 [PMID: 26912704 DOI: 10.1126/science.aad5440]
- 73 **Zhang C**, Wang D, Wang J, Wang L, Qiu W, Kume T, Dowell R, Yi R. Escape of hair follicle stem cells causes stem cell exhaustion during aging. *Nat Aging* 2021; **1**: 889-903 [PMID: 37118327 DOI: 10.1038/s43587-021-00103-w]
- 74 **Nakamura M**, Okano H, Blendy JA, Montell C, Musashi, a neural RNA-binding protein required for Drosophila adult external sensory organ development. *Neuron* 1994; **13**: 67-81 [PMID: 8043282 DOI: 10.1016/0896-6273(94)90460-X]
- 75 **Sakakibara S**, Imai T, Hamaguchi K, Okabe M, Aruga J, Nakajima K, Yasutomi D, Nagata T, Kurihara Y, Uesugi S, Miyata T, Ogawa M,

- Mikoshiha K, Okano H. Mouse-Musashi-1, a neural RNA-binding protein highly enriched in the mammalian CNS stem cell. *Dev Biol* 1996; **176**: 230-242 [PMID: [8660864](#) DOI: [10.1006/dbio.1996.0130](#)]
- 76 **Sakakibara S**, Nakamura Y, Satoh H, Okano H. Rna-binding protein Musashi2: developmentally regulated expression in neural precursor cells and subpopulations of neurons in mammalian CNS. *J Neurosci* 2001; **21**: 8091-8107 [PMID: [11588182](#) DOI: [10.1523/JNEUROSCI.21-20-08091.2001](#)]
- 77 **Ma X**, Tian Y, Song Y, Shi J, Xu J, Xiong K, Li J, Xu W, Zhao Y, Shuai J, Chen L, Plikus MV, Lengner CJ, Ren F, Xue L, Yu Z. Msi2 Maintains Quiescent State of Hair Follicle Stem Cells by Directly Repressing the Hh Signaling Pathway. *J Invest Dermatol* 2017; **137**: 1015-1024 [PMID: [28143780](#) DOI: [10.1016/j.jid.2017.01.012](#)]
- 78 **Chovatiya G**, Ghuwalewala S, Walter LD, Cosgrove BD, Tumber T. High-resolution single-cell transcriptomics reveals heterogeneity of self-renewing hair follicle stem cells. *Exp Dermatol* 2021; **30**: 457-471 [PMID: [33319418](#) DOI: [10.1111/exd.14262](#)]
- 79 **Joost S**, Zeisel A, Jacob T, Sun X, La Manno G, Lönnberg P, Linnarsson S, Kasper M. Single-Cell Transcriptomics Reveals that Differentiation and Spatial Signatures Shape Epidermal and Hair Follicle Heterogeneity. *Cell Syst* 2016; **3**: 221-237.e9 [PMID: [27641957](#) DOI: [10.1016/j.cels.2016.08.010](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: office@baishideng.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

