

World Journal of

Gastrointestinal Endoscopy

World J Gastrointest Endosc 2022 November 16; 14(11): 657-738



Contents**Monthly Volume 14 Number 11 November 16, 2022****OPINION REVIEW**

- 657** Current approaches and questions yet to be resolved for the prophylaxis of post-endoscopic retrograde cholangiopancreatography pancreatitis
Saito H, Fujimoto A, Oomoto K, Kadokawa Y, Tada S

MINIREVIEWS

- 667** Optimal traction direction in traction-assisted gastric endoscopic submucosal dissection
Nagata M

ORIGINAL ARTICLE**Retrospective Cohort Study**

- 672** Quality of colonoscopy performed by medical or surgical specialists and trainees in five Australian hospitals
Ow TW, Sukococheva OA, Tran V, Lin R, Lee SZ, Chu M, Angelica B, Rayner CK, Tse E, Iyngkaran G, Bampton PA

Retrospective Study

- 684** Effectiveness and safety of endoscopic resection for duodenal gastrointestinal stromal tumors: A single center analysis
Wang ZZ, Yan XD, Yang HD, Mao XL, Cai Y, Fu XY, Li SW

- 694** Impact of looping on premalignant polyp detection during colonoscopy

Toyoshima O, Nishizawa T, Yoshida S, Matsuno T, Arano T, Kondo R, Kinoshita K, Yasumi Y, Tsuji Y, Fujishiro M

- 704** Self-expanding metal stent placement and pathological alterations among obstructive colorectal cancer cases

Kosumi K, Mima K, Kanemitsu K, Tajiri T, Takematsu T, Sakamoto Y, Inoue M, Miyamoto Y, Mizumoto T, Kubota T, Miyanari N, Baba H

META-ANALYSIS

- 718** Antibiotic prophylaxis to prevent complications in endoscopic retrograde cholangiopancreatography: A systematic review and meta-analysis of randomized controlled trials

Merchan MFS, de Moura DTH, de Oliveira GHP, Proença IM, do Monte Junior ES, Ide E, Moll C, Sánchez-Luna SA, Bernardo WM, de Moura EGH

LETTER TO THE EDITOR

- 731** Minimally invasive colorectal surgery learning curve

Vanella S, Bottazzi EC, Farese G, Murano R, Noviello A, Palma T, Godas M, Crafa F

CORRECTION

- 737 Correction to "Laparoscopy-assisted resection of colorectal cancer with situs inversus totalis: A case report and literature review"

Chen W, Liang JL, Ye JW, Luo YX, Huang MJ

ABOUT COVER

Editorial Board Member of *World Journal of Gastrointestinal Endoscopy*, Manish Manrai, MD, DM, FRCP (Edinburgh), Professor, Department of Internal Medicine, Armed Forces Medical College, Pune, Maharashtra 411040, India. manishmanrai@yahoo.com

AIMS AND SCOPE

The primary aim of *World Journal of Gastrointestinal Endoscopy* (WJGE, *World J Gastrointest Endosc*) is to provide scholars and readers from various fields of gastrointestinal endoscopy with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGE mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal endoscopy and covering a wide range of topics including capsule endoscopy, colonoscopy, double-balloon enteroscopy, duodenoscopy, endoscopic retrograde cholangiopancreatography, endosonography, esophagoscopy, gastrointestinal endoscopy, gastroscopy, laparoscopy, natural orifice endoscopic surgery, proctoscopy, and sigmoidoscopy.

INDEXING/ABSTRACTING

The WJGE is now abstracted and indexed in Emerging Sources Citation Index (Web of Science), PubMed, PubMed Central, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 edition of Journal Citation Reports® cites the 2021 Journal Citation Indicator (JCI) for WJGE as 0.33.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Yi-Xuan Cai*; Production Department Director: *Xu Guo*; Editorial Office Director: *Yun-Xiaojiao Wu*.

NAME OF JOURNAL

World Journal of Gastrointestinal Endoscopy

ISSN

ISSN 1948-5190 (online)

LAUNCH DATE

October 15, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Anastasios Koulaouzidis, Bing Hu, Sang Chul Lee, Joo Young Cho

EDITORIAL BOARD MEMBERS

<https://www.wjnet.com/1948-5190/editorialboard.htm>

PUBLICATION DATE

November 16, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Submit a Manuscript: <https://www.f6publishing.com>*World J Gastrointest Endosc* 2022 November 16; 14(11): 667-671DOI: [10.4253/wjge.v14.i11.667](https://doi.org/10.4253/wjge.v14.i11.667)

ISSN 1948-5190 (online)

MINIREVIEWS

Optimal traction direction in traction-assisted gastric endoscopic submucosal dissection

Mitsuru Nagata

Specialty type: Gastroenterology and hepatology**Provenance and peer review:**
Invited article; Externally peer reviewed.**Peer-review model:** Single blind**Peer-review report's scientific quality classification**

Grade A (Excellent): 0

Grade B (Very good): B, B

Grade C (Good): 0

Grade D (Fair): 0

Grade E (Poor): 0

P-Reviewer: Guo JT, China; Li S, China**Received:** September 7, 2022**Peer-review started:** September 7, 2022**First decision:** September 26, 2022**Revised:** October 2, 2022**Accepted:** October 14, 2022**Article in press:** October 14, 2022**Published online:** November 16, 2022**Mitsuru Nagata**, Department of Endoscopy, Shonan Fujisawa Tokushukai Hospital, Fujisawa 251-0041, Kanagawa, Japan**Corresponding author:** Mitsuru Nagata, MD, Chief Doctor, Department of Endoscopy, Shonan Fujisawa Tokushukai Hospital, 1-5-1 Tsujidokandai, Fujisawa 251-0041, Kanagawa, Japan.
mitsuru10jp@yahoo.co.jp

Abstract

Various traction devices have been developed to secure a visual field and sufficient tension at the dissection plane during endoscopic submucosal dissection (ESD). However, few large-scale studies have investigated the effectiveness of traction devices in gastric ESD. Clip-with-line (CWL) is one such traction device that is widely used in cases of gastric ESD. The CONNECT-G trial was the first multicenter randomized controlled trial to compare conventional ESD with CWL-assisted ESD (CWL-ESD) for superficial gastric neoplasms. Overall, no significant intergroup difference was observed in terms of the gastric ESD procedure time. However, subgroup analysis according to lesion location revealed a significant reduction in the procedure time of gastric ESD for the lesion located at the greater curvature of the middle and upper third of the stomach in the CWL-ESD group. In this subgroup analysis, lesion location was categorized as follows: anterior wall, posterior wall, lesser curvature, and greater curvature of the upper, middle, and lower thirds of the stomach. However, the gastric ESD procedure time showed no significant difference, except for lesions located at the greater curvature of the upper and middle thirds of the stomach. The traction direction of CWL in the stomach was limited to the cardia and changed depending on the lesion location. Therefore, outcomes of the CONNECT-G trial suggest that the effectiveness of CWL was influenced by lesion location, i.e., traction direction. Further studies are warranted to investigate the optimal traction direction in gastric ESD.

Key Words: Endoscopic submucosal dissection; ESD; Traction device; Clip-with-line; Traction direction; Vertical traction

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Various traction devices have been developed for endoscopic submucosal dissection (ESD). However, few traction devices have been validated in large-scale studies thus far. The CONNECT-G trial was the first multicenter randomized controlled trial to compare conventional ESD with clip-with-line-assisted ESD for superficial gastric neoplasms. This study suggested that the effectiveness of traction devices in gastric ESD depends on the traction direction; in addition, the most optimal traction direction is vertical to the gastric wall.

Citation: Nagata M. Optimal traction direction in traction-assisted gastric endoscopic submucosal dissection. *World J Gastrointest Endosc* 2022; 14(11): 667-671

URL: <https://www.wjgnet.com/1948-5190/full/v14/i11/667.htm>

DOI: <https://dx.doi.org/10.4253/wjge.v14.i11.667>

INTRODUCTION

Endoscopic submucosal dissection (ESD) allows en bloc resection of superficial gastric neoplasms. However, gastric ESD is a challenging procedure. Surgeons can use their nondominant hand to generate traction for lesions while they resect using their dominant hand. Meanwhile, endoscopists cannot use their nondominant hand to generate traction because they cannot insert their hand into the stomach. Therefore, endoscopists occasionally cannot secure a visual field and sufficient tension at the dissection plane, resulting in a long ESD procedure time and a high perforation rate. Recently, many traction devices were reported to overcome these problems, but few large-scale studies investigated the effectiveness of traction devices in gastric ESD.

CONNECT-G TRIAL

The CONNECT-G trial was the first multicenter randomized controlled trial to compare conventional ESD with traction-assisted ESD for the treatment of superficial gastric neoplasms[1]. In this study, clip-with-line (CWL) was used as a traction device (Figure 1), and its traction direction is restricted to the direction where the line is drawn[2,3]. The primary endpoint was the mean gastric ESD procedure time, which was 58.1 min in the conventional ESD group and 60.7 min in the CWL-assisted ESD (CWL-ESD) group, with no significant difference ($P = 0.45$). R0 resection was not statistically significant in both groups (96.8% vs 97.8%, $P = 0.45$). However, the perforation rate was significantly lower in the CWL-ESD group (0.3% vs 2.2%, $P = 0.04$), suggesting that CWL may have improved the field of vision and reduced blind submucosal dissection.

For lesions located at the greater curvature of the middle and upper third of the stomach, the CWL-ESD group had a significantly shorter gastric ESD procedure time than the conventional ESD group (57.2 min vs 104.1 min, $P = 0.01$). This part of the stomach is a challenging area for conventional ESD because it is basically a gravitational lower side, so a mucosal flap is difficult to deploy, and the visual field tends to deteriorate due to fluid retention. Nevertheless, CWL-ESD is particularly useful in this area. In this subgroup analysis, lesion location was divided into the anterior wall, posterior wall, lesser curvature, and greater curvature of the upper, middle, and lower third of the stomach. However, no significant difference was found in the procedure time of gastric ESD, except for lesions located at the greater curvature of the middle and upper third of the stomach.


TRACTION DIRECTION OF CWL DIFFERS DEPENDING ON THE LESION LOCATION AND ENDOSCOPIC POSITION

The results of the CONNECT-G trial suggest that the effectiveness of CWL-ESD varies depending on the lesion location. Traction direction can be classified into five categories (Figure 2)[4]. Since CWL is a per-oral traction device, its traction direction is limited to the cardia and varies depending on the lesion location. Another consideration for the traction direction of CWL is the endoscopic position during submucosal dissection. Because of the large lumen of the stomach, there are two possible endoscopic positions: forward and retroflexed. Therefore, the traction direction also varies depending on the endoscopic position even if the lesion is in the same location. For example, for lesions located at the lesser curvature of the middle third of the stomach, CWL commonly provides a distal traction in retroflexed endoscopic position (Figure 3A) and proximal traction in forward endoscopic position (Figure 3B).



DOI: 10.4253/wjge.v14.i11.667 Copyright ©The Author(s) 2022.

Figure 1 A clip-with-line was made by tying a commercially available dental floss to the arm section of the hemoclip.




DOI: 10.4253/wjge.v14.i11.667 Copyright ©The Author(s) 2022.

Figure 2 Classification of the traction direction. A: Proximal traction; B: Diagonally proximal traction; C: Vertical traction; D: Diagonally distal traction; E: Distal traction. Citation: Reprinted from Mitsuru Nagata. Advances in traction methods for endoscopic submucosal dissection: What is the best traction method and traction direction? *World Journal of Gastroenterology* 2022; 28(1): 1-22. Copyright ©Mitsuru Nagata 2022. Published by Baishideng Publishing Group Inc.

WHAT IS THE OPTIMAL TRACTION DIRECTION IN GASTRIC ESD?

The optimal traction direction in gastric ESD was not yet fully investigated. However, several studies indicated that a vertical traction is the optimal traction direction. The CONNECT-G trial suggests that CWL is effective for lesions located at the greater curvature of the upper and middle third of the stomach, and vertical traction is frequently performed in this area from an anatomical point of view ([Figure 3C](#)). CWL can essentially only provide vertical traction for lesions located at the greater curvature of the stomach, but multidirectional traction devices, such as a spring-and-loop with clip



DOI: 10.4253/wjge.v14.i11.667 Copyright ©The Author(s) 2022.

Figure 3 Differences in traction direction depending on the lesion location in clip-with-line-assisted endoscopic submucosal dissection.

A: Distal traction; B: Proximal traction; C: Vertical traction. Citation: Reprinted from Mitsuru Nagata. Advances in traction methods for endoscopic submucosal dissection: What is the best traction method and traction direction? *World Journal of Gastroenterology* 2022; 28(1): 1–22. Copyright ©Mitsuru Nagata 2022. Published by Baishideng Publishing Group Inc.



DOI: 10.4253/wjge.v14.i11.667 Copyright ©The Author(s) 2022.

Figure 4 An S-O clip (Zeon Medical, Tokyo, Japan) was made of a 5 mm-long spring and 4 mm-long nylon loop on one side of the clip claws.

(Figure 4; SLC; S-O clip; Zeon Medical, Tokyo, Japan), may provide a vertical traction for lesions in other areas. The SLC allows the traction direction to be controlled in any direction. This clip was developed as a traction device to provide traction for colorectal ESD. Hence, we have devised a novel usage of the SLC with both forward and retroflexed endoscopic positions for gastric ESD[5,6]. A single-center randomized controlled trial comparing conventional ESD and SLC-assisted ESD (SLC-ESD) was conducted. In SLC-ESD, a vertical traction was selected using the multidirectional traction function. This study demonstrated that the median gastric ESD procedure time was significantly shorter in SLC-ESD than in conventional ESD (29.1 min vs 52.6 min; $P = 0.005$)[7]. However, SLC-ESD was not associated with a reduction in the gastric ESD procedure time for lesions > 20 mm. As submucosal dissection progresses, the distance between the SLC attachment site and the anchor site diminishes gradually, resulting in weaker traction force due to the spring shortening. For larger lesions, diagonally proximal traction may be preferable to vertical traction to maintain spring extension even as submucosal dissection progresses or an additional SLC should be considered when traction force becomes weaker. Overall, considering the results of these two randomized controlled trials, vertical traction may be the optimal traction direction for most cases of gastric ESD.

It is unclear whether other traction directions are effective in gastric ESD. Especially in distal traction, as the submucosal dissection progresses, the dissection plane falls toward a distal direction, which may be counterproductive because it may not provide an effective tension on the dissection plane. In CWL-ESD, a retroflexed endoscopic position occasionally results in a distal traction, and this position is common in gastric ESD. It is possible that a distal traction was provided for a relatively large number of cases in the CONNECT-G trial, and this could cause no significant difference in gastric ESD procedure time between conventional ESD and CWL-ESD in the total population of the CONNECT-G trial. However, the traction direction and endoscopic position were not reported, so this point should be

further investigated.

In CWL-ESD, combined with the pulley method[8,9], the traction direction can be controlled, and vertical traction can be obtained. Therefore, the pulley method may improve the gastric ESD procedure time in CWL-ESD. However, since the pulley method in gastric ESD has been reported mainly in case series studies or *ex-vivo* studies, its feasibility and effectiveness should be further investigated.

CONCLUSION

Vertical traction may be the optimal traction direction in traction-assisted ESD for gastric neoplasms. Further studies are needed to investigate the effectiveness of other traction directions.

FOOTNOTES

Author contributions: Nagata M has been associated with the conception, drafting of the article, and final approval of the article.

Conflict-of-interest statement: No financial relationships with a commercial entity producing health-care related products and/or services relevant to this article.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/Licenses/by-nc/4.0/>

Country/Territory of origin: Japan

ORCID number: Mitsuru Nagata 0000-0002-5697-5953.

S-Editor: Liu JH

L-Editor: A

P-Editor: Liu JH

REFERENCES

- 1 **Yoshida M**, Takizawa K, Suzuki S, Koike Y, Nonaka S, Yamasaki Y, Minagawa T, Sato C, Takeuchi C, Watanabe K, Kanzaki H, Morimoto H, Yano T, Sudo K, Mori K, Gotoda T, Ono H; CONNECT-G Study Group. Conventional vs traction-assisted endoscopic submucosal dissection for gastric neoplasms: a multicenter, randomized controlled trial (with video). *Gastrointest Endosc* 2018; **87**: 1231-1240 [PMID: 29233673 DOI: 10.1016/j.gie.2017.11.031]
- 2 **Oyama T**, Kikuchi Y, Shimaya S, Tomori T, Hotta K, Miyata Y, Yamada S. Endoscopic mucosal resection using a hooking knife (hooking EMR) [in Japanese]. *Stomach Intest* 2002; **37**: 1151-1161 [DOI: 10.11477/mf.1403104523]
- 3 **Oyama T**. Counter traction makes endoscopic submucosal dissection easier. *Clin Endosc* 2012; **45**: 375-378 [PMID: 23251884 DOI: 10.5946/ce.2012.45.4.375]
- 4 **Nagata M**. Advances in traction methods for endoscopic submucosal dissection: What is the best traction method and traction direction? *World J Gastroenterol* 2022; **28**: 1-22 [PMID: 35125817 DOI: 10.3748/wjg.v28.i1.1]
- 5 **Nagata M**. Modified attachment method using an S-O clip for gastric endoscopic submucosal dissection. *VideoGIE* 2019; **4**: 151-153 [PMID: 31032463 DOI: 10.1016/j.vgie.2018.12.015]
- 6 **Nagata M**. Internal traction method using a spring-and-loop with clip (S-O clip) allows countertraction in gastric endoscopic submucosal dissection. *Surg Endosc* 2020; **34**: 3722-3733 [PMID: 32350668 DOI: 10.1007/s00464-020-07590-9]
- 7 **Nagata M**, Fujikawa T, Munakata H. Comparing a conventional and a spring-and-loop with clip traction method of endoscopic submucosal dissection for superficial gastric neoplasms: a randomized controlled trial (with videos). *Gastrointest Endosc* 2021; **93**: 1097-1109 [PMID: 33058886 DOI: 10.1016/j.gie.2020.09.049]
- 8 **Li CH**, Chen PJ, Chu HC, Huang TY, Shih YL, Chang WK, Hsieh TY. Endoscopic submucosal dissection with the pulley method for early-stage gastric cancer (with video). *Gastrointest Endosc* 2011; **73**: 163-167 [PMID: 21030018 DOI: 10.1016/j.gie.2010.08.041]
- 9 **Ge PS**, Thompson CC, Jirapinyo P, Aihara H. Suture pulley countertraction method reduces procedure time and technical demand of endoscopic submucosal dissection among novice endoscopists learning endoscopic submucosal dissection: a prospective randomized *ex vivo* study. *Gastrointest Endosc* 2019; **89**: 177-184 [PMID: 30148993 DOI: 10.1016/j.gie.2018.08.032]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjnet.com>

