

# World Journal of *Gastrointestinal Endoscopy*

*World J Gastrointest Endosc* 2023 June 16; 15(6): 420-490



**EVIDENCE REVIEW**

- 420 Applications of endoscopic vacuum therapy in the upper gastrointestinal tract  
*Kouladouros K*

**MINIREVIEWS**

- 434 Flexible robotic endoscopy for treating gastrointestinal neoplasms  
*Kume K*
- 440 Endoscopic intraductal radiofrequency ablation for extrahepatic cholangiocarcinoma: An update (2023)  
*Inoue T, Yoneda M*

**ORIGINAL ARTICLE****Retrospective Cohort Study**

- 447 Role of endoscopic ultrasound for pre-intervention evaluation in early esophageal cancer  
*Kahlon S, Amar A, Butt Z, Urayama S*
- 458 Multicenter evaluation of recurrence in endoscopic submucosal dissection and endoscopic mucosal resection in the colon: A Western perspective  
*Wei MT, Zhou MJ, Li AA, Ofosu A, Hwang JH, Friedland S*
- 469 Endoscopic sleeve gastropasty in class III obesity: Efficacy, safety, and durability outcomes in 404 consecutive patients  
*Maselli DB, Hoff AC, Kucera A, Weaver E, Sebring L, Gooch L, Walton K, Lee D, Cratty T, Beal S, Nanduri S, Rease K, Gainey CS, Eaton L, Coan B, McGowan CE*

**Retrospective Study**

- 480 Prevalence and clinical risk factors for esophageal candidiasis in non-human immunodeficiency virus patients: A multicenter retrospective case-control study  
*Kimchy AV, Ahmad AI, Tully L, Lester C, Sanghavi K, Jennings JJ*

**ABOUT COVER**

Editor-in-Chief of *World Journal of Gastrointestinal Endoscopy*, Anastasios Koulaouzidis, MD, PhD, Professor, Department of Social Medicine and Public Health, Faculty of Health Sciences, Pomeranian Medical University, Szczecin EH16 4SA, Poland. akoulaouzidis@hotmail.com

**AIMS AND SCOPE**

The primary aim of *World Journal of Gastrointestinal Endoscopy* (WJGE, *World J Gastrointest Endosc*) is to provide scholars and readers from various fields of gastrointestinal endoscopy with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGE mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal endoscopy and covering a wide range of topics including capsule endoscopy, colonoscopy, double-balloon enteroscopy, duodenoscopy, endoscopic retrograde cholangiopancreatography, endosonography, esophagoscopy, gastrointestinal endoscopy, gastroscopy, laparoscopy, natural orifice endoscopic surgery, proctoscopy, and sigmoidoscopy.

**INDEXING/ABSTRACTING**

The WJGE is now abstracted and indexed in Emerging Sources Citation Index (Web of Science), PubMed, PubMed Central, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJGE as 2.0; IF without journal self cites: 1.9; 5-year IF: 3.3; Journal Citation Indicator: 0.28.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: Yi-Xuan Cai; Production Department Director: Xu Guo; Editorial Office Director: Jia-Ping Yan.

**NAME OF JOURNAL**

*World Journal of Gastrointestinal Endoscopy*

**ISSN**

ISSN 1948-5190 (online)

**LAUNCH DATE**

October 15, 2009

**FREQUENCY**

Monthly

**EDITORS-IN-CHIEF**

Anastasios Koulaouzidis, Bing Hu, Sang Chul Lee, Joo Young Cho

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/1948-5190/editorialboard.htm>

**PUBLICATION DATE**

June 16, 2023

**COPYRIGHT**

© 2023 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/gerinfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/gerinfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/gerinfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>



Retrospective Cohort Study

# Multicenter evaluation of recurrence in endoscopic submucosal dissection and endoscopic mucosal resection in the colon: A Western perspective

Mike T Wei, Margaret J Zhou, Andrew A Li, Andrew Ofosu, Joo Ha Hwang, Shai Friedland

**Specialty type:** Gastroenterology and hepatology

**Provenance and peer review:** Invited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): 0  
Grade B (Very good): 0  
Grade C (Good): C, C  
Grade D (Fair): D  
Grade E (Poor): 0

**P-Reviewer:** Kobayashi N, Japan; Tsou YK, Taiwan

**Received:** February 20, 2023

**Peer-review started:** February 20, 2023

**First decision:** April 13, 2023

**Revised:** May 12, 2023

**Accepted:** May 31, 2023

**Article in press:** May 31, 2023

**Published online:** June 16, 2023



**Mike T Wei, Margaret J Zhou, Andrew A Li, Joo Ha Hwang, Shai Friedland,** Department of Gastroenterology and Hepatology, Stanford University Medical Center, Palo Alto, CA 94306, United States

**Andrew Ofosu,** Department of Gastroenterology, University of Cincinnati College of Medicine, Cincinnati, OH 45267, United States

**Corresponding author:** Mike T Wei, MD, Clinical Assistant Professor, Department of Gastroenterology and Hepatology, Stanford University Medical Center, 300 Pasteur Drive, Palo Alto, CA 94306, United States. [mtwei@stanford.edu](mailto:mtwei@stanford.edu)

## Abstract

### BACKGROUND

While colon endoscopic mucosal resection (EMR) is an effective technique, removal of larger polyps often requires piecemeal resection, which can increase recurrence rates. Endoscopic submucosal dissection (ESD) in the colon offers the ability for *en bloc* resection and is well-described in Asia, but there are limited studies comparing ESD vs EMR in the West.

### AIM

To evaluate different techniques in endoscopic resection of large polyps in the colon and to identify factors for recurrence.

### METHODS

The study is a retrospective comparison of ESD, EMR and knife-assisted endoscopic resection performed at Stanford University Medical Center and Veterans Affairs Palo Alto Health Care System between 2016 and 2020. Knife-assisted endoscopic resection was defined as use of electrosurgical knife to facilitate snare resection, such as for circumferential incision. Patients  $\geq 18$  years of age undergoing colonoscopy with removal of polyp(s)  $\geq 20$  mm were included. The primary outcome was recurrence on follow-up.

### RESULTS

A total of 376 patients and 428 polyps were included. Mean polyp size was greatest in the ESD group (35.8 mm), followed by knife-assisted endoscopic resection (33.3 mm) and EMR (30.5 mm) ( $P < 0.001$ ). ESD achieved highest *en bloc*

resection (90.4%) followed by knife-assisted endoscopic resection (31.1%) and EMR (20.2%) ( $P < 0.001$ ). A total of 287 polyps had follow-up (67.1%). On follow-up analysis, recurrence rate was lowest in knife-assisted endoscopic resection (0.0%) and ESD (1.3%) and highest in EMR (12.9%) ( $P = 0.0017$ ). *En bloc* polyp resection had significantly lower rate of recurrence (1.9%) compared to non-*en bloc* (12.0%,  $P = 0.003$ ). On multivariate analysis, ESD (in comparison to EMR) adjusted for polyp size was found to significantly reduce risk of recurrence [adjusted hazard ratio 0.06 (95%CI: 0.01-0.57,  $P = 0.014$ )].

## CONCLUSION

In our study, EMR had significantly higher recurrence compared to ESD and knife-assisted endoscopic resection. We found factors including resection by ESD, *en bloc* removal, and use of circumferential incision were associated with significantly decreased recurrence. While further studies are needed, we have demonstrated the efficacy of ESD in a Western population.

**Key Words:** Endoscopic mucosal resection; Endoscopic submucosal dissection; Recurrence; Colonoscopy; Polypectomy

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** Endoscopic submucosal dissection is an effective and safe technique. Compared to endoscopic mucosal resection, we find that endoscopic submucosal dissection as well as knife-assisted endoscopic resection to achieve higher *en bloc* resection, circumferential incision, R0 resection as well as lower recurrence rate. While further studies are needed, we have demonstrated the efficacy of endoscopic submucosal dissection in a Western population.

**Citation:** Wei MT, Zhou MJ, Li AA, Ofosu A, Hwang JH, Friedland S. Multicenter evaluation of recurrence in endoscopic submucosal dissection and endoscopic mucosal resection in the colon: A Western perspective. *World J Gastrointest Endosc* 2023; 15(6): 458-468

**URL:** <https://www.wjgnet.com/1948-5190/full/v15/i6/458.htm>

**DOI:** <https://dx.doi.org/10.4253/wjge.v15.i6.458>

## INTRODUCTION

Large non-pedunculated colorectal polyps are currently removed primarily through endoscopic mucosal resection (EMR) or endoscopic submucosal dissection (ESD)[1]. ESD has been slowly adopted in the United States, limited in large part due to lack of experts, long training required and significantly increased time for resection compared to EMR[2]. As a result, there is limited data for Western experience in ESD. In the largest prospective multicenter study to date in North America, Draganov *et al* [3] identified 399 cases of ESD in the colorectum, identifying an *en bloc* resection rate of 87.2%, and recurrence rate of 2.7% (8 of 296)[3]. With the growing experience in North America in performing ESD, there is increased attention to performance outcomes of ESD compared to EMR. In this study, we seek to evaluate our experience of ESD compared to EMR at two tertiary centers in California.

## MATERIALS AND METHODS

### Study cohort

We performed a retrospective study evaluating endoscopic resection performed of polyps  $\geq 20$  mm at two centers (Stanford University Medical Center and Veterans Affairs Palo Alto Health Care System) by two practitioners (JHH and SF), between January 1, 2016, and December 31, 2020. Inclusion criteria included adults age  $\geq 18$  who presented for colonoscopy with endoscopic removal of polyp  $\geq 20$  mm in size. Exclusion criteria included age  $< 18$  and pregnancy.

### Definitions for endoscopic resection

Endoscopic resection was categorized as EMR, knife-assisted endoscopic resection and ESD. EMR was defined by hot or cold snare resection of the polyp with or without submucosal injection. Knife-assisted endoscopic resection was defined as use of electrosurgical knife to facilitate snare resection, such as for circumferential incision and minimal submucosal dissection with an ESD knife. ESD was defined as use

of electrosurgical knife for circumferential incision and submucosal dissection with the intention of performing a complete *en bloc* resection using the knife (Figure 1)[4]. *En bloc* resection was defined as removal of the polyp in its entirety in one singular piece. Determination of each technique is up to the discretion of the endoscopist. Knife-assisted endoscopic resection was performed when the endoscopist determined at the initial submucosal injection step that full ESD would be too dangerous, typically due to fibrosis or poor scope stability, but that there was a clinical benefit to utilizing an ESD knife to perform selected parts of the procedure.

Endoscopy was performed using high-definition video endoscopes (*e.g.* PCF-H190DL; GIF-1TH190). A transparent cap was attached to the tip of the endoscope for each procedure. Each polyp was carefully examined under both white light and narrow band imaging (NBI)[5] and evaluated to predict histopathological diagnosis and invasion depth. Polyps were characterized by Paris classification[6] as well as by Japan NBI Expert Team (JNET)[7,8]. Submucosal injection was performed using hydroxyethyl starch with dye, saline with dye, ORISE™ gel (Boston Scientific), Eleview™ liquid composition (Aries Pharmaceuticals), or EverLift™ (GI Supply). Lesion marking, mucosal incision, and submucosal dissection were performed using an DualKnife (Olympus), FlushKnife (Fujinon), Hybrid Knife (ERBE) or ProKnife (Boston Scientific) with an electrosurgical generator (ERBE Elektromedizin, Tübingen, Germany). In select cases, the resection site was closed with hemostatic clips, X-Tac (Apollo Endosurgery), or OverStitch (Apollo Endosurgery). Resected specimens were pinned on cork or foam board for better pathologic analysis. The specimens were fixed with formalin[1].

The size of the polyp was determined by using the snare as reference, or if the polyp was removed *en bloc*, was measured against a ruler when it was retrieved from the colon.

### Data collection

All procedures performed by SF (Stanford and Veterans Affairs Palo Alto Health Care System) and JHH (Stanford) between January 1, 2016 and December 31, 2020 were reviewed. Data collected included patient demographics (age, sex, race/ethnicity), sedation, bowel preparation, polyp size, location, Paris and JNET classification, history of prior resection, method of resection, *en bloc* removal of polyp, and pathology of the polyp. Bowel preparation was characterized as adequate or inadequate. 30-d complications recorded included bleeding with or without intervention, perforation, small bowel obstruction, abdominal pain, as well as complications unrelated to procedure. Follow-up endoscopic evaluation was measured for presence or absence of recurrence. Follow-up was reviewed up to December 31, 2022. Recurrence was defined as evidence of polyp in the area of the prior resection. During follow-up endoscopy, careful examination was performed in the area of the resection, with both white light and NBI, to evaluate for recurrence. When there was suspicion for recurrence, resection or biopsies were performed of the area. The primary outcome was recurrence on follow-up. Secondary outcomes included *en bloc* resection and complication rates

### Specimen histology

Specimen from knife-assisted endoscopic resection and ESD were spread and pinned onto cork or Styrofoam boards immediately following endoscopic resection. The specimens were fixed in 10% buffered formalin, paraffin embedded, and cut into 2-mm-thick slices, prior to evaluation by a pathologist.

### Statistical analysis

All analyses were performed with *P*-value < 0.05 considered significant. All tests were 2-tailed.  $\chi^2$  test was performed to compare the frequencies of categorical outcomes and student's *t*-test was performed to evaluate averages of normally distributed continuous variables. Cox regression analysis was performed to estimate unadjusted and adjusted hazard ratios (HR and aHR) relating potential confounders such as resection technique, age, sex, race, polyp location, prior resection attempt, polyp size, with polyp recurrence.

### Ethics statement

This study was performed under the approval of the Institutional Review Board at Stanford University, Stanford, California, United States.

## RESULTS


### Patient demographics

There were 376 patients included in the study, 122 of whom received ESD, 44 received knife-assisted endoscopic resection and 216 received EMR. A total of 38 patients had more than one  $\geq 20$  mm polyp removed. There were 6 patients who underwent two resection techniques (*e.g.* EMR and ESD). There was similar distribution in age, sex, race/ethnicity across the three categories of procedures (Table 1). Patients undergoing ESD had a higher likelihood of receiving the procedure under general anesthesia or

**Table 1 Patient characteristics by intervention, *n* (%)**

<i>N</i> = 376	ESD ( <i>n</i> = 122)	Knife-assisted endoscopic resection ( <i>n</i> = 44)	EMR ( <i>n</i> = 216)	<i>P</i> value
Mean age (mean ± SD)	66.9 (11.8)	64.5 (11.8)	66 (9.7)	0.452
Male (%)	78 (63.9)	23 (52.3)	130 (60.2)	0.395
Race/Ethnicity				0.113
White	78 (63.9)	28 (63.6)	144 (66.7)	
Asian	19 (15.6)	6 (13.6)	12 (5.6)	
African American	4 (3.3)	1 (2.3)	11 (5.1)	
Latino	14 (11.5)	6 (13.6)	25 (11.6)	
Other	7 (5.7)	3 (6.8)	24 (11.1)	
Sedation				< 0.001
General anesthesia	11 (9.0)	2 (4.5)	7 (3.2)	
Monitored anesthesia care	93 (76.2)	29 (65.9)	117 (54.2)	
Moderate sedation	17 (13.9)	13 (29.5)	82 (38.0)	
None	1 (0.8)	0 (0.0)	10 (4.6)	
Adequate bowel preparation	121 (99.2)	41 (93.2)	210 (97.2)	0.100

EMR: Endoscopic mucosal resection; ESD: Endoscopic submucosal dissection.



DOI: 10.4253/wjge.v15.i6.458 Copyright ©The Author(s) 2023.

**Figure 1 Endoscopic submucosal dissection.** A and B: Steps of endoscopic submucosal dissection includes careful surveillance of the polyp, with techniques including near focus and narrow band imaging; C: Mucosal incision is performed following by submucosal dissection, which is aided by a submucosal lifting agent (in this case a mixture of epinephrine, hetastarch, and indigo carmine); D-F: Following complete resection of the polyp, the complete resection bed is closed, with techniques including clips.

monitored anesthesia care (85.2%) compared to knife-assisted endoscopic resection (70.5%) and EMR (57.4%).

### Polyp resection, overall

A total of 428 polyps underwent endoscopic resection, with 258 by EMR and 125 by ESD (Supplementary Table 1). Polyps removed by ESD (35.8 mm) were larger compared to by knife-assisted

endoscopic resection (33.3 mm), which was larger than by EMR (30.5 mm) ( $P < 0.001$ ). ESD achieved the highest *en bloc* resection (90.4%) followed by knife-assisted endoscopic resection (31.1%) and EMR (20.2%) ( $P < 0.001$ ). There was no significant difference in proportion of polyps that had history of prior resection attempt in the three resection techniques. Non-neoplastic polyps were removed more frequently in EMR (5.8%) compared to ESD (0.0%), while cancer was removed more frequently with ESD (13.6%) compared to EMR (3.5%).

### Polyp resection, follow-up

EMR (69.0%) and knife-assisted endoscopic resection (71.1%) had greater proportion of patients that underwent follow-up compared to in the ESD group (61.6%), though this was not statistically significant ( $P = 0.266$ ). On evaluation of polyps that received follow-up evaluation (Table 2), ESD had highest rate of *en bloc* resection (89.7%) followed by knife-assisted endoscopic resection (25.0%), followed by EMR (15.2%) ( $P < 0.001$ ). A higher proportion (44.2%) of polyps undergoing ESD were identified in the rectum compared to knife-assisted endoscopic resection and EMR, while a higher percentage of polyps were removed in the right colon by knife-assisted endoscopic resection or EMR. A higher proportion (74.0%) removed by ESD were identified as Paris classification Is, compared to 56.3% for knife-assisted endoscopic resection and 36.0% for EMR. EMR had the longest mean follow-up (516.2 d) compared to ESD (456.8) and knife-assisted endoscopic resection (365.0), though this was not statistically significant ( $P = 0.061$ ). ESD (74.0%) and knife-assisted endoscopic resection (18.8%) had higher R0 resection compared to EMR (4.5%) ( $P < 0.001$ ). There was no recurrence in the knife-assisted endoscopic removal group (0/30). Recurrence rate was lowest in knife-assisted endoscopic resection (0.0%), followed by ESD (1.3%), and highest in EMR (12.9%) ( $P = 0.002$ ).

In categorizing polyps by presence of recurrence (Table 3), there was overall a low proportion of polyps with recurrence (8.4%). Polyps with recurrence had greater mean average size (37.4 vs 32.7 mm,  $P = 0.202$ ), though this was not statistically significant. Polyps with recurrence more often had non *en bloc* resection (91.7% vs 61.2%,  $P = 0.003$ ). Polyps with recurrence more often did not undergo circumferential incision (95.8% vs 62.7%,  $P = 0.001$ ). Of note, polyps removed with circumferential incision had higher proportion of *en bloc* removal (76.8% vs 14.9%,  $P < 0.0001$ ). Recurrence polyps had a higher proportion of polyps that had prior attempt at removal (17.6% vs 5.3%), though this was not statistically significant ( $P = 0.154$ ). There was no significant difference in pathology or mean follow-up between the two groups. Compared to no recurrence, polyps with recurrence had higher proportion of R1 (91.7% vs 67.7%) and lower proportion of R0 (4.2% vs 26.6%) ( $P = 0.041$ ).

### Procedural complications

Overall, there was a low patient complication rate [25 patients (6.6%)], with similar proportion of complication (6.5%-6.8%) among the three procedures (Table 4). There were 3 cases of perforation (two ESD and one knife-assisted endoscopic resection). One patient received knife-assisted endoscopic resection of a > 50 mm polyp in cecum involving the ileocecal valve. There was only partial lifting of the lesion with submucosal injection. Dense fibrosis was encountered, and as such the remainder of the resection was performed by piecemeal EMR. Following the procedure, the patient had abdominal pain, and was found to have pneumoperitoneum. The patient underwent exploratory laparotomy with resection of the terminal ileum and proximal colon. Pathology returned as tubular adenoma with focal high-grade dysplasia. In the second case, the patient had a fungating partially obstructing 50 mm mass in ascending colon. Following ESD, five hemostatic clips placed to close the wound. The patient had worsening abdominal pain following the procedure, and perforation was seen on computed tomography, leading to hemicolectomy. Pathology was consistent with tubulovillous adenoma. In the third case, a 30 mm fungating non-obstructing mass was found in the cecum, encasing the appendiceal orifice. A 40 mm specimen was resected *en bloc*. A single small perforation (< 2 mm) occurred, which was closed with a single clip followed by full mucosal closure with an Endoloop and clips. The patient recovered uneventfully.

### Predictors of recurrence

On univariate Cox regression, age, sex, race and polyp location were not significant risk factors. Relative to EMR, ESD was found to decrease risk of recurrence [hazard ratio (HR): 0.12 (95%CI: 0.02-0.92),  $P = 0.041$ ]. Completion of circumferential incision, *en bloc* resection as well as R0 resection were found to significantly reduce risk of recurrence (Table 5). On multivariable Cox regression adjusted for polyp size and type of resection (ESD vs EMR), ESD significantly reduced risk of recurrence [adjusted HR (aHR): 0.06 (95%CI: 0.01-0.57,  $P = 0.014$ )] (Table 6). In this analysis, we did not include *en bloc* resection, R0 resection, and presence of circumferential incision as these are factors closely tied with performance of ESD. When evaluating EMR compared to knife-assisted endoscopic resection combined with ESD, on multivariate analysis ESD and knife-assisted endoscopic resection also demonstrated significant decrease in risk of recurrence [aHR: 0.05 (95%CI: 0.01-0.45),  $P = 0.008$ ] (Supplementary Tables 2 and 3). Knife-assisted endoscopic resection was unable to be evaluated independently of ESD as there were no cases of recurrence.

Table 2 Characteristics of polyp with follow-up, by intervention, *n* (%)

<i>N</i> = 287	ESD ( <i>n</i> = 77)	Knife-assisted endoscopic resection ( <i>n</i> = 32)	EMR ( <i>n</i> = 178)	<i>P</i> value
Size of polyp, mm (mean ± SD)	37.2 (19.7)	32.7 (8.7)	31.4 (11.5)	0.010
<i>En bloc</i>	69 (89.7)	8 (25.0)	27 (15.2)	< 0.001
Location of polyp				< 0.001
Cecum	10 (13.0)	7 (21.9)	47 (26.4)	
Ascending	13 (16.9)	12 (37.5)	63 (35.4)	
Transverse	8 (10.4)	6 (18.8)	45 (25.3)	
Descending	2 (2.6)	4 (12.5)	12 (6.7)	
Sigmoid	10 (13.0)	1 (3.1)	5 (2.8)	
Rectum	34 (44.2)	2 (6.3)	6 (3.4)	
Paris classification				< 0.001
Is	57 (74.0)	18 (56.3)	64 (36.0)	
Ila	16 (20.8)	9 (28.1)	102 (57.3)	
Ilb	0 (0.0)	1 (3.1)	2 (1.1)	
Ila+c	2 (2.6)	1 (3.1)	2 (1.1)	
Ilc	0 (0.0)	1 (3.1)	0 (0.0)	
Isp	2 (2.6)	2 (6.3)	8 (4.5)	
Pathology				< 0.001
Non-neoplastic	0 (0.0)	1 (3.1)	10 (5.6)	
Neoplastic, no high-grade dysplasia	50 (64.9)	25 (78.1)	152 (85.4)	
High-grade dysplasia	17 (22.1)	6 (18.8)	12 (6.7)	
Cancer	10 (13.0)	0 (0.0)	4 (2.2)	
First follow-up, days (mean ± SD)	456.8 (326.1)	365.0 (230.2)	516.2 (377.7)	0.061
Recurrence	1 (1.3)	0 (0.0)	23 (12.9)	0.0017
Complete resection				< 0.001
R0	57 (74.0)	6 (18.8)	8 (4.5)	
R1	18 (23.4)	26 (81.3)	156 (87.6)	
Rx	2 (2.6)	0 (0.0)	14 (7.9)	

EMR: Endoscopic mucosal resection; ESD: Endoscopic submucosal dissection.

## DISCUSSION

The development of advanced polypectomy techniques has allowed patients to avoid colorectal surgeries. While ESD is frequently performed in Asia, it is not commonly performed elsewhere including in the West. However, there are several compelling arguments for performance of ESD over EMR in large ( $\geq 20$  mm) polyps. In a recent meta-analysis, Lim *et al*[9] found that ESD of polyps  $\geq 20$  mm was associated to higher *en bloc* resection [relative risk (RR): 1.9, 95%CI: 1.4-2.7;  $P < 0.001$ ] and lower recurrence (RR 0.19, 95%CI: 0.09-0.43;  $P < 0.001$ ) compared to EMR[9]. Given the benefits of ESD, this has culminated in a multicenter randomized controlled trial based in France led by Jacques *et al*[10] which found ESD to be superior to EMR in *en bloc* resection as well as decreased recurrence[10]. Given advantages seen with ESD, we performed the first North American study comparing ESD to EMR.

In our retrospective comparison of ESD, EMR and knife-assisted endoscopic resection, ESD was able to achieve the highest *en bloc* resection, followed by knife-assisted endoscopic resection; EMR had the lowest *en bloc* resection rate. Recurrence rate was lowest in the ESD (1.3%) and knife-assisted endoscopic resection group (0.0%), and highest in the EMR group (12.9%). On multivariate regression, we found that performance of ESD (in comparison to EMR) significantly decreased recurrence. Increased polyp size significantly increased risk of recurrence. We were able to achieve *en bloc* resection rate of 90.4% with ESD. This is comparable to work by Gupta *et al*[1], in which overall *en bloc* resection rate was

**Table 3 Comparison of recurrence with no recurrence**

	Recurrence, <i>n</i> (%)	No recurrence, <i>n</i> (%)	<i>P</i> value
<b>Size of polyp, mm (mean ± SD)</b>	37.4 (17.1)	32.7 (13.8)	0.202
<b>Procedure</b>			0.002
ESD	1/24 (4.2)	76/263 (28.9)	
Knife-assisted endoscopic removal	0/24 (0.0)	32/263 (12.2)	
EMR	23/24 (95.8)	155/263 (58.9)	
<b><i>En bloc</i> resection</b>			0.003
<i>En bloc</i>	2/24 (8.3)	102/263 (38.8)	
Non <i>en bloc</i>	22/24 (91.7)	161/263 (61.2)	
<b>Circumferential incision</b>			0.001
Yes	1/99 (1.0)	98/99 (99.0)	
No	23/188 (12.2)	165/188 (87.8)	
<b>Prior resection</b>			0.154
Prior attempt	3/24 (12.5) (17.6)	14/263 (5.3)	
No prior attempt	21/24 (87.5)	249/263 (94.7)	
<b>Pathology</b>			0.691
Non-neoplastic	0/24 (0.0)	11/263 (4.2)	
Neoplastic, no high-grade dysplasia	19/24 (79.2)	208/263 (79.1)	
High-grade dysplasia	4/24 (16.7)	31/263 (11.8)	
Cancer	1/24 (4.2)	13/263 (4.9)	
<b>First follow-up, days (mean ± SD)</b>	498.0 (406.9)	482.0 (348.7)	0.854
<b>Complete resection</b>			0.041
R0	1/24 (4.2)	70/263 (26.6)	
R1	22/24 (91.7)	178/263 (67.7)	
Rx	1/24 (4.2)	15/263 (5.7)	

EMR: Endoscopic mucosal resection; ESD: Endoscopic submucosal dissection.

**Table 4 Patient complications, *n* (%)**

<b><i>N</i> = 376</b>	<b>ESD (<i>n</i> = 122)</b>	<b>Knife-assisted endoscopic resection (<i>n</i> = 44)</b>	<b>EMR (<i>n</i> = 216)</b>
Complication	8 (6.6)	3 (6.8)	14 (6.5)
Bleeding without intervention	3 (2.5) <sup>1</sup>	1 (2.3) <sup>1</sup>	5 (2.3)
Bleeding with intervention	3 (2.5)	1 (2.3)	1 (0.5)
SBO/partial SBO	0 (0.0)	0 (0.0)	2 (0.9)
Bowel perforation	2 (1.6)	1 (2.3)	0 (0.0)
Abdominal pain	1 (0.8) <sup>1</sup>	1 (2.3) <sup>1</sup>	3 (1.4)
Unrelated complication	0 (0.0)	0 (0.0)	3 (1.4)

<sup>1</sup>One patient had both bleeding without intervention and abdominal pain.

SBO: Small bowel obstruction; EMR: Endoscopic mucosal resection; ESD: Endoscopic submucosal dissection.

73.1%, and the rate for the second half of their study was 84.6%. Similarly, our study had ESD recurrence rate of 1.3%, slightly lower than the 4.3% (*n* = 2) by Gupta *et al*[1]. Overall, there was low risk of complication across the three procedures. Under appropriate training, we feel the three procedures to

**Table 5 Univariate Cox regression evaluating predictors of recurrence, including endoscopic submucosal dissection versus endoscopic mucosal resection**

Covariates	Unadjusted hazard ratio (95%CI)	P value
<b>Treatment type</b>		0.041
EMR, pure	Reference	
ESD, pure	0.12 (0.02-0.92)	
<b>Age, per year</b>	1.04 (0.99-1.08)	0.109
<b>Sex</b>		0.898
Female	Reference	
Male	1.06 (0.46-2.40)	
<b>Race</b>		0.139
White	Reference	
Non-White	1.85 (0.82-4.19)	
<b>Polyp location</b>		0.376
Non-rectum	Reference	
Rectum	0.52 (0.12-2.22)	
<b>Prior resection attempt</b>	2.65 (0.76-9.29)	0.127
<b>Polyp size, by mm</b>	1.03 (1.01-1.05)	0.001
<b>Presence of circumferential incision</b>	0.12 (0.02-0.92)	0.041
<b>En bloc resection</b>	0.15 (0.03-0.63)	0.010
<b>JNET classification</b>		
Type 1	Reference	
Type 2A	3.07 (0.41-22.92)	0.273
Type 2B or 3	4.57 (0.41-50.74)	0.216
<b>R0 resection</b>	0.13 (0.02-0.93)	0.042

EMR: Endoscopic mucosal resection; ESD: Endoscopic submucosal dissection; JNET: Japan NBI Expert Team.

**Table 6 Multivariate Cox regression evaluating predictors of recurrence**

Covariates	Adjusted hazard ratio (95%CI)	P value
<b>Treatment type</b>		0.014
EMR, pure	Reference	
ESD, pure	0.06 (0.01-0.57)	
<b>Polyp size, by mm</b>	1.05 (1.02-1.07)	< 0.001

EMR: Endoscopic mucosal resection; ESD: Endoscopic submucosal dissection.

be safe techniques.

While operational proficiency is related to study outcome, in this study we try to evaluate the specific factors that lead to success in reducing polyp recurrence. Specifically, we look at factors such as *en bloc* resection and performance of circumferential incision. Circumferential incision was found to be associated with decreased recurrence. In one evaluation of ESD compared to hybrid ESD (circumferential mucosal incision followed by snare resection), hybrid ESD trended towards lower *en bloc* resection rate and complete resection rate compared to ESD, though this did not reach statistical significance. However, importantly, on surveillance of hybrid ESD by the Korean specialist ( $n = 21$ ) and United States novice practitioner ( $n = 9$ ), there was no recurrence in either group[4]. While this study was limited by overall low numbers, it provided early suggestion that circumferential incision alone may help improve

the outcomes of polyp resection compared to EMR. A major advantage of knife-assisted endoscopic resection over ESD is the relative technical simplicity; in particular, circumferential incision is a relatively safe technique that in our experience is easily taught to trainees with sufficient experience in routine colonoscopy. Over time, with increased experience and proficiency performing ESD, we expect that many endoscopists will choose ESD over knife-assisted endoscopic resection to maximize *en bloc* and R0 resection, but our data highlights the generally excellent long-term results of the knife-assisted technique.

While there is justifiable concern about the risk of perforation with ESD, and the 3 cases of perforation in this series were all in the ESD/knife-assisted endoscopic resection group rather than EMR, it is notable that the perforations occurred in very challenging cases where EMR was deemed not feasible, and surgery was the only other viable option. For large lesions involving greater than half the circumference of the lumen, the Japan Gastroenterological Endoscopy Society does not recommend piecemeal EMR, but rather ESD and consideration for surgery if ESD is not endoscopically feasible[8].

There were several limitations for our study. First, retrospective data from only two endoscopists were used. However, given the lack of ESD experts in the country, having 125 cases of ESD is relatively robust. In addition, while more EMR cases could have been achieved by including other endoscopists at the two hospitals included, this would potentially introduce more bias with variation in technique and approach to EMR as well as skill with polypectomy. Another concern is the limited follow-up (67.1%). A lot of the patients were referred for endoscopic removal but received follow-up with the referring provider. Despite reaching out to community providers, we only received limited response. Further, the retrospective nature of EMR and ESD studies introduce selection bias in the determination of which polyp to undergo EMR, ESD, or knife-assisted endoscopic resection. A randomized clinical trial would be ideal but is logistically challenging given the overall low frequency of these procedures.

## CONCLUSION

In this multicenter study evaluating ESD, knife-assisted endoscopic resection and EMR, ESD and knife-assisted endoscopic resection were able to achieve higher rates of *en bloc* resection and was able to achieve significantly lower risk of recurrence compared to EMR. Given the results of this study, ESD and knife-assisted endoscopic resection should be strongly considered when possible for polyps  $\geq 20$  mm to improve *en bloc* and curative resection and decrease risk of recurrence.

## ARTICLE HIGHLIGHTS

### Research background

Adoption of endoscopic submucosal dissection (ESD) has been slow in the United States, largely related to lack of experts, long training required and significant time for procedure compared to endoscopic mucosal resection (EMR).

### Research motivation

In this study, we seek to evaluate our experience of ESD compared to EMR in California.

### Research objectives

We evaluate ESD, knife-assisted endoscopic resection as well as EMR to identify factors for recurrence.

### Research methods

This was a retrospective comparison performed at two tertiary centers within California between 2016 and 2020. Adult patients that received colonoscopy with endoscopic removal of a polyp at least 20 mm in size were included. Primary outcome of interest was recurrence on follow-up.

### Research results

ESD achieved highest *en bloc* resection followed by knife-assisted endoscopic resection and EMR. On follow-up, recurrence rate was lowest in knife-assisted endoscopic resection (0.0%) and ESD (1.3%), while EMR had the highest recurrence rate (12.9%,  $P = 0.0017$ ).

### Research conclusions

In our study, we found that EMR had significantly higher recurrence compared to ESD or knife-assisted endoscopic resection.

### Research perspectives

We have demonstrated efficacy of ESD in a Western population.

## FOOTNOTES

**Author contributions:** Wei MT contributed to project conception, data collection, manuscript writing, data analysis, manuscript revision; Zhou MJ, Li AA and Ofosu A contributed to data analysis, manuscript revision; Hwang JH and Friedland S contributed to project conception, data collection, manuscript revision.

**Institutional review board statement:** This study was performed under the approval of the Institutional Review Board at Stanford University, Stanford, California, USA.

**Informed consent statement:** Because of retrospective study signed informed consent form is not needed.

**Conflict-of-interest statement:** Mike T. Wei: Consultant for Neptune Medical, AgilTx, Capsovision; Margaret Zhou and Andrew Ofosu: No conflicts; Andrew Li: Consultant for Neptune Medical; Joo Ha Hwang: Consultant for Olympus, Medtronic, Boston Scientific, Lumendi, Fujifilm, Noah Medical, Neptune Medical, and Micro-Tech; Shai Friedland: Consultant for Intuitive Surgical and Capsovision.

**Data sharing statement:** No additional data are available.

**STROBE statement:** The authors have read the STROBE Statement—checklist of items, and the manuscript was prepared and revised according to the STROBE Statement—checklist of items.

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** United States

**ORCID number:** Mike T Wei 0000-0003-4756-9010; Margaret J Zhou 0000-0002-9111-1659; Joo Ha Hwang 0000-0002-7534-230X; Shai Friedland 0000-0002-9782-000X.

**S-Editor:** Fan JR

**L-Editor:** A

**P-Editor:** Fan JR

## REFERENCES

- Gupta N, Rodríguez-Ruiz G, Siddiqui UD, Chapman CG, Donboli K, Hart J, Xiao SY, Waxman I. Endoscopic submucosal dissection for colorectal lesions: outcomes from a United States experience. *Surg Endosc* 2022; **36**: 236-243 [PMID: 33523276 DOI: 10.1007/s00464-020-08262-4]
- Herreros de Tejada A. ESD training: A challenging path to excellence. *World J Gastrointest Endosc* 2014; **6**: 112-120 [PMID: 24748918 DOI: 10.4253/wjge.v6.i4.112]
- Draganov PV, Aihara H, Karasik MS, Ngamruengphong S, Aadam AA, Othman MO, Sharma N, Grimm IS, Rostom A, Elmunzer BJ, Jawaid SA, Westerveld D, Perbtani YB, Hoffman BJ, Schlachterman A, Siegel A, Coman RM, Wang AY, Yang D. Endoscopic Submucosal Dissection in North America: A Large Prospective Multicenter Study. *Gastroenterology* 2021; **160**: 2317-2327.e2 [PMID: 33610532 DOI: 10.1053/j.gastro.2021.02.036]
- Yang DH, Kwak MS, Park SH, Ye BD, Byeon JS, Myung SJ, Yang SK, Kim HG, Friedland S. Endoscopic Mucosal Resection with Circumferential Mucosal Incision for Colorectal Neoplasms: Comparison with Endoscopic Submucosal Dissection and between Two Endoscopists with Different Experiences. *Clin Endosc* 2017; **50**: 379-387 [PMID: 28264251 DOI: 10.5946/ce.2016.058]
- Sano Y, Tanaka S, Kudo SE, Saito S, Matsuda T, Wada Y, Fujii T, Ikematsu H, Uraoka T, Kobayashi N, Nakamura H, Hotta K, Horimatsu T, Sakamoto N, Fu KI, Tsuruta O, Kawano H, Kashida H, Takeuchi Y, Machida H, Kusaka T, Yoshida N, Hirata I, Terai T, Yamano HO, Kaneko K, Nakajima T, Sakamoto T, Yamaguchi Y, Tamai N, Nakano N, Hayashi N, Oka S, Iwatate M, Ishikawa H, Murakami Y, Yoshida S, Saito Y. Narrow-band imaging (NBI) magnifying endoscopic classification of colorectal tumors proposed by the Japan NBI Expert Team. *Dig Endosc* 2016; **28**: 526-533 [PMID: 26927367 DOI: 10.1111/den.12644]
- The Paris endoscopic classification of superficial neoplastic lesions: esophagus, stomach, and colon: November 30 to December 1, 2002. *Gastrointest Endosc* 2003; **58**: S3-43 [PMID: 14652541 DOI: 10.1016/s0016-5107(03)02159-x]
- Inoue T, Nakagawa K, Yamasaki Y, Shichijo S, Kanesaka T, Maekawa A, Higashino K, Uedo N, Ishihara R, Takeuchi Y. Underwater endoscopic mucosal resection versus endoscopic submucosal dissection for 20-30 mm colorectal polyps. *J Gastroenterol Hepatol* 2021; **36**: 2549-2557 [PMID: 33724540 DOI: 10.1111/jgh.15494]
- Tanaka S, Kashida H, Saito Y, Yahagi N, Yamano H, Saito S, Hisabe T, Yao T, Watanabe M, Yoshida M, Saitoh Y, Tsuruta O, Sugihara KI, Igarashi M, Toyonaga T, Ajioka Y, Kusunoki M, Koike K, Fujimoto K, Tajiri H. Japan Gastroenterological Endoscopy Society guidelines for colorectal endoscopic submucosal dissection/endoscopic mucosal resection. *Dig Endosc* 2020; **32**: 219-239 [PMID: 31566804 DOI: 10.1111/den.13545]

- 9 **Lim XC**, Nistala KRY, Ng CH, Lin SY, Tan DJH, Ho KY, Chong CS, Muthiah M. Endoscopic submucosal dissection vs endoscopic mucosal resection for colorectal polyps: A meta-analysis and meta-regression with single arm analysis. *World J Gastroenterol* 2021; **27**: 3925-3939 [PMID: [34321855](#) DOI: [10.3748/wjg.v27.i25.3925](#)]
- 10 **Jacques J**, Wallenhorst T, Chevaux JB, Lépilliez DV, Chaussade S, Rivory J, Legros R, Schaefer M, Leblanc S, Rostain F, Barret M, Belle A, Crepin S, Magne J, Albouys J, Preux PM, Lepetit H, Dahan M, Ponchon T, Pioche M. Endoscopic submucosal dissection (ESD) vs piece-meal endoscopic mucosal resection (PM-EMR) for large laterally spreading lesions: French randomized controlled trial RESECT-COLON (Abstract). *Gastrointest Endosc* 2022; **95** Suppl 6 [DOI: [10.1016/j.gie.2022.04.384](#)]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

**Telephone:** +1-925-3991568

**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

**Help Desk:** <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

