

# World Journal of *Hepatology*

*World J Hepatol* 2019 July 27; 11(7): 562-612



**REVIEW**

- 562 Rise of sodium-glucose cotransporter 2 inhibitors in the management of nonalcoholic fatty liver disease  
*Dokmak A, Almeqdadi M, Trivedi H, Krishnan S*

**MINIREVIEWS**

- 574 Role of innovative 3D printing models in the management of hepatobiliary malignancies  
*Bangeas P, Tsioukas V, Papadopoulos VN, Tsoulfas G*

**ORIGINAL ARTICLE****Retrospective Study**

- 586 Epidemiology and outcomes of acute liver failure in Australia  
*Hey P, Hanrahan TP, Sinclair M, Testro AG, Angus PW, Peterson A, Warrillow S, Bellomo R, Perini MV, Starkey G, Jones RM, Fink M, McClure T, Gow P*

**SYSTEMATIC REVIEW**

- 596 Spontaneous fungal peritonitis: Micro-organisms, management and mortality in liver cirrhosis-A systematic review  
*Tariq T, Irfan FB, Farishta M, Dykstra B, Sieloff EM, Desai AP*

**CASE REPORT**

- 607 Wilson disease developing osteoarthritic pain in severe acute liver failure: A case report  
*Kido J, Matsumoto S, Sugawara K, Nakamura K*

**ABOUT COVER**

Associate Editor of *World Journal of Hepatology*, Yong-Ping Yang, MD, Professor, Center of Therapeutic Research for Hepatocellular Carcinoma, the Fifth Medical Center of Chinese PLA General Hospital, Beijing 100039, China

**AIMS AND SCOPE**

*World Journal of Hepatology (World J Hepatol, WJH, online ISSN 1948-5182, DOI: 10.4254)*, is a peer-reviewed open access academic journal that aims to guide clinical practice and improve diagnostic and therapeutic skills of clinicians.

The *WJH* covers topics concerning liver biology/pathology, cirrhosis and its complications, liver fibrosis, liver failure, portal hypertension, hepatitis B and C and inflammatory disorders, steatohepatitis and metabolic liver disease, hepatocellular carcinoma, etc. Priority publication will be given to articles concerning diagnosis and treatment of hepatology diseases. The following aspects are covered: Clinical diagnosis, laboratory diagnosis, etc.

We encourage authors to submit their manuscripts to *WJH*. We will give priority to manuscripts that are supported by major national and international foundations and those that are of great basic and clinical significance.

**INDEXING/ABSTRACTING**

The *WJH* is now abstracted and indexed in PubMed, PubMed Central, Emerging Sources Citation Index (Web of Science), Scopus, China National Knowledge Infrastructure (CNKI), China Science and Technology Journal Database (CSTJ), and Superstar Journals Database.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Responsible Electronic Editor: *Yan-Liang Zhang*

Proofing Production Department Director: *Yun-Xiaojuan Wu*

**NAME OF JOURNAL**

*World Journal of Hepatology*

**ISSN**

ISSN 1948-5182 (online)

**LAUNCH DATE**

October 31, 2009

**FREQUENCY**

Monthly

**EDITORS-IN-CHIEF**

Ke-Qin Hu, Koo Jeong Kang, Nikolaos T Pysopoulos

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/1948-5182/editorialboard.htm>

**EDITORIAL OFFICE**

Ya-Juan Ma, Director

**PUBLICATION DATE**

July 27, 2019

**COPYRIGHT**

© 2019 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>

## Retrospective Study

## Epidemiology and outcomes of acute liver failure in Australia

Penelope Hey, Timothy P Hanrahan, Marie Sinclair, Adam G Testro, Peter W Angus, Adam Peterson, Stephen Warrillow, Rinaldo Bellomo, Marcos V Perini, Graham Starkey, Robert M Jones, Michael Fink, Tess McClure, Paul Gow

**ORCID number:** Penelope Hey (0000-0002-5219-0382); Timothy P Hanrahan (0000-0002-3682-7400); Marie Sinclair (0000-0003-0657-3048); Adam G Testro (0000-0001-6776-3115); Peter W Angus (0000-0001-8505-2317); Adam Peterson (0000-0002-8052-292X); Stephen Warrillow (0000-0002-7240-4106); Rinaldo Bellomo (0000-0002-1650-8939); Marcos V Perini (0000-0002-0165-1564); Graham Starkey (0000-0002-4285-1343); Robert M Jones (0000-0002-8415-5565); Michael Fink (0000-0002-9684-4691); Tess McClure (0000-0003-2810-7443); Paul Gow (0000-0001-6505-7233).

**Author contributions:** All authors have contributed to this manuscript and have agreed on the content; Hey P, Hanrahan T and Gow P were involved in the study design; Hey P, Hanrahan T and Peterson A were involved with data collection and analysis; Hanrahan H provided statistical support; Hey P, Gow P, Hanrahan T, Sinclair M, Testro A, Warrillow S, Bellomo R, Perini M, Starkey G, Jones R, Fink M and McClure T were involved with data interpretation, drafting and revising the work; All authors provided approval for the final version to be published.

**Institutional review board statement:** This study was approved through the Austin Health Human Research Ethics Committee.

**Informed consent statement:**

Penelope Hey, Timothy P Hanrahan, Marie Sinclair, Adam G Testro, Peter W Angus, Adam Peterson, Marcos V Perini, Graham Starkey, Robert M Jones, Tess McClure, Paul Gow, Liver Transplant Unit, Austin Health, Heidelberg 3084, Australia

Penelope Hey, Marie Sinclair, Adam G Testro, Peter W Angus, Stephen Warrillow, Rinaldo Bellomo, Marcos V Perini, Graham Starkey, Robert M Jones, Tess McClure, Paul Gow, The University of Melbourne, Melbourne 3052, Australia

Stephen Warrillow, Rinaldo Bellomo, Department of Intensive Care, Austin Health, Melbourne 3084, Australia

Michael Fink, Department of Surgery, Austin Health, Melbourne 3084, Australia

**Corresponding author:** Penelope Hey, MBBS, BMedSci, Liver Transplant Unit, Austin Health, 145 Studley Road, Heidelberg 3084, Australia. [penelope.hey@austin.org.au](mailto:penelope.hey@austin.org.au)

**Telephone:** +61-3-94965353

**Fax:** +61-3-94963487

## Abstract

### BACKGROUND

Acute liver failure (ALF) is a life-threatening syndrome with varying aetiologies requiring complex care and multidisciplinary management. Its changing incidence, aetiology and outcomes over the last 16 years in the Australian context remain uncertain.

### AIM

To describe the changing incidence, aetiology and outcomes of ALF in South Eastern Australia.

### METHODS

The database of the Victorian Liver Transplant Unit was interrogated to identify all cases of ALF in adults (> 16 years) in adults hospitalised between January 2002 and December 2017. Overall, 169 patients meeting criteria for ALF were identified. Demographics, aetiology of ALF, rates of transplantation and outcomes were collected for all patients. Transplant free survival and overall survival (OS) were assessed based on survival to discharge from hospital. Results were compared to data from a historical cohort from the same unit from 1988-2001.

### RESULTS

Paracetamol was the most common aetiology of acute liver failure, accounting for

Patients were not required to give informed consent to the study because the analysis used anonymous clinical data collected as a clinical Audit.

**Conflict-of-interest statement:** All authors declare no conflict-of-interest related to this article.

**Data sharing statement:** No additional data are available.

**Open-Access:** This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

**Manuscript source:** Unsolicited manuscript

**Received:** April 23, 2019

**Peer-review started:** April 23, 2019

**First decision:** June 5, 2019

**Revised:** June 19, 2019

**Accepted:** July 4, 2019

**Article in press:** July 5, 2019

**Published online:** July 27, 2019

**P-Reviewer:** Auzinger G, He SQ, Zheng YW

**S-Editor:** Ma YJ

**L-Editor:** A

**E-Editor:** Zhang YL



50% of cases, with an increased incidence compared with the historical cohort ( $P = 0.046$ ). Viral hepatitis and non-paracetamol drug or toxin induced liver injury accounted for 15% and 10% of cases respectively. Transplant free survival (TFS) improved significantly compared to the historical cohort (52% vs 38%,  $P = 0.032$ ). TFS was highest in paracetamol toxicity with spontaneous recovery in 72% of cases compared to 31% of non-paracetamol ALF ( $P < 0.001$ ). Fifty-nine patients were waitlisted for emergency liver transplantation. Nine of these died while waiting for an organ to become available. Forty-two patients (25%) underwent emergency liver transplantation with a 1, 3 and 5 year survival of 81%, 78% and 72% respectively.

## CONCLUSION

Paracetamol toxicity is the most common aetiology of ALF in South-Eastern Australia with a rising incidence over 30 years. TFS has improved, however it remains low in non-paracetamol ALF.

**Key words:** Liver failure; Acute; Paracetamol; Australia; Victoria; Liver transplant

©The Author(s) 2019. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core tip:** Acute liver failure (ALF) is a life-threatening syndrome with varying aetiologies based on geographic location. Paracetamol is the most common cause of ALF in South Eastern Australia with a rising incidence over 30 years. Despite this, transplantation for paracetamol induced ALF is lower than other large centres at 4% with a comparable overall survival. Non-paracetamol ALF however portends a poor prognosis with less than one third surviving without emergency liver transplantation.

**Citation:** Hey P, Hanrahan TP, Sinclair M, Testro AG, Angus PW, Peterson A, Warrillow S, Bellomo R, Perini MV, Starkey G, Jones RM, Fink M, McClure T, Gow P. Epidemiology and outcomes of acute liver failure in Australia. *World J Hepatol* 2019; 11(7): 586-595

**URL:** <https://www.wjgnet.com/1948-5182/full/v11/i7/586.htm>

**DOI:** <https://dx.doi.org/10.4254/wjh.v11.i7.586>

## INTRODUCTION

Acute liver failure (ALF) is a clinical syndrome characterised by severe liver injury associated with the development of coagulopathy and hepatic encephalopathy in the absence of known pre-existing liver disease<sup>[1]</sup>. The causes of ALF differ markedly based on geographic location<sup>[2]</sup>. In developing countries viral hepatitis is the most common aetiology, whereas in the developed world, the majority of ALF cases are due to paracetamol poisoning and / or other drug reactions<sup>[3-5]</sup>.

Historically, ALF was associated with low rates of spontaneous survival and in regions with transplant programs, the majority of eligible patients underwent emergency liver transplantation (ELT)<sup>[6]</sup>. However, more recently, transplant-free survival (TFS) has improved<sup>[3,7,8]</sup>. In the United States, rates of liver transplantation for ALF have been reported to be as low as 20% with an overall survival to hospital discharge of 75%<sup>[8]</sup>. For non-paracetamol aetiologies, however, TFS is less than 30%, and ELT has an established survival benefit<sup>[9]</sup>. In contrast, for paracetamol induced ALF, TFS is much higher and the survival advantage of transplantation is less clear<sup>[10]</sup>.

The Victorian Liver Transplant Unit (VLTU) provides liver transplantation and quaternary hepatology services for a population of 6.7 million in South-East Australia, representing 27% of the total Australian population<sup>[11]</sup>. Essentially all cases of ALF are referred to the VLTU, with only very few patients either not referred or dying prior to transfer. King's College Criteria is used to determine suitability for liver transplantation.

The primary aim of this study was to report the aetiology, incidence and outcomes of all adult cases of ALF presenting to the VLTU over the last 16 years. Our secondary aim was to compare current data to historical data by the same unit<sup>[12]</sup>. We hypothesised that the incidence and underlying causes of ALF across the population would be unchanged, but that outcomes would be improved despite relatively low utilization of ELT.

## MATERIALS AND METHODS

All adult patients with ALF aged greater than 16 years managed at the VLTU between January 01, 2002 and December 31, 2017 were included in this study. ALF was defined as acute liver injury with the development of coagulopathy and hepatic encephalopathy within 26 wk of onset of symptoms, in the absence of known chronic liver disease<sup>[43]</sup>. Data were extracted from the VLTU database for comparison with a previously published historical cohort<sup>[42]</sup>. Patients with previous liver transplantation and primary non-function of the graft were excluded.

Information was collected from the unit database and crosschecked against medical records. The aetiology of ALF was determined by the treating team based on clinical history, paracetamol levels, viral and autoimmune serology, metabolic testing and, if available, histology from liver biopsy or explant specimens. Indeterminate ALF, also known as non-A non-B hepatitis or seronegative hepatitis, was diagnosed when all other aetiologies had been excluded.

Demographics, aetiology of ALF, rates of transplantation and outcomes were collected for all patients. Transplant free survival and overall survival (OS) were assessed based on survival to discharge from hospital. Patients who recovered without the need for liver transplantation were typically discharged back to the care of the referring health service and therefore, long-term outcomes are unknown. Medical records of patients who undergo liver transplantation, however, are regularly updated and were used to calculate post-transplant survival.

This study was approved through the Austin Health Human Research Ethics Committee.

### Statistical analysis

The incidence of ALF was calculated based on annual local population data available from the Australian Bureau of Statistics. Where data was normally distributed (by Shapiro-Wilk analysis), two-tailed student *t*-tests were used to compare for significance. Data not meeting normality criteria were reported as median (IQR) and compared using the Mann-Whitney *U* test. To assess differences in endpoints and changes in aetiology over time, data were first analysed by comparison of the two cohorts, and then as a continuous set. For categorical data, comparison was made using chi-square test. Time-series analysis was completed using linear regression.

## RESULTS

Records of 221 patients who had a diagnosis of acute liver injury on the liver transplant database were reviewed. Twenty-one patients were excluded due to the absence of hepatic encephalopathy. Twenty-two cases were excluded as they had underlying cirrhosis with acute decompensation and four cases because they had been transplanted at other centres. Five patients were excluded because they suffered ALF secondary to primary graft non-function after liver transplantation for non-ALF indications.

One hundred and sixty-nine patients met the inclusion criteria. One hundred and thirty-two cases (78%) were female and 37 (22%) were male. The median age at presentation was 41 years [31;52] for females and 37 years [27;49] for males. The median rate of referral over the study period was 9 cases per year, or 1.6 cases per million population per year. Twenty-two patients (13%) presented directly to our centre, 118 cases (70%) were referred from other metropolitan hospitals and 29 (17%) from regional and rural hospitals. One hundred and fifty-seven patients (92.9%) required support in the intensive care unit during their admission.

### Aetiology

Paracetamol toxicity was the most common cause of ALF accounting for a total of 84 cases (50%) (Table 1). The median rate of paracetamol induced ALF referred to the centre over the 16-year study period was 0.7 cases per million population per year<sup>[41]</sup>. Non-paracetamol drug or toxin induced liver injury was the cause of ALF in 17 patients (10%). Viral hepatitis was the cause of ALF in 26 cases (15%). This included 20 cases of fulminant hepatitis B virus (HBV) infection and two from reactivated chronic HBV following systemic chemotherapy. Hepatitis A virus (HAV) was identified in four cases. Other viral aetiologies included herpes simplex virus-2 (HSV) and varicella zoster virus both in isolated cases.

Fulminant autoimmune hepatitis was diagnosed in nine patients (5%). The diagnosis was made based on a combination of positive auto-antibody testing, elevated IgG levels and histology results either during the episode of ALF or after recovery. ALF secondary to malignancy was identified in four patients. Veno-

**Table 1** Drugs and toxins implicated in cases of acute liver failure managed at the Victorian Liver Transplant Unit from 2002-2017

Drug/Toxins	Cases of ALF	Waitlisted for ELT	ELT
	(n = 101)	(n = 21)	(n = 13)
Paracetamol	84	11	3
Antibiotics	4	2	2
Amoxicillin/clavulanate (2), clarithromycin, isoniazid			
Infliximab	2	2	2
Illicit drugs	4	1	1
LSD, injected buprenorphine, amphetamines, MDMA			
NSAIDs	1	1	1
Amanita phalloides	1	1	1
Herbal medicines			
Black cohosh herb, kava kava	2	2	2
Other			
Moxonidine, fenofibrate, chlorambucil	3	1	1

ALF: Acute liver failure; ELT: Emergency liver transplantation; NSAIDs: Non-steroid anti-inflammatory drugs; LSD: Lysergic acid diethylamide; MDMA: Methylenedioxyamphetamine; NSAIDs: Non-steroidal anti-inflammatory drugs

occlusive disease caused ALF in one case following allogeneic stem cell transplantation. Wilson's disease was the cause of ALF in two cases. Severe ischemic hepatitis due to hepatic artery injury and extensive portal vein thrombosis resulted in ALF in four patients.

ALF occurred in five patients during pregnancy. Aetiologies included acute fatty liver disease of pregnancy, HSV hepatitis and pregnancy-precipitated liver failure from a urea-cycle disorder<sup>[14]</sup>. Unintentional paracetamol poisoning causing ALF occurred in two cases in the context of hyperemesis and malnutrition. One of these cases presented with foetal death in utero in the third trimester of pregnancy, however in all other pregnancies, the infants survived. Twenty-one cases (12%) were classified as indeterminate ALF after exclusion of other aetiologies.

### Survival

Over the study period the TFS was 52% and OS to discharge from hospital was 72%. TFS was significantly higher in patients with paracetamol induced ALF at 74% compared to non-paracetamol aetiologies at 31% ( $P < 0.0001$ ). OS however did not differ significantly at 77% and 67% for paracetamol and non-paracetamol aetiologies respectively ( $P = 0.13$ ). ALF caused by indeterminate hepatitis had the lowest TFS at 10%.

### Transplantation

Fifty-nine patients (35%) were waitlisted for transplantation (Figure 1). Eight patients improved and were delisted, however one subsequently died of sepsis. One patient was delisted following an intraoperative finding of ischemic gut and eight patients died while waiting for a donor liver (14%). 42 patients (25%) underwent liver transplantation for ALF. Rates of liver transplantation were highest in patients with indeterminate hepatitis at 67%. Only three patients with paracetamol induced ALF were transplanted (4%).

After medical and psychosocial assessment, liver transplantation was contraindicated in 24 patients. Of these, fifteen cases were deemed medically unsuitable for transplantation. Medical contraindications included uncontrolled sepsis, ischemic bowel, intracranial events, active malignancy and known severe coronary artery disease. Only three (20%) of these patients recovered and survived to discharge. Six patients had psychiatric or psychosocial contraindications and three patients had concurrent heavy alcohol intake rendering them unsuitable for transplantation. Paracetamol was the cause of ALF in eight of these patients (89%). Four (44%) recovered and were discharged from hospital.

For patients undergoing liver transplantation for ALF, 1, 3 and 5-year survival was 81%, 78% and 72% respectively. Seven patients (17%) died within 30 d of transplantation. The aetiology of ALF for such patients included indeterminate hepatitis (2), AIH (2), drug and toxins (2) and hepatic arterial injury leading to




Figure 1 Flowchart of outcomes of patients meeting criteria for acute liver failure.

ischemic hepatitis (1). Causes of death post operatively included sepsis in three cases, cardiovascular or cerebral events in three patients and hepatic artery thrombosis in one case.

**Comparison over time**

Compared to historical data published by the unit from 1988-2001<sup>[12]</sup>, there was no difference in age at presentation or gender distribution (Table 2). Paracetamol poisoning remained the most common aetiology over the two time periods. There was a significant increase in the rate of paracetamol-induced ALF in this current data set compared to the historical cohort (36% vs 50%,  $P = 0.046$ ). When presented as continuous data, linear regression analysis also identified a significant increase in paracetamol-induced ALF per capita across the 30 years ( $R^2 = 0.275$ ;  $F(1,28) = 10.643$ ,  $P = 0.003$ ) (Figure 2). Overall ALF referrals to the unit also increased significantly over the time period ( $R^2 = 0.178$ ;  $F(1,28) = 6.074$ ,  $P = 0.020$ ). Comparing this dataset to historical data (1988-2001) also identified no differences in rates of viral hepatitis.

TFS to discharge from hospital improved in the present study compared to the historical cohort from 38% to 52% ( $P = 0.032$ ). An improvement in TFS was observed in both paracetamol toxicity (69% vs 74%,  $P = 0.614$ ) and non-paracetamol aetiologies (20% vs 31%,  $P = 0.160$ ) although neither met significance (Table 3). Overall, there was no statistically significant difference in hospital survival (63% to 72%,  $P = 0.122$ ) or transplantation rates for ALF (33% to 25%,  $P = 0.206$ ).

**DISCUSSION**

This cohort study represents the largest modern series of patients presenting with ALF in Australia. As the sole liver transplant unit responsible for a population of 6.7 million, the VLTU captured both the incidence and outcomes of this rare but life-threatening syndrome. We identified paracetamol toxicity as the commonest cause of ALF, responsible for 50% of presentations. We also found an increasing rate of paracetamol induced ALF referrals to our centre over a 30-year period. Finally, we found that transplant-free survival for both paracetamol toxicity and non-paracetamol aetiologies has improved, but that spontaneous survival for non-paracetamol ALF remains low. This is particularly the case for indeterminate hepatitis and non-paracetamol drug-induced liver injury, highlighting the need for early referral to a transplant centre.

Our reported prevalence of paracetamol-induced ALF is similar to series from the United Kingdom and United States with rates ranging from 37%-79%<sup>[3,4,15]</sup>. However, of concern, this is the only cohort to show an overall steady increase in its incidence. These results differ from local Victorian data demonstrating a decrease in hospital admissions for non-ALF paracetamol poisoning from 2000 to 2007<sup>[16]</sup>. In the United States, rates of ALF due to paracetamol poisoning have also fallen following the introduction of paracetamol sales restrictions in 1998<sup>[3,17]</sup>. These regulations limited pharmacy sales to a maximum of 32 tablets per pack. While restrictions in non-pharmacy sales of paracetamol have been in place in Australia since 2013, packets of

**Table 2 Demographics and aetiology of acute liver failure managed at the Victorian Liver Transplant Unit in a historical cohort<sup>†</sup> compared to current series**

	1988-2001 <sup>†</sup> (n = 80)		2002-2017 (n = 169)		P value
	n	%	n	%	
Referral rate <sup>‡</sup> , median [IQR]	1.2 [0.6;1.6]		1.6 [1.3;1.7]		0.020
Age, yr, median [IQR]	36 [27.0;48.0]		40 [30.0;52.0]		0.168
Gender					
Male	16	20.0	37	21.9	
Female	64	80.0	132	78.1	0.733
Aetiology					
Paracetamol	29	36.3	84	49.7	0.046
Viral hepatitis	11	13.8	26	15.3	0.734
Hepatitis B	8	10.0	20	11.8	0.669
Hepatitis A	3	3.8	4	2.4	0.538
Varicella zoster	0	0.0	1	0.1	0.491
Herpes simplex	0	0.0	1	0.1	0.491
Non-paracetamol drug/toxin	5	6.3	17	10.1	0.323
Indeterminate	27	33.8	21	12.4	<0.001
Autoimmune hepatitis	0	0.0	9	5.3	0.036
Other	8	10.0	12	7.1	0.432
Waitlisted for transplantation	35	43.8	59	33.5	0.179
Waitlist mortality	9	25.7	8	13.6	0.139
Liver transplantation	26	32.5	42	24.9	0.206
Transplant-free survival	30	37.5	88	52.1	0.032
Overall hospital survival	50	62.5	122	72.2	0.122


<sup>†</sup>Gow *et al*<sup>[12]</sup>, 2001;<sup>‡</sup>Referral rate per million population per year. IQR: Interquartile range.

up to 100 tablets are still readily available in pharmacies. Risk factors for ALF from paracetamol include prolonged fasting and malnutrition often resulting in inadvertent toxicity<sup>[18]</sup> and combination pain relief products can also result in accidental overdose<sup>[16]</sup>. Improved awareness and public health strategies to address these factors may curtail this trend in life-threatening paracetamol toxicity.

Viral hepatitis was the second most common cause of ALF in this series. HBV accounted for 12% of all cases of ALF with no change compared to the historical cohort. This is in contrast to the declining incidence of HBV-induced ALF in other developed countries<sup>[3,19,20]</sup>. This may in part reflect the fact that many older Australians are still not vaccinated against the virus<sup>[21]</sup>. Also, Australia has high rates of immigration from countries in Asia where HBV remains endemic. Fulminant flares of HBV in the setting of inappropriately ceasing anti-viral therapy was a common background scenario in this patient group. Additionally, two patients in this cohort developed ALF from HBV infection during treatment of haematological malignancies with systemic chemotherapy. In both cases, reactivation of HBV was fatal. This emphasises the importance of early identification of patients at risk, monitoring and prophylaxis with nucleoside analogues where appropriate<sup>[22]</sup>.

Hepatitis A was the cause of ALF in just four (2%) of our cases. HAV vaccination is not routine in Australia and is recommended only in high-risk populations including men who have sex with men and travellers to endemic areas. However, outbreaks do occur in a predominantly non-immune population<sup>[23]</sup>. Of note, three of the HAV induced ALF cases occurred in the same year, coinciding with a large food-borne outbreak of HAV in Victoria<sup>[24]</sup>. While HAV and HBV were uncommon causes of ALF, mortality and morbidity were high with less than 40% of this patient population surviving without transplantation.

In this series we demonstrated that patients with paracetamol induced ALF have high rates of spontaneous survival and are unlikely to require transplantation. Indeed, rates of transplantation for paracetamol poisoning in this cohort are lower than reported by other large centres at 8%-25% but with comparable overall survival<sup>[3,15,18]</sup>.



**Figure 2** Rate of paracetamol induced acute liver failure per capita.

The role of ELT in paracetamol induced ALF has been questioned for more than 25 years<sup>[25]</sup>. King's College Hospital have been the world leaders in defining patients with paracetamol induced ALF who will die without transplantation and these criteria have been debated and refined over the last 30 years<sup>[1,10,26-28]</sup>. More recently the King's group have questioned whether transplantation plays any role in paracetamol induced ALF<sup>[28]</sup>. Our own policy is to consider transplantation only for the small subgroup of patients who have progressive coagulopathy and hyperlactatemia despite 48-72 hours of supportive treatment.

Over the 30-year period under review, TFS for ALF improved significantly. This finding has been observed in other large cohorts<sup>[3,4,8,15]</sup>. Early recognition of ALF and referral to specialised transplant centres may be one factor that has improved TFS. Supportive care for ALF within intensive care units has also evolved. This is particularly the case in the monitoring and management of cerebral oedema. For the last decade, we have had a protocolised approach to the management of this patient group with the aim of minimizing deaths from cerebral oedema. This combination approach, termed "quadruple-H therapy" comprises hyperventilation, hypernatremia, hypothermia and hemodiafiltration<sup>[29]</sup>. This multimodal approach has minimised the use of invasive intracranial pressure monitoring devices. Death attributed to cerebral oedema occurred in just one patient (0.6%) in this series.

Despite improvements in supportive care, transplantation still plays a pivotal role in this condition. Organ allocation in Australia is typically State based. However, patients with ALF who are ventilated in intensive care are classified as a category 1 and are allocated the next suitable organ within Australia or New Zealand<sup>[30]</sup>. Despite this approach, waitlist mortality remained high. There is often a brief window period where patients with ALF can be transplanted before they develop refractory multiple organ failure. Therefore, timely availability of a suitable donor organ for waitlisted patients remains critical.

This is the largest Australian cohort to report on the changing incidence and aetiology of ALF. We identified a concerning rise in paracetamol toxicity causing ALF. However, despite the high incidence of paracetamol induced ALF, we report on our very low use of ELT in this group of patients despite a comparable overall survival to other large international centres.

This study has several limitations. Firstly, this is a retrospective study, which interrogated prospectively recorded data. The diagnosis of ALF requires the presence of hepatic encephalopathy, which relied on the adequate documentation of this clinical finding in medical records. Additionally, these data did not include patients who were treated at other centres. It is possible that death prior to transfer, suicidal intent, substance abuse or significant comorbidities deemed unsuitable for transplantation may have prevented some patients being referred to the VLTU.

In conclusion, paracetamol toxicity remains the most common aetiology of ALF, with increasing rates over time, highlighting the need for public health measures to reduce this preventable cause. TFS has improved which may reflect advances in

**Table 3 Outcomes of acute liver failure managed at the Victorian Liver Transplant Unit in a historical cohort<sup>†</sup> compared to the current series**

Aetiology	TFS (%)				P value	OS (%)				P value
	1988-2001 <sup>1</sup>		2002-2017			1988-2001 <sup>1</sup>		2002-2017		
	n	%	n	%	n	%	n	%		
Paracetamol	20	69.0	62	73.8	0.614	21	72.4	65	77.4	0.589
Non-paracetamol	10	19.6	26	30.6	0.160	29	56.9	57	67.1	0.230
Viral hepatitis	2	18.2	10	38.4	0.228	5	45.5	18	69.2	0.173
HBV	2	25.0	7	35.0	0.609	4	50.0	14	70.0	0.318
HAV	0	0.0	3	75.0	0.047	1	0.33	3	75.0	0.270
Drug/toxin induced	3	60.0	5	29.4	0.211	4	80.0	12	70.6	0.678
Autoimmune hepatitis	0	0.0	3	33.3	-	0	0.0	6	66.7	-
Indeterminate hepatitis	5	18.5	2	9.5	0.381	15	55.5	14	66.7	0.435
Other	0	0.0	6	50.0	0.017	5	62.5	7	58.3	0.852

<sup>1</sup>Gow *et al*<sup>[12]</sup>, 2001. TFS: Transplant-free survival; OS: Overall hospital survival; HBV: Hepatitis B virus; HAV: Hepatitis A virus.

supportive care measures. Regardless, the majority of cases of non-paracetamol ALF still require transplantation and therefore early referral to a specialised transplant centre remains imperative.

## ARTICLE HIGHLIGHTS

### Research background

Acute liver failure (ALF) is a rare clinical syndrome with varying aetiologies based on geographic location. This condition is associated with high morbidity and mortality, and emergency liver transplantation is often life-saving.

### Research motivation

In Australia, published data from 1988-2001 demonstrated that paracetamol toxicity was the major cause of ALF, followed by non-A non-B hepatitis. An updated analysis of aetiologies and outcomes in an Australian context is therefore required.

### Research objections

This study aimed to provide a description of the aetiologies and outcomes of acute liver failure presenting to a large Australian liver transplant centre. We also aimed to describe changes over the past thirty years since the availability of liver transplantation for this condition.

### Research methods

This is a retrospective cohort study of all patients admitted to the Victorian Liver Transplant Unit from 2001-2017. Data were compared to previous published series from the unit from 1988-2001, and as continuous data, to assess changes in aetiologies and outcomes over the past 30 years.

### Research results

Paracetamol toxicity accounted for half of all cases of ALF, with a rise in the incidence of this condition over the past 30 years. Despite this observation, rates of liver transplantation for this condition are low at 4%, with an excellent overall survival. Rates of emergency liver transplantation were highest in indeterminate hepatitis and non-paracetamol drug induced liver injury. Transplant-free survival improved in this cohort compared to the historical cohort, however there was no significant change in overall survival.

### Research conclusions

Paracetamol represents the major cause of ALF in South-Eastern Australia with a concerning rise in its incidence over the past 30 years. Transplant-free survival has improved but remains low for ALF due to non-paracetamol causes.

### Research perspectives

This study shows a concerning rise in the incidence of paracetamol induced ALF in Australia, raising important questions regarding awareness and public health strategies to curb this rise. Larger multi-centre studies are required to confirm this observation. Transplant-free survival

improved in this population similar to reports from other large international series, highlighting advances in supportive care.

## REFERENCES

- O'Grady JG, Alexander GJ, Hayllar KM, Williams R. Early indicators of prognosis in fulminant hepatic failure. *Gastroenterology* 1989; **97**: 439-445 [PMID: 2490426]
- Ichai P, Samuel D. Etiology and prognosis of fulminant hepatitis in adults. *Liver Transpl* 2008; **14** Suppl 2: S67-S79 [PMID: 18825677 DOI: 10.1002/lt.21612]
- Bernal W, Hyyrylainen A, Gera A, Audimoolam VK, McPhail MJ, Auzinger G, Rela M, Heaton N, O'Grady JG, Wendon J, Williams R. Lessons from look-back in acute liver failure? A single centre experience of 3300 patients. *J Hepatol* 2013; **59**: 74-80 [PMID: 23439263 DOI: 10.1016/j.jhep.2013.02.010]
- Ostapowicz G, Fontana RJ, Schiødt FV, Larson A, Davern TJ, Han SH, McCashland TM, Shakil AO, Hay JE, Hyman L, Crippin JS, Blei AT, Samuel G, Reisch J, Lee WM; U. S. Acute Liver Failure Study Group. Results of a prospective study of acute liver failure at 17 tertiary care centers in the United States. *Ann Intern Med* 2002; **137**: 947-954 [PMID: 12484709]
- Khuroo MS, Kamili S. Aetiology and prognostic factors in acute liver failure in India. *J Viral Hepat* 2003; **10**: 224-231 [PMID: 12753342]
- Schiødt FV, Atillasoy E, Shakil AO, Schiff ER, Caldwell C, Kowdley KV, Stribling R, Crippin JS, Flamm S, Somberg KA, Rosen H, McCashland TM, Hay JE, Lee WM. Etiology and outcome for 295 patients with acute liver failure in the United States. *Liver Transpl Surg* 1999; **5**: 29-34 [PMID: 9873089]
- Koch DG, Tillman H, Durkalski V, Lee WM, Reuben A. Development of a Model to Predict Transplant-free Survival of Patients With Acute Liver Failure. *Clin Gastroenterol Hepatol* 2016; **14**: 1199-1206.e2 [PMID: 27085756 DOI: 10.1016/j.cgh.2016.03.046]
- Reuben A, Tillman H, Fontana RJ, Davern T, McGuire B, Stravitz RT, Durkalski V, Larson AM, Liou I, Fix O, Schilsky M, McCashland T, Hay JE, Murray N, Shaikh OS, Ganger D, Zaman A, Han SB, Chung RT, Smith A, Brown R, Crippin J, Harrison ME, Koch D, Munoz S, Reddy KR, Rossaro L, Satyanarayana R, Hassanein T, Hanje AJ, Olson J, Subramanian R, Karvellas C, Hameed B, Sherker AH, Robuck P, Lee WM. Outcomes in Adults With Acute Liver Failure Between 1998 and 2013: An Observational Cohort Study. *Ann Intern Med* 2016; **164**: 724-732 [PMID: 27043883 DOI: 10.7326/m15-2211]
- Kremers WK, van IJperen M, Kim WR, Freeman RB, Harper AM, Kamath PS, Wiesner RH. MELD score as a predictor of pretransplant and posttransplant survival in OPTN/UNOS status 1 patients. *Hepatology* 2004; **39**: 764-769 [PMID: 14999695]
- O'Grady J. Timing and benefit of liver transplantation in acute liver failure. *J Hepatol* 2014; **60**: 663-670 [PMID: 24211740 DOI: 10.1016/j.jhep.2013.10.024]
- Australian Bureau of Statistics. 3101.0 - Australian Demographic Statistics, Dec 2017. [Cited 2018 Mar. 3] Available from: <http://www.abs.gov.au/AUSSTATS/abs@.nsf/DetailsPage/3101.0Dec%202017>
- Gow PJ, Jones RM, Dobson JL, Angus PW. Etiology and outcome of fulminant hepatic failure managed at an Australian liver transplant unit. *J Gastroenterol Hepatol* 2004; **19**: 154-159 [PMID: 14731124]
- Lee WM, Stravitz RT, Larson AM. Introduction to the revised American Association for the Study of Liver Diseases Position Paper on acute liver failure 2011. *Hepatology* 2012; **55**: 965-967 [PMID: 22213561 DOI: 10.1002/hep.25551]
- Sinclair M, Ket S, Testro A, Gow PJ, Angus PW. Acute hepatic decompensation precipitated by pregnancy-related catabolic stress: a rare mimic of acute liver failure. *Obstet Gynecol* 2014; **123**: 480-483 [PMID: 24413235]
- Donnelly MC, Davidson JS, Martin K, Baird A, Hayes PC, Simpson KJ. Acute liver failure in Scotland: changes in aetiology and outcomes over time (the Scottish Look-Back Study). *Aliment Pharmacol Ther* 2017; **45**: 833-843 [PMID: 28097670 DOI: 10.1111/apt.13943]
- Sood S, Howell J, Sundararajan V, Angus PW, Gow PJ. Paracetamol overdose in Victoria remains a significant health-care burden. *J Gastroenterol Hepatol* 2013; **28**: 1356-1360 [PMID: 23489151 DOI: 10.1111/jgh.12196]
- Hawton K, Simkin S, Deeks J, Cooper J, Johnston A, Waters K, Arundel M, Bernal W, Gunson B, Hudson M, Suri D, Simpson K. UK legislation on analgesic packs: before and after study of long term effect on poisonings. *BMJ* 2004; **329**: 1076 [PMID: 15516343 DOI: 10.1136/bmj.38253.572581.7C]
- Larson AM, Polson J, Fontana RJ, Davern TJ, Lalani E, Hyman LS, Reisch JS, Schiødt FV, Ostapowicz G, Shakil AO, Lee WM; Acute Liver Failure Study Group. Acetaminophen-induced acute liver failure: results of a United States multicenter, prospective study. *Hepatology* 2005; **42**: 1364-1372 [PMID: 16317692 DOI: 10.1002/hep.20948]
- Manka P, Verheyen J, Gerken G, Canbay A. Liver Failure due to Acute Viral Hepatitis (A-E). *Visc Med* 2016; **32**: 80-85 [PMID: 27413724 DOI: 10.1159/000444915]
- Daniels D, Grytdal S, Wasley A; Centers for Disease Control and Prevention (CDC). Surveillance for acute viral hepatitis - United States, 2007. *MMWR Surveill Summ* 2009; **58**: 1-27 [PMID: 19478727]
- The Kirby Institute. HIV, viral hepatitis and sexually transmissible infections in Australia: annual surveillance report 2017. Sydney: Kirby Institute, UNSW Sydney. Available from: <https://kirby.unsw.edu.au/report/annual-surveillance-report-hiv-viral-hepatitis-and-stis-australia-2017>
- European Association For The Study Of The Liver. EASL clinical practice guidelines: Management of chronic hepatitis B virus infection. *J Hepatol* 2012; **57**: 167-185 [PMID: 22436845 DOI: 10.1016/j.jhep.2012.02.010]
- Wasley A, Fiore A, Bell BP. Hepatitis A in the era of vaccination. *Epidemiol Rev* 2006; **28**: 101-111 [PMID: 16775039 DOI: 10.1093/epirev/mxj012]
- Donnan EJ, Fielding JE, Gregory JE, Lalor K, Rowe S, Goldsmith P, Antoniou M, Fullerton KE, Knope K, Copland JG, Bowden DS, Tracy SL, Hogg GG, Tan A, Adamopoulos J, Gaston J, Vally H. A multistate outbreak of hepatitis A associated with semidried tomatoes in Australia, 2009. *Clin Infect Dis* 2012; **54**: 775-781 [PMID: 22238166 DOI: 10.1093/cid/cir949]
- Gow PJ, Angus PW, Smallwood RA. Transplantation in patients with paracetamol-induced fulminant hepatic failure. *Lancet* 1997; **349**: 651-652 [PMID: 9057756 DOI: 10.1016/S0140-6736(05)61600-5]
- Craig DG, Ford AC, Hayes PC, Simpson KJ. Systematic review: prognostic tests of paracetamol-induced acute liver failure. *Aliment Pharmacol Ther* 2010; **31**: 1064-1076 [PMID: 20180786 DOI: 10.1111/jgh.12196]

- 10.1111/j.1365-2036.2010.04279.x]
- 27 **Bailey B**, Amre DK, Gaudreault P. Fulminant hepatic failure secondary to acetaminophen poisoning: a systematic review and meta-analysis of prognostic criteria determining the need for liver transplantation. *Crit Care Med* 2003; **31**: 299-305 [PMID: 12545033 DOI: 10.1097/01.ccm.0000034674.51554.4c]
- 28 **O'Grady JG**. Transplant in haste, repent at your leisure? *Liver Transpl* 2015; **21**: 570-571 [PMID: 25824853 DOI: 10.1002/lt.24124]
- 29 **Warrillow SJ**, Bellomo R. Preventing cerebral oedema in acute liver failure: the case for quadruple-H therapy. *Anaesth Intensive Care* 2014; **42**: 78-88 [PMID: 24471667 DOI: 10.1177/0310057X1404200114]
- 30 The Transplantation Society of Australia and New Zealand: Clinical Guidelines for Organ Transplantation from Deceased Donors May 2017. Available from: <https://www.tsanz.com.au/organallocationguidelines/documents/ClinicalGuidelinesV1.1May2017.pdf>



Published By Baishideng Publishing Group Inc  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA  
Telephone: +1-925-2238242  
Fax: +1-925-2238243  
E-mail: [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
Help Desk: <https://www.f6publishing.com/helpdesk>  
<https://www.wjgnet.com>

