

# World Journal of *Hepatology*

*World J Hepatol* 2021 December 27; 13(12): 1816-2200



### OPINION REVIEW

- 1816** Non-alcoholic fatty liver disease in irritable bowel syndrome: More than a coincidence?  
*Purssell H, Whorwell PJ, Athwal VS, Vasant DH*

### REVIEW

- 1828** Liver-side of inflammatory bowel diseases: Hepatobiliary and drug-induced disorders  
*Mazza S, Soro S, Verga MC, Elvo B, Ferretti F, Cereatti F, Drago A, Grassia R*
- 1850** Gastrointestinal and hepatic side effects of potential treatment for COVID-19 and vaccination in patients with chronic liver diseases  
*Law MF, Ho R, Law KWT, Cheung CKM*
- 1875** Genotype E: The neglected genotype of hepatitis B virus  
*Ingasia LAO, Wose Kinge C, Kramvis A*

### MINIREVIEWS

- 1892** One stop shop approach for the diagnosis of liver hemangioma  
*Sandulescu LD, Urhut CM, Sandulescu SM, Ciurea AM, Cazacu SM, Iordache S*
- 1909** Liver function in COVID-19 infection  
*Przekop D, Gruszevska E, Chrostek L*
- 1919** Potential role of noninvasive biomarkers during liver fibrosis  
*Kaur N, Goyal G, Garg R, Tapasvi C, Chawla S, Kaur R*
- 1936** Imaging evaluation of the liver in oncology patients: A comparison of techniques  
*Freitas PS, Janicas C, Veiga J, Matos AP, Herédia V, Ramalho M*
- 1956** Liver manifestations and complications in inflammatory bowel disease: A review  
*Gaspar R, Branco CC, Macedo G*
- 1968** Dengue hemorrhagic fever and the liver  
*Leowattana W, Leowattana T*
- 1977** Artificial Intelligence in hepatology, liver surgery and transplantation: Emerging applications and frontiers of research  
*Veerankutty FH, Jayan G, Yadav MK, Manoj KS, Yadav A, Nair SRS, Shabeerali TU, Yeldho V, Sasidharan M, Rather SA*
- 1991** De novo and recurrence of metabolic dysfunction-associated fatty liver disease after liver transplantation  
*Han MAT, Olivo R, Choi CJ, Pyrsopoulos N*

- 2005** Liver dysfunction as a cytokine storm manifestation and prognostic factor for severe COVID-19

*Taneva G, Dimitrov D, Velikova T*

- 2013** COVID-19 and the liver: A brief and core review

*Kayaaslan B, Guner R*

- 2024** Newer variants of progressive familial intrahepatic cholestasis

*Vinayagamoorthy V, Srivastava A, Sarma MS*

- 2039** Deep learning in hepatocellular carcinoma: Current status and future perspectives

*Ahn JC, Qureshi TA, Singal AG, Li D, Yang JD*

## ORIGINAL ARTICLE

### Basic Study

- 2052** Gut dysbiosis and systemic inflammation promote cardiomyocyte abnormalities in an experimental model of steatohepatitis

*Longo L, Rampelotto PH, Filippi-Chiela E, de Souza VEG, Salvati F, Cerski CT, da Silveira TR, Oliveira CP, Uribe-Cruz C, Álvares-da-Silva MR*

### Case Control Study

- 2071** Leukocyte cell-derived chemotaxin-2 and fibroblast growth factor 21 in alcohol-induced liver cirrhosis

*Sak JJ, Prystupa A, Kiciński P, Luchowska-Kocot D, Kurys-Denis E, Bis-Wencel H*

### Retrospective Study

- 2081** Biliary complications in recipients of living donor liver transplantation: A single-centre study

*Guirguis RN, Nashaat EH, Yassin AE, Ibrahim WA, Saleh SA, Bahaa M, El-Meteini M, Fathy M, Dabbous HM, Montasser IF, Salah M, Mohamed GA*

- 2104** Liver function tests and metabolic-associated fatty liver disease: Changes in upper normal limits, does it really matter?

*Forlano R, Mullish BH, Dhar A, Goldin RD, Thursz M, Manousou P*

- 2113** Use of oral vancomycin in children with autoimmune liver disease: A single centre experience

*Di Giorgio A, Tulone A, Nicastro E, Norsa L, Sonzogni A, D'Antiga L*

- 2128** Trends of alcoholic liver cirrhosis readmissions from 2010 to 2018: Rates and healthcare burden associated with readmissions

*Kichloo A, El-Amir Z, Dahiya DS, Wani F, Singh J, Solanki D, Edigin E, Eseaton P, Mehboob A, Shaka H*

### Observational Study

- 2137** New stem cell autophagy surrogate diagnostic biomarkers in early-stage hepatocellular carcinoma in Egypt: A pilot study

*Yosef T, Ibrahim WA, Matboli M, Swilam AA, El-Nakeep S*

- 2150** Determination of "indeterminate score" measurements in lean nonalcoholic fatty liver disease patients from western Saudi Arabia

*Khayyat YM*

- 2161** Managing liver transplantation during the COVID-19 pandemic: A survey among transplant centers in the Southeast United States

*Gonzalez AJ, Kapila N, Thomas E, Pinna A, Tzakis A, Zervos XB*

### Prospective Study

- 2168** Accuracy of virtual chromoendoscopy in differentiating gastric antral vascular ectasia from portal hypertensive gastropathy: A proof of concept study

*Al-Taei AM, Cubillan MP, Hinton A, Sobotka LA, Befeler AS, Hachem CY, Hussan H*

- 2179** Non-alcoholic steatohepatitis in liver transplant recipients diagnosed by serum cytokeratin 18 and transient elastography: A prospective study

*Alhinai A, Qayyum-Khan A, Zhang X, Samaha P, Metrakos P, Deschenes M, Wong P, Ghali P, Chen TY, Sebastiani G*

### CASE REPORT

- 2192** Rare primary mature teratoma of the liver: A case report

*Kovalenko YA, Zharikov YO, Kiseleva YV, Goncharov AB, Shevchenko TV, Gurmikov BN, Kalinin DV, Zhao AV*

**ABOUT COVER**

Editorial Board Member of *World Journal of Hepatology*, Manuel Luis Rodríguez-Perálvarez, MD, PhD, Consultant Hepatologist and Associate Professor of Medicine, Department of Hepatology and Liver Transplantation, Reina Sofia University Hospital, Córdoba 14014, Spain. ropeml@hotmail.com

**AIMS AND SCOPE**

The primary aim of *World Journal of Hepatology* (*WJH*, *World J Hepatol*) is to provide scholars and readers from various fields of hepatology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

*WJH* mainly publishes articles reporting research results and findings obtained in the field of hepatology and covering a wide range of topics including chronic cholestatic liver diseases, cirrhosis and its complications, clinical alcoholic liver disease, drug induced liver disease autoimmune, fatty liver disease, genetic and pediatric liver diseases, hepatocellular carcinoma, hepatic stellate cells and fibrosis, liver immunology, liver regeneration, hepatic surgery, liver transplantation, biliary tract pathophysiology, non-invasive markers of liver fibrosis, viral hepatitis.

**INDEXING/ABSTRACTING**

The *WJH* is now abstracted and indexed in PubMed, PubMed Central, Emerging Sources Citation Index (Web of Science), Scopus, China National Knowledge Infrastructure (CNKI), China Science and Technology Journal Database (CSTJ), and Superstar Journals Database. The 2021 edition of Journal Citation Reports® cites the 2020 Journal Citation Indicator (JCI) for *WJH* as 0.61. The *WJH*'s CiteScore for 2020 is 5.6 and Scopus CiteScore rank 2020: Hepatology is 24/62.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: *Xu Guo*; Production Department Director: *Xiang Li*; Editorial Office Director: *Xiang Li*.

**NAME OF JOURNAL**

*World Journal of Hepatology*

**ISSN**

ISSN 1948-5182 (online)

**LAUNCH DATE**

October 31, 2009

**FREQUENCY**

Monthly

**EDITORS-IN-CHIEF**

Nikolaos Pyrsopoulos, Ke-Qin Hu, Koo Jeong Kang

**EDITORIAL BOARD MEMBERS**

<https://www.wjnet.com/1948-5182/editorialboard.htm>

**PUBLICATION DATE**

December 27, 2021

**COPYRIGHT**

© 2021 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>



## Deep learning in hepatocellular carcinoma: Current status and future perspectives

Joseph C Ahn, Touseef Ahmad Qureshi, Amit G Singal, Debiao Li, Ju-Dong Yang

**ORCID number:** Joseph C Ahn 0000-0001-6994-2870; Touseef Ahmad Qureshi 0000-0002-6683-4556; Amit G Singal 0000-0002-1172-3971; Debiao Li 0000-0001-8560-8231; Ju-Dong Yang 0000-0001-7834-9825.

**Author contributions:** Yang JD devised the project and the main conceptual ideas for the review; Ahn JC conducted the literature search and identified relevant studies to be included in the review; Qureshi TA and Li D provided the technical expertise on artificial intelligence and deep learning; Ahn JC drafted the manuscript; Yang JD and Singal AG revised the manuscript critically for important intellectual content; and all authors approved the final version to be published.

**Conflict-of-interest statement:** Dr. Yang provides a consulting service for Exact Sciences and Gilead; Dr. Singal has been on advisory boards and served as a consultant for Genentech, Bayer, Eisai, BMS, Exelixis, AstraZeneca, and TARGET RWE. No other potential conflicts of interest relevant to this article exist.

**Country/Territory of origin:** United States

**Specialty type:** Gastroenterology and hepatology

**Joseph C Ahn**, Gastroenterology and Hepatology, Mayo Clinic, Rochester, MN 55904, United States

**Touseef Ahmad Qureshi, Debiao Li**, Biomedical Imaging Research Institute, Cedars-Sinai Medical Center, Los Angeles, CA 90048, United States

**Amit G Singal**, Internal Medicine, University of Texas Southwestern Medical Center, Dallas, TX 75390, United States

**Ju-Dong Yang**, Karsh Division of Gastroenterology and Hepatology, Cedars-Sinai Medical Center, Los Angeles, CA 90048, United States

**Corresponding author:** Ju-Dong Yang, MD, MS, Assistant Professor, Karsh Division of Gastroenterology and Hepatology, Cedars-Sinai Medical Center, 8900 Beverly Blvd, Los Angeles, CA 90048, United States. [judong.yang@cshs.org](mailto:judong.yang@cshs.org)

### Abstract

Hepatocellular carcinoma (HCC) is among the leading causes of cancer incidence and death. Despite decades of research and development of new treatment options, the overall outcomes of patients with HCC continue to remain poor. There are areas of unmet need in risk prediction, early diagnosis, accurate prognostication, and individualized treatments for patients with HCC. Recent years have seen an explosive growth in the application of artificial intelligence (AI) technology in medical research, with the field of HCC being no exception. Among the various AI-based machine learning algorithms, deep learning algorithms are considered state-of-the-art techniques for handling and processing complex multimodal data ranging from routine clinical variables to high-resolution medical images. This article will provide a comprehensive review of the recently published studies that have applied deep learning for risk prediction, diagnosis, prognostication, and treatment planning for patients with HCC.

**Key Words:** Hepatocellular carcinoma; Artificial intelligence; Deep learning

©The Author(s) 2021. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** There are emerging roles for deep learning technology in the field of hepato-

**Provenance and peer review:**

Invited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): A, A

Grade B (Very good): 0

Grade C (Good): 0

Grade D (Fair): D

Grade E (Poor): 0

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

**Received:** May 28, 2021

**Peer-review started:** May 28, 2021

**First decision:** July 6, 2021

**Revised:** July 19, 2021

**Accepted:** November 15, 2021

**Article in press:** November 15, 2021

**Published online:** December 27, 2021

**P-Reviewer:** Gumbs A, Shafqat S

**S-Editor:** Wang JJ

**L-Editor:** Webster JR

**P-Editor:** Wang JJ



cellular carcinoma (HCC) including HCC risk prediction, as well as diagnosis, prognostication, and treatment planning leveraging readily available data from radiologic and histopathologic medical images. This article will provide a comprehensive review of the recently published studies that have applied deep learning for risk prediction, diagnosis, prognostication, and treatment planning for patients with HCC.

**Citation:** Ahn JC, Qureshi TA, Singal AG, Li D, Yang JD. Deep learning in hepatocellular carcinoma: Current status and future perspectives. *World J Hepatol* 2021; 13(12): 2039-2051

**URL:** <https://www.wjgnet.com/1948-5182/full/v13/i12/2039.htm>

**DOI:** <https://dx.doi.org/10.4254/wjh.v13.i12.2039>

## INTRODUCTION


Hepatocellular carcinoma (HCC) is an aggressive primary liver cancer that develops in the setting of chronic parenchymal liver diseases, and is among the top causes of cancer incidence and mortality worldwide[1,2]. While the burden of HCC has been declining with effective antiviral therapy against hepatitis B virus (HBV) and hepatitis C virus (HCV), HCC incidence related to metabolic syndrome will likely continue to rise due to the dramatic increase in the prevalence of non-alcoholic fatty liver disease (NAFLD) in the general population[3]. Decades of HCC research led to the development of a screening protocol, non-invasive diagnostic modalities based on imaging, and various treatment modalities including surgical, locoregional and systemic therapies[4,5]. However, the overall outcomes of patients with HCC continue to remain poor and there are areas of significant unmet need in risk prediction, early detection, accurate prognostication, and individualized treatments for patients with HCC.

Patients with HCC generate enormous amounts of health data. While promising for researchers, ensuring that such high volumes of data are turned into actionable knowledge can be a significant challenge. Artificial intelligence (AI) is thought to be capable of synthesizing and analyzing multimodal data with superhuman degrees of accuracy or reliability, and recent years have seen a rapid growth in the application of AI to many fields of medicine including hepatology[6]. This "AI revolution" over the past decade has been possible due to the advent of deep learning technology. Deep learning algorithms can process a broad spectrum of medical data from structured numeric data such as vital signs and laboratory values, high dimensional data from multi-omics studies, as well as digitized high-resolution images from various radiologic and histopathologic studies. This review aims to provide an overview as well as highlight examples of the many potential applications of deep learning to improve the care of patients with HCC.

## AI, MACHINE LEARNING, AND DEEP LEARNING

AI-based approaches provide a variety of methods for a range of tasks and clinical application including image classification, organ and lesion segmentation, accurate extraction of key imaging features and measurements, tumor detection, stratification of high-risk subjects, prediction of disease and treatment outcome (Figure 1). Advancements in AI in recent years, particularly in the realm of medical image processing and analysis, offer an enormous range of automated tools for extracting precise measurements of biomarkers, revealing complex features, quantifying tissue characteristics and performing radiomics for deep analysis of raw imaging data.

The term "artificial intelligence" encompasses a broad range of technology that enables machines to perform tasks typically thought to require human reasoning and problem-solving skills[7]. "Machine learning" is a branch of AI in which computer algorithms train on sample data to build a mathematical model that makes predictions or decisions without being explicitly programmed to do so[8]. Machine learning algorithms can be broadly divided into supervised and unsupervised learning. Supervised learning algorithms train on sample data with labeled outcome data, and their goal is to learn the relationship between the input data and the outcomes to make



**Figure 1** Schematic representation of the relationships between the terms artificial intelligence, machine learning, and deep learning, and how deep learning can utilize multimodal data to improve care for patients with hepatocellular carcinoma.

accurate predictions about the outcome when provided with a new set of input data [9]. Examples of supervised learning algorithms include traditional techniques such as linear regression and logistic regression, as well as more sophisticated techniques including support vector machines, random forest and gradient boosting. On the other hand, unsupervised learning algorithms train on unlabeled sample data and analyze the underlying structure or distribution within the data to discover new clusters or patterns [10]. Examples of unsupervised learning algorithms include K-means and principle component analysis among many others.

Among the various AI-based machine learning algorithms, artificial neural networks (ANNs) consist of layers of interconnected mathematical formulas that enable them to analyze complex non-linear relationships [11]. “Deep learning (DL)” refers to highly complex AI models utilizing multiple layers of ANNs and has recently emerged as a state-of-the-art AI technique for analyzing complex, high-dimensional healthcare data. Some of the commonly used DL techniques include convolutional neural networks (CNNs) and recurrent neural networks (RNNs) [12]. CNNs have connective patterns resembling those of an animal visual cortex and can detect inherent spatial features of high dimensional images. RNNs have connections forming a directed graph along a temporal sequence, and therefore can be highly useful in time series prediction.

It is crucial to recognize that any AI-based machine learning algorithms require external validation in an independent dataset as models could be overfitted and end up overestimating the performance. In this review article, the performance characteristics of the various DL models are from the validation cohorts, and not the original derivation cohorts used to train the algorithms.

## HCC CLINICAL DATA

Despite multiple available risk prediction tools for HCC, none have been rigorously validated or endorsed by major liver societies. Currently, HCC surveillance is recommended for patients with cirrhosis and high risk patients with chronic HBV infection [13]. Accurate prediction models utilizing more specific risk factors for HCC development at individual levels would allow health systems to implement targeted screening strategies. Ioannou *et al* [14] trained a RNN to predict HCC development within 3 years using 4 baseline variables and 27 longitudinal variables from 48151 patients with HCV-related cirrhosis in the national Veterans Health Administration. The RNN model significantly outperformed logistic regression and exhibited an area under the curve (AUC) of 0.759 among all samples and an AUC of 0.806 among patients with sustained virologic response. Phan *et al* [15] surveyed 1 million random samples from Taiwan’s National Health Insurance Research Database between 2002 to 2010 to predict liver cancer among patients with viral hepatitis. The disease history of



each patient was transformed into a  $108 \times 998$  matrix and applied to a CNN, which predicted liver cancer with an AUC of 0.886 and an accuracy of 0.980. Another study by Nam *et al*[16] constructed a deep neural network to predict 3-year and 5-year incidence of HCC in 424 patients with HBV-related cirrhosis on entecavir therapy. When applied to an external validation cohort of 316 patients, the DL model achieved a Harrell's C-index of 0.782 and significantly outperformed 6 previously reported models based on traditional modeling. The same group also developed another DL model called the AI-based Model of Recurrence after Liver Transplantation (MoRAL-AI) to predict HCC recurrence after liver transplantation using variables such as tumor diameter, age, alpha-fetoprotein (AFP), and prothrombin time[17]. The MoRAL-AI showed significantly better predictive performance compared to conventional models such as the Milan, UCSF, up-to-seven, and Kyoto criteria (C-index = 0.75 *vs* 0.64, 0.62, 0.50, 0.50, respectively;  $P < 0.001$ ).

## HCC MULTI-OMICS

Serum AFP has been widely used as a predictive and prognostic biomarker for HCC [18], but AFP has limited sensitivity for detecting early-stage HCC and its levels do not reliably correlate with disease progression[19]. Recent advances in multi-omics related to HCC are expected to address this unmet need for novel biomarkers. Multi-omics refers to an approach to biological analysis which utilizes data sets from multiple "omics", such as the genome, epigenome, transcriptome, proteome, metabolome and microbiome. Multi-omics experiments generate an enormous amount of information, and various machine learning techniques including DL that can help with the computational challenges of processing and analyzing such high dimensional data. Xie *et al*[20] used gene expression profiling of peripheral blood to build an ANN model that classifies HCC patients from a control group. Using a nine-gene expression system, the ANN was able to distinguish HCC patients from controls with an AUC of 0.943, 98% sensitivity, and 85% specificity, although it should be noted that the control group was healthy individuals rather than patients with cirrhosis, which could have overestimated the performance of the model. Choi *et al*[21] proposed a novel network-based DL method to identify prognostic gene signatures *via* G2Vec, a modified Word2Vec model originally used for natural language processing (NLP). When applied to gene expression data for HCC from the Cancer Genome Atlas (TCGA), G2Vec showed superior prediction accuracy for patient outcomes compared to existing gene selection methods and was able to identify two distinct gene modules significantly associated with HCC prognosis. Chaudhary *et al*[22] used RNA sequencing, miRNA, and methylation data of 360 HCC patients from TCGA to build an autoencoder, which is an unsupervised feed-forward neural network. Using this DL model, they were able to distinguish patients with survival differences and identify specific mutations and pathways as predictors of aggressive tumor behavior.

## RADIOLOGY

### HCC diagnosis and segmentation

In recent years, there have been remarkable advances in the application of AI for the interpretation of medical imaging, primarily due to the use of DL algorithms using CNN[23]. CNN algorithms trained on ultrasound, computed tomography (CT), or magnetic resonance imaging (MRI) images have shown excellent performances in detection of lesions, classification of lesions, segmentation of organs or anatomic structures, and imaging reconstruction[24].

In 2012, Streba *et al*[25] prospectively studied contrast-enhanced ultrasound images of 112 patients to train an ANN that classified five different types of liver tumors. The ANN showed promising performances with accuracies of 94.5% in the training set and 87.1% in the testing set. In 2017, Hassan *et al*[26] reported using the stacked sparse auto-encoder, an unsupervised DL technique, to segment and classify liver lesions on ultrasound images with a classification accuracy of 97.2%. Additionally, Bharti *et al*[27] built a CNN using echotexture and roughness of liver surface on 754 segmented ultrasound images, which differentiated between normal liver, chronic liver disease, cirrhosis, and HCC with a classification accuracy of 96.6%. Schmauch *et al*[28] also created a CNN which detects and characterizes benign and malignant focal liver lesions on 2-D ultrasound images from 367 patients from various institutions. When

applied to a new dataset of 177 patients, the model achieved a weighted mean AUC of 0.891. Recently, Brehar *et al*[29] conducted a study comparing CNN's performance for HCC detection on ultrasound images against conventional machine learning algorithms including multi-layer perceptron, support vector machines, random forest and AdaBoost. The CNN achieved an AUC of 0.95% with 91.0% accuracy, 94.4% sensitivity, and 88.4% specificity and significantly outperformed the conventional machine learning algorithms. Beyond detecting the actual presence of HCC on ultrasound images, studies have also attempted to predict the risk of future HCC development based on analyzing the ultrasound images of liver parenchyma in patients without HCC. For example, Jin *et al*[30] performed a DL radiomics analysis on 2-D shear wave elastography and corresponding B-mode ultrasound images of 434 chronic HBV patients, which predicted 5-year HCC development with AUC of 0.900 in the test cohort.

In addition to ultrasound images, cross-sectional imaging from CT or MRI studies serve as an extremely abundant and promising source of data for DL. In 2018, Yasaka *et al*[31] used CT image sets of liver masses from 460 patients to train a CNN that can classify liver lesions into five categories of: (1) HCC; (2) Other malignant tumors; (3) Indeterminate masses; (4) Hemangiomas; and (5) Cysts with a median AUC of 0.92. Shi *et al*[32] showed that incorporation of a CNN enabled identification of HCC using a three-phase CT imaging protocol with a diagnostic accuracy similar to that of a four-phase protocol, which would allow patients to receive lower doses of radiation. Segmentation of HCC, liver parenchyma, and other organs on CT scan is very important for determination of tumor extent and treatment planning, but manual contouring of the images is highly time-consuming and subject to inter-observer variability. The 2017 International Conference On Medical Image Computing Computer Assisted Intervention called for a Liver Tumor Segmentation Benchmark (LITS) challenge, encouraging researchers to develop automatic segmentation algorithms to segment liver lesions using 200 CT scans (training: 130; testing: 70) provided by clinical sites around the world. Several teams participating in the challenge have developed DL algorithms with promising performances for HCC segmentation using CT images[33-37]. Beyond the LITS challenge, there are ongoing research efforts to improve segmentation using different architectures of DL networks [38-42].

Hamm *et al*[43] used MRI images from 494 patients to train a CNN which can classify hepatic lesions into six different categories. When applied to random cases in the test set, the CNN outperformed expert radiologists (90% sensitivity and 98% specificity *vs* 82.5% sensitivity and 96.5% specificity) and especially for HCC detection (90% sensitivity *vs* 60%-70% sensitivity). The same group conducted additional studies to make their CNN interpretable by generating highlighted feature maps corresponding to liver lesions[44]. Wu *et al*[45] built a CNN using multiphase MRI images and achieved an AUC of 0.95 for distinguishing Liver Imaging Reporting and Data System (LI-RADS) grade 3 from LI-RADS 4 and 5 lesions for HCC diagnosis. Zhen *et al* [46] also trained a CNN model combining unenhanced MRI images and clinical variables from 1210 patients with liver tumors, which demonstrated diagnostic performances on par with three experienced radiologists using enhanced MRI images.

### **HCC prognostication, treatment planning, and response to treatment**

In addition to serving as accurate and efficient tools for diagnosis of HCC, DL models utilizing radiology data can also be used for prognostication, treatment planning, and assessing tumor response to therapy. Vascular invasion is a key prognostic element in patients with HCC. Recent studies developed CNN models with promising ability to detect microvascular invasion on MRI images of HCC patients undergoing surgical resection[47-49]. An *et al*[50] used an unsupervised CNN-based deformable image registration technique to assess the relationship between ablative margins and local tumor progression in 141 patients with single HCC who underwent microwave ablation, and demonstrated that patients with ablative margins < 5 mm were at significantly higher risk of local tumor progression. Liu *et al*[51] developed a DL radiomics model to predict responses to trans-arterial chemoembolization (TACE) using ultrasound images of 130 HCC patients, which accurately predicted TACE response with an AUC of 0.93. The same group also assessed their ultrasound-based DL radiomics model to predict 2-year progression-free survival among 419 HCC patients and facilitate optimized treatment selection. Peng *et al*[52] trained a residual CNN model to predict response to TACE using CT images from 562 patients with intermediate-stage HCC undergoing TACE, which showed accuracies of 85.1% and 82.8% in two external validation cohorts. Another study developed a DL score for disease-specific survival by using CT images in a cohort of 243 patients with HCC

treated with TACE, with a higher score predicting poor prognosis [hazard ratio (HR): 3.01; 95% cumulative incidence (CI): 2.02-4.50][53]. Finally, Zhang *et al*[54] built a DL-based model predicting overall survival using CT images from 201 patients with unresectable HCC treated with TACE and sorafenib, which achieved superior predictive performance compared to the clinical nomogram (C-index of 0.730 *vs* 0.679,  $P = 0.023$ ).

## HCC PATHOLOGY

Automated interpretation of histopathologic images from liver biopsy is another major area of medical imaging in patients with HCC where DL can be utilized. In addition to effectively replicating the human pathologists' jobs of diagnosing and grading HCC, DL models can help identify and analyze additional complex imaging features and patterns which are related to specific mutations and disease prognosis. Lin *et al*[55] used images from multiphoton microscopy of 113 HCC patients to train a CNN with over 90% accuracy for determining HCC differentiation. Kiani *et al*[56] developed a CNN-based "Liver Cancer Assistant" which accurately differentiated hematoxylin and eosin (H&E) images of HCC and cholangiocarcinoma and helped improve the diagnostic performance of nine pathologists. Liao *et al*[57] used TCGA dataset for training a CNN that distinguished HCC from adjacent normal tissues with perfect performance (AUC: 1.00) and predicted the presence of specific somatic mutations with AUCs over 0.70. Wang *et al*[58] trained a CNN for automated segmentation and classification of individual nuclei at single-cell levels on H&E-stained tissue sections of HCC tumors from TCGA, and performed feature extraction to identify 246 quantitative image features. Then, a clustering analysis by an unsupervised learning approach identified three distinct histologic subtypes which were independent of previously established genomic clusters and had different prognosis. Chen *et al*[59] trained a CNN for automatic grading of HCC tumors on histopathological H&E images, which showed 96% accuracy for benign and malignant classification and 89.6% accuracy for the degree of tumor differentiation, and predicted the presence of specific genetic mutations.

Lu *et al*[60] applied three pre-trained CNN models to extract imaging features from HCC histopathology and performed Cox proportional hazards analysis to predict overall survival and disease-free survival, and observed significant correlations between the imaging features and established biological pathways. Saillard *et al*[61] used two DL algorithms based on whole-slide digitized histological slides from 194 patients with HCC to predict the survival of patients treated by surgical resection. When tested on an independent validation set from TCGA, both DL models had a higher discriminatory power than a score combining all baseline variables associated with survival. Shi *et al*[62] built an interpretable DL framework using pathologic images from 1445 patients with HCC and developed a "tumor risk score" which showed prognostic performances independent of and superior to clinical staging systems and stratified patients into five groups of different prognosis. A recent study by Yamashita *et al*[63] developed a histopathology-based DL based system which stratified patients with risk scores for postsurgical recurrence of HCC.

## FUTURE DIRECTION

There are several key issues to address before DL-based AI models can be universally implemented in real world clinical practice settings. Due to their complexity, DL models are traditionally considered to be "black-box" models, meaning humans cannot understand how the DL models make their predictions. Interpretability of the DL models are crucial for physicians to accept and trust them in everyday clinical practice, and for troubleshooting and improving the models for rare cases. This is being addressed by recent developments in various "explainable AI" techniques but currently there is no clear consensus on the best methodology. Another potential limitation is the generalizability of the individual DL algorithms. Concerns have been raised that AI algorithms developed at highly specialized academic medical centers using their own patients' data may over-represent certain groups of patients and not accurately reflect the real-world population of patients seen at local community hospitals. Finally, AI models, like other prediction models, are often not publicly available, limiting external validation. Independent validation of the proposed model and comparison to old models are as important as deriving new models. Large-scale,

**Table 1 Studies applying deep learning for hepatocellular carcinoma**

Study	Cohort	Data source	Deep learning	Input	Output	Main findings
Predicting HCC risk using clinical variables						
Ioannou <i>et al</i> [14] 2020	48151 HCV cirrhosis (T: 90%, V: 10%)	VHA database	RNN	Clinical variables	Risk of HCC development	RNN predicted HCC development with AUC of 0.759, and AUC of 0.806 among those who achieved SVR
Phan <i>et al</i> [15] 2020	6052 HBV and HCV (T: 70%, V: 30%)	Taiwanese NHIRD	CNN	Disease history data	Risk of HCC development	CNN achieved an accuracy of 0.980 and AUC of 0.886 for predicting HCC development among viral hepatitis patients
Nam <i>et al</i> [16] 2020	T: 424 HBV cirrhosis; V: 316 HBV cirrhosis	2 Korean centers	ResNet	Clinical variables	Risk of HCC development	DL model achieved an accuracy of 0.763 and AUC of 0.782 in the validation cohort and outperformed previous models
Nam <i>et al</i> [17] 2020	T: 349 LT recipients; V: 214 LT recipients	3 Korean LT centers	ResNet	Clinical variables	Recurrent HCC after LT	DL model significantly outperformed conventional models in prediction of post-T HCC recurrence with AUC of 0.75
Multi-omics-based HCC diagnosis and prognostication						
Xie <i>et al</i> [20] 2018	T: 133 HCC/54 HV; V: 52 HCC/34 HV	1 center in China	ANN	Gene expression	HCC detection	ANN using nine genes had an AUC of 0.943, 98% sensitivity, and 85% specificity for classifying HCC
Choi <i>et al</i> [21] 2018	135 HCC (10-fold CV)	TCGA	G2Vec	Gene expression	HCC prognosis	G2Vec showed significantly higher prediction accuracy for patient outcomes compared to existing gene selection tools
Chaudhary <i>et al</i> [22] 2018	T: 360 HCC; V: 220, 221, 166, 40, 27 HCC	TCGA; 5 external datasets	Auto-encoder	RNA-seq, miRNA-seq, methylation	HCC prognosis	DL model distinguished groups with survival differences and identified mutations and pathways predicting aggressive tumor behavior
Radiology-based HCC diagnosis/prediction						
Streba <i>et al</i> [25] 2012	112 FLL (10-fold CV)	1 center in Romania	ANN	US images	FLL type	ANN had 87.12% testing accuracy, 93.2% sensitivity, and 89.7% specificity for classifying 5 classes of liver lesions
Hassan <i>et al</i> [26] 2017	110 FLL (10-fold CV)	1 center in Egypt	Auto-encoder	US images	FLL type	The proposed system had 97.2% accuracy, 98% sensitivity, and 95.70% specificity for classifying liver lesions
Bharti <i>et al</i> [27] 2018	24 normal, 25 CLD, 25 cirrhosis, 20 HCC	1 center in India	CNN	US images	Liver stages	CNN achieved 96.6% classification accuracy for differentiating normal liver, CLD, cirrhosis, and HCC
Schmauch <i>et al</i> [28] 2019	T: 367 FLL; V: 177 FLL	Centers in France	ResNet	US images	FLL type	DL model reached mean AUC of 0.935 for focal liver lesion detection and 0.916 for focal liver lesion characterization
Breher <i>et al</i> [29] 2020	T: 200 HCC; V: 68 HCC	1 center in Romania	CNN	US images	HCC detection	CNN achieved AUC of 0.95, accuracy of 0.91, 94.4% sensitivity and 88.4% specificity for HCC detection
Jin <i>et al</i> [30] 2021	434 HBV (3:1:1 split)	1 center in China	DL radiomics	US images	Risk of HCC development	DL radiomics model predicted 5-yr HCC development risk with AUC of 0.900 in the test set
Yasaka <i>et al</i> [31] 2018	T: 460 liver masses; V: 100 liver masses	1 center in Japan	CNN	CT images	Liver mass type	CNN classified liver lesions into five categories with a median AUC of 0.92
Shi <i>et al</i> [32] 2020	449 FLL; (T: 80%, V: 20%)	1 center in China	CNN	CT images	FLL type	CNN applied to three-phase CT protocol images achieved AUC of 0.925 for differentiating HCC

						from other FLLs
Hamm <i>et al</i> [43] 2019	T: 434 FLL; V: 60 FLL	1 center in United States	CNN	MRI images	FLL type	CNN achieved 90% sensitivity and 98% specificity for classifying FLLs and AUC of 0.992 for HCC classification
Wang <i>et al</i> [44] 2019	T: 434 FLL; V: 60 FLL	1 center in United States	CNN	MRI images	FLL type	Interpretable DL system achieved 76.5% PPV and 82.9% sensitivity for identifying correct radiological features
Wu <i>et al</i> [45] 2020	89 liver tumors; (60: 20: 20)	1 center in United States	CNN	MRI images	LI-RADS grading	CNN achieved AUC of 0.95, 90% accuracy, 100% sensitivity and 83.5% PPV for LI-RADS grading of liver tumors
Zhen <i>et al</i> [46] 2020	T: 1210 liver tumors; V: 201 liver tumors	1 center in China	CNN	MRI images	Liver tumor type	CNN combined with clinical data showed AUC of 0.985 for classifying HCC with 91.9% agreement with pathology
Radiology-based HCC prognostication, treatment planning, and response to treatment						
Zhang <i>et al</i> [47] 2021	T: 158 HCC; V: 79 HCC	1 center in China	CNN	MRI images	MVI in HCC	CNN achieved AUC of 0.72, 55% sensitivity, and 81% specificity for preoperative MVI in HCC patients
Wang <i>et al</i> [48] 2020	T: 60 HCC; V: 40 HCC	1 center in China	CNN	MRI images	MVI in HCC	Fusion of deep features from MRI images yielded AUC of 0.79 for MVI prediction in HCC patients
Jiang <i>et al</i> [49] 2021	405 HCC; (T: 80%, V: 20%)	1 center in China	CNN	CT images	MVI in HCC	CNN achieved AUC of 0.906 for prediction of MVI. Mean survival was significantly better in the group without MVI
An <i>et al</i> [50] 2020	141 single HCC resect MWA	1 center in China	CNN	MRI images	Ablative margin	Deep learning model accurately estimated ablative margins and risk of local tumor progression
Liu <i>et al</i> [51] 2020	T: 89 HCC resect TACE; V: 41 HCC rec. TACE	1 center in China	CNN	Ultrasound images	Response to TACE	Deep learning radiomics model predicted tumor response to TACE with AUC of 0.93
Peng <i>et al</i> [52] 2020	T: 562 HCC resect TACE; V: 227 HCC rec. TACE	3 centers in China	CNN	CT images	Response to TACE	Deep learning model had accuracies of 85.1% and 82.8% for predicting TACE response in 2 validation cohorts
Liu <i>et al</i> [53] 2020	243 HCC resect TACE (6:1:3 split)	1 center in China	CNN	CT images	Post-TACE survival	Higher DL score was an independent prognostic factor and predicted overall survival with AUCs of 0.85-0.90
Zhang <i>et al</i> [54] 2020	201 HCC resect TACE + sorafenib (T: 120, V: 81)	3 centers in China	CNN	CT images	OS on TACE + sorafenib	Deep learning signature achieved C-index of 0.714 for predicting OS in HCC patients receiving TACE + sorafenib
Histopathology-based HCC diagnosis, subtyping, and outcome predictions						
Lin <i>et al</i> [55] 2019	113 HCC	1 center in China	CNN	Histopath images	HCC differentiation	CNN achieved an accuracy of 0.941 for determining HCC differentiation on multiphoton microscopy
Kiani <i>et al</i> [56] 2020	70 WSI (35 HCC, 35 CC)	TCGA	CNN	Histopath images	HCC <i>vs</i> CC	CNN-based "Liver Cancer Assistant" accurately differentiated HCC <i>vs</i> cholangiocarcinoma
Liao <i>et al</i> [57] 2020	T: 491 HCC; V: 455 HCC	TCGA; 1 center in China	CNN	Histopath images	HCC detection, mutations	CNN distinguished HCC from adjacent tissues with AUC of 1.00 and predicted specific mutations with AUC over 0.70
Wang <i>et al</i> [58] 2020	T: 99 HCC; V: 205 HCC	TCGA	CNN	Histopath images	Histological HCC subtype	Unsupervised clustering identified 3 histological subtypes complementing molecular pathways and



						prognostic value
Chen <i>et al</i> [59] 2020	T: 402 HCC/89 normal; V: 67 HCC/34 normal	GDC portal; 1 center in China	CNN	Histopath images	HCC grade mutations	CNN achieved 89.6% accuracy for tumor differentiation stage and predicted presence of specific gene mutations
Lu <i>et al</i> [60] 2020	421 HCC/105 normal (6-fold CV)	GDC portal	CNN	Histopath images	HCC prognosis	Pre-trained CNN predicted OS using pathology images and identified HCC subgroups with different prognosis
Saillard <i>et al</i> [61] 2020	T: 194 HCC; V: 328 HCC	1 French center TCGA	CNN	Histopath images	Survival after HCC resection	CNN models using pathology images predicted survival with C-index 0.75-0.78 and outperformed conventional models
Shi <i>et al</i> [62] 2021	T: 1125 HCC; V: 320 HCC	1 center in China; TCGA	CNN	Histopath images	HCC outcomes	Deep learning-based “tumor risk score” was superior to clinical staging and stratified 5 groups of different prognosis
Yamashita <i>et al</i> [63] 2021	T: 36 WSI; V: 30 WSI	1 center in United States; TCGA	CNN	Histopath images	Post-surgical recurrence	CNN risk scores outperformed TNM system for predicting recurrence and identified high- and low-risk subgroups

ANN: Artificial neural network; AUC: Area under the curve; CC: Cholangiocarcinoma; CNN: Convolutional neural network; CV: Cross-validation; FLL: Focal liver lesion; GDC: Genomic Data Commons; HBV: Hepatitis B virus; HCC: Hepatocellular carcinoma; HCV: Hepatitis C virus; HV: Healthy volunteers; LT: Liver transplant; MVI: Microvascular invasion; MWA: Microwave ablation; OS: Overall survival; PFS: Progression-free survival; RFA: Radio-frequency ablation; RNN: Recurrent neural network; SR: Surgical resection; STS-net: Spatial transformed similarity network; SVR: Sustained virologic response; T: Training; TCGA: The Cancer Genome Atlas; V: Validation; VHA: Veterans Health Administration; WSI: Whole slide image; CT: Computed tomography; MRI: Magnetic resonance imaging; NHIRD: National Health Insurance Research Database; TNM: Tumor, Nodes, Metastasis; TACE: Trans-arterial chemoembolization; LI-RADS: Liver Imaging Reporting and Data System.

prospective, multi-centered studies involving diverse populations with external validation will be necessary before DL algorithms can be widely accepted.

A currently under-explored, but highly promising and exciting area for the application of DL is the field of autonomous robotics. In a recent editorial, Gumbs *et al* [64] state that while the current form of robotic surgery seems like a form of minimally invasive surgery, the true power of robotic surgery exists in its potential to create autonomous actions. Recently, a DL-based surgical instrument tracking algorithm was able to closely track the instruments during robotic surgery and evaluate the surgeons' performance, demonstrating that DL algorithms can learn the correct steps of robotic surgery [65]. With the help of DL and other AI technologies, it may be possible to imagine a future where fully autonomous robots perform resection of large, complex HCC in ways that no human surgeons can mimic. However, there are significant barriers before the idea of fully autonomous robotic surgery can become a reality, including the current technical limitations of autonomous surgical robotics, as well as the hesitation of patients and providers to fully trust autonomous robots to perform invasive operations. “Explainability” of the DL algorithms will be critical here, as humans would need to be able to understand and correct every single mistake that an autonomous robot makes during surgery. Therefore, for the foreseeable future, DL will most likely remain as a helpful, adjunctive tool to assist human surgeons.

## CONCLUSION

This review has provided a comprehensive overview of various ways in which DL algorithms can be employed to assist medical providers and enhance the care of patients with HCC (Table 1). DL algorithms not only can efficiently and accurately replicate the same jobs performed by human physicians, but more importantly can help discover novel biologic pathways and disease subgroups with clinical significance by processing and analyzing complex high-dimensional data in ways impossible for the human brain.

Despite some important limitations to overcome, application of state-of-the-art AI technologies such as DL for the care of patients with HCC is no longer a futuristic idea but is rapidly becoming a reality. Most of the studies covered in this review were published within the past two years, and the number of studies utilizing DL continues

to increase exponentially. We anticipate that DL algorithms will soon take a major role in the diagnosis, prognostication, and treatment of patients with HCC.

## REFERENCES

- 1 **Bray F**, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018; **68**: 394-424 [PMID: 30207593 DOI: 10.3322/caac.21492]
- 2 **Yang JD**, Hainaut P, Gores GJ, Amadou A, Plymoth A, Roberts LR. A global view of hepatocellular carcinoma: trends, risk, prevention and management. *Nat Rev Gastroenterol Hepatol* 2019; **16**: 589-604 [PMID: 31439937 DOI: 10.1038/s41575-019-0186-y]
- 3 **Stepanova M**, De Avila L, Afendy M, Younossi I, Pham H, Cable R, Younossi ZM. Direct and Indirect Economic Burden of Chronic Liver Disease in the United States. *Clin Gastroenterol Hepatol* 2017; **15**: 759-766.e5 [PMID: 27464590 DOI: 10.1016/j.cgh.2016.07.020]
- 4 **Llovet JM**, Kelley RK, Villanueva A, Singal AG, Pikarsky E, Roayaie S, Lencioni R, Koike K, Zucman-Rossi J, Finn RS. Hepatocellular carcinoma. *Nat Rev Dis Primers* 2021; **7**: 6 [PMID: 33479224 DOI: 10.1038/s41572-020-00240-3]
- 5 **Yang JD**, Heimbach JK. New advances in the diagnosis and management of hepatocellular carcinoma. *BMJ* 2020; **371**: m3544 [PMID: 33106289 DOI: 10.1136/bmj.m3544]
- 6 **Ahn JC**, Connell A, Simonetto DA, Hughes C, Shah VH. Application of Artificial Intelligence for the Diagnosis and Treatment of Liver Diseases. *Hepatology* 2021; **73**: 2546-2563 [PMID: 33098140 DOI: 10.1002/hep.31603]
- 7 **Haenlein M**, Kaplan A. A Brief History of Artificial Intelligence: On the Past, Present, and Future of Artificial Intelligence. *Calif Manage Rev* 2019 [DOI: 10.1177/0008125619864925]
- 8 **Koza JR**, Bennett FH, Andre D, Keane MA. Automated Design of Both the Topology and Sizing of Analog Electrical Circuits Using Genetic Programming. In: Gero JS, Sudweeks F. Artificial Intelligence in Design '96. Dordrecht: Springer Netherlands, 1996: 151-170
- 9 **Cunningham P**, Cord M, Delany SJ. Supervised Learning. In: Cord M, Cunningham P. Machine Learning Techniques for Multimedia: Case Studies on Organization and Retrieval. Berlin, Heidelberg: Springer Berlin Heidelberg, 2008: 21-49
- 10 **Ghahramani Z**, Unsupervised Learning. In: Bousquet O, von Luxburg U, Rätsch G. Advanced Lectures on Machine Learning: ML Summer Schools 2003, Canberra, Australia, February 2 - 14, 2003, Tübingen, Germany, August 4 - 16, 2003, Revised Lectures. Berlin, Heidelberg: Springer Berlin Heidelberg, 2004: 72-112
- 11 **Esteva A**, Robicquet A, Ramsundar B, Kuleshov V, DePristo M, Chou K, Cui C, Corrado G, Thrun S, Dean J. A guide to deep learning in healthcare. *Nat Med* 2019; **25**: 24-29 [PMID: 30617335 DOI: 10.1038/s41591-018-0316-z]
- 12 **Puttagunta M**, Ravi S. Medical image analysis based on deep learning approach. *Multimed Tools Appl* 2021; 1-34 [PMID: 33841033 DOI: 10.1007/s11042-021-10707-4]
- 13 **Marrero JA**, Kulik LM, Sirlin CB, Zhu AX, Finn RS, Abecassis MM, Roberts LR, Heimbach JK. Diagnosis, Staging, and Management of Hepatocellular Carcinoma: 2018 Practice Guidance by the American Association for the Study of Liver Diseases. *Hepatology* 2018; **68**: 723-750 [PMID: 29624699 DOI: 10.1002/hep.29913]
- 14 **Ioannou GN**, Tang W, Beste LA, Tincopa MA, Su GL, Van T, Tapper EB, Singal AG, Zhu J, Waljee AK. Assessment of a Deep Learning Model to Predict Hepatocellular Carcinoma in Patients With Hepatitis C Cirrhosis. *JAMA Netw Open* 2020; **3**: e2015626 [PMID: 32870314 DOI: 10.1001/jamanetworkopen.2020.15626]
- 15 **Phan DV**, Chan CL, Li AA, Chien TY, Nguyen VC. Liver cancer prediction in a viral hepatitis cohort: A deep learning approach. *Int J Cancer* 2020; **147**: 2871-2878 [PMID: 32761609 DOI: 10.1002/ijc.33245]
- 16 **Nam JY**, Sinn DH, Bae J, Jang ES, Kim JW, Jeong SH. Deep learning model for prediction of hepatocellular carcinoma in patients with HBV-related cirrhosis on antiviral therapy. *JHEP Rep* 2020; **2**: 100175 [PMID: 33117971 DOI: 10.1016/j.jhepr.2020.100175]
- 17 **Nam JY**, Lee JH, Bae J, Chang Y, Cho Y, Sinn DH, Kim BH, Kim SH, Yi NJ, Lee KW, Kim JM, Park JW, Kim YJ, Yoon JH, Joh JW, Suh KS. Novel Model to Predict HCC Recurrence after Liver Transplantation Obtained Using Deep Learning: A Multicenter Study. *Cancers (Basel)* 2020; **12** [PMID: 33003306 DOI: 10.3390/cancers12102791]
- 18 **Behne T**, Copur MS. Biomarkers for hepatocellular carcinoma. *Int J Hepatol* 2012; **2012**: 859076 [PMID: 22655201 DOI: 10.1155/2012/859076]
- 19 **Tzartzeva K**, Obi J, Rich NE, Parikh ND, Marrero JA, Yopp A, Waljee AK, Singal AG. Surveillance Imaging and Alpha Fetoprotein for Early Detection of Hepatocellular Carcinoma in Patients With Cirrhosis: A Meta-analysis. *Gastroenterology* 2018; **154**: 1706-1718.e1 [PMID: 29425931 DOI: 10.1053/j.gastro.2018.01.064]
- 20 **Xie H**, Xue YQ, Liu P, Zhang PJ, Tian ST, Yang Z, Guo Z, Wang HM. Multi-parameter gene expression profiling of peripheral blood for early detection of hepatocellular carcinoma. *World J Gastroenterol* 2018; **24**: 371-378 [PMID: 29391759 DOI: 10.3748/wjg.v24.i3.371]
- 21 **Choi J**, Oh I, Seo S, Ahn J. G2Vec: Distributed gene representations for identification of cancer

- prognostic genes. *Sci Rep* 2018; **8**: 13729 [PMID: 30213980 DOI: 10.1038/s41598-018-32180-0]
- 22 **Chaudhary K**, Poirion OB, Lu L, Garmire LX. Deep Learning-Based Multi-Omics Integration Robustly Predicts Survival in Liver Cancer. *Clin Cancer Res* 2018; **24**: 1248-1259 [PMID: 28982688 DOI: 10.1158/1078-0432.CCR-17-0853]
  - 23 **Chartrand G**, Cheng PM, Vorontsov E, Drozdal M, Turcotte S, Pal CJ, Kadoury S, Tang A. Deep Learning: A Primer for Radiologists. *Radiographics* 2017; **37**: 2113-2131 [PMID: 29131760 DOI: 10.1148/rg.2017170077]
  - 24 **Azer SA**. Deep learning with convolutional neural networks for identification of liver masses and hepatocellular carcinoma: A systematic review. *World J Gastrointest Oncol* 2019; **11**: 1218-1230 [PMID: 31908726 DOI: 10.4251/wjgo.v11.i12.1218]
  - 25 **Streba CT**, Ionescu M, Gheonea DI, Sandulescu L, Ciurea T, Saftoiu A, Vere CC, Rogoveanu I. Contrast-enhanced ultrasonography parameters in neural network diagnosis of liver tumors. *World J Gastroenterol* 2012; **18**: 4427-4434 [PMID: 22969209 DOI: 10.3748/wjg.v18.i32.4427]
  - 26 **Hassan TM**, Elmogy M, Sallam E-S. Diagnosis of Focal Liver Diseases Based on Deep Learning Technique for Ultrasound Images. *Arab J Sci Eng* 2017; **42**: 3127-3140 [DOI: 10.1007/s13369-016-2387-9]
  - 27 **Bharti P**, Mittal D, Ananthasivan R. Preliminary Study of Chronic Liver Classification on Ultrasound Images Using an Ensemble Model. *Ultrason Imaging* 2018; **40**: 357-379 [PMID: 30015593 DOI: 10.1177/0161734618787447]
  - 28 **Schmauch B**, Herent P, Jehanno P, Dehaene O, Saillard C, Aubé C, Luciani A, Lassau N, Jégou S. Diagnosis of focal liver lesions from ultrasound using deep learning. *Diagn Interv Imaging* 2019; **100**: 227-233 [PMID: 30926443 DOI: 10.1016/j.diii.2019.02.009]
  - 29 **Brehar R**, Mitrea DA, Vancea F, Marita T, Nedevschi S, Lupsor-Platon M, Rotaru M, Badea RI. Comparison of Deep-Learning and Conventional Machine-Learning Methods for the Automatic Recognition of the Hepatocellular Carcinoma Areas from Ultrasound Images. *Sensors (Basel)* 2020; **20** [PMID: 32485986 DOI: 10.3390/s20113085]
  - 30 **Jin J**, Yao Z, Zhang T, Zeng J, Wu L, Wu M, Wang J, Wang Y, Yu J, Zheng R. Deep learning radiomics model accurately predicts hepatocellular carcinoma occurrence in chronic hepatitis B patients: a five-year follow-up. *Am J Cancer Res* 2021; **11**: 576-589 [PMID: 33575088]
  - 31 **Yasaka K**, Akai H, Abe O, Kiryu S. Deep Learning with Convolutional Neural Network for Differentiation of Liver Masses at Dynamic Contrast-enhanced CT: A Preliminary Study. *Radiology* 2018; **286**: 887-896 [PMID: 29059036 DOI: 10.1148/radiol.2017170706]
  - 32 **Shi W**, Kuang S, Cao S, Hu B, Xie S, Chen S, Chen Y, Gao D, Zhu Y, Zhang H, Liu H, Ye M, Sirlin CB, Wang J. Deep learning assisted differentiation of hepatocellular carcinoma from focal liver lesions: choice of four-phase and three-phase CT imaging protocol. *Abdom Radiol (NY)* 2020; **45**: 2688-2697 [PMID: 32232524 DOI: 10.1007/s00261-020-02485-8]
  - 33 **Alirr OI**. Deep learning and level set approach for liver and tumor segmentation from CT scans. *J Appl Clin Med Phys* 2020; **21**: 200-209 [PMID: 33113290 DOI: 10.1002/acm2.13003]
  - 34 **Chlebus G**, Schenk A, Moltz JH, van Ginneken B, Hahn HK, Meine H. Automatic liver tumor segmentation in CT with fully convolutional neural networks and object-based postprocessing. *Sci Rep* 2018; **8**: 15497 [PMID: 30341319 DOI: 10.1038/s41598-018-33860-7]
  - 35 **Deng Z**, Guo Q, Zhu Z. Dynamic Regulation of Level Set Parameters Using 3D Convolutional Neural Network for Liver Tumor Segmentation. *J Healthc Eng* 2019; **2019**: 4321645 [PMID: 30918620 DOI: 10.1155/2019/4321645]
  - 36 **Wardhana G**, Naghibi H, Sirmacek B, Abayazid M. Toward reliable automatic liver and tumor segmentation using convolutional neural network based on 2.5D models. *Int J Comput Assist Radiol Surg* 2021; **16**: 41-51 [PMID: 33219906 DOI: 10.1007/s11548-020-02292-y]
  - 37 **Li X**, Chen H, Qi X, Dou Q, Fu CW, Heng PA. H-DenseUNet: Hybrid Densely Connected UNet for Liver and Tumor Segmentation From CT Volumes. *IEEE Trans Med Imaging* 2018; **37**: 2663-2674 [PMID: 29994201 DOI: 10.1109/TMI.2018.2845918]
  - 38 **Ahn SH**, Yeo AU, Kim KH, Kim C, Goh Y, Cho S, Lee SB, Lim YK, Kim H, Shin D, Kim T, Kim TH, Youn SH, Oh ES, Jeong JH. Comparative clinical evaluation of atlas and deep-learning-based auto-segmentation of organ structures in liver cancer. *Radiat Oncol* 2019; **14**: 213 [PMID: 31775825 DOI: 10.1186/s13014-019-1392-z]
  - 39 **Almotairi S**, Kareem G, Aouf M, Almutairi B, Salem MA. Liver Tumor Segmentation in CT Scans Using Modified SegNet. *Sensors (Basel)* 2020; **20** [PMID: 32164153 DOI: 10.3390/s20051516]
  - 40 **Ayalew YA**, Fante KA, Mohammed MA. Modified U-Net for liver cancer segmentation from computed tomography images with a new class balancing method. *BMC Biomed Eng* 2021; **3**: 4 [PMID: 33641679 DOI: 10.1186/s42490-021-00050-y]
  - 41 **Budak Ü**, Guo Y, Tanyildizi E, Şengür A. Cascaded deep convolutional encoder-decoder neural networks for efficient liver tumor segmentation. *Med Hypotheses* 2020; **134**: 109431 [PMID: 31669758 DOI: 10.1016/j.mehy.2019.109431]
  - 42 **Chen Y**, Wang K, Liao X, Qian Y, Wang Q, Yuan Z, Heng PA. Channel-Unet: A Spatial Channel-Wise Convolutional Neural Network for Liver and Tumors Segmentation. *Front Genet* 2019; **10**: 1110 [PMID: 31827487 DOI: 10.3389/fgene.2019.01110]
  - 43 **Hamm CA**, Wang CJ, Savic LJ, Ferrante M, Schobert I, Schlachter T, Lin M, Duncan JS, Weinreb JC, Chapiro J, Letzen B. Deep learning for liver tumor diagnosis part I: development of a convolutional neural network classifier for multi-phasic MRI. *Eur Radiol* 2019; **29**: 3338-3347 [PMID: 31016442 DOI: 10.1007/s00330-019-06205-9]

- 44 **Wang CJ**, Hamm CA, Savic LJ, Ferrante M, Schobert I, Schlachter T, Lin M, Weinreb JC, Duncan JS, Chapiro J, Letzen B. Deep learning for liver tumor diagnosis part II: convolutional neural network interpretation using radiologic imaging features. *Eur Radiol* 2019; **29**: 3348-3357 [PMID: [31093705](#) DOI: [10.1007/s00330-019-06214-8](#)]
- 45 **Wu Y**, White GM, Cornelius T, Gowdar I, Ansari MH, Supanich MP, Deng J. Deep learning LI-RADS grading system based on contrast enhanced multiphase MRI for differentiation between LR-3 and LR-4/LR-5 liver tumors. *Ann Transl Med* 2020; **8**: 701 [PMID: [32617321](#) DOI: [10.21037/atm.2019.12.151](#)]
- 46 **Zhen SH**, Cheng M, Tao YB, Wang YF, Juengpanich S, Jiang ZY, Jiang YK, Yan YY, Lu W, Lue JM, Qian JH, Wu ZY, Sun JH, Lin H, Cai XJ. Deep Learning for Accurate Diagnosis of Liver Tumor Based on Magnetic Resonance Imaging and Clinical Data. *Front Oncol* 2020; **10**: 680 [PMID: [32547939](#) DOI: [10.3389/fonc.2020.00680](#)]
- 47 **Zhang Y**, Lv X, Qiu J, Zhang B, Zhang L, Fang J, Li M, Chen L, Wang F, Liu S, Zhang S. Deep Learning With 3D Convolutional Neural Network for Noninvasive Prediction of Microvascular Invasion in Hepatocellular Carcinoma. *J Magn Reson Imaging* 2021; **54**: 134-143 [PMID: [33559293](#) DOI: [10.1002/jmri.27538](#)]
- 48 **Wang G**, Jian W, Cen X, Zhang L, Guo H, Liu Z, Liang C, Zhou W. Prediction of Microvascular Invasion of Hepatocellular Carcinoma Based on Preoperative Diffusion-Weighted MR Using Deep Learning. *Acad Radiol* 2020 [PMID: [33303346](#) DOI: [10.1016/j.acra.2020.11.014](#)]
- 49 **Jiang YQ**, Cao SE, Cao S, Chen JN, Wang GY, Shi WQ, Deng YN, Cheng N, Ma K, Zeng KN, Yan XJ, Yang HZ, Huan WJ, Tang WM, Zheng Y, Shao CK, Wang J, Yang Y, Chen GH. Preoperative identification of microvascular invasion in hepatocellular carcinoma by XGBoost and deep learning. *J Cancer Res Clin Oncol* 2021; **147**: 821-833 [PMID: [32852634](#) DOI: [10.1007/s00432-020-03366-9](#)]
- 50 **An C**, Jiang Y, Huang Z, Gu Y, Zhang T, Ma L, Huang J. Assessment of Ablative Margin After Microwave Ablation for Hepatocellular Carcinoma Using Deep Learning-Based Deformable Image Registration. *Front Oncol* 2020; **10**: 573316 [PMID: [33102233](#) DOI: [10.3389/fonc.2020.573316](#)]
- 51 **Liu F**, Liu D, Wang K, Xie X, Su L, Kuang M, Huang G, Peng B, Wang Y, Lin M, Tian J. Deep Learning Radiomics Based on Contrast-Enhanced Ultrasound Might Optimize Curative Treatments for Very-Early or Early-Stage Hepatocellular Carcinoma Patients. *Liver Cancer* 2020; **9**: 397-413 [PMID: [32999867](#) DOI: [10.1159/000505694](#)]
- 52 **Peng J**, Kang S, Ning Z, Deng H, Shen J, Xu Y, Zhang J, Zhao W, Li X, Gong W, Huang J, Liu L. Residual convolutional neural network for predicting response of transarterial chemoembolization in hepatocellular carcinoma from CT imaging. *Eur Radiol* 2020; **30**: 413-424 [PMID: [31332558](#) DOI: [10.1007/s00330-019-06318-1](#)]
- 53 **Liu QP**, Xu X, Zhu FP, Zhang YD, Liu XS. Prediction of prognostic risk factors in hepatocellular carcinoma with transarterial chemoembolization using multi-modal multi-task deep learning. *EClinicalMedicine* 2020; **23**: 100379 [PMID: [32548574](#) DOI: [10.1016/j.eclinm.2020.100379](#)]
- 54 **Zhang L**, Xia W, Yan ZP, Sun JH, Zhong BY, Hou ZH, Yang MJ, Zhou GH, Wang WS, Zhao XY, Jian JM, Huang P, Zhang R, Zhang S, Zhang JY, Li Z, Zhu XL, Gao X, Ni CF. Deep Learning Predicts Overall Survival of Patients With Unresectable Hepatocellular Carcinoma Treated by Transarterial Chemoembolization Plus Sorafenib. *Front Oncol* 2020; **10**: 593292 [PMID: [33102242](#) DOI: [10.3389/fonc.2020.593292](#)]
- 55 **Lin H**, Wei C, Wang G, Chen H, Lin L, Ni M, Chen J, Zhuo S. Automated classification of hepatocellular carcinoma differentiation using multiphoton microscopy and deep learning. *J Biophotonics* 2019; **12**: e201800435 [PMID: [30868728](#) DOI: [10.1002/jbio.201800435](#)]
- 56 **Kiani A**, Uyumazturk B, Rajpurkar P, Wang A, Gao R, Jones E, Yu Y, Langlotz CP, Ball RL, Montine TJ, Martin BA, Berry GJ, Ozawa MG, Hazard FK, Brown RA, Chen SB, Wood M, Allard LS, Ylagan L, Ng AY, Shen J. Impact of a deep learning assistant on the histopathologic classification of liver cancer. *NPJ Digit Med* 2020; **3**: 23 [PMID: [32140566](#) DOI: [10.1038/s41746-020-0232-8](#)]
- 57 **Liao H**, Long Y, Han R, Wang W, Xu L, Liao M, Zhang Z, Wu Z, Shang X, Li X, Peng J, Yuan K, Zeng Y. Deep learning-based classification and mutation prediction from histopathological images of hepatocellular carcinoma. *Clin Transl Med* 2020; **10**: e102 [PMID: [32536036](#) DOI: [10.1002/ctm2.102](#)]
- 58 **Wang H**, Jiang Y, Li B, Cui Y, Li D, Li R. Single-Cell Spatial Analysis of Tumor and Immune Microenvironment on Whole-Slide Image Reveals Hepatocellular Carcinoma Subtypes. *Cancers (Basel)* 2020; **12** [PMID: [33260561](#) DOI: [10.3390/cancers12123562](#)]
- 59 **Chen M**, Zhang B, Topatana W, Cao J, Zhu H, Juengpanich S, Mao Q, Yu H, Cai X. Classification and mutation prediction based on histopathology H&E images in liver cancer using deep learning. *NPJ Precis Oncol* 2020; **4**: 14 [PMID: [32550270](#) DOI: [10.1038/s41698-020-0120-3](#)]
- 60 **Lu L**, Daigle BJ Jr. Prognostic analysis of histopathological images using pre-trained convolutional neural networks: application to hepatocellular carcinoma. *PeerJ* 2020; **8**: e8668 [PMID: [32201640](#) DOI: [10.7717/peerj.8668](#)]
- 61 **Saillard C**, Schmauch B, Laifa O, Moarii M, Toldo S, Zaslavskiy M, Pronier E, Laurent A, Amaddeo G, Regnault H, Sommacale D, Ziolo M, Pawlotsky JM, Mulé S, Luciani A, Wainrib G, Clozel T, Courtiol P, Calderaro J. Predicting Survival After Hepatocellular Carcinoma Resection Using Deep Learning on Histological Slides. *Hepatology* 2020; **72**: 2000-2013 [PMID: [32108950](#) DOI: [10.1002/hep.31207](#)]
- 62 **Shi JY**, Wang X, Ding GY, Dong Z, Han J, Guan Z, Ma LJ, Zheng Y, Zhang L, Yu GZ, Wang XY, Ding ZB, Ke AW, Yang H, Wang L, Ai L, Cao Y, Zhou J, Fan J, Liu X, Gao Q. Exploring prognostic

- indicators in the pathological images of hepatocellular carcinoma based on deep learning. *Gut* 2021; **70**: 951-961 [PMID: [32998878](#) DOI: [10.1136/gutjnl-2020-320930](#)]
- 63 **Yamashita R**, Long J, Saleem A, Rubin DL, Shen J. Deep learning predicts postsurgical recurrence of hepatocellular carcinoma from digital histopathologic images. *Sci Rep* 2021; **11**: 2047 [PMID: [33479370](#) DOI: [10.1038/s41598-021-81506-y](#)]
- 64 **Gumbs AA**, Perretta S, d'Allemagne B, Chouillard E. What is Artificial Intelligence Surgery? *Art Int Surg* 2021; **1**: 1-10 [DOI: [10.20517/ais.2021.01](#)]
- 65 **Lee D**, Yu HW, Kwon H, Kong HJ, Lee KE, Kim HC. Evaluation of Surgical Skills during Robotic Surgery by Deep Learning-Based Multiple Surgical Instrument Tracking in Training and Actual Operations. *J Clin Med* 2020; **9** [PMID: [32585953](#) DOI: [10.3390/jcm9061964](#)]





Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

**Telephone:** +1-925-3991568

**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

**Help Desk:** <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

