World Journal of *Hepatology*

World J Hepatol 2024 March 27; 16(3): 300-489





Published by Baishideng Publishing Group Inc

World Journal of Hepatology

Contents

Monthly Volume 16 Number 3 March 27, 2024

EDITORIAL

300 Is there a need for universal double reflex testing of HBsAg-positive individuals for hepatitis D infection? Abbas Z. Abbas M

REVIEW

- 304 Non-alcoholic fatty liver disease and sleep disorders Bu LF, Xiong CY, Zhong JY, Xiong Y, Li DM, Hong FF, Yang SL
- 316 Amebic liver abscess: An update Kumar R, Patel R, Priyadarshi RN, Narayan R, Maji T, Anand U, Soni JR

MINIREVIEWS

331 Advances in discovery of novel investigational agents for functional cure of chronic hepatitis B: A comprehensive review of phases II and III therapeutic agents

Lam R. Lim JK

- 344 Protein succinylation, hepatic metabolism, and liver diseases Liu S, Li R, Sun YW, Lin H, Li HF
- 353 Hepatocellular carcinoma immune microenvironment and check point inhibitors-current status Gupta T, Jarpula NS

ORIGINAL ARTICLE

Case Control Study

Associations of PNPLA3 and LEP genetic polymorphisms with metabolic-associated fatty liver disease in 366 Thai people living with human immunodeficiency virus

Choochuay K, Kunhapan P, Puangpetch A, Tongsima S, Srisawasdi P, Sobhonslidsuk A, Sungkanuparph S, Biswas M, Sukasem Č

Retrospective Cohort Study

379 Comparison of fungal vs bacterial infections in the medical intensive liver unit: Cause or corollary for high mortality?

Khan S, Hong H, Bass S, Wang Y, Wang XF, Sims OT, Koval CE, Kapoor A, Lindenmeyer CC

393 Lean body mass index is a marker of advanced tumor features in patients with hepatocellular carcinoma deLemos AS, Zhao J, Patel M, Kooken B, Mathur K, Nguyen HM, Mazhar A, McCarter M, Burney H, Kettler C, Chalasani N, Gawrieh S



Contents

Monthly Volume 16 Number 3 March 27, 2024

Retrospective Study

405 Prediction model for hepatitis B e antigen seroconversion in chronic hepatitis B with peginterferon-alfa treated based on a response-guided therapy strategy

Zhang PX, Zheng XW, Zhang YF, Ye J, Li W, Tang QQ, Zhu J, Zou GZ, Zhang ZH

418 Retrospective study of the incidence, risk factors, treatment outcomes of bacterial infections at uncommon sites in cirrhotic patients

Schneitler S, Schneider C, Casper M, Lammert F, Krawczyk M, Becker SL, Reichert MC

Observational Study

428 Palliative long-term abdominal drains vs large volume paracenteses for the management of refractory ascites in end-stage liver disease

Kaur S, Motta RV, Chapman B, Wharton V, Collier JD, Saffioti F

Basic Study

439 Comprehensive prognostic and immune analysis of sterol O-acyltransferase 1 in patients with hepatocellular carcinoma

Gan CJ, Zheng Y, Yang B, Cao LM

SYSTEMATIC REVIEWS

452 Update in lean metabolic dysfunction-associated steatotic liver disease

Sato-Espinoza K, Chotiprasidhi P, Huaman MR, Díaz-Ferrer J

META-ANALYSIS

465 Influence of nonalcoholic fatty liver disease on response to antiviral treatment in patients with chronic hepatitis B: A meta-analysis

Liu SY, Wang D, Liu J, Yang LP, Chen GY

477 Prognostic value of neutrophil-to-lymphocyte ratio in end-stage liver disease: A meta-analysis Cai XH, Tang YM, Chen SR, Pang JH, Chong YT, Cao H, Li XH



Contents

Monthly Volume 16 Number 3 March 27, 2024

ABOUT COVER

Editorial Board Member of World Journal of Hepatology, Marcello Maida, MD, Doctor, Gastroenterology and Endoscopy Unit, S.Elia - Raimondi Hospital, Caltanissetta, Caltanissetta 93100, Italy. marcello.maida@hotmail.it

AIMS AND SCOPE

The primary aim of World Journal of Hepatology (WJH, World J Hepatol) is to provide scholars and readers from various fields of hepatology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJH mainly publishes articles reporting research results and findings obtained in the field of hepatology and covering a wide range of topics including chronic cholestatic liver diseases, cirrhosis and its complications, clinical alcoholic liver disease, drug induced liver disease autoimmune, fatty liver disease, genetic and pediatric liver diseases, hepatocellular carcinoma, hepatic stellate cells and fibrosis, liver immunology, liver regeneration, hepatic surgery, liver transplantation, biliary tract pathophysiology, non-invasive markers of liver fibrosis, viral hepatitis.

INDEXING/ABSTRACTING

The WJH is now abstracted and indexed in PubMed, PubMed Central, Emerging Sources Citation Index (ESCI), Scopus, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJH as 2.4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Yi-Xuan Cai; Production Department Director: Xiang Li; Cover Editor: Xiang Li.

NAME OF JOURNAL	INSTRUCTIONS TO AUTHORS			
World Journal of Hepatology	https://www.wjgnet.com/bpg/gerinfo/204			
ISSN	GUIDELINES FOR ETHICS DOCUMENTS			
ISSN 1948-5182 (online)	https://www.wjgnet.com/bpg/GerInfo/287			
LAUNCH DATE	GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH			
October 31, 2009	https://www.wjgnet.com/bpg/gerinfo/240			
FREQUENCY	PUBLICATION ETHICS			
Monthly	https://www.wjgnet.com/bpg/GerInfo/288			
EDITORS-IN-CHIEF	PUBLICATION MISCONDUCT			
Nikolaos Pyrsopoulos, Ke-Qin Hu, Koo Jeong Kang	https://www.wjgnet.com/bpg/gerinfo/208			
EXECUTIVE ASSOCIATE EDITORS-IN-CHIEF	POLICY OF CO-AUTHORS			
Shuang-Suo Dang	https://www.wjgnet.com/bpg/GerInfo/310			
EDITORIAL BOARD MEMBERS	ARTICLE PROCESSING CHARGE			
nttps://www.wjgnet.com/1948-5182/editorialboard.htm	https://www.wignet.com/bpg/gerinfo/242			
PUBLICATION DATE	STEPS FOR SUBMITTING MANUSCRIPTS			
March 27, 2024	https://www.wjgnet.com/bpg/GerInfo/239			
COPYRIGHT	ONLINE SUBMISSION			
© 2024 Baishideng Publishing Group Inc	https://www.f6publishing.com			
PUBLISHING PARTNER	PUBLISHING PARTNER's OFFICIAL WEBSITE			
Department of Infectious Diseases, the Second Affiliated Hospital of Xi'an liaotong University	http://2yuan.xjtu.edu.cn/Html/Departments/Main/Index_21148.html			

© 2024 Baishideng Publishing Group Inc. All rights reserved. 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA E-mail: office@baishideng.com https://www.wjgnet.com



W J H World Journal of Henatology Hepatology

Submit a Manuscript: https://www.f6publishing.com

World J Hepatol 2024 March 27; 16(3): 331-343

DOI: 10.4254/wjh.v16.i3.331

ISSN 1948-5182 (online)

MINIREVIEWS

Advances in discovery of novel investigational agents for functional cure of chronic hepatitis B: A comprehensive review of phases II and III therapeutic agents

Robert Lam, Joseph K Lim

Specialty type: Gastroenterology and hepatology

Provenance and peer review: Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0 Grade B (Very good): 0 Grade C (Good): C Grade D (Fair): D Grade E (Poor): 0

P-Reviewer: Janicko M, Slovakia; Kapritsou M, Greece

Received: November 27, 2023 Peer-review started: November 27, 2023 First decision: January 5, 2024 Revised: January 23, 2024 Accepted: February 29, 2024 Article in press: February 29, 2024 Published online: March 27, 2024



Robert Lam, Joseph K Lim, Section of Digestive Diseases, Yale University School of Medicine, New Haven, CT 06520, United States

Corresponding author: Joseph K Lim, MD, Professor, Section of Digestive Diseases, Yale University School of Medicine, Yale Liver Center, 333 Cedar Street, LMP 1080, New Haven, CT 06520, United States. joseph.lim@yale.edu

Abstract

Chronic hepatitis B virus (HBV) infection affects over 295 million people globally and an estimated 1.6 million people in the United States. It is associated with significant morbidity and mortality due to cirrhosis, liver failure, and liver cancer. Antiviral therapy with oral nucleos(t)ide analogues is associated with high rates of virologic suppression, which in turn has been associated with a decreased risk of liver complications. However, current antiviral regimens are limited by concerns with adverse effects, adherence, resistance, long-term treatment, and ongoing risk for liver events. Novel investigational agents are currently in development and are targeted at achieving functional cure with sustained hepatitis B surface antigen (HBsAg) loss and suppression of HBV DNA. Herein we review key evidence from phases II and III trials defining the efficacy and safety profiles for key investigational agents for functional cure of chronic hepatitis B, including core/capsid inhibitors, entry inhibitors, RNA interference (siRNA/ASO), HBsAg inhibitors, Toll-like receptor agonists, checkpoint inhibitors, and therapeutic vaccines.

Key Words: Hepatitis B virus; Treatment; Clinical trials; RNA interference; Entry inhibitors; Core inhibitors; Immunomodulators

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Novel investigational agents targeting functional cure [sustained hepatitis B surface antigen (HBsAg) loss and undetectable hepatitis B virus (HBV) DNA] are currently in clinical trial development. Herein we review key evidence from phases 2 and 3 trials defining the efficacy and safety profiles for key investigational agents, including core/capsid inhibitors, entry inhibitors, RNA interference (siRNA/ASO), HBsAg inhibitors, Toll-like receptor agonists, checkpoint inhibitors, and therapeutic vaccines.

Citation: Lam R, Lim JK. Advances in discovery of novel investigational agents for functional cure of chronic hepatitis B: A comprehensive review of phases II and III therapeutic agents. World J Hepatol 2024; 16(3): 331-343 URL: https://www.wjgnet.com/1948-5182/full/v16/i3/331.htm DOI: https://dx.doi.org/10.4254/wjh.v16.i3.331

INTRODUCTION

Chronic hepatitis B virus (HBV) infection is a global health problem with more than 295 million people infected worldwide and as many as 1.6 million people infected in the United States[1,2]. Complications of chronic HBV infection, including cirrhosis, hepatocellular carcinoma (HCC), and liver failure, may take years to develop and have led to more than 800000 deaths each year[3-5].

While the ideal goal of HBV therapy would be a complete sterilizing cure, such a therapy does not exist because it is difficult to directly target the covalently closed circular DNA (cccDNA) in the hepatocyte nucleus and the integration of HBV DNA into the host genome. The best that can be achieved with current therapies is a functional cure where there is loss of HBV surface antigen (HBsAg) with undetectable HBV DNA after 6 months off therapy. This is an important endpoint given its association with reduced liver necroinflammation, reduced risk of HCC, increased liver fibrosis regression, normalization of alanine aminotransferase (ALT) levels, reduced risk of liver cirrhosis, and increased survival [6-11].

Current FDA-approved therapies include pegylated interferons (PEGIFN α) and nucleos(t)ide analogues (NA)[12,13]. PEGIFN α are administered as subcutaneous injections on a once weekly dosing schedule for one year. They exert antiviral and immunomodulatory activities by enhancing cccDNA degradation and modifying cccDNA transcription. PEGIFNα therapy has higher rates of HBsAg loss and HBV e-antigen (HBeAg) seroconversion than NA, but are associated with poor tolerability and risk for depression [14,15]. In contrast, NA are administered orally every day. They suppress HBV replication by causing chain termination when incorporated into HBV DNA undergoing reverse transcription. Early NA such as Lamivudine and Adefovir had high rates of antiviral resistance with only a few years of treatment[16]. NA currently used in clinical practice, namely, Tenofovir disoproxil fumarate (TDF) and Entecavir (ETV), have potent antiviral activity and a high barrier to resistance^[17]. Compared to PEGIFNα therapy, NA are well-tolerated, but require a long term duration of maintenance therapy[18]. Both PEG-IFNα and NA therapies are unable to eliminate the HBV because they do not directly target cccDNA and integrated HBV DNA. Consequently, cccDNA persists, which enables transcription of RNA and translation of viral HBV proteins, such as HBsAg, to continue[19,20].

Rates of a functional cure with PEGIFN α and NA therapies are low. With PEGIFN α treatment, HBsAg loss has only been reported in approximately 7% of both HBeAg positive and negative patients after a year of treatment[21]. HBsAg loss is even lower in patients receiving NA, with only 0.3%-5% of HBeAg negative patients and 1.4% of HBeAg positive patients achieving HBsAg loss after treatment for 5-7 years[22,23]. Given the limitations of existing HBV therapies, there is great interest in novel HBV therapeutics that can lead to the following outcomes: Functional cure, improvement in quality of life, and preventing progression of chronic HBV infection to cirrhosis, HCC, and HBV-related mortality. Herein, this review will focus on novel HBV therapies in active phases II and III clinical trial development.

METHODOLOGY

This paper is a narrative review. Investigational agents for treatment of chronic HBV infection under active phases II and III development were identified using the National Institutes of Health Clinical Trials directory^[24]. This directory includes details regarding the study design, population, treatment arms, and sponsoring pharmaceutical company for all publicly supported clinical studies. Information from this website was incorporated in the development of Table 1, which summarizes information about investigational agents in phases II and III trials without published study results. A PubMed search was conducted for each investigational agent under active phases II and II development. Data was retrieved from published original research articles and conference abstracts. The sponsoring pharmaceutical company website for each investigational agent was reviewed for published presentation slides from international liver meetings.

FUTURE THERAPY FOR HEPATITIS B

A summary of the novel therapies in phases II and III development with study data are listed in Table 2, while therapies



Table 1 Novel therapeutic agents for treatment of chronic hepatitis B infection in phase II or III development								
Drug sponsor	Phase							
Janssen Pharmaceutics	П							
Assembly Biosciences	П							
Gilead Sciences	III							
Ionis Pharmaceuticals	III							
Vir Biotechnology	П							
Vir Biotechnology	Ш							
GlaxoSmithKline Pharmaceuticals	Ш							
Arbutus Biopharma	Ш							
Replicor	Ш							
Gilead Sciences	Ш							
Gilead Sciences	Ш							
Gilead Sciences	Ш							
Brii Biosciences	п							
	itis B infection in phase II or III development Drug sponsor Janssen Pharmaceutics Assembly Biosciences Gilead Sciences Vir Biotechnology Vir Biotechnology GlaxoSmithKline Pharmaceuticals Arbutus Biopharma Replicor Gilead Sciences Gilead Sciences Gilead Sciences Gilead Sciences Brii Biosciences							

Table 2 Summary of novel investigational agents in phase II trials for treatment of chronic hepatitis B infection

Drug name (therapeutic class)	Drug sponsor	Phase	Trial ID	Study design	Study population	Sample size	Intervention and control	Primary outcome
RG6346 (siRNA); RO7020531 (TLR-7 agonist)	Hoffman-La Roche	П	NCT04225715	Randomized, open-label, parallel assignment	Chronic HBV infection patients on established NA monotherapy for ≥ 12 months, HBV DNA < 20 IU/mL, ALT ≤ 1.5 ULN	280	Control arm: NA; Experi- mental arms: (1) CpAM (RO7049389) + TLR-7 agonist (RO7020531) + NA; (2) siRNA (RG6346) + NA; (3) siRNA (RG6346) + PEG- IFN + NA; (4) siRNA (RG6346) + TLR (RO7020531) + NA; (5) siRNA (RG6346) + PD-L1 LNA (R07191863) + NA	Percentage of participants with HBsAg loss at 24 wk after end of treatment
GC1102 (HBsAg neutralizing antibody)	Green Cross Corporation	Ш	NCT03801798	Double-blind, randomized, placebo- controlled, parallel-group	Chronic HBV infection patients on NA \ge 24 wk before screening	42	Control arm: NA + placebo; Experimental arm: NA + GC1102	Proportion of participants with ≥ 1 log IU/mL reduction in HBsAg titer

CpAM: Core Protein Allosteric Modulator; HBV: Hepatitis B Virus; HBsAg: Hepatitis B surface antigen; LLOQ: Lower limit of quantification; NA: Nucleos(t)ide therapy; TLR: Toll-like receptor; SiRNA: Short interfering RNA; ULN: Upper limit of normal.

without study data are listed in Table 1.

Core/capsid inhibitors

Capsid allosteric modulators directly target the destabilization of HBV core protein, resulting in the formation of abnormal capsids or morphologically normal capsids lacking genetic material[25]. This prevents further release and spread of HBV to other hepatocytes.



Znishideng® WJH | https://www.wjgnet.com

JNJ56136379 (JNJ-6379): JNJ56136379 (also known as JNJ-6379) targets the HBV capsid assembly process needed for HBV replication. It accelerates the rate of HBV capsid assembly to form empty, morphologically intact viral capsids and has a secondary mechanism of inhibiting de novo cccDNA[26].

The JADE study was a randomized, partially blinded, placebo-controlled phase II study evaluating the efficacy and safety of JNJ-6379 in 232 adults with non-cirrhotic, chronic HBV infection. Participants were virally suppressed or not on active treatment at the time of entry into the clinical trial. Participants were randomized to receive JNJ-6379, either as monotherapy or in combination with NA (ETV or TDF), and then compared to a control group of placebo plus NA. Dosing of JNJ-6379 at 75 mg and 250 mg daily was investigated. Overall, JNJ6379 did not show a clear benefit over NA monotherapy. The primary endpoint of a 1 log IU/mL mean decrease in HBsAg from baseline to week 24 for the JNJ-6379 treatment groups and control was not achieved. Specifically, the mean change in HBsAg compared to baseline for the JNJ-6379 treatment groups ranged from -0.41 to 0.11 log IU/mL. Among participants who were HBeAg-positive at the start of the study, there was also a limited reduction of HBeAg of 0.49 and 0.70 log IU/mL for JNJ-6379 75 mg and 250 mg plus NA treatment groups, respectively. Over the 24-wk follow-up period, the use of JNJ6379 both as monotherapy and in combination with NA led to a marked reduction in both HBV DNA (mean JNJ-6379 75 mg or 250 mg plus NA HBV DNA reduction of 5.53 and 5.58 log IU/mL, respectively, compared with 5.21 log IU/mL in the placebo plus NA group) and HBV RNA (mean JNJ-6379 75 mg and 250 mg plus NA HBV RNA reduction was 2.96 and 3.15 log IU/mL, respectively, compared with 1.33 log IU/mL in the placebo plus NA group). Both doses of JNJ6379 were safe and welltolerated^{26]}.

ABI-H0731 (Vebicorvir): ABI-H0731 (Vebicorvir/VBR) is an orally administered small molecule that disrupts HBV replication by inducing altered, non-functional core protein assembly^[27].

One of the phase II studies evaluated the efficacy and safety of VBR in combination with ETV for treatment-naïve, noncirrhotic, HBeAg positive study participants. In this double-blind, randomized, and placebo-controlled study, participants either received a combination of once daily VBR 300 mg daily and ETV 0.5 mg daily, or a combination of placebo and ETV 0.5 mg daily. The study revealed that the combination of VBR and ETV was safe and well-tolerated, and augmented a reduction of HBV DNA and RNA. The primary endpoint was achieved as there was a significantly greater mean log reduction in HBV DNA from baseline with VBR plus ETV combination therapy as compared to placebo plus ETV therapy at both treatment weeks 12 (-4.45 log IU/mL with VBR and ETV vs -3.3 log IU/mL with placebo and ETV) and 24 (-5.33 log IU/mL with VBR and ETV vs -4.2 log IU/mL with placebo and ETV). Furthermore, a greater proportion of patients had normalized ALT levels by treatment week 24 among the VBR and ETV combination therapy group (12/13 participants) as compared to the placebo and ETV therapy group (5/12 participants). No resistance breakthrough occurred with the use of VBR. The study demonstrated that VBR can be combined with current NA therapy to enhance anti-viral activity in treatment-naïve patients with chronic HBV infection[27].

Another phase II study evaluated the efficacy and safety of combination VBR and NA therapy as compared to NA monotherapy in non-cirrhotic, chronic HBV participants who were virally suppressed by NA for at least 6 months. The 73 enrolled study participants were randomized to receive VBR 300 mg daily plus NA or matching placebo plus NA for 24 wk. Results showed that there was no difference between the two groups for the change in HBsAg or HBeAg from baseline to treatment week 24. Of note, the combination of VBR plus NA led to a more marked reduction of HBV DNA and pregenomic RNA at week 24 from baseline compared to the placebo plus NA group, irrespective of HBeAg status. Among patients with detectable HBV DNA at baseline, there were a greater proportion of patients in the VBR plus NA group (29/35 HBeAg+ patients, 16/17 HBeAg- patients) compared to the placebo plus NA group (17/59 HBeAg+ patients, 10/14 HBeAg- patients) who achieved undetectable HBV DNA levels at week 24. VBR was found to be safe and well-tolerated. This clinical study provided further support that even greater levels of viral suppression can occur with the addition of a VBR core inhibitor to existing NA therapies, although the clinical significance of this is yet to be investigated[28].

Entry inhibitors

Entry inhibitors target the function of HBV surface proteins or host receptors to prevent HBV entry into the host cell required for infection[29].

Bulevirtide (formerly Myrcludex): Bulevirtide is a synthetic myristoylated peptide entry inhibitor that competitively binds and blocks a hepatocyte surface protein, sodium taurocholate cotransporting polypeptide (NTCP) receptor, such that HBsAg is unable to enter the hepatocyte^[29]. Hepatitis D virus (HDV) uses the same NTCP receptor as HBV, so Bulevirtide has been also used to prevent co-infection by HDV[30]. Increases in bile acid level are expected since NTCP plays a role in bile transport[31].

The MYR-201 study was a phase Ib/IIa, randomized, open-label study investigating the safety and efficacy of Bulevirtide with regard to the HBV and HDV virologic response and tolerability. The study featured 24 participants randomized to receive either Bulevirtide for 24 wk followed by PEGIFNα-2a weekly for 48 wk (Bulevirtide cohort), 2 mg Bulevirtide daily plus PEGIFNα-2a weekly for 24 wk followed by 24 wk of PEGIFNα-2a alone (Bulevirtide-IFN cohort), or PEGIFNα-2a weekly alone for 48 wk (IFN cohort). Study results revealed that HBsAg levels remained unchanged compared to baseline throughout the study in all treatment groups. ALT normalized in 6/8 patients in the Bulevirtide cohort compared to only 1/15 patient in the Bulevirtide-IFN and IFN cohorts. Notably, mean HBV DNA was significantly reduced by $10^{1.28}$ copies/mL at week 24 from baseline in the Bulevirtide-IFN cohort, with 6/7 patients showing a $\geq 1 \log 1$ decline. There was a non-significant decline of the HBV DNA from baseline in the IFN and Bulevirtide cohorts. This was the first proof-of-concept study showing that Bulevirtide was safe and well-tolerated, and could enhance viral suppression when used in combination with PEGIFN α -2a[32].



Another phase II, multicenter, open-label study, known as MYR-202, randomized patients into four groups: 2 mg subcutaneous Bulevirtide daily with TDF daily, 5 mg subcutaneous Bulevirtide daily with TDF daily, 10 mg subcutaneous Bulevirtide daily with TDF daily, or TDF alone for a total of 24 wk. Therapeutic impact on HBsAg was investigated as a secondary endpoint. There was no significant change in HBsAg concentration from baseline in any of the treatment groups throughout the treatment and follow-up period. Like the MYR-201 study, Bulevirtide in the MYR-202 study was well-tolerated. Common treatment-related adverse events included elevations in asymptomatic bile salt levels and ALT levels[33].

MYR-203 assessed the safety and efficacy of Bulevirtide alone or in combination with PEGIFN α for 48 wk. Treatment arms included PEGIFN α alone weekly, 2 mg Bulevirtide daily plus PEGIFN α weekly, 5 mg Bulevirtide daily plus PEGIFN α weekly, or 2 mg Bulevirtide daily for a total of 48 wk. This was then followed by a treatment free period of 24 wk. By weeks 48 and 72, there was a > 1 log reduction from baseline or undetectable HBsAg levels in the Bulevirtide plus PEGIFN α combination groups, but not in the monotherapy groups. Specifically, by 72 wk, 6/15 participants in the combination arm of the 2 mg Bulevirtide plus PEGIFN α group and 2/15 participants in the 5 mg Bulevirtide plus PEGIFN α group achieved either a > 1 log IU/mL decline or undetectable levels of HBsAg. MYR-203 study findings demonstrated a potential role of combination Bulevirtide and PEGIFN α therapy in future HBV cure given that it led to a large proportion of patients achieving HBsAg loss[34].

The MYR-204 multicenter, randomized phase II trial studied the safety and efficacy of Bulevirtide administered subcutaneously at 2 mg or 10 mg daily dosing in combination with PEGIFN α weekly compared to Bulevirtide 10 mg monotherapy over 48 wk. Interim data at the 24-wk mark showed that a > 1 log IU/mL decline in HBsAg levels from baseline was achieved only in the Bulevirtide and PEGIFN α combination groups (10/100 participants) and the PEGIFN α alone group (1/24 participants). There was a modest decline in HBv DNA from baseline in the groups that received Bulevirtide (mean HBV DNA change ranged from -0.3 to -0.7 log IU/mL)[35].

The MYR-301 trial was the first phase III multicenter, randomized, parallel design study of Bulevirtide monotherapy at 2 and 10 mg daily dosing compared to no active anti-HDV treatment for 48 wk, defined as delayed treatments. For HBV efficacy endpoints at week 48, no patient in any group experienced HBsAg loss and changes in HBsAg from baseline were minimal. Only a small decline in HBV DNA levels was observed with Bulevirtide treatment. No severe adverse effects were observed in patients receiving Bulevirtide that led to discontinuation of the drug[36].

Small interfering RNAs

Small interfering RNAs (siRNAs) are short RNA molecules that hybridize to specific viral mRNA sequences and target bound mRNA for degradation[37]. Effectively, siRNA prevents the expression of HBV proteins needed for replication.

Bepirovirsen (GSK3228836): Bepirovirsen is an antisense oligonucleotide that targets all HBV RNA, including mRNA and pregenomic RNA, and designates it for degradation[38].

One of the two phase II randomized controlled trials evaluated the safety, tolerability, and antiviral activity of Bepirovirsen. The study enrolled 24 treatment-naïve participants and 7 participants receiving stable NA therapy with chronic HBV infection. Patients who were treatment-naïve were randomized to receive placebo or Bepirovirsen at a dose of 150 mg or 300 mg. Patients on stable NA therapy were randomized to receive placebo or Bepirovirsen at a dose of 300 mg. Bepirovirsen was administered twice weekly for 2 wk and then once weekly for another 2 wk, after which patients were followed for 26 wk to assess for a change in HBsAg levels from baseline. After 4 wk of treatment with 300 mg Bepirovirsen for treatment-naïve patients, there was a significant decrease in HBsAg levels and HBV DNA from baseline compared to placebo; this was not observed in the Bepirovirsen 150 mg group. Specifically, among treatment-naïve subjects, there was a mean 1.56 log IU/mL reduction in HBsAg in the Bepirovirsen 300 mg group from baseline to day 29, as compared to a 0.5 log IU/mL reduction in the Bepirovirsen 150 mg group and < 0.07 log IU/mL reduction in the placebo group. The timing of HBsAg reduction in responders occurred rapidly after 4 wk of therapy. Bepirovirsen was found to have a favorable safety profile and treatment response, which encouraged its use in a larger study cohort[39].

The B-Clear Trial was a phase IIb randomized controlled study investigating the efficacy and safety of Bepirovirsen in 457 enrolled participants with chronic HBV infection when used for 12 and 24 wk. Results revealed that 6/68 participants and 7/70 participants who received 24 wk of Bepirovirsen once weekly with and without NA therapy, respectively, achieved HBsAg and HBV DNA loss that persisted for 24 wk following the end of the treatment period. While there were similar results of HBsAg loss irrespective of NA therapy use or HBeAg status, HBsAg loss among patients who were HBeAg-positive only occurred in those receiving NA therapy. The study also showed that levels of HBsAg at baseline can be predictive of response to therapy. Specifically, receiver operating characteristic curve analysis revealed that baseline HBsAg level < 3000 IU/mL was the cutoff level associated with functional cure when treated with Bepirovirsen. Common adverse events observed more commonly in the study cohort receiving Bepirovirsen compared to placebo included injection site reactions, pyrexia, fatigue, and increased ALT levels. A brief increase in HBV DNA observed after stopping Bepirovirsen raised potential concerns about the durability of treatment response; however, these blips in HBV DNA levels were postulated to be due to spontaneous release of virions from the hepatic reservoir. Durability of treatment response will be investigated in future studies with longer follow-up time[40].

VIR-2218: VIR-2218 is a triantennary N-acetyl galactosamine (GalNAc) conjugated siRNA that targets the X region of the HBV genome[41]. As the X region contains overlapping HBV gene templates, the use of a single siRNA can effectively silence all HBV RNA production in this region. VIR-2218 can also suppress the X-mediated upregulation of cccDNA transcription.

VIR-2218-1001 was a two-part, phase I/II, randomized, double-blind, and placebo-controlled study. The first part of the study evaluated the safety and tolerability of a single dose of VIR-2218 at six dosing levels administered to healthy

Zaisbideng® WJH | https://www.wjgnet.com

adult volunteers. The second part of the study evaluated the safety and therapeutic effect across various increase doses of VIR-2218 given 4 wk apart. Study participants were non-cirrhotic adults with chronic HBV infection on NA therapy for at least 6 months and HBV DNA < 90 IU/mL. In both parts of the study, VIR-2218 was well-tolerated across all dose levels with only mild adverse events, commonly headache, injection site reactions, and mild ALT elevations. The study found a dose-dependent reduction in HBsAg in all VIR-2218 treatment groups compared to placebo by the 48-wk follow-up. A total of 12/24 participants across the VIR-2218 cohorts as compared to none in the placebo group achieved a reduction of HBsAg levels to < 100 IU/mL. The greatest mean reduction of HBsAg (-1.65 log IU/mL) occurred at week 20 for those receiving the 200 mg VIR-2218. While no participants had serum HBsAg loss or anti-HBs seroconversion by week 48, an HBsAg level < 100 IU/mL has been associated with a significantly higher chance of HBsAg loss[42]. This study demonstrated that VIR-2218 is well-tolerated with antiviral effects that could potentially lead to functional cure[43].

Another phase II trial investigated the safety and efficacy of VIR-2218 alone and in combination with PEGIFNα in noncirrhotic participants with chronic HBV infection. Inclusion criteria included NA therapy for at least 2 months, HBsAg > 50 IU/mL, and HBV DNA < 90 IU/mL. Preliminary data revealed that VIR-2218 was generally well-tolerated both alone and in combination with PEGIFNa. Adverse events that occurred were more consistent with known effects of PEGIFNa, such as mild ALT elevations and reductions in neutrophil and platelet levels. Four of 13 study participants treated with VIR-2218 combined with PEGIFNa for a longer duration of 48 wk achieved HBsAg seroclearance and anti-HBs seroconversion. Patients in this longer duration combination group also had the largest mean HBsAg reduction of 2.9 log IU/mL at the end of therapy. While the study is still ongoing with longer follow-up time, the preliminary results demonstrate that the antiviral effect of VIR-2218 may be potentiated by PEGIFNa and they show promise as a future combination therapy[44].

VIR-3434: VIR-3434 is a subcutaneously administered monoclonal antibody that targets an antigenic loop of HBsAg to block HBV cell entry[45]. In addition to clearing HBsAg, it can stimulate T cells for a vaccinal effect.

The MARCH trial was a phase II study that evaluated the safety, tolerability, and antiviral activity of VIR-2218 and VIR-3434 either as monotherapy or as combination therapy. Study participants were virally-suppressed, non-cirrhotic adults on NA therapy with chronic HBV infection. Both VIR-2218 and VIR-3434 were well-tolerated with mild adverse effects. The study was instrumental in showing that the combination of VIR-2218 and VIR-3434 Led to a marked mean HBsAg decline of > 2.5 log IU/mL in all cohorts. In fact, most participants were able to achieve an HBsAg level < 10 IU/mL. VIR-3434 has an additive effect to VIR-2218 in achieving a greater HBsAg reduction compared to monotherapy; this is consistent with their established complimentary mechanisms of action on HBV replication[46].

JNJ-73753989 (JNJ-3989, formerly ARO-HBV): JNJ-3989 is composed of two siRNAs which target both the S gene and X gene of the HBV. Consequently, it impairs the production of HBV RNA transcripts which are essential for replication [47].

A phase IIa clinical trial assessed the safety and efficacy of JNJ-3989 both with and without JNJ-6379 in 84 recruited participants with chronic HBV infection who were treatment-naïve or on chronic NA-suppressive therapy. All participants received an NA throughout the study. JNJ-3989 was well-tolerated across all doses throughout the study period. By day 112, there was an HBsAg reduction of $\geq 1 \log IU/mL$ from baseline in 39/40 participants who received 100 to 400 mg of JNJ-3989 every 4 wk in combination with an NA daily. Also, 30/40 patients achieved HBsAg < 100 IU/mL by day 112. A dose-dependent relationship was seen with higher doses of JNJ-3989 achieving higher levels of HBsAg reductions. More frequent dosing intervals did not change the magnitude and rate of response compared to dosing of JNJ-3989 every 4 wk. All 12 patients in the triple combination of JNJ-3989, JNJ-6379, and NA therapy achieved a \geq 1 log IU/mL HBsAg reduction from baseline to the nadir. The HBsAg reduction was also durable - 15/19 participants maintained a \geq 1 log HBsAg reduction for nearly 336 d after their last JNJ-3989 dose. This trial provided support that JNJ-3989 can be used safely in combination with an NA and that JNJ-6379 is an efficacious and durable HBV therapy[48].

The REEF-1 study was a large multicenter, double-blinded, randomized, phase IIb clinical trial studying the efficacy and safety of combination therapies of JNJ6379, JNJ-3989, and NA at various doses. The study featured non-cirrhotic adults with chronic HBV infection who were either treatment-naïve or virologically suppressed on NA therapy. The primary endpoint was the proportion of patients who met NA stopping criteria, as defined as ALT < 3 × upper limit of normal, HBV DNA less than the lower limit of quantitation, HBeAg negativity, and HBsAg < 10 IU/mL by week 48. Over the course of 48 wk, JNJ-3989 in combination with NA therapy led to a robust, dose-dependent response for meeting NA stopping criteria as well as reducing HBsAg and HBV RNA levels. In fact, 94/96 patients in the combination JNJ3989 200 mg every 4 wk and NA group had a \geq 1 log IU/mL HBsAg decline with a mean decline of 2.6 log IU/mL. Most patients did not reach the NA stopping criteria for two reasons: Failure to achieve HBsAg < 10 IU for those who were HBeAgnegative at baseline, or not achieving HBeAg seronegative status for patients who were HBeAg-positive at baseline. JNJ-3989 in combination with NA was safe and well-tolerated. Overall, REEF-1 showed that the combination of novel therapies, involving JNJ-3989 and/or JNJ6379, with established NA therapies is insufficient to achieve functional cure, but can achieve substantial HBsAg reductions[49].

Arbutus-729 (AB-729 or Imdusiran): AB-729 is a subcutaneously administered, GalNAc-conjugated RNA interference agent that blocks all RNA transcripts and reduces all HBV viral antigens [50]. It has an immunostimulatory component by enhancing HBV-specific T cell responses following repeat dosing[51].

In the AB-729-001 phase II study, healthy subjects and those with chronic HBV infection were subjected to single and repeat doses of AB-729 at various doses (60 or 90 mg of AB-729) and frequencies (every 4, 8, or 12 wk). ABI-729 with repeat dosing was found to be safe and well-tolerated. The most frequent adverse events included injection site events and asymptomatic ALT elevations which were Grade 2 in severity or lower. There was a robust and persistent decline in HBsAg in most subjects across cohorts regardless of dose, dosing interval, or HBeAg status; there was a mean reduction of HBsAg by 1.5 log IU/mL from baseline to 24 wk after the last dose. In fact, 26/34 participants achieved HBsAg < 100

IU/mL at some point in the study. As well, there was a sustained reduction in HBsAg and HBV DNA in 7 of 9 patients even after discontinuation of both AB-729 and NA-therapy. Only one subject seroconverted at week 48. These study findings demonstrated that AB-729 may be considered as a potential therapy for achieving functional cure of chronic HBV infection[52].

The AB-729-201 trial was a randomized, open-label, multicenter, phase IIa study which evaluated the safety, tolerability, and antiviral activity of AB-729 with PEGIFNα. The 43 non-cirrhotic, HBeAg-negative subjects had virally suppressed chronic HBV infection and were on stable NA therapy for at least 12 months prior. Patients received 4 doses of ABI-729 60 mg every 8 wk, and at week 24 were randomized to either of two treatment combinations (AB-729 + NA + PEGIFNα or NA + IFN) and at two treatment durations (12 wk vs 14 wk) followed by another 24 wk of follow-up where patients were evaluated to stop NA therapy. Preliminary results showed that by week 24 of treatment, there was a mean HBsAg decline of 1.6 log IU/mL across all cohorts. As well, 38 of 41 subjects achieved HBsAg levels < 100 IU/mL at some point during the treatment period. The interim data also showed that AB-729 with and without IFN was safe and welltolerated with most treatment related adverse events unrelated to AB-729 therapy [53].

HBsAg inhibition

HBsAg is a main surface protein on the envelope of the new HBV virion and subviral particles that maintains chronic infection via immune exhaustion[54]. HBsAg loss is one primary component required for functional cure[55]. HBsAg inhibitors disrupt the secretory processes involved in translocating HBsAg to the surface and effectively decrease HBsAg availability[56].

REP 2139/REP 2165 (Replicor): REP2139 is a nucleic acid polymer (NAP) that stops the assembly of subviral particles in hepatocytes and blocks the release of HBsAg[57]. REP2165 is a biologically equivalent variant of REP2139 with equivalent HBV antiviral activity in vivo. However, it has accelerated clearance which may be useful in cases requiring high frequency dosing for patients with slow rates of HBsAg clearance[58].

REP401 was an open-label phase 2 study evaluating the safety and efficacy of the combination therapy TDF, PEGIFN α , and either REP2139 or REP2165. Participants had chronic HBV infection and were HBeAg-negative. Patients received 24 wk of TDF therapy, followed by 24 wk of a control backbone therapy (TDF and PEGIFN α) or combination triple therapy (TDF, PEGIFNα, and either REP2139 or REP2165). Then participants were monitored for a treatment-free period of 48 wk. The addition of either REP2139 or REP2165 to TDF and PEGIFNα was safe and well-tolerated. Use of REP2139/REP2165 did not affect PEGIFNα-induced thrombocytopenia and neutropenia. Notably, there was a significantly more frequent and greater increase in asymptomatic transaminase levels among patients receiving an NAP which correlated with an initial decrease in HBsAg levels. From weeks 25 to 48, the combination triple therapy led to a rapid 4 to 6 log IU/mL decline in HBsAg in 15/20 patients by week 35. By week 48, HBsAg was not detected in 10 of 20 patients and HBsAg seroconversion was achieved in 11/20 patients, all with HBsAg < 1 log IU/mL. In contrast, the control backbone therapy group had an HBsAg decline > 1 log IU/mL in only 3 of 20 patients with no HBsAg seroconversion observed. Both the triple combination group and control group achieved a similar HBV DNA decline with 18 of 40 participants achieving HBV DNA less than the lower limit of quantification by week 48. In the 48-wk follow period, functional cure persisted in 14 of the 40 patients. Within the triple combination therapy group, there was no difference in response between REP2139 and REP2165 with regards to HBsAg, hepatitis B surface antibody (anti-HBs), and HBV DNA levels. REP401 showed that the addition of REP2139 or REP2165 to TDF and PEGIFNα therapy did not affect tolerability and increased rate of functional cure both during and after therapy[59].

REP301 was an open-label, nonrandomized, phase II trial investigating the use of REP2139 with PEGIFNa-2a in adults with chronic HBV infection. These participants were HBeAg-positive, anti-hepatitis D antigen-positive, and HDV RNApositive, and had an HBsAg levels > 1000 IU/mL. Study subjects received intravenous (IV) REP2139 once weekly for 15 wk, followed by a combination of IV REP2139 and subcutaneous PEGIFNα-2a once weekly for another 15 wk, and then finally, PEGIFNα-2a for 33 wk. By the end of treatment, 6/12 subjects had HBsAg < 50 IU/mL, 6/12 subjects had HBsAb > 10 mIU/mL, and 9/12 subjects had suppressed HBV DNA < 10 IU/mL. The response was durable to 1 year of followup: 5/6 patients maintained HBsAg suppression < 50 IU/mL, all 6/6 patients maintained HbsAb > 10 mIU/mL, and 7/9 patients had HBV DNA < 10 IU/mL. Use of both REP2139 and PEGIFNα-2a was safe and well-tolerated. The most frequent adverse events with REP2139 monotherapy were pyrexia and chills, while the introduction of PEGIFNα-2a led to asymptomatic transient elevations in ALT and aspartate aminotransferase (AST). REP301 underscored that combination REP2139 therapy with PEGIFNα-2a has robust and durable HBV and HDV antiviral effects even after completion of therapy[60].

Toll-like receptor agonists

Toll-like receptor (TLR) agonists act as immunomodulators to enhance the immune response against chronic HBV infection[61]. They induce the production of interferons, cytokines, and chemokines which upregulate antiviral effects [62].

Vesatolimod (GS-9620): Vesatolimod selectively activates TLR-7 found in gut-associated plasmacytoid dendritic cells and B lymphocytes to upregulate T and B cell responses[63].

The first phase II double-blind, randomized, placebo-controlled study evaluated the safety, efficacy, and pharmacodynamics of Vesatolimod in virally-suppressed, non-cirrhotic patients with chronic HBV infection. The 162 participants were randomized to receive weekly dosed placebo or Vesatolimod (1 mg, 2 mg, or 4 mg) for various treatment durations (4, 8, 12, and 48 wk). Vesatolimod was safe and well-tolerated at all doses with no clinically significant adverse events or lab derangements in the cohorts. Although the biological activity of Vesatolimod was verified with a dose-dependent

pharmacodynamic induction of the biomarker ISG15, no significant HBsAg decline from baseline was observed in any of the cohorts by week 48[64].

The second phase II study evaluated the safety and efficacy of Vesatolimod on patients with non-cirrhotic, chronic HBV infection who were not on oral antiviral treatment for at least 3 months. Additionally, patients had HBV DNA \geq 2000 IU/mL. In this multicenter, double blind, randomized, placebo-controlled study, patients were randomized to receive weekly placebo or oral Vesatolimod (1 mg, 2 mg, or 4 mg) for 12 wk. All subjects also received TDF of 300 mg daily for 48 wk. Vesatolimod was safe and well-tolerated. None of the patients achieved HBsAg loss or HBsAb seroconversion in any of the cohorts, and there was no significant difference in the decline of HBsAg among the Vesatolimod treatment groups compared to placebo. Only three total patients in the Vesatolimod groups had HBeAg loss and HBeAb seroconversion at week 48. There was no significant difference in HBV DNA decline among the Vesatolimod groups compared to placebo. Like the first study, a pharmacodynamic response was verified with a consistent dose-dependent induction of ISG15 biomarker level[65].

Selgantolimod (GS-9688): Selgantolimod is a selective TLR-8 agonist with antiviral activity against chronic HBV infection. It leads to the production of proinflammatory cytokines, chemokines, and interferons that initiate an innate and adaptive immune response against HBV[66].

A phase II, randomized, double-blind, placebo-controlled, multicenter study investigated the safety and efficacy of Selgantolimod in virally suppressed individuals on antiviral therapy with chronic HBV infection. Patients were randomized to receive once weekly placebo or oral Selgantolimod dosed at 1.5 mg or 3 mg for a total of 24 wk while continuing oral NA agents. Only one of the 48 participants in the 1.5 mg Selgantolimod group achieved the primary endpoint of $a \ge 1 \log IU/mL$ decline in HBsAg from baseline to week 24. As compared to placebo where no participants achieved HBsAg or HBeAg loss, 2 of the 39 subjects with HBeAg negative status achieved HBsAg loss and 3 of the 39 subjects had HBeAg loss in the Selgantolimod groups. The largest HBsAg reductions during the study occurred in patients who received Selgantolimod. In fact, HBsAg declines persisted even after treatment cessation. Selgantolimod was safe and generally well-tolerated with the most common adverse events including nausea, vomiting, and headache[67].

Therapeutic vaccinations

Therapeutic vaccinations present HBV vaccine antigens in a non-infective form to antigen presenting cells to stimulate a CD4 and CD8-mediated T cell response against HBV[68]. In comparison to preventative vaccines, therapeutic vaccinations are given during ongoing infection.

GS-4774: GS-4774 is a vaccine composed of heat-inactivated yeast cells expressing HBsAg, hepatitis B core antigen, and HBV-encoded oncogene X protein as a single fusion protein. Inoculation of individuals with GS-4774 as a subcutaneous injection elicits a significant T cell response^[69]

A phase II study evaluated the safety, tolerability, and efficacy of GS-4774 in non-cirrhotic patients with chronic HBV infection who were virally suppressed with oral antiviral therapy for at least a year. Subjects were randomized to receive either oral antivirals alone or a combination therapy of oral antivirals plus GS-4772 (dosed as 2, 10, or 40 yeast units) subcutaneously every 4 wk until week 20. Subjects continued oral antivirals for the remainder of the study to week 24 and then followed to week 48. No significant difference in mean HBsAg decline was found from baseline to week 24 or week 48 between any of the GS-4772 combinations therapy groups compared to oral antivirals alone. No patient experienced loss of HBsAg. Combination therapy of GS-4774 and antivirals was found to be safe and well-tolerated - there was no virologic breakthrough or treatment discontinuations in any patient and injection site reactions were the most common adverse event. The study showed that GS-4774 has limited efficacy for functional cure of chronic HBV infection among virally suppressed patients[70].

Another phase II, open-label, multicenter study evaluated the safety and efficacy of GS-4774 in combination with TDF in patients who were treatment-naïve. Inclusion criteria included positive HBsAg serology for at least 6 months, HBV DNA levels ≥ 2000 IU/mL, and no use of antiviral therapy within 3 months of study screening. Subjects were randomized to receive oral TDF alone or in combination with GS-4774 (dosed 2, 10, or 40 yeast units) every 4 wk until week 20. GS-4774 was safe and well-tolerated. There was no significant decrease in levels of HBsAg from baseline to weeks 24 and 48 among treatment groups despite a strong immune stimulatory effect on CD8+ T cells[71].

BRII-179: BRII-179 is a virus-like therapeutic vaccine expressing Pre-S1, Pre-S2, and S HBV surface antigens which stimulates an HBV specific T and B cell-mediated response[72].

In a randomized, open-label phase Ib/IIa study, the safety, antiviral activity, and immunogenicity of subcutaneouslyadministered BRII-179 at 20 mcg and 40 mcg doses with and without PEGIFN α was evaluated in subjects with noncirrhotic, chronic HBV infection. Subjects did not have detectable levels of HBsAg and were on NA for at least 6 months prior to the study. Results showed that both doses of BRII-179 were safe and well-tolerated with no severe adverse events. Limited HBsAg reductions (< 0.2 log HBsAg IU/mL) from baseline were observed after 4 doses of BRI-179 in both dosing groups. BRII-179 was found to be immunogenic: All BRII-179 treatment groups had increased HBsAb levels by at least > 30%, as compared to NA therapy alone which elicited no detectable anti-HBs response[72].

BRII-179 was also studied in combination with VIR-2218 for treating chronic HBV infection. An ongoing phase II study with interim results compared the combination of BRII-179 and VIR-2218 to VIR-2218 alone. Subjects were virally suppressed on an NA for at least 12 months and had HBV DNA less than the lower limit of quantification. Patients were followed to week 40. Interim results showed that BRII-179 in combination with VIR-2218 was safe and well-tolerated with mild adverse events, most commonly an injection site reaction. Although no significant difference in mean HBsAg reduction from baseline was found between combination therapy and VIR-2218 alone, the combination of BRII-179 and VIR-2218 led to a potent increase in anti-HBs level of more than 100 IU/L in more than 40% of the subjects compared to



none in the VIR-2218 alone. Final results will evaluate the long-term therapeutic and immune response to BRII-179 and VIR-2218 combination therapies^[73].

Anti-programmed cell death ligand-1

In chronic HBV infection, there is upregulation of programmed cell death ligand-1 (PD-L1) which is responsible for T-cell exhaustion and persistence of HBV viral disease [74]. The goal of checkpoint inhibitor therapy that blocks PD-L1 is to restore the function of HBV-specific T cells[75].

ASC22 (Envafolimab): ASC22 is a subcutaneously administered immunotherapy that blocks the programmed cell death protein 1 (PD-1)/PD-L1 pathway to restore T cell function. A phase IIb, randomized, single-blind, multicenter clinical trial was conducted to assess the efficacy and safety of ASC22 in subjects with chronic HBV infection who were virally suppressed on NA. Included subjects had HBsAg < 10000 IU/mL, HBV DNA < 20 IU/mL, and ALT/AST less than 2 × upper limit of normal, and were HBeAg-negative. Subjects were randomized to recieve either 1 mg/kg or 2 mg/kg subcutaneously-administered ASC22 every 2 wk in combination with an NA for 24 wk or placebo with NA. Both groups then received an additional 24 wk of NA therapy. Interim results of the combination therapy group with 1 mg/kg ASC22 and NA showed a more significant HBsAg reduction as compared to placebo and NA therapy, especially among patients with a baseline HbsAg level ≤ 100 IU/mL. This response was durably sustained - 3 of the 7 patients with baseline HBsAg ≤ 100 IU/mL in the ASC22 treatment group was able to sustain an HBsAg loss lower than the lower limit of quantification (0.05 IU/mL) by the end of the follow-up period. ASC22 1 mg/kg combined with NA for up to 24 wk was also safe and well-tolerated. Low-grade ALT flares were observed in 10/48 patients from the ASC22 group compared to none in the placebo group; these ALT flares also tended to occur more frequently in patients with a more significant HBsAg reduction. Thus, ALT flares may be a marker to monitor treatment response [76,77].

CONCLUSION

Current antiviral therapy with PEGIFN α and NA have low rates of functional cure and have limitations with regards to adverse effects, adherence, resistance, long-term treatment, and ongoing risk for liver events. Innovative clinical trials have been key in the development of novel therapies with a diverse range of mechanisms that strive to achieve the goal of functional cure (sustained HBsAg loss and undetectable HBV DNA 24 wk post-treatment). Based on available phases 2 and 3 data, it appears that single agent approaches (e.g., RNAi alone) are unlikely to result in HBsAg loss and therefore agents combining HBsAg lowering antivirals (e.g., RNAi and monoclonal antibody) +/- immunomodulator +/- NA may be required. Combination regimens with two drug (RNAi plus NA with bepirovirsen) or three drug approaches (RNAi plus immunomodulator plus NA with VIR-2218/ PEGIFN α /NA) have demonstrated proof of principle that functional cure can be achieved. Future randomized controlled trials in larger representative cohorts (HBeAg-positive/negative, NA-naïve/experienced, low vs high HBsAg titer) are needed to further confirm the efficacy/safety profiles of functional curative regimens and predictors of virologic response.

FOOTNOTES

Author contributions: Lam R collected the data; Lam R and Lim JK wrote and revised the manuscript.

Conflict-of-interest statement: Robert Lam reports no conflict of interest; Joseph K Lim has received research funding (to Yale University) from Gilead, Intercept, Inventiva, Novo Nordisk, Pfizer, and Viking.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: https://creativecommons.org/Licenses/by-nc/4.0/

Country/Territory of origin: United States

ORCID number: Joseph K Lim 0000-0003-1037-5773.

S-Editor: Liu JH L-Editor: Wang TQ P-Editor: Zheng XM

REFERENCES

- Lim JK, Nguyen MH, Kim WR, Gish R, Perumalswami P, Jacobson IM. Prevalence of Chronic Hepatitis B Virus Infection in the United 1 States. Am J Gastroenterol 2020; 115: 1429-1438 [PMID: 32483003 DOI: 10.14309/ajg.00000000000651]
- 2 Cui F, Blach S, Manzengo Mingiedi C, Gonzalez MA, Sabry Alaama A, Mozalevskis A, Séguy N, Rewari BB, Chan PL, Le LV, Doherty M,



Luhmann N, Easterbrook P, Dirac M, de Martel C, Nayagam S, Hallett TB, Vickerman P, Razavi H, Lesi O, Low-Beer D. Global reporting of progress towards elimination of hepatitis B and hepatitis C. Lancet Gastroenterol Hepatol 2023; 8: 332-342 [PMID: 36764320 DOI: 10.1016/S2468-1253(22)00386-7

- Lok AS, McMahon BJ. Chronic hepatitis B: update 2009. Hepatology 2009; 50: 661-662 [PMID: 19714720 DOI: 10.1002/hep.23190] 3
- McMahon BJ. The natural history of chronic hepatitis B virus infection. Hepatology 2009; 49: S45-S55 [PMID: 19399792 DOI: 4 10.1002/hep.22898]
- Polaris Observatory Collaborators. Global prevalence, treatment, and prevention of hepatitis B virus infection in 2016: a modelling study. 5 Lancet Gastroenterol Hepatol 2018; 3: 383-403 [PMID: 29599078 DOI: 10.1016/S2468-1253(18)30056-6]
- Chu CM, Liaw YF. Hepatitis B surface antigen seroclearance during chronic HBV infection. Antivir Ther 2010; 15: 133-143 [PMID: 6 20386068 DOI: 10.3851/IMP1497]
- 7 Arase Y, Ikeda K, Suzuki F, Suzuki Y, Saitoh S, Kobayashi M, Akuta N, Someya T, Hosaka T, Sezaki H, Kumada H. Long-term outcome after hepatitis B surface antigen seroclearance in patients with chronic hepatitis B. Am J Med 2006; 119: 71.e9-71.16 [PMID: 16431195 DOI: 10.1016/j.amjmed.2005.02.033]
- Liu J, Yang HI, Lee MH, Lu SN, Jen CL, Batrla-Utermann R, Wang LY, You SL, Hsiao CK, Chen PJ, Chen CJ; R. E.V.E.A.L.-HBV Study 8 Group. Spontaneous seroclearance of hepatitis B seromarkers and subsequent risk of hepatocellular carcinoma. Gut 2014; 63: 1648-1657 [PMID: 24225939 DOI: 10.1136/gutjnl-2013-305785]
- Moucari R, Marcellin P. [HBsAg seroclearance: prognostic value for the response to treatment and the long-term outcome]. Gastroenterol 9 *Clin Biol* 2010; **34** Suppl 2: S119-S125 [PMID: 21095515 DOI: 10.1016/S0399-8320(10)70031-2]
- 10 Yuen MF, Wong DK, Fung J, Ip P, But D, Hung I, Lau K, Yuen JC, Lai CL. HBsAg Seroclearance in chronic hepatitis B in Asian patients: replicative level and risk of hepatocellular carcinoma. Gastroenterology 2008; 135: 1192-1199 [PMID: 18722377 DOI: 10.1053/j.gastro.2008.07.008]
- Yuen MF, Wong DK, Sablon E, Tse E, Ng IO, Yuan HJ, Siu CW, Sander TJ, Bourne EJ, Hall JG, Condreay LD, Lai CL. HBsAg 11 seroclearance in chronic hepatitis B in the Chinese: virological, histological, and clinical aspects. Hepatology 2004; 39: 1694-1701 [PMID: 15185311 DOI: 10.1002/hep.20240]
- Zhang W, Zhang D, Dou X, Xie Q, Jiang J, Chen X, Ren H. Consensus on Pegylated Interferon Alpha in Treatment of Chronic Hepatitis B. J 12 Clin Transl Hepatol 2018; 6: 1-10 [PMID: 29577026 DOI: 10.14218/JCTH.2017.00073]
- Kim SS, Cheong JY, Cho SW. Current Nucleos(t)ide Analogue Therapy for Chronic Hepatitis B. Gut Liver 2011; 5: 278-287 [PMID: 13 21927654 DOI: 10.5009/gnl.2011.5.3.278]
- Craxi A, Cooksley WG. Pegylated interferons for chronic hepatitis B. Antiviral Res 2003; 60: 87-89 [PMID: 14638403 DOI: 14 10.1016/j.antiviral.2003.08.015]
- 15 Yang JF, Kao YH, Dai CY, Huang JF, Hsieh MY, Lin ZY, Chen SC, Wang LY, Chuang WL, Yu ML. Comparison of adverse effects related to pegylated interferon-based therapy for patients with chronic hepatitis B and chronic hepatitis C in Taiwan. Hepatol Int 2010; 4: 732-740 [PMID: 21286344 DOI: 10.1007/s12072-010-9208-0]
- Ryu SH, Chung YH. [Resistance to adefovir in patients with chronic hepatitis B]. Korean J Hepatol 2006; 12: 484-492 [PMID: 17237626] 16
- 17 Lam YF, Yuen MF, Seto WK, Lai CL. Current Antiviral Therapy of Chronic Hepatitis B: Efficacy and Safety. Curr Hepat Rep 2011; 10: 235-243 [PMID: 22131901 DOI: 10.1007/s11901-011-0109-z]
- Choi MS, Yoo BC. Management of chronic hepatitis B with nucleoside or nucleotide analogues: a review of current guidelines. Gut Liver 18 2010; 4: 15-24 [PMID: 20479908 DOI: 10.5009/gnl.2010.4.1.15]
- 19 Yang HC, Kao JH. Persistence of hepatitis B virus covalently closed circular DNA in hepatocytes: molecular mechanisms and clinical significance. Emerg Microbes Infect 2014; 3: e64 [PMID: 26038757 DOI: 10.1038/emi.2014.64]
- Martinez MG, Villeret F, Testoni B, Zoulim F. Can we cure hepatitis B virus with novel direct-acting antivirals? Liver Int 2020; 40 Suppl 1: 20 27-34 [PMID: 32077597 DOI: 10.1111/liv.14364]
- Flink HJ, van Zonneveld M, Hansen BE, de Man RA, Schalm SW, Janssen HL; HBV 99-01 Study Group. Treatment with Peg-interferon 21 alpha-2b for HBeAg-positive chronic hepatitis B: HBsAg loss is associated with HBV genotype. Am J Gastroenterol 2006; 101: 297-303 [PMID: 16454834 DOI: 10.1111/j.1572-0241.2006.00418.x]
- Chang TT, Lai CL, Kew Yoon S, Lee SS, Coelho HS, Carrilho FJ, Poordad F, Halota W, Horsmans Y, Tsai N, Zhang H, Tenney DJ, Tamez 22 R, Iloeje U. Entecavir treatment for up to 5 years in patients with hepatitis B e antigen-positive chronic hepatitis B. Hepatology 2010; 51: 422-430 [PMID: 20049753 DOI: 10.1002/hep.23327]
- Hadziyannis SJ, Tassopoulos NC, Heathcote EJ, Chang TT, Kitis G, Rizzetto M, Marcellin P, Lim SG, Goodman Z, Ma J, Brosgart CL, 23 Borroto-Esoda K, Arterburn S, Chuck SL; Adefovir Dipivoxil 438 Study Group. Long-term therapy with adefovir dipivoxil for HBeAgnegative chronic hepatitis B for up to 5 years. Gastroenterology 2006; 131: 1743-1751 [PMID: 17087951 DOI: 10.1053/j.gastro.2006.09.020]
- National Institutes of Health. U.S. Government, 2023. Available from: https://clinicaltrials.gov 24
- Fanning GC, Zoulim F, Hou J, Bertoletti A. Therapeutic strategies for hepatitis B virus infection: towards a cure. Nat Rev Drug Discov 2019; 25 18: 827-844 [PMID: 31455905 DOI: 10.1038/s41573-019-0037-0]
- Berke JM, Dehertogh P, Vergauwen K, Van Damme E, Mostmans W, Vandyck K, Pauwels F. Capsid Assembly Modulators Have a Dual 26 Mechanism of Action in Primary Human Hepatocytes Infected with Hepatitis B Virus. Antimicrob Agents Chemother 2017; 61 [PMID: 28584155 DOI: 10.1128/AAC.00560-17]
- Sulkowski MS, Agarwal K, Ma X, Nguyen TT, Schiff ER, Hann HL, Dieterich DT, Nahass RG, Park JS, Chan S, Han SB, Gane EJ, Bennett 27 M, Alves K, Evanchik M, Yan R, Huang Q, Lopatin U, Colonno R, Ma J, Knox SJ, Stamm LM, Bonacini M, Jacobson IM, Ayoub WS, Weilert F, Ravendhran N, Ramji A, Kwo PY, Elkhashab M, Hassanein T, Bae HS, Lalezari JP, Fung SK, Yuen MF. Safety and efficacy of vebicorvir administered with entecavir in treatment-naïve patients with chronic hepatitis B virus infection. J Hepatol 2022; 77: 1265-1275 [PMID: 35697332 DOI: 10.1016/j.jhep.2022.05.027]
- Yuen MF, Agarwal K, Ma X, Nguyen TT, Schiff ER, Hann HL, Dieterich DT, Nahass RG, Park JS, Chan S, Han SB, Gane EJ, Bennett M, 28 Alves K, Evanchik M, Yan R, Huang Q, Lopatin U, Colonno R, Ma J, Knox SJ, Stamm LM, Bonacini M, Jacobson IM, Ayoub WS, Weilert F, Ravendhran N, Ramji A, Kwo PY, Elkhashab M, Hassanein T, Bae HS, Lalezari JP, Fung SK, Sulkowski MS. Safety and efficacy of vebicorvir in virologically suppressed patients with chronic hepatitis B virus infection. J Hepatol 2022; 77: 642-652 [PMID: 35460726 DOI: 10.1016/j.jhep.2022.04.005
- 29 Yardeni D, Koh C. Bulevirtide for HBV and HDV infections. Drugs Today (Barc) 2021; 57: 433-448 [PMID: 34268531 DOI: 10.1358/dot.2021.57.7.3283861]



- Ferenci P, Reiberger T, Jachs M. Treatment of Chronic Hepatitis D with Bulevirtide-A Fight against Two Foes-An Update. Cells 2022; 11 30 [PMID: 36428959 DOI: 10.3390/cells11223531]
- Deterding K, Xu C, Port K, Dietz-Fricke C, Xun J, Maasoumy B, Cornberg M, Wedemeyer H. Bile acid increase during bulevirtide treatment 31 of hepatitis D is not associated with a decline in HDV RNA. J Viral Hepat 2023; 30: 597-606 [PMID: 36924318 DOI: 10.1111/jvh.13831]
- Bogomolov P, Alexandrov A, Voronkova N, Macievich M, Kokina K, Petrachenkova M, Lehr T, Lempp FA, Wedemeyer H, Haag M, Schwab 32 M, Haefeli WE, Blank A, Urban S. Treatment of chronic hepatitis D with the entry inhibitor myrcludex B: First results of a phase Ib/IIa study. J Hepatol 2016; 65: 490-498 [PMID: 27132170 DOI: 10.1016/j.jhep.2016.04.016]
- Wedemeyer H, Schöneweis K, Bogomolov P, Blank A, Voronkova N, Stepanova T, Sagalova O, Chulanov V, Osipenko M, Morozov V, 33 Geyvandova N, Sleptsova S, Bakulin IG, Khaertynova I, Rusanova M, Pathil A, Merle U, Bremer B, Allweiss L, Lempp FA, Port K, Haag M, Schwab M, Zur Wiesch JS, Cornberg M, Haefeli WE, Dandri M, Alexandrov A, Urban S. Safety and efficacy of bulevirtide in combination with tenofovir disoproxil fumarate in patients with hepatitis B virus and hepatitis D virus coinfection (MYR202): a multicentre, randomised, parallel-group, open-label, phase 2 trial. Lancet Infect Dis 2023; 23: 117-129 [PMID: 36113537 DOI: 10.1016/S1473-3099(22)00318-8]
- 34 Wedemeyer H, Schöneweis K, Bogomolov PO, Voronkova N, Chulanov VP, Stepanova T, Bremer B, Allweiss L, Dandri M, Burhenne J, Haefeli WE, Ciesek S, Dittmer U, Alexandrov A, Urban S. GS-13-Final results of a multicenter, open-label phase 2 clinical trial (MYR203) to assess safety and efficacy of myrcludex B in cwith PEG-interferon Alpha 2a in patients with chronic HBV/HDV co-infection. J Hepatol 2019 [DOI: 10.1016/S0618-8278(19)30141-0]
- Asselah T, Arama SS, Bogomolov P, Bourliere M, Fontaine H, Gherlan S, Gorodin V, Hilleret M-N, Lazar S, Mamonova N. Safety and 35 efficacy of bulevirtide monotherapy and in combination with peginterferon alfa-2a in patients with chronic hepatitis delta: 24 wk interim data of MYR204 phase 2b study. J Hepatol 2021; 75 (Suppl 2): S291
- Wedemeyer H, Aleman S, Brunetto M, Blank A, Andreone P, Bogomolov P, Chulanov V, Mamonova N, Geyvandova N, Viacheslav M, 36 Sagalova O, Stepanova T, Manuilov D, Suri V, An Q, Flaherty J, Osinusi A, Schulze zur Wiesch J, Cornberg M, Lampertico P. Efficacy and safety of bulevirtide monotherapy given at 2 mg or 10 mg dose level once daily for treatment of chronic hepatitis delta: week 48 primary end point results from a phase 3 randomized, multicenter, parallel design study. J Hepatol 2022; 77: S4-S5 [DOI: 10.1016/S0168-8278(22)00433-0]
- Mak LY, Seto WK, Yuen MF. Novel Antivirals in Clinical Development for Chronic Hepatitis B Infection. Viruses 2021; 13 [PMID: 37 34207458 DOI: 10.3390/v13061169]
- Mak LY, Hui RW, Fung J, Seto WK, Yuen MF. Bepirovirsen (GSK3228836) in chronic hepatitis B infection: an evaluation of phase II 38 progress. Expert Opin Investig Drugs 2023; 32: 971-983 [PMID: 37902953 DOI: 10.1080/13543784.2023.2277389]
- Yuen MF, Heo J, Jang JW, Yoon JH, Kweon YO, Park SJ, Tami Y, You S, Yates P, Tao Y, Cremer J, Campbell F, Elston R, Theodore D, Paff 39 M, Bennett CF, Kwoh TJ. Safety, tolerability and antiviral activity of the antisense oligonucleotide bepirovirsen in patients with chronic hepatitis B: a phase 2 randomized controlled trial. Nat Med 2021; 27: 1725-1734 [PMID: 34642494 DOI: 10.1038/s41591-021-01513-4]
- Yuen MF, Lim SG, Plesniak R, Tsuji K, Janssen HLA, Pojoga C, Gadano A, Popescu CP, Stepanova T, Asselah T, Diaconescu G, Yim HJ, 40 Heo J, Janczewska E, Wong A, Idriz N, Imamura M, Rizzardini G, Takaguchi K, Andreone P, Arbune M, Hou J, Park SJ, Vata A, Cremer J, Elston R, Lukić T, Quinn G, Maynard L, Kendrick S, Plein H, Campbell F, Paff M, Theodore D; B-Clear Study Group. Efficacy and Safety of Bepirovirsen in Chronic Hepatitis B Infection. N Engl J Med 2022; 387: 1957-1968 [PMID: 36346079 DOI: 10.1056/NEJMoa2210027]
- Gupta SV, Fanget MC, MacLauchlin C, Clausen VA, Li J, Cloutier D, Shen L, Robbie GJ, Mogalian E. Clinical and Preclinical Single-Dose 41 Pharmacokinetics of VIR-2218, an RNAi Therapeutic Targeting HBV Infection. Drugs R D 2021; 21: 455-465 [PMID: 34741731 DOI: 10.1007/s40268-021-00369-w]
- Tseng TC, Liu CJ, Su TH, Wang CC, Chen CL, Chen PJ, Chen DS, Kao JH. Serum hepatitis B surface antigen levels predict surface antigen 42 loss in hepatitis B e antigen seroconverters. Gastroenterology 2011; 141: 517-525, 525.e1 [PMID: 21672542 DOI: 10.1053/j.gastro.2011.04.046]
- Gane E, Lim YS, Kim JB, Jadhav V, Shen L, Bakardjiev AI, Huang SA, Cathcart AL, Lempp FA, Janas MM, Cloutier DJ, Kaittanis C, Sepp-43 Lorenzino L, Hinkle G, Taubel J, Haslett P, Milstein S, Anglero-Rodriguez YI, Hebner CM, Pang PS, Yuen MF. Evaluation of RNAi therapeutics VIR-2218 and ALN-HBV for chronic hepatitis B: Results from randomized clinical trials. J Hepatol 2023; 79: 924-932 [PMID: 37290591 DOI: 10.1016/j.jhep.2023.05.023]
- Gane E, Lim YS, Tangkijvanich P, O'Beirne J, Lim TH, Bakardjiev A, Ding X, Connolly L, Huang S, Kim J. Preliminary safety and antiviral 44 activity of VIR-2218, an X-targeting HBV RNAi therapeutic, in chronic hepatitis B patients. Proceedings of the Digital International Liver Congress (Digital ILC 2020; 2020. Elsevier BV [DOI: 10.1016/S0168-8278(20)30647-4]
- Lempp FA, Volz T, Cameroni E, Benigni F, Zhou J, Rosen LE, Noack J, Zatta F, Kaiser H, Bianchi S, Lombardo G, Jaconi S, Vincenzetti L, 45 Imam H, Soriaga LB, Passini N, Belnap DM, Schulze A, Lütgehetmann M, Telenti A, Cathcart AL, Snell G, Purcell LA, Hebner CM, Urban S, Dandri M, Corti D, Schmid MA. Potent broadly neutralizing antibody VIR-3434 controls hepatitis B and D virus infection and reduces HBsAg in humanized mice. J Hepatol 2023; 79: 1129-1138 [PMID: 37459920 DOI: 10.1016/j.jhep.2023.07.003]
- Gane E, Jucov A, Dobryanska M, Yoon KT, Lim TH, Arizpe A, Cloutier D, Shen L, Gupta SV, Lau AH. Safety, tolerability, and antiviral 46 activity of the siRNA VIR-2218 in combination with the investigational neutralizing monoclonal antibody VIR-3434 for the treatment of chronic hepatitis B virus infection: preliminary results from the phase 2 march trial. Proceedings of the Hepatology; 2022. WILEY 111 RIVER ST, HOBOKEN 07030-5774, NJ USA: S18-S19
- Li H, Niu X, Zhang Y, Zhang D, Wang L, Miao Y, Jiang Y, Ji J, Chen Q, Wu X, Ediage EN, Kakuda TN, Biermer M. Pharmacokinetics, 47 Safety, and Tolerability of the siRNA JNJ-73763989 in Healthy Chinese Adult Participants. Clin Pharmacol Drug Dev 2023; 12: 175-180 [PMID: 36415122 DOI: 10.1002/cpdd.1197]
- Yuen MF, Locarnini S, Lim TH, Strasser SI, Sievert W, Cheng W, Thompson AJ, Given BD, Schluep T, Hamilton J, Biermer M, Kalmeijer R, 48 Beumont M, Lenz O, De Ridder F, Cloherty G, Ka-Ho Wong D, Schwabe C, Jackson K, Lai CL, Gish RG, Gane E. Combination treatments including the small-interfering RNA JNJ-3989 induce rapid and sometimes prolonged viral responses in patients with CHB. J Hepatol 2022; 77: 1287-1298 [PMID: 35870702 DOI: 10.1016/j.jhep.2022.07.010]
- 49 Yuen MF, Asselah T, Jacobson IM, Brunetto MR, Janssen HLA, Takehara T, Hou JL, Kakuda TN, Lambrecht T, Beumont M, Kalmeijer R, Guinard-Azadian C, Mayer C, Jezorwski J, Verbinnen T, Lenz O, Shukla U, Biermer M; REEF-1 Study Group. Efficacy and safety of the siRNA JNJ-73763989 and the capsid assembly modulator JNJ-56136379 (bersacapavir) with nucleos(t)ide analogues for the treatment of chronic hepatitis B virus infection (REEF-1): a multicentre, double-blind, active-controlled, randomised, phase 2b trial. Lancet Gastroenterol Hepatol 2023; 8: 790-802 [PMID: 37442152 DOI: 10.1016/S2468-1253(23)00148-6]
- Hui RW, Mak LY, Seto WK, Yuen MF. RNA interference as a novel treatment strategy for chronic hepatitis B infection. Clin Mol Hepatol 50



2022; 28: 408-424 [PMID: 35172540 DOI: 10.3350/cmh.2022.0012]

- Paratala B, Park JJ, Ganchua SC, Gane E, Yuen R, Lee AC, Moore C, Lam AM, Sevinsky H, Sims K. Inhibition of hepatitis B surface 51 antigen in chronic hepatitis B subjects by RNA interference therapeutic AB-729 is accompanied by upregulation of HBV-specific T cell activation markers. Proceedings of the The International Liver Congress 2021 (ILC 2021); 2021. Elsevier BV
- 52 Yuen R, Berliba E, Sukeepaisarnjaroen W, Holmes J, Leerapun A, Tangkijvanich P, Strasser S, Jucov A, Gane E, Thi E. Long-term suppression maintained after cessation of AB-729 treatment and comparable on-treatment response observed in HBeAg+ subjects. Poster Presentation (SAT443). J Hepatol 2022 [DOI: 10.1016/S0168-8278(22)02045-1]
- Yuen M-F, Heo J, Nahass RG, Wong GL-H, Burda T, Bhamidimarri KR, Hu T-H, Nguyen T, Lim Y-S, Chen C-Y. LBP-38-Preliminary safety 53 and antiviral activity of AB-729 combination treatment with pegylated interferon alfa-2a in virally suppressed, HBeAg-negative subjects with chronic HBV infection. J Hepatol 2023; 78: S125-S125 [DOI: 10.1016/S0168-8278(23)00618-9]
- Carman WF. The clinical significance of surface antigen variants of hepatitis B virus. J Viral Hepat 1997; 4 Suppl 1: 11-20 [PMID: 9097273 54 DOI: 10.1111/j.1365-2893.1997.tb00155.x]
- 55 Cornberg M, Lok AS, Terrault NA, Zoulim F; 2019 EASL-AASLD HBV Treatment Endpoints Conference Faculty. Guidance for design and endpoints of clinical trials in chronic hepatitis B - Report from the 2019 EASL-AASLD HBV Treatment Endpoints Conference. Hepatology 2019 [PMID: 31713892 DOI: 10.1002/hep.31030]
- Baugh SDP. Inhibiting the Secretion of Hepatitis B Surface Antigen (HBsAg) to Treat Hepatitis B Infection- a Review. Infect Disord Drug 56 Targets 2017; 17: 24-35 [PMID: 28056752 DOI: 10.2174/1871526517666170104113730]
- Vaillant A. REP 2139: Antiviral Mechanisms and Applications in Achieving Functional Control of HBV and HDV Infection. ACS Infect Dis 57 2019; 5: 675-687 [PMID: 30199230 DOI: 10.1021/acsinfecdis.8b00156]
- 58 Vaillant A. Editorial: In vitro mechanistic evaluation of nucleic acid polymers: A cautionary tale. Mol Ther Nucleic Acids 2022; 28: 168-174 [PMID: 35402067 DOI: 10.1016/j.omtn.2022.03.002]
- 59 Bazinet M, Pântea V, Placinta G, Moscalu I, Cebotarescu V, Cojuhari L, Jimbei P, Iarovoi L, Smesnoi V, Musteata T, Jucov A, Dittmer U, Krawczyk A, Vaillant A. Safety and Efficacy of 48 Weeks REP 2139 or REP 2165, Tenofovir Disoproxil, and Pegylated Interferon Alfa-2a in Patients With Chronic HBV Infection Naïve to Nucleos(t)ide Therapy. Gastroenterology 2020; 158: 2180-2194 [PMID: 32147484 DOI: 10.1053/j.gastro.2020.02.058]
- Bazinet M, Pântea V, Cebotarescu V, Cojuhari L, Jimbei P, Albrecht J, Schmid P, Le Gal F, Gordien E, Krawczyk A, Mijočević H, 60 Karimzadeh H, Roggendorf M, Vaillant A. Safety and efficacy of REP 2139 and pegylated interferon alfa-2a for treatment-naive patients with chronic hepatitis B virus and hepatitis D virus co-infection (REP 301 and REP 301-LTF): a non-randomised, open-label, phase 2 trial. Lancet Gastroenterol Hepatol 2017; 2: 877-889 [PMID: 28964701 DOI: 10.1016/S2468-1253(17)30288-1]
- Ma Z, Cao Q, Xiong Y, Zhang E, Lu M. Interaction between Hepatitis B Virus and Toll-Like Receptors: Current Status and Potential 61 Therapeutic Use for Chronic Hepatitis B. Vaccines (Basel) 2018; 6 [PMID: 29337856 DOI: 10.3390/vaccines6010006]
- Isogawa M, Robek MD, Furuichi Y, Chisari FV. Toll-like receptor signaling inhibits hepatitis B virus replication in vivo. J Virol 2005; 79: 62 7269-7272 [PMID: 15890966 DOI: 10.1128/jvi.79.11.7269-7272.2005]
- Riddler SA, Para M, Benson CA, Mills A, Ramgopal M, DeJesus E, Brinson C, Cyktor J, Jacobs J, Koontz D, Mellors JW, Laird GM, Wrin T, 63 Patel H, Guo S, Wallin J, Boice J, Zhang L, Humeniuk R, Begley R, German P, Graham H, Geleziunas R, Brainard DM, SenGupta D. Vesatolimod, a Toll-like Receptor 7 Agonist, Induces Immune Activation in Virally Suppressed Adults Living With Human Immunodeficiency Virus-1. Clin Infect Dis 2021; 72: e815-e824 [PMID: 33043969 DOI: 10.1093/cid/ciaa1534]
- 64 Janssen HLA, Brunetto MR, Kim YJ, Ferrari C, Massetto B, Nguyen AH, Joshi A, Woo J, Lau AH, Gaggar A, Subramanian GM, Yoshida EM, Ahn SH, Tsai NCS, Fung S, Gane EJ. Safety, efficacy and pharmacodynamics of vesatolimod (GS-9620) in virally suppressed patients with chronic hepatitis B. J Hepatol 2018; 68: 431-440 [PMID: 29104121 DOI: 10.1016/j.jhep.2017.10.027]
- Agarwal K, Ahn SH, Elkhashab M, Lau AH, Gaggar A, Bulusu A, Tian X, Cathcart AL, Woo J, Subramanian GM, Andreone P, Kim HJ, 65 Chuang WL, Nguyen MH. Safety and efficacy of vesatolimod (GS-9620) in patients with chronic hepatitis B who are not currently on antiviral treatment. J Viral Hepat 2018; 25: 1331-1340 [PMID: 29851204 DOI: 10.1111/jvh.12942]
- Ayithan N, Ghosh A, Dwivedi A, Wallin JJ, Tan SK, Chen D, Kottilil S, Poonia B. Oral Selective TLR8 Agonist Selgantolimod Induces 66 Multiple Immune Cell Responses in Humans. Viruses 2021; 13 [PMID: 34960669 DOI: 10.3390/v13122400]
- 67 Gane EJ, Dunbar PR, Brooks AE, Zhang F, Chen D, Wallin JJ, van Buuren N, Arora P, Fletcher SP, Tan SK, Yang JC, Gaggar A, Kottilil S, Tang L. Safety and efficacy of the oral TLR8 agonist selgantolimod in individuals with chronic hepatitis B under viral suppression. J Hepatol 2023; 78: 513-523 [PMID: 38133554 DOI: 10.1016/j.jhep.2022.09.027]
- Cargill T, Barnes E. Therapeutic vaccination for treatment of chronic hepatitis B. Clin Exp Immunol 2021; 205: 106-118 [PMID: 33969474 68 DOI: 10.1111/cei.13614]
- Gaggar A, Coeshott C, Apelian D, Rodell T, Armstrong BR, Shen G, Subramanian GM, McHutchison JG. Safety, tolerability and 69 immunogenicity of GS-4774, a hepatitis B virus-specific therapeutic vaccine, in healthy subjects: a randomized study. Vaccine 2014; 32: 4925-4931 [PMID: 25045824 DOI: 10.1016/j.vaccine.2014.07.027]
- Lok AS, Pan CQ, Han SH, Trinh HN, Fessel WJ, Rodell T, Massetto B, Lin L, Gaggar A, Subramanian GM, McHutchison JG, Ferrari C, Lee 70 H, Gordon SC, Gane EJ. Randomized phase II study of GS-4774 as a therapeutic vaccine in virally suppressed patients with chronic hepatitis B. J Hepatol 2016; 65: 509-516 [PMID: 27210427 DOI: 10.1016/j.jhep.2016.05.016]
- Boni C, Janssen HLA, Rossi M, Yoon SK, Vecchi A, Barili V, Yoshida EM, Trinh H, Rodell TC, Laccabue D, Alfieri A, Brillo F, Fisicaro P, 71 Acerbi G, Pedrazzi G, Andreone P, Cursaro C, Margotti M, Santoro R, Piazzolla V, Brunetto MR, Coco B, Cavallone D, Zhao Y, Joshi A, Woo J, Lau AH, Gaggar A, Subramanian GM, Massetto B, Fung S, Ahn SH, Ma X, Mangia A, Ferrari C. Combined GS-4774 and Tenofovir Therapy Can Improve HBV-Specific T-Cell Responses in Patients With Chronic Hepatitis. Gastroenterology 2019; 157: 227-241.e7 [PMID: 30930022 DOI: 10.1053/j.gastro.2019.03.044]
- 72 Ma H, Lim TH, Leerapun A, Weltman M, Jia J, Lim YS, Tangkijvanich P, Sukeepaisarnjaroen W, Ji Y, Le Bert N, Li D, Zhang Y, Hamatake R, Tan N, Li C, Strasser SI, Ding H, Yoon JH, Stace NH, Ahmed T, Anderson DE, Yan L, Bertoletti A, Zhu Q, Yuen MF. Therapeutic vaccine BRII-179 restores HBV-specific immune responses in patients with chronic HBV in a phase Ib/IIa study. JHEP Rep 2021; 3: 100361 [PMID: 34661089 DOI: 10.1016/j.jhepr.2021.100361]
- Yuen MF, Wong GLH, Douglas M, Ma H, Zhu C, Ji Y, Liu W, Chen X, Zhu Q. Preliminary Safety and Efficacy of the Combination Therapy 73 of BRII-835 (VIR-2218) and BRII-179 (VBI-2601) Treating Chronic HBV Infection. Proceedings of the 32nd Annual Conference of the Asian Pacific Association for the Study of the Liver (APASL) 2023; Taipei. Available from: https://www.vbivaccines.com/wp-content/uploads/2023/



03/APASL-2023-Oral-Presentation.pdf

- Féray C, López-Labrador FX. Is PD-1 blockade a potential therapy for HBV? JHEP Rep 2019; 1: 142-144 [PMID: 32040093 DOI: 74 10.1016/j.jhepr.2019.07.007]
- Bertoletti A, Le Bert N. Immunotherapy for Chronic Hepatitis B Virus Infection. Gut Liver 2018; 12: 497-507 [PMID: 29316747 DOI: 75 10.5009/gnl17233]
- Wang G, Cui Y, Xie Y, Mao Q, Xie Q, Gu Y, Chen X, Hu G, Yang Y, He HW. Effects of Subcutaneous PD-L1 Antibody ASC22 76 (Envafolimab) Plus nucleos(t)ide analogs on HBsAg reduction in patients with chronic hepatitis B infection are correlated with pre-treatment HBsAg level. Proceedings of the Asian Pacific Association for the Study of the Liver Annual Meeting 2023; Taipei. Available from: https:// www.prnewswire.com/news-releases/subcutaneously-administered-pd-11-antibody-asc22-envafolimab-is-safe-and-well-tolerated-in-phase-iiahbv-study-301186235.html
- Wang G. ALT flares were linked to HBsAg reduction, seroclearance and seroconversion: interim results from a phase IIb study in chronic 77 hepatitis B patients with 24-week treatment of subcutaneous PD-L1 antibody ASC22 (Envafolimab) plus nucleos(t)ide analogs. Proceedings of the EASL International Liver Congress 2022 [DOI: 10.1016/S0168-8278(22)00538-4]





Published by Baishideng Publishing Group Inc 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA Telephone: +1-925-3991568 E-mail: office@baishideng.com Help Desk: https://www.f6publishing.com/helpdesk https://www.wjgnet.com

