

World Journal of *Hepatology*

World J Hepatol 2024 April 27; 16(4): 490-660



EDITORIAL

- 490 Erdafitinib and checkpoint inhibitors for first-line and second-line immunotherapy of hepatic, gastrointestinal, and urinary bladder carcinomas: Recent concept
Wishahi M
- 494 Progress of mitochondrial and endoplasmic reticulum-associated signaling and its regulation of chronic liver disease by Chinese medicine
Zheng Y, Zheng YH, Wang JH, Zhao TJ, Wang L, Liang TJ
- 506 Subclinical hepatitis E virus genotype 1 infection: The concept of “dynamic human reservoir”
Shrestha A, Basnet S, KC S
- 511 Metabolic dysfunction-associated steatotic liver disease: A silent pandemic
Samanta A, Sen Sarma M

REVIEW

- 517 Spectrum of COVID-19 induced liver injury: A review report
Singh L, Kumar A, Rai M, Basnet B, Rai N, Khanal P, Lai KS, Cheng WH, Asaad AM, Ansari S
- 537 Multifaceted roles of lymphatic and blood endothelial cells in the tumor microenvironment of hepatocellular carcinoma: A comprehensive review
Li JJ, Mao JX, Zhong HX, Zhao YY, Teng F, Lu XY, Zhu LY, Gao Y, Fu H, Guo WY
- 550 Quantitative hepatitis B core antibody and quantitative hepatitis B surface antigen: Novel viral biomarkers for chronic hepatitis B management
Leowattana W, Leowattana P, Leowattana T
- 566 Molecular mechanism of nanomaterials induced liver injury: A review
Das SK, Sen K, Ghosh B, Ghosh N, Sinha K, Sil PC

ORIGINAL ARTICLE**Case Control Study**

- 601 Expression and clinical significance of short-chain fatty acids in patients with intrahepatic cholestasis of pregnancy
Ren SJ, Feng JT, Xiang T, Liao CL, Zhou YP, Xuan RR

Retrospective Cohort Study

- 612 *Klebsiella pneumoniae* infections after liver transplantation: Drug resistance and distribution of pathogens, risk factors, and influence on outcomes
Guo L, Peng P, Peng WT, Zhao J, Wan QQ

Retrospective Study

- 625 Development and validation of a nomogram for predicting in-hospital mortality of intensive care unit patients with liver cirrhosis

Tang XW, Ren WS, Huang S, Zou K, Xu H, Shi XM, Zhang W, Shi L, Lü MH

Prospective Study

- 640 Prospective study of hepatitis B and D epidemiology and risk factors in Romania: A 10-year update

Iacob S, Gheorghe L, Onica M, Huiban L, Pop CS, Brisc C, Sirli R, Ester C, Brisc CM, Diaconu S, Rogoveanu I, Sandulescu L, Vuletic D, Trifan A

SYSTEMATIC REVIEWS

- 650 Relative carcinogenicity of tacrolimus vs mycophenolate after solid organ transplantation and its implications for liver transplant care

Liu D, Youssef MM, Grace JA, Sinclair M

ABOUT COVER

Peer Reviewer of *World Journal of Hepatology*, Somchai Amornytin, MD, Professor, Staff Physician, Department of Anesthesiology and Siriraj Gastrointestinal Endoscopy Center, Faculty of Medicine, Siriraj Hospital, Mahidol University, Bangkok 10700, Thailand. somchai.amo@mahidol.ac.th

AIMS AND SCOPE

The primary aim of *World Journal of Hepatology (WJH, World J Hepatol)* is to provide scholars and readers from various fields of hepatology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJH mainly publishes articles reporting research results and findings obtained in the field of hepatology and covering a wide range of topics including chronic cholestatic liver diseases, cirrhosis and its complications, clinical alcoholic liver disease, drug induced liver disease autoimmune, fatty liver disease, genetic and pediatric liver diseases, hepatocellular carcinoma, hepatic stellate cells and fibrosis, liver immunology, liver regeneration, hepatic surgery, liver transplantation, biliary tract pathophysiology, non-invasive markers of liver fibrosis, viral hepatitis.

INDEXING/ABSTRACTING

The *WJH* is now abstracted and indexed in PubMed, PubMed Central, Emerging Sources Citation Index (ESCI), Scopus, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for *WJH* as 2.4.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Yi-Xuan Cai, Production Department Director: Xiang Li, Cover Editor: Xiang Li.

NAME OF JOURNAL

World Journal of Hepatology

ISSN

ISSN 1948-5182 (online)

LAUNCH DATE

October 31, 2009

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Nikolaos Pylsopoulos, Ke-Qin Hu, Koo Jeong Kang

EXECUTIVE ASSOCIATE EDITORS-IN-CHIEF

Shuang-Suo Dang

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/1948-5182/editorialboard.htm>

PUBLICATION DATE

April 27, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

PUBLISHING PARTNER

Department of Infectious Diseases, the Second Affiliated Hospital of Xi'an Jiaotong University

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

POLICY OF CO-AUTHORS

<https://www.wjgnet.com/bpg/GerInfo/310>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

PUBLISHING PARTNER'S OFFICIAL WEBSITE

http://2yuan.xjtu.edu.cn/Html/Departments/Main/Index_21148.html

Relative carcinogenicity of tacrolimus vs mycophenolate after solid organ transplantation and its implications for liver transplant care

Dorothy Liu, Mark M Youssef, Josephine A Grace, Marie Sinclair

Specialty type: Transplantation

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0

Grade B (Very good): B

Grade C (Good): 0

Grade D (Fair): 0

Grade E (Poor): 0

P-Reviewer: Chauhan S, United States

Received: December 16, 2023

Peer-review started: December 16, 2023

First decision: January 17, 2024

Revised: January 30, 2024

Accepted: March 19, 2024

Article in press: March 19, 2024

Published online: April 27, 2024



Dorothy Liu, Josephine A Grace, Marie Sinclair, Department of Gastroenterology, Austin Health, Melbourne 3084, Victoria, Australia

Dorothy Liu, Marie Sinclair, Victorian Liver Transplant Unit, Austin Health, Melbourne 3084, Victoria, Australia

Dorothy Liu, Mark M Youssef, Josephine A Grace, Marie Sinclair, Department of Medicine, University of Melbourne, Melbourne 3084, Victoria, Australia

Corresponding author: Dorothy Liu, B-BMed, MD, FRACP, Doctor, Department of Gastroenterology, Austin Health, 145 Studley Road, Heidelberg, Melbourne 3084, Victoria, Australia. dorothy.liu2@austin.org.au

Abstract

BACKGROUND

De novo malignancy is a leading cause of late morbidity and mortality in liver transplant recipients. Cumulative immunosuppression has been shown to contribute to post-transplant malignancy (PTM) risk. There is emerging evidence on the differential carcinogenic risk profile of individual immunosuppressive drugs, independent of the net effect of immunosuppression. Calcineurin inhibitors such as tacrolimus may promote tumorigenesis, whereas mycophenolic acid (MPA), the active metabolite of mycophenolate mofetil, may limit tumour progression. Liver transplantation (LT) is relatively unique among solid organ transplantation in that immunosuppression monotherapy with either tacrolimus or MPA is often achievable, which makes careful consideration of the risk-benefit profile of these immunosuppression agents particularly relevant for this cohort. However, there is limited clinical data on this subject in both LT and other solid organ transplant recipients.

AIM

To investigate the relative carcinogenicity of tacrolimus and MPA in solid organ transplantation.

METHODS

A literature search was conducted using MEDLINE and Embase databases using the key terms "solid organ transplantation", "tacrolimus", "mycophenolic acid", and "carcinogenicity", in order to identify relevant articles published in English between 1st January 2002 to 11th August 2022. Related terms, synonyms and explosion of MeSH terms, Boolean operators and truncations were also utilised in

the search. Reference lists of retrieved articles were also reviewed to identify any additional articles. Excluding duplicates, abstracts from 1230 records were screened by a single reviewer, whereby 31 records were reviewed in detail. Full-text articles were assessed for eligibility based on pre-specified inclusion and exclusion criteria.

RESULTS

A total of 6 studies were included in this review. All studies were large population registries or cohort studies, which varied in transplant era, type of organ transplanted and immunosuppression protocol used. Overall, there was no clear difference demonstrated between tacrolimus and MPA in *de novo* PTM risk following solid organ transplantation. Furthermore, no study provided a direct comparison of carcinogenic risk between tacrolimus and MPA monotherapy in solid organ transplantation recipients.

CONCLUSION

The contrasting carcinogenic risk profiles of tacrolimus and MPA demonstrated in previous experimental studies, and its application in solid organ transplantation, is yet to be confirmed in clinical studies. Thus, the optimal choice of immunosuppression drug to use as maintenance monotherapy in LT recipients is not supported by a strong evidence base and remains unclear.

Key Words: Immunosuppression; Solid organ transplantation; Liver transplantation; Carcinogenicity; Tacrolimus; Mycophenolate

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Cumulative immunosuppression exposure is an important risk factor for the development of post-transplant malignancy. There is emerging evidence on the differential carcinogenic risk profile of individual immunosuppressive drugs, independent of the net immunosuppression effect. This review demonstrates that the evidence on the relative carcinogenicity of tacrolimus and mycophenolic acid, the two agents most commonly used as maintenance monotherapy in liver transplant patients, remains unclear. Further studies are required to determine the clinical relevance of previous experimental findings to enable physicians to tailor immunosuppression regimens to minimize individual malignancy risk in solid organ transplantation.

Citation: Liu D, Youssef MM, Grace JA, Sinclair M. Relative carcinogenicity of tacrolimus vs mycophenolate after solid organ transplantation and its implications for liver transplant care. *World J Hepatol* 2024; 16(4): 650-660

URL: <https://www.wjgnet.com/1948-5182/full/v16/i4/650.htm>

DOI: <https://dx.doi.org/10.4254/wjh.v16.i4.650>

INTRODUCTION

Liver transplantation (LT) remains the only curative treatment for end-stage liver disease and some cases of hepatocellular carcinoma, with an overall median survival of 20 years[1]. Despite improvements in short-term survival with the decline in rates of rejection and graft failure with the advent of modern immunosuppression regimens, long-term complications including post-transplant malignancy (PTM), have risen. Liver transplant recipients incur a 2- to 3-fold increase in rates of *de novo* malignancy compared to the general population[2,3]. Indeed, PTM has become a leading cause of late mortality in LT recipients[4,5].

The cumulative exposure to immunosuppression and direct carcinogenicity of individual agents may contribute to the development of PTM[6]. Tacrolimus and mycophenolic acid (MPA) are the most commonly used backbone immunosuppressants post-LT, and are also utilised as maintenance monotherapy in 42% of LT recipients in the United States due to the relatively immune tolerant microenvironment of the liver[7,8]. Experimental data have demonstrated multiple pro-oncogenic effects of tacrolimus, whereas MPA may be protective against tumour growth and progression[9]. This systematic review aims to compare the relative carcinogenicity of tacrolimus and MPA in solid organ transplantation to assist clinicians in making informed decisions regarding choice of immunosuppression regimens for patients.

PTM immunology

The development of PTM is a consequence of complex interactions between genetic, lifestyle and transplant factors (Figure 1). The central role of the immune system in cancer surveillance is highlighted by the increased malignancy risk that results from congenital and acquired immunodeficiencies, as well as the efficacy of immunotherapy for a growing number of malignancies such as hepatocellular carcinoma, melanoma, and renal cell carcinoma[6].

An intact immune system prevents oncogenesis through 3 main mechanisms. Firstly, the immune system eliminates or suppresses viral infections to prevent virus-induced tumours, as seen in the role of Epstein Barr virus infections in the development of early post-transplant lymphoproliferative disorder (PTLD)[6,10]. Secondly, inflammation resolution and

Post-transplant malignancy		
Genetic factors	Lifestyle factors	Transplant factors
Age Gender Ethnicity Family history of cancer Previous history of cancer	Smoking Alcohol Obesity Oncogenic viruses Sun exposure Radiation exposure	Organ type Cancer in explant Immunosuppression drug type, dose, and duration

Figure 1 Risk factors for post-transplant malignancy.

pathogen elimination prevents the establishment of a pro-inflammatory environment conducive to tumourigenesis[6,10]. Thirdly, cells of the innate and adaptive immune system can identify and eliminate tumour cells based on the expression of tumour-specific antigens and danger signals[6,10]. Chronic immunosuppression exposure disrupts the integrity of cancer immunosurveillance. Furthermore, animal studies have suggested that tumours developing in an immunocompromised host are more immunogenic compared to an immunocompetent host, enabling tumour cells to evade immune recognition and destruction[10]. Unsurprisingly, the incidence of PTM is as high as 20% in solid organ transplant recipients after 10 years of cumulative immunosuppression exposure[6].

Potential mechanisms of carcinogenicity of individual immunosuppression drugs

Individual immunosuppression drugs may also have direct carcinogenic effects, resulting in DNA damage and gene expression changes that promote cancer progression independent of the effects of overall immunosuppression exposure.

Tacrolimus

Calcineurin inhibitors (CNI) such as tacrolimus and cyclosporine suppress T cell activation and proliferation by inhibiting *interleukin-2* gene transcription (Figure 2)[11]. The reduced rate of cellular rejection and resultant improved graft and patient survival associated with tacrolimus-based immunosuppression has led to tacrolimus being the CNI of choice following solid organ transplantation[12]. However, experimental data suggest tacrolimus may promote cancer progression by creating a tumour-permissive microenvironment independent of its immunosuppressive effects.

Tacrolimus has a dose-dependent effect on the production of transforming growth factor $\beta 1$ (TGF- $\beta 1$), a cytokine implicated in tumour growth, metastatic spread and development of biologically aggressive cancers[13,14]. The microenvironment is further altered by TGF- $\beta 1$ through inhibition of anti-tumour immune responses and promotion of extracellular matrix production and angiogenesis[15].

The direct effect of tacrolimus on tumour angiogenesis is not fully understood and may be tissue-dependent. *In vivo* studies have demonstrated tacrolimus enhanced lymphangiogenesis and invasion of hepatocellular carcinoma *via* increased vascular endothelial growth factor (VEGF)-C expression[16]. However, tacrolimus may also hinder angiogenesis through its indirect inhibition of nuclear factor of activated T cells, which has a critical role in mediating angiogenesis through its stimulation of VEGF and secreted frizzled-related protein 2[17,18]. This anti-angiogenic effect has an emerging therapeutic role in rheumatoid arthritis, breast cancer, corneal neovascularisation and hypertrophic scars [17,18].

Tacrolimus exposure may lead to alterations in gene expression that promote cancer development and progression. Tacrolimus has been found to activate the proto-oncogene, Ras, in human renal epithelial cells and renal cancer cells, contributing to renal cancer development[19]. Notably, the activation of Ras is critical for VEGF over-expression and subsequent angiogenesis[19]. Tacrolimus can also interfere with proline-oxidase and p53-mediated apoptosis, thus promoting tumour growth[20].

Experimental data on cyclosporine has similarly demonstrated its oncogenic effects through the over-expression of TGF β and VEGF, impaired repair of radiation-induced DNA damage and promotion of apoptosis[21-25]. The shared mechanism of action between tacrolimus and cyclosporine possibly reflects a class-effect of CNIs on malignancy risk.

MPA

Mycophenolate mofetil is a key component of backbone immunosuppression following LT, allowing for CNI de-escalation or cessation and minimisation of renal and metabolic dysfunction. The active metabolite, MPA, inhibits inosine monophosphate dehydrogenase (IMPDH) which is a crucial enzyme involved in *de novo* guanosine nucleotide and DNA synthesis (Figure 2)[26]. This leads to the preferential depletion of lymphocytes due their dependency on *de novo* purine synthesis[26]. There is currently no experimental data linking MPA to increased carcinogenicity risk independent of its effects associated with overall immunosuppression. On the contrary, MPA has *in vitro* and *in vivo* anti-neoplastic properties which may confer a reduced risk of PTM.

MPA has been shown to inhibit the growth of a variety of *in vivo* tumour cell lines[27-31]. Upregulation of peroxisome proliferative-activated receptor gamma by MPA prevents tumour cell differentiation[32]. Reduced expression of adhesion molecules on lymphocytes and endothelial cells interferes with adhesion receptor-dependent tumour dissemination[33-36]. Furthermore, increased expression of subtypes of adhesion receptors from the $\beta 1$ integrin family may induce re-




Figure 2 Mechanism of action of commonly used immunosuppression drugs following solid organ transplantation. APC: Antigen presenting cell; IL: Interleukin; MHC: Major histocompatibility complex; mTOR: Mammalian target of rapamycin; NFAT: Nuclear factor of activated T cells.

differentiation of tumour cells towards a lower invasive phenotype[36]. However, some cancer types have been found to be resistant to the anti-neoplastic properties of MPA[28,37].

The anti-neoplastic properties of MPA may also have a therapeutic potential. The enzyme IMPDH, the target of MPA, is over-expressed in cancer cells[26]. Furthermore, MPA-mediated inhibition of IMPDH has been demonstrated to induce tumour cell apoptosis, however these findings are yet to be confirmed *in vivo*[38].

Other immunosuppression agents

Azathioprine is a purine analogue that is incorporated into cellular DNA, where it inhibits purine nucleotide synthesis and interferes with RNA synthesis and metabolism (Figure 2)[15]. It is well known that azathioprine is a risk factor for the development of PTM, in particular, non-melanoma skin cancer. Multiple studies have demonstrated the synergistic effect between ultraviolet A radiation and the azathioprine metabolite, 6-thioguanine, in the generation of mutagenic oxidative DNA damage[39,40]. The carcinogenic effects of azathioprine have limited its use in transplantation in favour of MPA.

Sirolimus and everolimus inhibit mammalian target of rapamycin (mTOR), which subsequently downregulates cyclin-dependent kinases and mRNAs required for cell cycle progression, thus preventing interleukin-2-mediated lymphocyte proliferation (Figure 2)[9]. *In vivo* studies have shown mTOR inhibitors precipitate tumour cell cycle progression arrest and subsequent apoptosis[41,42]. Impaired VEGF production and signalling also restricts tumour angiogenesis and metastatic spread[43-45]. Interestingly, the simultaneous administration of sirolimus in these models can reverse the pro-angiogenic effects of cyclosporine[43-45]. The potential dual immunosuppressive and anti-neoplastic properties of mTOR inhibitors has led to its increasing utilisation in the transplantation setting.

MATERIALS AND METHODS

Multiple population and cohort studies have investigated the role of tacrolimus and MPA in the development of *de novo* PTM, however a direct causal relationship is difficult to establish. As the two most commonly used drugs for maintenance monotherapy post-LT, the oncogenic risk profile of tacrolimus and MPA warrants further review.

Preferred Reporting Items for Systematic Reviews and Meta-analysis (PRISMA) guidelines were utilised to shortlist relevant articles for this narrative review to minimise bias. A comprehensive literature search was conducted through MEDLINE and Embase electronic databases between 1st January 2002 to 11th August 2022. This time period was selected to include relevant literature since the introduction and clinical use of MPA. The following terms were used, including synonyms and closely related words, as Medical Subject Headings (MeSH) and text words: "Solid Organ Transplantation", "Tacrolimus", "Mycophenolic Acid", and "Carcinogenicity". Explosion of MeSH terms, Boolean operators and truncations were also utilised throughout the search. Further articles were identified through reference lists of published systematic reviews in the area. Excluding duplicates, abstracts from 1230 records were screened by a single

reviewer, whereby 31 records were deemed appropriate for full-text review. Full-text articles were assessed for eligibility based on inclusion and exclusion criteria, leaving 6 studies for inclusion in this review (Table 1, Figure 3).

RESULTS

The 6 studies included in this review are summarised in Table 2. All studies were large population-based registries or cohort studies that analysed PTM risk in the presence or absence of tacrolimus or MPA use. No studies included data on individual drug dosages, plasma levels or duration to assess for cumulative drug exposure. There was heterogeneity amongst the studied populations in type of organ transplanted, transplantation era and immunosuppression regimens used. No studies provided a direct comparative risk of PTM with tacrolimus or MPA monotherapy.

Cutaneous and non-cutaneous malignancy

A Taiwanese population-based study evaluated risk factors for *de novo* cutaneous and non-cutaneous malignancy in 7852 liver, heart, and kidney transplant recipients[46]. Among 2127 liver transplant recipients, 111 (5.2%) malignancies were recorded during the mean follow-up period of 4.2 years[46]. Despite the majority of liver transplant recipients using tacrolimus (77.3%) or MPA (99.0%), neither immunosuppressant was associated with PTM risk[46].

Among 687 heart transplant patients, 31 (4.5%) *de novo* malignancies were reported[46]. Immunosuppression therapy was also not associated with PTM risk in this cohort[46]. However, the smaller number of malignancy outcomes may have contributed to attenuated risk estimates.

De novo malignancy was diagnosed in 470 out of 5038 (9.3%) kidney transplant recipients[46]. The use of MPA was an independent risk factor for PTM in kidney transplant recipients, compared to no MPA use [adjusted hazard ratio (HR): 1.5, 95% confidence interval (95%CI): 1.2-1.8; $P < 0.001$][46]. MPA exposure was also a risk factor for *de novo* transitional cell carcinoma (adjusted HR: 1.7, 95%CI: 1.2-2.4; $P < 0.01$) and renal cell carcinoma (adjusted HR: 1.7, 95%CI: 1.1-2.8; $P < 0.05$) in a sub-analysis of kidney transplant recipients without hypertension or diabetes as an underlying cause for renal failure[46].

On the contrary, a smaller Taiwanese population cohort study of 642 kidney transplant recipients did not demonstrate an association between MPA or tacrolimus exposure, and the development of 54 (8.4%) *de novo* malignancies[47]. However, the study's primary endpoint of hospitalisation due to malignancy as the primary coded diagnosis, likely underestimated the incidence of *de novo* PTM from the exclusion of malignancies coded as secondary diagnoses or those diagnosed in the community.

Differences in immunosuppression regimens and cumulative exposure to individual drugs may also contribute to the conflicting findings of the aforementioned studies, however this data was not available for analysis. Additionally, lifestyle factors known to influence malignancy risk such as smoking and alcohol consumption, were not included in either study.

Cutaneous malignancy

Three studies investigated the relationship between immunosuppression and post-transplant cutaneous malignancy.

A population-based study in the United Kingdom investigated the development of post-transplant melanoma and non-melanoma skin cancers in 2852 liver, kidney, pancreas, heart, and lung transplant recipients, compared to 13527 matched controls from the general population[48]. Among 437 liver transplant recipients, 19 (4.3%) skin cancers were diagnosed during the 6.2 year median follow-up period[48]. Liver transplant recipients had the lowest incidence of skin cancer compared to other solid organ transplant recipients [Incidence rate ratio (IRR): 4.34, 95%CI: 2.48-7.58, $P = 0.00$], possibly reflecting lower immunosuppression requirements and relative immune privilege[48]. Neither tacrolimus nor MPA use was associated with the development of *de novo* cutaneous malignancy across all solid organ transplantation[48]. However, these findings are limited by small outcome numbers. Additionally, the complex interaction between immunosuppression agents and other risk factors for skin cancer including smoking status and ultraviolet light exposure was not considered.

An American study compared 170 kidney, kidney/pancreas, and heart transplant recipients with *de novo* cutaneous squamous cell carcinoma (SCC) to 324 matched recipient controls[49]. Risk factors such as smoking status, family history of skin cancer and personal history of pre-cancerous skin lesions were adjusted for, however the cancer group were significantly older than the non-cancer group despite matching. In azathioprine naïve patients, MPA use was associated with lower cutaneous SCC risk, independent of tacrolimus exposure (OR: 0.52, 95%CI: 0.32-0.84)[49]. Current and previous MPA use was also inversely associated with the development of multiple cutaneous SCCs (previous MPA use: OR: 0.53, 95%CI: 0.3-0.94; current MPA use: OR: 0.52, 95%CI: 0.29-0.94)[49]. Conversely, cyclosporine-naïve patients treated with tacrolimus had no significant difference in cutaneous SCC risk compared to no tacrolimus use, when adjusted for MPA exposure[49]. Although the authors considered individual immunosuppression exposure risk in the clinical context of changing multi-drug regimens, this was limited by potential recall bias associated with self-reported questionnaires used to obtain immunosuppression data.

Finally, *de novo* lip SCC was evaluated in a large Australian and New Zealand registry study of 8162 kidney transplant patients[50]. Mycophenolate use was associated with reduced risk of SCC of the lower vermilion of the lip in univariate (IRR: 0.28, 95%CI: 0.12-0.69, $P = 0.006$), but not multivariate (IRR: 0.85, 95%CI: 0.28-2.60, $P = 0.774$) analyses[50]. There was no difference between tacrolimus use *vs* no use in the risk of lip SCC of the lower vermilion (IRR: 2.07, 95%CI: 0.45-9.50, $P = 0.35$)[50]. Of note, the study included patients transplanted between 1982 and 2003, with less use of tacrolimus (2/121, 1.7%) and MPA (5/121, 4.1%) during this transplant era, compared to cyclosporine and azathioprine, respectively.

Table 1 Inclusion and exclusion criteria utilised for literature search strategy

Inclusion criteria	Exclusion criteria
Involve human solid organ transplant recipients	Presents risk data on only one of the immunosuppressant medications
Independent malignancy risk analysis related to both immunosuppressants mycophenolic acid and tacrolimus	Does not specify type of immunosuppression
Contains a group of participants exposed to tacrolimus or mycophenolic acid, exclusive of the other	Mean follow up less than one year (given the slow growing nature of malignancy)
Greater than 100 participants	Not published in English
Greater than 5 cases of malignancy	Full text not available
Randomised controlled trials and observational studies	Systematic reviews and meta-analyses

This study was likely underpowered to draw conclusions between tacrolimus and MPA exposure and risk of SCC of the lower vermillion of the lip.

PTLD

A large population registry in France evaluated risk factors for PTLT occurrence in kidney and kidney/pancreas transplant recipients over a 10-year period[51]. Compared to 21170 control kidney transplant recipients, 327 cases of PTLT were recorded and 181 cases were included in the final analysis[51]. Tacrolimus and MPA use were not associated with overall PTLT risk, even when simultaneous kidney pancreas transplant recipients were excluded[51]. However, tacrolimus and MPA were negatively associated with graft site PTLT (tacrolimus: HR: 0.33, 95%CI: 0.16-0.68; MPA: HR: 0.44; 95%CI: 0.23-0.86), which may be attributed to fewer episodes of acute rejection and less immunosuppression exposure in this subgroup[51].

DISCUSSION

With long-term survival now commonplace following LT, there is an increasing need to improve non-hepatic health to avoid complications including metabolic derangements, renal impairment and *de novo* malignancy. *De novo* PTM accounts for approximately 16.4% of late deaths following LT[12,52]. Although immunosuppression exposure is a well-known contributor of PTM risk, there remains uncertainty regarding the carcinogenic effect of specific immunosuppression drugs, alone or in combination. This is the first narrative review that compares the relative carcinogenicity of tacrolimus and MPA in solid organ transplant recipients.

Existing *in vitro* and *in vivo* experimental data have portrayed a contrasting carcinogenic risk profile between tacrolimus and MPA. Tacrolimus promotes oncogenesis and tumour growth in its surrounding microenvironment with the activation of proto-oncogenes, production of TGF- β and inhibition of apoptosis[13,19,20]. The data on MPA is limited but suggests possible inhibition of tumour cell differentiation and prevention of vascular spread through alteration of cellular adhesion molecule expression[6]. However, there is currently no human data that directly compares the carcinogenic effects of tacrolimus and MPA in LT or other solid organ transplantation.

This review included a small number of studies that did not demonstrate a clear difference between tacrolimus and MPA in *de novo* PTM risk following solid organ transplantation. Our findings are in keeping with a recent systematic review and meta-analysis of kidney, liver, heart, and lung transplant recipients, whereby the risk of *de novo* malignancy did not differ between patients who received MPA and patients who received tacrolimus (OR: 0.88, 95%CI: 0.69-1.14, $P = 0.33$)[53]. However, the relationship between immunosuppression exposure and *de novo* PTM risk may vary based on transplant type. In liver transplant recipients, cumulative tacrolimus exposure has been associated with the development of PTM[54,55], although the high tacrolimus doses utilised in these studies are no longer aimed for in routine clinical practice. Furthermore, the conversion from CNI-based immunosuppression to MPA monotherapy post-LT results in either similar or lower rates of PTM[56,57]. Whether the reduction in PTM risk found in these studies is due to the effects of MPA or the reduction in tacrolimus exposure, is unknown. Thus, the differential carcinogenic risk profile of tacrolimus and MPA found in previous experimental studies is yet to be replicated in the clinical setting. Further clarification with large prospective studies is required.

There are inherent practical and financial difficulties in designing studies to compare the relative risk of *de novo* PTM between tacrolimus and MPA. Large prospective population-based studies of prolonged follow-up duration are required to ensure adequate statistical power. Variables that influence PTM risk such as age, gender, ethnicity, and smoking should be identified. However, there may be unidentifiable confounders that are difficult to capture, owing to the complex interaction between genetic, lifestyle and disease factors in oncogenesis. Population-based registries often rely on International Classification of Diseases coding for data collection, which can lead under-representation of malignancy incidence due to miscoding. Finally, longitudinal recording of drug dose, plasma levels and duration is required to capture changes in immunosuppression regimens frequently seen in routine clinical practice. The accurate calculation of cumulative immunosuppression exposure minimises drug exposure misclassification bias seen in current transplant

Table 2 Summary of studies comparing tacrolimus to mycophenolic acid in solid organ transplantation

Ref.	Transplant era	Organ	n	De novo malignancy	Time to malignancy (median)	TAC vs no TAC (OR/HR/IRR)	MPA vs no MPA (OR/HR/IRR)
Van Leeuwen <i>et al</i> [50], 2009	1982 to 2003	Kidney	8162	203 lip SCC; 121 lip SCC of the lower vermillion in first transplant)	6.1 yr	SCC of the lower vermillion of lip during first transplant [IRR: 2.07, 95%CI: 0.45-9.50 ($P = 0.35$)]	SCC of the lower vermillion of lip during first transplant [IRR: 0.85, 95%CI: 0.28-2.60 ($P = 0.77$)]
Caillard <i>et al</i> [51], 2012	1998 to 2007	Kidney \pm pancreas	21351	181 PTLD; 43 graft PTLD	Not specified	Any PTLD [aHR: 0.66, 95%CI: 0.36-1.22 ($P = 0.19$)]; Graft PTLD [HR: 0.33, 95%CI: 0.16-0.68 ($P = 0.003$)*]	Any PTLD [aHR: 1.22, 95%CI: 0.74-2.02 ($P = 0.44$)]; Graft PTLD [HR: 0.44, 95%CI: 0.23-0.86 ($P = 0.015$)*]
Hsiao <i>et al</i> [47], 2014	2000 to 2008	Kidney	642	54 non-cutaneous malignancy	3.9 yr	HR: 1.99, 95%CI: 0.66-6.00 ($P = 0.22$)	HR: 1.00, 95%CI: 0.40-2.45 ($P = 0.99$)
Coghill <i>et al</i> [49], 2016	1995 to 2010	Kidney \pm pancreas, and heart	2004	170 SCC	9.0 yr	Single SCC (OR: 1.11, 95%CI: 0.48-2.60)	Single SCC (OR: 0.52, 95%CI: 0.32-0.84*)
Yeh <i>et al</i> [46], 2020	1997 to 2011	Liver, kidney, and heart	7852	612 cutaneous and non-cutaneous malignancy	Not specified	cHR (heart): 0.6, 95%CI: 0.1-2.7; cHR (kidney): 1.5, 95%CI: 0.8-2.6; aHR (liver): 0.6, 95%CI: 0.2-1.7	cHR (heart): 1.6, 95%CI: 0.7-3.3; aHR (kidney): 1.5, 95%CI: 1.2-1.8 ($P < 0.001$)*; cHR (liver): 1.5, 95%CI: 0.9-2.5
Gibson <i>et al</i> [48], 2021	2010 to 2018	Liver, kidney \pm pancreas, heart, and lung	2852	242 cutaneous malignancy	4.7 yr	IRR: 0.83, 95%CI: 0.55-1.25 ($P = 0.37$)	IRR: 0.78, 95%CI: 0.54-1.12 ($P = 0.18$)

* $P < 0.05$.

aHR: Adjusted hazard ratio; cHR: Crude hazard ratio; HR: Hazard ratio; IRR: Incidence rate ratio; MPA: Mycophenolic acid; OR: Odds ratio; PTLD: Post-transplant lymphoproliferative disorder; SCC: Squamous cell carcinoma; TAC: Tacrolimus; 95%CI: 95% confidence interval.

cohort analyses that presume an unvarying drug regimen.

Immunosuppression minimisation is an important strategy to reduce PTM risk given the limited clinical data surrounding individual agents. There are currently no clear guidelines regarding immunosuppression drug choice to minimise PTM risk following LT. European LT guidelines state CNI-related *de novo* PTM risk may be due to dosage, and that there is no evidence to suggest MPA contributes to *de novo* PTM development[58]. In our centre, there is a preference for MPA, alone or in combination with everolimus, due to improved renal outcomes and experimental data suggesting higher PTM risk with tacrolimus. Overall, the choice of immunosuppression needs to be individualised based on recipient characteristics, liver disease aetiology, and alloimmune risk.

Routine cancer surveillance for all transplant recipients is recommended in addition to immunosuppression minimisation. Strict cancer surveillance strategies may lead to earlier cancer detection rates and improved non-cutaneous cancer patient survival in LT recipients[59,60]. As non-melanoma skin cancer is the leading cause of PTM in LT recipients, annual skin examinations by a dermatologist are recommended from 5 years or more after LT[5,61]. Recipients with primary sclerosing cholangitis and inflammatory bowel disease require annual colonoscopies for colorectal cancer surveillance[61]. Age and gender based cancer surveillance for all LT recipients is also recommended.




Figure 3 Search strategy utilised for article selection.

CONCLUSION

The clinical relevance of previous experimental studies on the relative carcinogenicity of tacrolimus and MPA, and its application in solid organ transplantation, is yet to be confirmed. Consequently, the choice of immunosuppressive agent to use as maintenance monotherapy in LT patients is not currently supported by a strong evidence base and remains unclear. Further studies are required to enable physicians to tailor immunosuppression regimens to minimise individual malignancy risk.

ARTICLE HIGHLIGHTS

Research background

Many liver transplant (LT) recipients are able to be maintained on long-term immunosuppressive monotherapy, most commonly with either tacrolimus or mycophenolate. In experimental studies, tacrolimus is associated with increased carcinogenicity, whereas mycophenolic acid (MPA) may have anti-neoplastic properties. However, there is minimal clinical data comparing the relative carcinogenicity of tacrolimus and MPA in LT or other solid organ transplant recipients.

Research motivation

Post-transplant malignancy (PTM) is a leading cause of late mortality in LT recipients. Thus, a clinically relevant difference in the carcinogenic risk profile between tacrolimus and MPA will affect the choice of immunosuppressive agent used as maintenance monotherapy in LT patients.

Research objectives

To determine the relative carcinogenicity of tacrolimus and MPA in solid organ transplantation.

Research methods

A systematic review was conducted using PRISMA guidelines with relevant articles published between 1st January 2002 to 11th August 2022 retrieved from MEDLINE and Embase databases for review.

Research results

A total of 6 studies were included in this systematic review, which did not demonstrate a clear difference between tacrolimus and MPA in the development of *de novo* PTM following solid organ transplantation.

Research conclusions

The relative carcinogenicity of tacrolimus and MPA, and its clinical relevance in solid organ transplantation, remains unclear.

Research perspectives

This review highlights the need for further large, population-based prospective studies to further assess the carcinogenic profiles of tacrolimus and MPA, to assist physicians in the choice of immunosuppressive agent to use as maintenance monotherapy in LT patients.

FOOTNOTES

Author contributions: Liu D was involved in data interpretation, drafting of the original and revised manuscripts, and prepared some of the figures and tables; Youssef MM performed data acquisition and prepared some of the figures and tables; Sinclair M was responsible for the study design; Grace JA and Sinclair M were involved in data interpretation and critical revision of the manuscript; and all authors have read and approved the final manuscript.

Conflict-of-interest statement: The authors have no conflicts of interest to report.

PRISMA 2009 Checklist statement: The authors have read the PRISMA 2009 Checklist, and the manuscript was prepared and revised according to the PRISMA 2009 Checklist.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Australia

ORCID number: Dorothy Liu 0000-0002-7802-2803; Josephine A Grace 0000-0002-8435-4740; Marie Sinclair 0000-0003-0657-3115.

S-Editor: Chen YL

L-Editor: A

P-Editor: Yu HG

REFERENCES

- 1 **McCaughan GW**, Munn SR. Liver transplantation in Australia and New Zealand. *Liver Transpl* 2016; **22**: 830-838 [PMID: 27028552 DOI: 10.1002/lt.24446]
- 2 **Herrero JI**. De novo malignancies following liver transplantation: impact and recommendations. *Liver Transpl* 2009; **15** Suppl 2: S90-S94 [PMID: 19877025 DOI: 10.1002/lt.21898]
- 3 **Rodríguez-Perálvarez M**, De la Mata M, Burroughs AK. Liver transplantation: immunosuppression and oncology. *Curr Opin Organ Transplant* 2014; **19**: 253-260 [PMID: 24685671 DOI: 10.1097/MOT.000000000000069]
- 4 **Gelson W**, Hoare M, Dawwas MF, Vowler S, Gibbs P, Alexander G. The pattern of late mortality in liver transplant recipients in the United Kingdom. *Transplantation* 2011; **91**: 1240-1244 [PMID: 21516069 DOI: 10.1097/TP.0b013e31821841ba]
- 5 **Chandok N**, Watt KD. Burden of de novo malignancy in the liver transplant recipient. *Liver Transpl* 2012; **18**: 1277-1289 [PMID: 22887956 DOI: 10.1002/lt.23531]
- 6 **Buell JF**, Gross TG, Woodle ES. Malignancy after transplantation. *Transplantation* 2005; **80**: S254-S264 [PMID: 16251858 DOI: 10.1097/01.tp.0000186382.81130.ba]
- 7 **Abrol N**, Jadlowiec CC, Taner T. Revisiting the liver's role in transplant alloimmunity. *World J Gastroenterol* 2019; **25**: 3123-3135 [PMID: 31333306 DOI: 10.3748/wjg.v25.i25.3123]
- 8 **Wiesner RH**, Fung JJ. Present state of immunosuppressive therapy in liver transplant recipients. *Liver Transpl* 2011; **17** Suppl 3: S1-S9 [PMID: 21850697 DOI: 10.1002/lt.22410]
- 9 **Guba M**, Graeb C, Jauch KW, Geissler EK. Pro- and anti-cancer effects of immunosuppressive agents used in organ transplantation. *Transplantation* 2004; **77**: 1777-1782 [PMID: 15223891 DOI: 10.1097/01.Tp.0000120181.89206.54]
- 10 **Schreiber RD**, Old LJ, Smyth MJ. Cancer immunoediting: integrating immunity's roles in cancer suppression and promotion. *Science* 2011; **331**: 1565-1570 [PMID: 21436444 DOI: 10.1126/science.1203486]
- 11 **Panackel C**, Mathew JF, Fawas N M, Jacob M. Immunosuppressive Drugs in Liver Transplant: An Insight. *J Clin Exp Hepatol* 2022; **12**: 1557-1571 [PMID: 36340316 DOI: 10.1016/j.jceh.2022.06.007]
- 12 **Rana A**, Ackah RL, Webb GJ, Halazun KJ, Vierling JM, Liu H, Wu MF, Yoeli D, Kueht M, Mindikoglu AL, Sussman NL, Galván NT, Cotton RT, O'Mahony CA, Goss JA. No Gains in Long-term Survival After Liver Transplantation Over the Past Three Decades. *Ann Surg* 2019; **269**: 20-27 [PMID: 29303806 DOI: 10.1097/SLA.0000000000002650]
- 13 **Maluccio M**, Sharma V, Lagman M, Vyas S, Yang H, Li B, Suthanthiran M. Tacrolimus enhances transforming growth factor-beta1 expression and promotes tumor progression. *Transplantation* 2003; **76**: 597-602 [PMID: 12923450 DOI: 10.1097/01.Tp.0000081399.75231.3b]

- 14 **Suthanthiran M**, Hojo M, Maluccio M, Boffa DJ, Luan FL. Post-transplantation malignancy: a cell autonomous mechanism with implications for therapy. *Trans Am Clin Climatol Assoc* 2009; **120**: 369-388 [PMID: 19768190]
- 15 **Gutierrez-Dalmau A**, Campistol JM. Immunosuppressive therapy and malignancy in organ transplant recipients: a systematic review. *Drugs* 2007; **67**: 1167-1198 [PMID: 17521218 DOI: 10.2165/00003495-200767080-00006]
- 16 **Zhou S**, Tan C, Dai Z, Zhu H, Xu M, Zhou Z, Wang W, Zhao Y, Fu X, Zhou J, Fan J. Tacrolimus enhances the invasion potential of hepatocellular carcinoma cells and promotes lymphatic metastasis in a rat model of hepatocellular carcinoma: involvement of vascular endothelial growth factor-C. *Transplant Proc* 2011; **43**: 2747-2754 [PMID: 21911157 DOI: 10.1016/j.transproceed.2011.06.040]
- 17 **Siamakpour-Reihani S**, Caster J, Bandhu Nepal D, Courtwright A, Hilliard E, Usary J, Ketelsen D, Darr D, Shen XJ, Patterson C, Klauber-Demore N. The role of calcineurin/NFAT in SFRP2 induced angiogenesis--a rationale for breast cancer treatment with the calcineurin inhibitor tacrolimus. *PLoS One* 2011; **6**: e20412 [PMID: 21673995 DOI: 10.1371/journal.pone.0020412]
- 18 **Shen Y**, Jin R, Liang X, Deng Z, He J, Ding Y, Ding F, Lu L, Liu F, Yang J. Angiogenesis modulation-mediated inhibitory effects of tacrolimus on hypertrophic scar formation. *Microvasc Res* 2023; **145**: 104446 [PMID: 36270418 DOI: 10.1016/j.mvr.2022.104446]
- 19 **Datta D**, Contreras AG, Basu A, Dormond O, Flynn E, Briscoe DM, Pal S. Calcineurin inhibitors activate the proto-oncogene Ras and promote protumorigenic signals in renal cancer cells. *Cancer Res* 2009; **69**: 8902-8909 [PMID: 19903851 DOI: 10.1158/0008-5472.CAN-09-1404]
- 20 **Rivera A**, Maxwell SA. The p53-induced gene-6 (proline oxidase) mediates apoptosis through a calcineurin-dependent pathway. *J Biol Chem* 2005; **280**: 29346-29354 [PMID: 15914462 DOI: 10.1074/jbc.M504852200]
- 21 **Hojo M**, Morimoto T, Maluccio M, Asano T, Morimoto K, Lagman M, Shimbo T, Suthanthiran M. Cyclosporine induces cancer progression by a cell-autonomous mechanism. *Nature* 1999; **397**: 530-534 [PMID: 10028970 DOI: 10.1038/17401]
- 22 **Shihab FS**, Bennett WM, Isaac J, Yi H, Andoh TF. Nitric oxide modulates vascular endothelial growth factor and receptors in chronic cyclosporine nephrotoxicity. *Kidney Int* 2003; **63**: 522-533 [PMID: 12631117 DOI: 10.1046/j.1523-1755.2003.00757.x]
- 23 **Herman M**, Weinstein T, Korzets A, Chagnac A, Ori Y, Zevin D, Malachi T, Gafer U. Effect of cyclosporin A on DNA repair and cancer incidence in kidney transplant recipients. *J Lab Clin Med* 2001; **137**: 14-20 [PMID: 11150019 DOI: 10.1067/mLc.2001.111469]
- 24 **Morisaki T**, Matsunaga H, Beppu K, Ihara E, Hirano K, Kanaide H, Mori M, Katano M. A combination of cyclosporin-A (CsA) and interferon-gamma (INF-gamma) induces apoptosis in human gastric carcinoma cells. *Anticancer Res* 2000; **20**: 3363-3373 [PMID: 11131636]
- 25 **Nomura T**, Yamamoto H, Mimata H, Shitashige M, Shibasaki F, Miyamoto E, Nomura Y. Enhancement by cyclosporin A of taxol-induced apoptosis of human urinary bladder cancer cells. *Urol Res* 2002; **30**: 102-111 [PMID: 12086014 DOI: 10.1007/s00240-002-0239-4]
- 26 **Kaltenborn A**, Schrem H. Mycophenolate mofetil in liver transplantation: a review. *Ann Transplant* 2013; **18**: 685-696 [PMID: 24346057 DOI: 10.12659/AOT.889299]
- 27 **Williams RH**, Lively DH, DeLong DC, Cline JC, Sweeny MJ. Mycophenolic acid: antiviral and antitumor properties. *J Antibiot (Tokyo)* 1968; **21**: 463-464 [PMID: 4303502 DOI: 10.7164/antibiotics.21.463]
- 28 **Tressler RJ**, Garvin LJ, Slate DL. Anti-tumor activity of mycophenolate mofetil against human and mouse tumors in vivo. *Int J Cancer* 1994; **57**: 568-573 [PMID: 8181860 DOI: 10.1002/ijc.2910570421]
- 29 **Yokoyama I**, Hayashi S, Kobayashi T, Negita M, Yasutomi M, Uchida K, Takagi H. Immunosuppressive drugs and their effect on experimental tumor growth. *Transpl Int* 1995; **8**: 251-255 [PMID: 7546145 DOI: 10.1007/bf00346876]
- 30 **Sweeney MJ**, Gerzon K, Harris PN, Holmes RE, Poore GA, Williams RH. Experimental antitumor activity and preclinical toxicology of mycophenolic acid. *Cancer Res* 1972; **32**: 1795-1802 [PMID: 4509039]
- 31 **Koehl GE**, Wagner F, Stoeltzing O, Lang SA, Steinbauer M, Schlitt HJ, Geissler EK. Mycophenolate mofetil inhibits tumor growth and angiogenesis in vitro but has variable antitumor effects in vivo, possibly related to bioavailability. *Transplantation* 2007; **83**: 607-614 [PMID: 17353782 DOI: 10.1097/01.tp.0000253756.69243.65]
- 32 **Zheng ZH**, Yang Y, Lu XH, Zhang H, Shui XX, Liu C, He XB, Jiang Q, Zhao BH, Si SY. Mycophenolic acid induces adipocyte-like differentiation and reversal of malignancy of breast cancer cells partly through PPAR γ . *Eur J Pharmacol* 2011; **658**: 1-8 [PMID: 21349264 DOI: 10.1016/j.ejphar.2011.01.068]
- 33 **Heemann U**, Azuma H, Hamar P, Schmid C, Tilney N, Philipp T. Mycophenolate mofetil inhibits lymphocyte binding and the upregulation of adhesion molecules in acute rejection of rat kidney allografts. *Transpl Immunol* 1996; **4**: 64-67 [PMID: 8762015 DOI: 10.1016/s0966-3274(96)80039-6]
- 34 **Blaheta RA**, Leckel K, Wittig B, Zenker D, Oppermann E, Harder S, Scholz M, Weber S, Encke A, Markus BH. Mycophenolate mofetil impairs transendothelial migration of allogeneic CD4 and CD8 T-cells. *Transplant Proc* 1999; **31**: 1250-1252 [PMID: 10083559 DOI: 10.1016/s0041-1345(98)01984-8]
- 35 **Leckel K**, Beecken WD, Jonas D, Oppermann E, Coman MC, Beck KF, Cinatl J, Hailer NP, Auth MK, Bechstein WO, Shipkova M, Blaheta RA. The immunosuppressive drug mycophenolate mofetil impairs the adhesion capacity of gastrointestinal tumour cells. *Clin Exp Immunol* 2003; **134**: 238-245 [PMID: 14616783 DOI: 10.1046/j.1365-2249.2003.02290.x]
- 36 **Engl T**, Makarević J, Relja B, Natshah I, Müller I, Beecken WD, Jonas D, Blaheta RA. Mycophenolate mofetil modulates adhesion receptors of the beta1 integrin family on tumor cells: impact on tumor recurrence and malignancy. *BMC Cancer* 2005; **5**: 4 [PMID: 15644133 DOI: 10.1186/1471-2407-5-4]
- 37 **Dun B**, Sharma A, Teng Y, Liu H, Purohit S, Xu H, Zeng L, She JX. Mycophenolic acid inhibits migration and invasion of gastric cancer cells via multiple molecular pathways. *PLoS One* 2013; **8**: e81702 [PMID: 24260584 DOI: 10.1371/journal.pone.0081702]
- 38 **Guidicelli G**, Chaigne-Delalande B, Dilhuydy MS, Pinson B, Mahfouf W, Pasquet JM, Mahon FX, Pourquier P, Moreau JF, Legembre P. The necrotic signal induced by mycophenolic acid overcomes apoptosis-resistance in tumor cells. *PLoS One* 2009; **4**: e5493 [PMID: 19430526 DOI: 10.1371/journal.pone.0005493]
- 39 **O'Donovan P**, Perrett CM, Zhang X, Montaner B, Xu YZ, Harwood CA, McGregor JM, Walker SL, Hanaoka F, Karran P. Azathioprine and UVA light generate mutagenic oxidative DNA damage. *Science* 2005; **309**: 1871-1874 [PMID: 16166520 DOI: 10.1126/science.1114233]
- 40 **Kelly GE**, Meikle W, Sheil AG. Effects of immunosuppressive therapy on the induction of skin tumors by ultraviolet irradiation in hairless mice. *Transplantation* 1987; **44**: 429-434 [PMID: 3629691 DOI: 10.1097/00007890-198709000-00021]
- 41 **Majewski M**, Korecka M, Joergensen J, Fields L, Kossev P, Schuler W, Shaw L, Wasik MA. Immunosuppressive TOR kinase inhibitor everolimus (RAD) suppresses growth of cells derived from posttransplant lymphoproliferative disorder at allograft-protecting doses. *Transplantation* 2003; **75**: 1710-1717 [PMID: 12777861 DOI: 10.1097/01.Tp.0000063934.89714.19]
- 42 **Nepomuceno RR**, Balatoni CE, Natkunam Y, Snow AL, Krams SM, Martinez OM. Rapamycin inhibits the interleukin 10 signal transduction pathway and the growth of Epstein Barr virus B-cell lymphomas. *Cancer Res* 2003; **63**: 4472-4480 [PMID: 12907620]

- 43 **Guba M**, von Breitenbuch P, Steinbauer M, Koehl G, Flegel S, Hornung M, Bruns CJ, Zuelke C, Farkas S, Anthuber M, Jauch KW, Geissler EK. Rapamycin inhibits primary and metastatic tumor growth by antiangiogenesis: involvement of vascular endothelial growth factor. *Nat Med* 2002; **8**: 128-135 [PMID: 11821896 DOI: 10.1038/nm0202-128]
- 44 **Luan FL**, Ding R, Sharma VK, Chon WJ, Lagman M, Suthanthiran M. Rapamycin is an effective inhibitor of human renal cancer metastasis. *Kidney Int* 2003; **63**: 917-926 [PMID: 12631072 DOI: 10.1046/j.1523-1755.2003.00805.x]
- 45 **Boffa DJ**, Luan F, Thomas D, Yang H, Sharma VK, Lagman M, Suthanthiran M. Rapamycin inhibits the growth and metastatic progression of non-small cell lung cancer. *Clin Cancer Res* 2004; **10**: 293-300 [PMID: 14734482 DOI: 10.1158/1078-0432.ccr-0629-3]
- 46 **Yeh CC**, Khan A, Muo CH, Yang HR, Li PC, Chang CH, Chen TL, Jeng LB, Liao CC. De Novo Malignancy After Heart, Kidney, and Liver Transplant: A Nationwide Study in Taiwan. *Exp Clin Transplant* 2020; **18**: 224-233 [PMID: 32133940 DOI: 10.6002/ect.2019.0210]
- 47 **Hsiao FY**, Hsu WW. Epidemiology of post-transplant malignancy in Asian renal transplant recipients: a population-based study. *Int Urol Nephrol* 2014; **46**: 833-838 [PMID: 24009082 DOI: 10.1007/s11255-013-0544-6]
- 48 **Gibson JAG**, Cordaro A, Dobbs TD, Griffiths R, Akbari A, Whitaker S, Hutchings HA, Lyons RA, Whitaker IS. The association between immunosuppression and skin cancer in solid organ transplant recipients: a control-matched cohort study of 2,852 patients. *Eur J Dermatol* 2021; **31**: 712-721 [PMID: 34427560 DOI: 10.1684/ejd.2021.4108]
- 49 **Coghill AE**, Johnson LG, Berg D, Resler AJ, Leca N, Madeleine MM. Immunosuppressive Medications and Squamous Cell Skin Carcinoma: Nested Case-Control Study Within the Skin Cancer after Organ Transplant (SCOT) Cohort. *Am J Transplant* 2016; **16**: 565-573 [PMID: 26824445 DOI: 10.1111/ajt.13596]
- 50 **van Leeuwen MT**, Grulich AE, McDonald SP, McCredie MR, Amin J, Stewart JH, Webster AC, Chapman JR, Vajdic CM. Immunosuppression and other risk factors for lip cancer after kidney transplantation. *Cancer Epidemiol Biomarkers Prev* 2009; **18**: 561-569 [PMID: 19190169 DOI: 10.1158/1055-9965.EPI-08-0919]
- 51 **Caillard S**, Lamy FX, Quelen C, Dantal J, Lebranchu Y, Lang P, Velten M, Moulin B; French Transplant Centers. Epidemiology of posttransplant lymphoproliferative disorders in adult kidney and kidney pancreas recipients: report of the French registry and analysis of subgroups of lymphomas. *Am J Transplant* 2012; **12**: 682-693 [PMID: 22226336 DOI: 10.1111/j.1600-6143.2011.03896.x]
- 52 **Fuochi E**, Anastasio L, Lynch EN, Campani C, Dragoni G, Milani S, Galli A, Innocenti T. Main factors influencing long-term outcomes of liver transplantation in 2022. *World J Hepatol* 2023; **15**: 321-352 [PMID: 37034235 DOI: 10.4254/wjh.v15.i3.321]
- 53 **Zwart ES**, Yüksel E, Pannekoek A, de Vries R, Mebius RE, Kazemier G. De Novo Carcinoma after Solid Organ Transplantation to Give Insight into Carcinogenesis in General-A Systematic Review and Meta-Analysis. *Cancers (Basel)* 2021; **13** [PMID: 33807849 DOI: 10.3390/cancers13051122]
- 54 **Rodríguez-Perálvarez M**, Colmenero J, González A, Gastaca M, Curell A, Caballero-Marcos A, Sánchez-Martínez A, Di Maira T, Herrero JI, Almohalla C, Lorente S, Cuadrado-Lavín A, Pascual S, López-Garrido MÁ, González-Grande R, Gómez-Orellana A, Alejandro R, Zamora-Olaya J, Bernal-Bellido C; Chronic immunosuppression, cancer Spanish consortium. Cumulative exposure to tacrolimus and incidence of cancer after liver transplantation. *Am J Transplant* 2022; **22**: 1671-1682 [PMID: 35286761 DOI: 10.1111/ajt.17021]
- 55 **Carenco C**, Assenat E, Faure S, Duny Y, Danan G, Bismuth M, Herrero A, Jung B, Ursic-Bedoya J, Jaber S, Larrey D, Navarro F, Pageaux GP. Tacrolimus and the risk of solid cancers after liver transplant: a dose effect relationship. *Am J Transplant* 2015; **15**: 678-686 [PMID: 25648361 DOI: 10.1111/ajt.13018]
- 56 **Schmeding M**, Kiessling A, Neuhaus R, Heidenhain C, Bahra M, Neuhaus P, Neumann UP. Mycophenolate mofetil monotherapy in liver transplantation: 5-year follow-up of a prospective randomized trial. *Transplantation* 2011; **92**: 923-929 [PMID: 21832958 DOI: 10.1097/TP.0b013e31822d880d]
- 57 **Aguiar D**, Martínez-Urbistondo D, D'Avola D, Iñarrairaegui M, Pardo F, Rotellar F, Sangro B, Quiroga J, Herrero JI. Conversion from Calcineurin Inhibitor-Based Immunosuppression to Mycophenolate Mofetil in Monotherapy Reduces Risk of De Novo Malignancies After Liver Transplantation. *Ann Transplant* 2017; **22**: 141-147 [PMID: 28302995 DOI: 10.12659/aot.901556]
- 58 **European Association for the Study of the Liver**. EASL Clinical Practice Guidelines: Liver transplantation. *J Hepatol* 2016; **64**: 433-485 [PMID: 26597456 DOI: 10.1016/j.jhep.2015.10.006]
- 59 **Finkenstedt A**, Graziadei IW, Oberaigner W, Hilbe W, Nachbaur K, Mark W, Margreiter R, Vogel W. Extensive surveillance promotes early diagnosis and improved survival of de novo malignancies in liver transplant recipients. *Am J Transplant* 2009; **9**: 2355-2361 [PMID: 19663894 DOI: 10.1111/j.1600-6143.2009.02766.x]
- 60 **Herrero JI**, Alegre F, Quiroga J, Pardo F, Iñarrairaegui M, Sangro B, Rotellar F, Montiel C, Prieto J. Usefulness of a program of neoplasia surveillance in liver transplantation. A preliminary report. *Clin Transplant* 2009; **23**: 532-536 [PMID: 19681977 DOI: 10.1111/j.1399-0012.2008.00927.x]
- 61 **Lucey MR**, Terrault N, Ojo L, Hay JE, Neuberger J, Blumberg E, Teperman LW. Long-term management of the successful adult liver transplant: 2012 practice guideline by the American Association for the Study of Liver Diseases and the American Society of Transplantation. *Liver Transpl* 2013; **19**: 3-26 [PMID: 23281277 DOI: 10.1002/lt.23566]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: office@baishideng.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

