

Effect of anesthetics on gastric damage using two models of portal hypertension

Paula RS Câmara, Gisele P Moi, José Geraldo P Ferraz, José Murilo R Zeitune

Paula RS Câmara, Gisele P Moi, UNIVAG, University Center of Várzea Grande, Mato Grosso, Brazil

Gisele P Moi, Biochemistry Laboratory, Dental School of Piracicaba, State University of Campinas, Piracicaba, SP, Brazil

José Geraldo P Ferraz, Division of Gastroenterology, Faculty of Medicine, University of Calgary, Calgary, AB T2N 1N4, Canada

Paula RS Câmara, José Murilo R Zeitune, Division of Gastroenterology, Faculty of Medicine, State University of Campinas, Brazil

Author contributions: Câmara PRS, Ferraz JGP, and Zeitune JMR designed the research; Câmara PRS performed the research; Câmara PRS, Moi GP, Ferraz JGP and Zeitune JMR analyzed the data; Câmara PRS and Moi GP wrote the paper; Ferraz JGP and Zeitune JMR reviewed it; All the authors wrote this paper.

Supported by a Doctoral Grant from the Fundação de Amparo à Pesquisa do Estado do São Paulo (FAPESP), Brazil

Correspondence to: Paula RS Câmara, PhD, UNIVAG, University Center of Várzea Grande, Avenida Dom Orlando Chaves, 2655, Várzea Grande-MT, 78118-000, Brazil. endothelin@hotmail.com

Telephone: +55-65-36886000 **Fax:** +55-65-36886015

Received: January 20, 2010 **Revised:** February 16, 2010

Accepted: February 23, 2010

Published online: August 6, 2010

increased under KX anesthesia while GBF was reduced.

CONCLUSION: The use of KX anesthesia in experimental procedures involving cirrhotic rats (but not those with pure portal hypertension) is preferable to SP anesthesia.

© 2010 Baishideng. All rights reserved.

Key words: Sodium pentobarbitone; Portal hypertension; Cirrhosis; Gastric damage; Ketamine/xylazine

Peer reviewer: Reidar Fossmark, MD, PhD, Department of Gastroenterology and Hepatology, St. Olav's Hospital, Olav Kyrre's gate 17, Trondheim N-7006, Norway

Câmara PRS, Moi GP, Ferraz JGP, Zeitune JMR. Effect of anesthetics on gastric damage using two models of portal hypertension. *World J Gastrointest Pharmacol Ther* 2010; 1(4): 81-86 Available from: URL: <http://www.wjgnet.com/2150-5349/full/v1/i4/81.htm> DOI: <http://dx.doi.org/10.4292/wjgpt.v1.i4.81>

Abstract

AIM: To investigate the effect of sodium pentobarbitone (SP) or ketamine/xylazine (KX) anesthetics on acute gastric injury.

METHODS: Portal hypertension was induced by bile duct ligation (BDL) or portal vein stenosis (PVS). Ethanol (EtOH)-induced gastric damage was assessed using *ex vivo* gastric chamber experiments. Gastric blood flow (GBF) was also measured by laser doppler flowmetry.

RESULTS: EtOH-induced gastric damage was reduced in BDL rats under KX anesthesia in comparison to those under SP anesthesia. GBF dysfunction in fasted BDL rats was partially restored under KX anesthesia. In contrast, in fasted PVS rats, EtOH-induced gastric damage was

INTRODUCTION

The use of anesthetics can differentially affect a number of physiological parameters (cardiovascular, metabolic and hemodynamic) in experimental models^[1-3]. As such, the use of certain anesthetics can affect the outcome of studies and different therapeutic interventions may be influenced by different anesthetic approaches. Ketamine is usually used as a short-acting anesthetic and analgesic agent that induces a trance-like anesthetic state known as dissociative anesthesia in both animals and humans^[4]. Xylazine is considered safe when used alone or in combination with other anesthetics and analgesics such as ketamine or isoflurane in animal research. Co-administration of ketamine with xylazine (KX) is a routine anesthetic regimen for domestic and laboratory animals including mice and rats^[5]. The sedative and muscle-relaxing properties of xylazine are beneficial in reducing side

effects of ketamine such as tremor and muscle rigidity^[6,7]. Recent research has revealed that KX may influence some physiological responses to surgical procedures or drug effects in small laboratory animals. In comparison to other anesthetics, it has been reported that KX increases the infarct size in cerebral ischemia^[8], induces hyperglycemia in fed rodents^[9], reduces TNF- α expression in rat spleen^[10] and influences lipopolysaccharide-induced endotoxemia^[11]. Some anesthetics have cardio- or renoprotective effects which may be relevant to designing ischemia-reperfusion protocols^[12,13]. They can affect the systemic hemodynamic^[14] and may induce moderate to severe tissue damage^[15].

The laboratorial use of anesthetic agents such as ketamine associated with xylazine (KX) and sodium pentobarbitone (SP) could induce alterations of gastric blood flow (GBF) in portal hypertensive rats. In addition, special attention to GBF during anesthesia in the cirrhotic rats' model could be important since the anesthetic agents could be worsening the gastric mucosa damage. This study aimed to evaluate the KX and SP anesthetic effects on GBF and ethanol (EtOH)-induced gastric damage in portal hypertensive rats.

MATERIALS AND METHODS

Animals

Male Sprague-Dawley rats (200-250 g) provided by the Animal House of the State University of Campinas were used in all experiments. They were housed in plastic cages and had free access to water and standard pellet chow. The experimental protocols were approved by the Ethical Principles in Animal Research adopted by the Brazilian College for Animal Experimentation.

Induction of portal hypertension

Portal hypertension was induced in male Sprague-Dawley rats anesthetized with either a mix of ketamine plus xylazine (100:10 mg/kg; ip) or SP (25-50 mg/kg; ip) using two procedures. Prehepatic portal hypertension was provoked by portal vein stenosis (PVS) according to Chojkier *et al.*^[16] (1981). The portal vein was isolated and a stenosis created by placing a single ligature of a 3-0 silk around both the portal vein and a 20-gauge blunt-tipped needle. The needle was then removed from the ligature, creating a calibrated constriction of the portal vein. In another experimental group, biliary cirrhosis with intrahepatic portal hypertension was induced by common bile duct ligation (BDL) as described by Lee *et al.*^[17] (1986). Briefly, BDL was performed by isolating the common bile duct and placing two 4-0 sutures proximal/distal to the porta hepatis. Next, a 5-10 mm segment between ligatures was resected. The abdomen was then sutured with 4-0 silk and the animals were allowed to recover. Controls had a sham operation. Experiments were performed 2^[16] or 4^[17] wk following surgery when pure portal hypertension and liver cirrhosis were established respectively.

Ethanol-induced gastric damage

Gastric resistance to a 40% EtOH-induced injury was

investigated using an *ex vivo* gastric chamber preparation. Rats were fasted for 16 h prior to the procedure. Rats were anesthetized by SP or ketamine plus xylazine and placed over a heating pad connected to a rectal probe (Harvard Apparatus, Holliston, MS) to maintain body temperature at 37°C. The stomach was exposed following a midline laparotomy and opened with an incision along the greater curvature. It was pinned over a plexiglass platform and clamped with a plexiglass cylinder. The experiments consisted of six periods of 10 min each. The mucosa was bathed with phosphate buffered saline for two periods followed by 40% ethanol for one period and then HCl at pH 1.5 for three periods. All solutions were at 37°C when added to the chamber and continuously stirred at 200 r/m. The stomach was photographed at the end of experiments and damage (percent total glandular mucosa) analyzed using computerized planimetry by an observer blinded to the different treatment groups^[18].

Gastric blood flow measurement

GBF was measured using laser-Doppler flowmetry and the *ex vivo* gastric chamber preparation^[18]. A pencil probe (type N, penetration 1 mm, Transonic, Ithaca, NY) connected to a flowmeter (BLF 21D, Transonic, Ithaca, NY) was placed over the corpus. The preparation was equilibrated for 5 min. Basal GBF was then recorded for 5 min and experiments performed according to the protocol described previously^[17]. GBF was recorded throughout the 60 min and maximum changes in flow were expressed as percentage change over basal^[19].

Drugs/chemicals

All drugs were of analytical grade. SP (Hypnol[®], Cristalia), Xylazine (Rompun[®], Bayer) and ketamine (Ketalar[®], Parke-Davis) were used as clinically available preparations.

Statistical analysis

Data are expressed as mean \pm SEM and comparisons among groups ($n = 5$) were analyzed by one-way ANOVA followed by the Student's Newman-Keul's test for multiple comparisons. Statistical significance was considered when $P < 0.05$.

RESULTS

Effects of anesthetics on ethanol-induced injury in rats with portal hypertension

In control animals under SP or KX anesthesia, topical application of ethanol produced minimal gastric damage (20% \pm 3%; $n = 5$, Figure 1). This was associated with a reduction in GBF followed by a hyperemic response when the mucosa was bathed with HCl (Figure 2A).

In BDL rats under SP anesthesia, the EtOH-induced damage increased over the control value to 76% \pm 4% ($n = 5$; Figure 1). However significant ($P < 0.001$) protection in gastric damage was obtained when BDL rats were anesthetized with KX (50% \pm 6%; $n = 5$; Figure 1). This was associated with the development of a hyperemic response to ethanol in BDL.




Figure 1 Gastric damage in control and portal hypertensive rats. Ethanol (EtOH)-induced gastric damage was significantly increased in cirrhotic bile duct ligation (BDL) rats compared to controls when sodium pentobarbitone (SP) anesthetic was administered. In BDL rats, ketamine/xylazine (KX) anesthesia partially restored the resistance of the portal hypertensive gastric mucosa to ethanol-induced damage. An inverse effect was obtained from KX and SP anesthetics in portal vein stenosis rats. Asterisks denote significant differences between the two anesthesia treatments in BDL and portal vein stenosis rats ($n = 5$; ^a $P < 0.001$; ^b $P < 0.05$). PVS: Portal vein stenosis.

In contrast to SP anesthesia, higher EtOH-induced gastric damage was observed in PVS rats under KX anesthesia ($16\% \pm 3\%$ for KX *vs* $5\% \pm 2\%$ for SP; $n = 5$; Figure 1).

Effects of anesthetics on gastric blood flow

The effects of anesthetics on GBF responses submitted to 40% EtOH-induced injury were investigated in control and portal hypertensive rats. Control animals under SP or KX anesthesia did not show a difference in hyperemic responses ($n = 5$, Figure 2A). In BDL rats under SP anesthesia, any changes in GBF were observed throughout the experiments. On the other hand, KX anesthesia partially restored GBF dysfunction in BDL rats, contributing to the reduced gastric damage observed ($n = 5$, Figure 2B).

In contrast, KX anesthesia lowered GBF in PVS rats. This was associated with reduced hyperemic response to topical application of 40% ethanol over the gastric mucosa. However, under SP anesthesia GBF was increased in PVS rats ($n = 5$, Figure 2C).

DISCUSSION

This study demonstrates that sodium pentobarbitone (SP) and Ketamine/Xylazine (KX) anesthetics have significantly different effects on GBF in fasted portal hypertensive rats. The cirrhotic rats (e.g. BDL rats) treated under KX anesthesia had less gastric mucosal damage from ethanol stimulus. While the specific process by which KX anesthetics affect GBF is unknown, previous studies demonstrate that glucose levels are improved under treatment with KX anesthetics *via* α_2 -adrenoceptor activation^[19] and in turn the glucose acts to impair vagal activity in the stomach^[20]. Therefore, it suggests that the increase of glucose serves to increase the pressure of GBF.

Physiological changes in the blood glucose concentration have been reported to modulate gastrointestinal motor function^[21,22], particularly stomach motor function^[20]. For example, hyperglycemia has been documented to exert an

inhibitory effect on gastric function under a number of experimental conditions and in a wide range of species.

The mechanism responsible for hyperglycemia-induced inhibition of gastric motility remains unclear. In terms of mechanism, it has been proposed by several investigators that glucose acts to impair vagal activity to the stomach^[23-25]. In addition, there are some studies showing that systemically administered glucose produces a decrease in efferent activity of the gastric vagus nerve^[20,26,27]. Moreover, it has been proposed that glucose acts in the hepatic portal area to inhibit hepatic afferent nerve traffic to the brain which results in a reflex reduction in efferent activity of the gastric vagus nerve^[26,28]. It has also been proposed that glucose acts in the brain to alter control of vagal mediated gastric motility^[25,27,29,30].

Xylazine is analogue of clonidine and its effect as a α_2 -adrenergic agonist is known. It has been reported that activation of α_2 -adrenoceptor can increase, decrease or not change the levels of glucagon. Increased glucagon levels associated with decreased insulin levels may account for hyperglycemic effect of α_2 -adrenoceptor agonist^[31,32]. In rats, xylazine increases urine flow rate by activating of the α_2 -adrenergic receptors in hypothalamic paraventricular nucleus which in turn decreases vasopressin release^[33]. Saha *et al*^[9] (2005) demonstrated that KX anesthetic increases blood glucose in fasted rats. It suggests that the acute hyperglycemia effect of KX reflects, in part, α_2 -adrenoceptor-dependent changes of glucoregulatory hormones including insulin, growth hormone, adrenocorticotrophic hormone and corticosterone^[34,35]. Ketamine also induces changes in GBF. Rodrigues *et al*^[36] (2006) demonstrated that KX anesthesia reduced the constrictor effect of noradrenaline in mesenteric arterioles when compared to chloral hydrate anesthesia. The reduction may be explained by a direct effect of ketamine on vascular smooth muscle cells causing relaxation^[37,38]. Hence, this inhibition of contraction can be caused by agonists. It has been demonstrated in isolated aorta^[39] and in the mesenteric artery^[40]. These effects are primarily caused by the reduc-




Figure 2 Gastric blood flow responses by topical application of ethanol (40%) in control (A), bile duct ligation (B) and portal vein stenosis (C) rats. A: Under both sodium pentobarbitone and ketamine/xylazine anesthesia, a reduction in gastric blood flow (GBF) was observed in control rats followed by hyperemic response of the gastric microcirculation ($n = 5$); B: GBF responses were absent in bile duct ligation rats under sodium pentobarbitone anesthesia whereas ketamine/xylazine anesthesia partially restored GBF responses to ethanol ($n = 5$); C: GBF rates increased in portal vein stenosis rats under sodium pentobarbitone anesthesia while under ketamine/xylazine anesthesia was reduced ($n = 5$).

tion of calcium flow into cells by the inhibition of voltage dependent L-type calcium channels^[41]. Besides, the effect of ketamine reduces the endothelial release of nitric oxide (NO)^[42]. Despite noradrenaline promoting vasoconstriction, this effect was reduced in the KX-anesthetized rats. It occurred independent of the NO release reduction that it promoted by ketamine^[36].

The data of the research suggest that administration of KX in BDL rats partially restored the hyperemic gastric response to topical ethanol administration. In contrast, when SP was administered in BDL rats, any changes in hyporesponsive GBF followed topical ethanol adminis-

tration was observed.

The gastric mucosa of the PVS rats group responded differently to anesthetic treatments. Naturally, the PVS rats group has elevated GBF because of the presence of excessive vasoactive mediators^[43]. However, under KX anesthesia it was observed that the GBF is reduced in PVS rats whereas the high GBF of PVS rats under SP anesthesia was not altered. The high GBF protected the gastric mucosa from damage stimulus caused by topical ethanol administration.

PVS and common BDL are models used to evaluate portal hypertension in rats. However, the effects of ethanol on portal hypertensive gastric mucosa are different between these models. GBF differs between the models under SP anesthesia. Thus it is suspected that links exist between liver damage, the pentobarbital administration and levels of vasodilator mediators. Pentobarbital has important vascular effects in rats with portal hypertension and cirrhosis.

Studies have shown different results under the two models of portal hypertensive gastropathy: in the BDL model there is evidence of alterations of basal hemodynamic parameters which augment the gastric lesions^[17,19,44]; these lesions were reduced in the PVS model. The low damage resistance in the BDL group could be due to the fact that, at the same time, the liver is involved and the pentobarbital anesthesia was used.

In animal models, it is generally assumed that the physiological parameters under general anesthesia represent the basal state of the animal (before institution of the disease model). However, the anesthetic category can variably affect cardiovascular, neurohumoral and behavioral parameters. Similarly, if the fed and fasted states of animals are chosen arbitrarily, it may cause several changes in physiological parameters. Therefore, it is possible that the anesthetic category agents and/or fed and fasted states of animals used in the studies with different therapeutic interventions can influence the results.

This study suggests that are significant differences in the GBF and gastric damage after intraperitoneal injection of the anesthetic agents between PVS and common BDL models. In the latter model, the rats under SP anesthesia did not show any changes in GBF although it increased gastric damage. On the other hand, ketamine associated with xilazine (KX) anesthesia was able to partially restore GBF dysfunction in BDL rats, contributing to the reduced gastric damage.

The use of KX anesthesia in BDL model should be preferred to SP anesthesia, demonstrating that the choice of appropriate anesthetics should avoid misleading interpretation of experimental data.

Data from this research suggest that the choice of appropriate anesthetics in experimental models of portal hypertension present extreme relevance and should be taken in to consideration before the completion of these studies in order to avoid misleading interpretation of the data.

COMMENTS

Background

Portal vein obstruction can cause portal hypertension and other clinical disorders

such as severe hemorrhage. So, it is very important to prevent hemorrhages in gastrointestinal surgical procedures. In addition, many kinds of drugs have vasoactive actions such as vasodilatation or vasoconstriction. Anesthetics are vasoactive drugs and should be used carefully in gastrointestinal surgical procedures due to side effects on gastric mucosal blood flow. In this work, the better anesthetic to use in surgery of the gastric mucosa of portal hypertensive rats was suggested.

Research frontiers

Ketamine/xylazine (KX) and sodium pentobarbitone (SP) are two anesthetics for experimental and medical use. In the area of preventing gastric hemorrhage and simultaneously reduce gastric damage, the effects of these anesthetics on gastric blood flow (GBF) response to 40% ethanol-induced injury were investigated in control and portal hypertensive rats using laser doppler flowmetry and gastric chamber techniques.

Innovations and breakthroughs

In order to reduce gastric damage and improve GBF from portal hypertensive rats, the rat gastric mucosa was submitted to ethanol stimulus under two different kinds of anesthetics. Bleeding and gastric damage in bile duct ligation (BDL) rats were reduced under KX anesthesia but not in pure portal hypertension. On the other hand, bleeding and gastric damage were increased in BDL rats under SP anesthesia while the GBF was reduced. The present study shows that the use of inappropriate anesthetic can aggravate gastric damage in portal hypertensive rats.

Applications

The results of this study suggest that the use of KX in experimental procedures involving cirrhotic rats is preferable to SP anesthesia, demonstrating that the choice of appropriate anesthetics should avoid misleading interpretations of experimental data.

Terminology

Portal Hypertensive Gastropathy (PHG) is a subclinical entity from portal hypertension and it can cause severe hemorrhage. BDL and portal vein stenosis are two experimental models to induce portal hypertension in rats. KX and SP are two anesthetics for experimental and medical use.

Peer review

This is a good descriptive study in which the authors analyze the effect of two different kinds of anesthetic drugs on gastric damage induced by ethanol in rats. The results are interesting and suggest that KX is a potential anesthetic that could be the better choice to reduce gastric hemorrhage in surgical procedures in PHG.

REFERENCES

- 1 **Wixson SK**, White WJ, Hughes HC Jr, Lang CM, Marshall WK. The effects of pentobarbital, fentanyl-droperidol, ketamine-xylazine and ketamine-diazepam on arterial blood pH, blood gases, mean arterial blood pressure and heart rate in adult male rats. *Lab Anim Sci* 1987; **37**: 736-742
- 2 **Rastaldo R**, Penna C, Pagliaro P. Comparison between the effects of pentobarbital or ketamine/nitrous oxide anesthesia on metabolic and endothelial components of coronary reactive hyperemia. *Life Sci* 2001; **69**: 729-738
- 3 **Brookes ZL**, Brown NJ, Reilly CS. Response of the rat cremaster microcirculation to hemorrhage in vivo: differential effects of intravenous anesthetic agents. *Shock* 2002; **18**: 542-548
- 4 **Bergman SA**. Ketamine: review of its pharmacology and its use in pediatric anesthesia. *Anesth Prog* 1999; **46**: 10-20
- 5 **Sumitra M**, Manikandan P, Rao KV, Nayeem M, Manohar BM, Puvanakrishnan R. Cardiorespiratory effects of diazepam-ketamine, xylazine-ketamine and thiopentone anesthesia in male Wistar rats—a comparative analysis. *Life Sci* 2004; **75**: 1887-1896
- 6 **Wright M**. Pharmacologic effects of ketamine and its use in veterinary medicine. *J Am Vet Med Assoc* 1982; **180**: 1462-1471
- 7 **Lei H**, Grinberg O, Nwaigwe CI, Hou HG, Williams H, Swartz HM, Dunn JF. The effects of ketamine-xylazine anesthesia on cerebral blood flow and oxygenation observed using nuclear magnetic resonance perfusion imaging and electron paramagnetic resonance oximetry. *Brain Res* 2001; **913**: 174-179
- 8 **Kawai N**, Keep RF, Betz AL. Hyperglycemia and the vascular effects of cerebral ischemia. *Stroke* 1997; **28**: 149-154
- 9 **Saha JK**, Xia J, Grondin JM, Engle SK, Jakubowski JA. Acute hyperglycemia induced by ketamine/xylazine anesthesia in rats: mechanisms and implications for preclinical models. *Exp Biol Med* (Maywood) 2005; **230**: 777-784
- 10 **Bette M**, Schlimme S, Mutters R, Menendez S, Hoffmann S, Schulz S. Influence of different anaesthetics on pro-inflammatory cytokine expression in rat spleen. *Lab Anim* 2004; **38**: 272-279
- 11 **Helmer KS**, Cui Y, Chang L, Dewan A, Mercer DW. Effects of ketamine/xylazine on expression of tumor necrosis factor-alpha, inducible nitric oxide synthase, and cyclo-oxygenase-2 in rat gastric mucosa during endotoxemia. *Shock* 2003; **20**: 63-69
- 12 **Saha DC**, Saha AC, Malik G, Astiz ME, Rackow EC. Comparison of cardiovascular effects of tiletamine-zolazepam, pentobarbital, and ketamine-xylazine in male rats. *J Am Assoc Lab Anim Sci* 2007; **46**: 74-80
- 13 **Lee HT**, Ota-Setlik A, Fu Y, Nasr SH, Emala CW. Differential protective effects of volatile anesthetics against renal ischemia-reperfusion injury in vivo. *Anesthesiology* 2004; **101**: 1313-1324
- 14 **Janssen BJ**, De Celle T, Debets JJ, Brouns AE, Callahan MF, Smith TL. Effects of anesthetics on systemic hemodynamics in mice. *Am J Physiol Heart Circ Physiol* 2004; **287**: H1618-H1624
- 15 **Smiler KL**, Stein S, Hrapkiewicz KL, Hibben JR. Tissue response to intramuscular and intraperitoneal injections of ketamine and xylazine in rats. *Lab Anim Sci* 1990; **40**: 60-64
- 16 **Chojkier M**, Groszmann RJ. Measurement of portal-systemic shunting in the rat by using gamma-labeled microspheres. *Am J Physiol* 1981; **240**: G371-G375
- 17 **Lee SS**, Braillon A, Girod C, Geoffroy P, Lebrec D. Haemodynamic rebound phenomena after abrupt cessation of propranolol therapy in portal hypertensive rats. *J Hepatol* 1986; **3**: 38-41
- 18 **Ferraz JG**, Tigley AW, Appleyard CB, Wallace JL. TNF-alpha contributes to the pathogenesis of ethanol-induced gastric damage in cirrhotic rats. *Am J Physiol* 1997; **272**: G809-G814
- 19 **Camara PR**, Ferraz GJ, Franco-Penteado CF, Sbragia-Neto L, Meirelles LR, Teixeira SA, Muscara MN, Velloso LA, Antunes E, Ferraz JG. Ablation of primary afferent neurons by neonatal capsaicin treatment reduces the susceptibility of the portal hypertensive gastric mucosa to ethanol-induced injury in cirrhotic rats. *Eur J Pharmacol* 2008; **589**: 245-250
- 20 **Shi M**, Jones AR, Niedringhaus MS, Pearson RJ, Biehl AM, Ferreira M Jr, Sahibzada N, Verbalis JG, Gillis RA. Glucose acts in the CNS to regulate gastric motility during hypoglycemia. *Am J Physiol Regul Integr Comp Physiol* 2003; **285**: R1192-R1202
- 21 **de Boer SY**, Masclee AA, Lamers CB. Effect of hyperglycemia on gastrointestinal and gallbladder motility. *Scand J Gastroenterol Suppl* 1992; **194**: 13-18
- 22 **Jones KL**, Kong MF, Berry MK, Rayner CK, Adamson U, Horowitz M. The effect of erythromycin on gastric emptying is modified by physiological changes in the blood glucose concentration. *Am J Gastroenterol* 1999; **94**: 2074-2079
- 23 **Samsom M**, Akkermans LM, Jebbink RJ, van Isselt H, van Berge-Henegouwen GP, Smout AJ. Gastrointestinal motor mechanisms in hyperglycaemia induced delayed gastric emptying in type I diabetes mellitus. *Gut* 1997; **40**: 641-646
- 24 **Ishiguchi T**, Nakajima M, Sone H, Tada H, Kumagai AK, Takahashi T. Gastric distension-induced pyloric relaxation: central nervous system regulation and effects of acute hyperglycaemia in the rat. *J Physiol* 2001; **533**: 801-813
- 25 **Ishiguchi T**, Tada H, Nakagawa K, Yamamura T, Takahashi T. Hyperglycemia impairs antro-pyloric coordination and delays gastric emptying in conscious rats. *Autonomic Neuroscience: Basic and Clinical* 2002; **95**: 112-120
- 26 **Sakaguchi T**, Yamaguchi K. Changes in efferent activities of the gastric vagus nerve by administration of glucose in the portal vein. *Experientia* 1979; **35**: 875-876
- 27 **Hirano T**, Niiijima A. Effects of 2-deoxy-D-glucose, glucose and insulin on efferent activity in gastric vagus nerve. *Exper-*

- entia* 1980; **36**: 1197-1198
- 28 **Sakaguchi T**, Shimojo E. Inhibition of gastric motility induced by hepatic portal injections of D-glucose and its anomers. *J Physiol* 1984; **351**: 573-581
 - 29 **Sakaguchi T**, Taguchi T, Okuda J. Different effects of D-glucose anomers on enhanced motility of the stomach. *Biochem Int* 1983; **7**: 299-305
 - 30 **Ferreira M Jr**, Browning KN, Sahibzada N, Verbalis JG, Gillis RA, Travagli RA. Glucose effects on gastric motility and tone evoked from the rat dorsal vagal complex. *J Physiol* 2001; **536**: 141-152
 - 31 **Saito M**, Saitoh T, Inoue S. Alpha 2-adrenergic modulation of pancreatic glucagon secretion in rats. *Physiol Behav* 1992; **51**: 1165-1171
 - 32 **Velliquette RA**, Ernsberger P. The role of I(1)-imidazoline and alpha(2)-adrenergic receptors in the modulation of glucose metabolism in the spontaneously hypertensive obese rat model of metabolic syndrome X. *J Pharmacol Exp Ther* 2003; **306**: 646-657
 - 33 **Cabral AD**, Kapusta DR, Kenigs VA, Varner KJ. Central alpha2-receptor mechanisms contribute to enhanced renal responses during ketamine-xylazine anesthesia. *Am J Physiol* 1998; **275**: R1867-R1874
 - 34 **Kasuya E**, Hodate K, Matsumoto M, Sakaguchi M, Hashizume T, Kanematsu S. The effects of xylazine on plasma concentrations of growth hormone, insulin-like growth factor-I, glucose and insulin in calves. *Endocr J* 1996; **43**: 145-149
 - 35 **Hampshire J**, Altszuler N. Clonidine or xylazine as provocative tests for growth hormone secretion in the dog. *Am J Vet Res* 1981; **42**: 1073-1076
 - 36 **Rodrigues SF**, de Oliveira MA, Martins JO, Sannomiya P, de Cássia Tostes R, Nigro D, Carvalho MH, Fortes ZB. Differential effects of chloral hydrate- and ketamine/xylazine-induced anesthesia by the s.c. route. *Life Sci* 2006; **79**: 1630-1637
 - 37 **Longnecker DE**, Miller FN, Harris PD. Small artery and vein response to ketamine HCl in the bat wing. *Anesth Analg* 1974; **53**: 64-68
 - 38 **Wendling WW**, Daniels FB, Chen D, Harakal C, Carlsson C. Ketamine directly dilates bovine cerebral arteries by acting as a calcium entry blocker. *J Neurosurg Anesthesiol* 1994; **6**: 186-192
 - 39 **Yamanaka I**, Dowdy EG. The effects of ketamine on spiral-cut strips of rabbit aorta. *Anesthesiology* 1974; **40**: 222-227
 - 40 **Kanmura Y**, Kajikuri J, Itoh T, Yoshitake J. Effects of ketamine on contraction and synthesis of inositol 1,4,5-trisphosphate in smooth muscle of the rabbit mesenteric artery. *Anesthesiology* 1993; **79**: 571-579
 - 41 **Yamazaki M**, Ito Y, Kuze S, Shibuya N, Momose Y. Effects of ketamine on voltage-dependent Ca²⁺ currents in single smooth muscle cells from rabbit portal vein. *Pharmacology* 1992; **45**: 162-169
 - 42 **Miyawaki I**, Nakamura K, Terasako K, Toda H, Kakuyama M, Mori K. Modification of endothelium-dependent relaxation by propofol, ketamine, and midazolam. *Anesth Analg* 1995; **81**: 474-479
 - 43 **Ohta M**, Kishihara F, Hashizume M, Kawanaka H, Tomikawa M, Higashi H, Tanoue K, Sugimachi K. Increased prostacyclin content in gastric mucosa of cirrhotic patients with portal hypertensive gastropathy. *Prostaglandins Leukot Essent Fatty Acids* 1995; **53**: 41-45
 - 44 **Lee SS**, Girod C, Valla D, Geoffroy P, Lebrec D. Effects of pentobarbital sodium anesthesia on splanchnic hemodynamics of normal and portal-hypertensive rats. *Am J Physiol* 1985; **249**: G528-G532

S- Editor Li LF L- Editor Roemmele A E- Editor Yang C