

Anticancer actions of PPAR γ ligands: Current state and future perspectives in human lung cancer

Shou Wei Han, Jesse Roman

Shou Wei Han, Jesse Roman, Division of Pulmonary, Critical Care and Sleep Disorders Medicine, Department of Medicine, University of Louisville School of Medicine, Louisville, KY 40202, United States

Jesse Roman, Louisville Veterans Affairs Medical Center, Louisville, KY, 40202, United States

Author contributions: Han SW wrote the paper; Roman J edited the paper.

Supported by The National Institutes of Health CA123104 (Han SW) and CA116812 (Roman J)

Correspondence to: Shou Wei Han, MD, PhD, Associate Professor, Division of Pulmonary, Critical Care and Sleep Disorders Medicine, Department of Medicine, University of Louisville School of Medicine, CTR Building, Room 524, 505 Hancock St., Louisville, KY 40202, United States. sw.han@louisville.edu

Telephone: +1-505-8528468 Fax: +1-505-8526233

Received: March 17, 2010 Revised: March 23, 2010

Accepted: March 24, 2010

Published online: March 26, 2010

Abstract

Peroxisome proliferator-activated receptors (PPARs) are ligand-dependent nuclear transcription factors and members of the nuclear receptor superfamily. Of the three PPARs identified to date (PPAR γ , PPAR β/δ , and PPAR α), PPAR γ has been studied the most, in part because of the availability of PPAR γ agonists (also known as PPAR γ ligands) and its significant effects on the management of several human diseases including type 2 diabetes, metabolic syndrome, cardiovascular disease and cancers. PPAR γ is expressed in many tumors including lung cancer, and its function has been linked to the process of lung cancer development, progression and metastasis. Studies performed in gynogenic and xenograft models of lung cancer showed decreased tumor growth and metastasis in animals treated with PPAR γ ligands. Furthermore, data are emerging from retrospective clinical studies that suggest a protective role for PPAR γ ligands on the incidence of lung cancer. This review summarizes the

research being conducted in this area and focuses on the mechanisms and potential therapeutic effects of PPAR γ ligands as a novel anti-lung cancer treatment strategy.

© 2010 Baishideng. All rights reserved.

Key words: Gene expression and regulation; Human lung cancer; Ligands; Peroxisome proliferator-activated receptor γ ; Signaling pathways; Therapy

Peer reviewers: Jongsun Park, Associate Professor, Department of Pharmacology, Cancer Research Institute, College of Medicine, Chungnam National University, 6 Munhwa-dong, Jung-gu, Taejeon 301-131, South Korea; Johan Lennartsson, PhD, Ludwig Institute for Cancer Research, Uppsala University, Box 595, SE-751 24 Uppsala, Sweden

Han SW, Roman J. Anticancer actions of PPAR γ ligands: Current state and future perspectives in human lung cancer. *World J Biol Chem* 2010; 1(3): 31-40 Available from: URL: <http://www.wjgnet.com/1949-8454/full/v1/i3/31.htm> DOI: <http://dx.doi.org/10.4331/wjbc.v1.i3.31>

INTRODUCTION

Lung carcinoma is the most common malignant tumor in the world, and is the leading cause of carcinoma death in the United States^[1]. This malignancy causes more deaths than the next three most common cancers combined (colon, breast and prostate). The expected 5-year survival rate for all patients in whom lung cancer is diagnosed is less than 13% compared to 65% for colon, 89% for breast, and 99% for prostate cancer although incremental and significant advances in available systemic treatments have taken place in the last decade to improve survival rates and to provide better palliation for patients with non-small-cell (NSCLC) and small-cell lung carcinoma (SCLC). Cigarette smoking is strongly correlated with the onset of lung cancer and effective tobacco con-

trol efforts have resulted in substantial declines in tobacco use and tobacco-related cancer deaths in the United States^[2]. Clinical approaches such as chemo- and radiotherapies have shown only a modest improvement in survival of patients with advanced NSCLC and limited-stage SCLC. However, agents targeting specific kinases and growth factor receptors show promise. For example, Erlotinib, an epidermal growth factor receptor (EGFR) tyrosine kinase inhibitor, and bevacizumab, a humanized monoclonal antibody that recognizes and blocks vascular endothelial growth factor A are currently used in medical practice and have resulted in improved survival^[3]. Unfortunately, targeted therapies which are initially effective in certain small subpopulations of patients, eventually fail to control the tumor. More recently, genomic and proteomic studies have unveiled a means for the molecular profiling of tumor tissue from patients with NSCLC, and could allow tailoring of therapy. Although there are still significant challenges to implementing genomic and proteomic testing in clinical practice, the rapid development of newer technologies provides hope for overcoming these barriers^[4]. The limitations in efficacy and safety associated with available treatments for lung cancer especially NSCLC underscore the need for novel agents with improved efficacy and safety profiles. Therefore, understanding and searching for novel molecular mechanisms responsible for lung cancer initiation and proliferation are needed to identify new targets for therapy.

Since their discovery in 1990, peroxisome proliferator-activated receptors (also known as PPARs) have emerged as potential targets for anti-cancer therapies. Although originally cloned in an attempt to identify the molecular mediators of peroxisome proliferation in the liver of rodents, PPARs are now recognized as versatile members of the ligand-activated nuclear hormone receptor superfamily of transcription factors that includes receptors for steroids, thyroid hormone, retinoic acid, and vitamin D, among others^[5]. PPARs are considered to play key roles in diverse physiological processes ranging from lipid metabolism to inflammation, and have been implicated in diseases such as cancer, atherosclerosis, and diabetes^[5,6]. Although information about the function of PPARs in lung is scarce, data implicating these molecules in key processes in lung biology are rapidly emerging.

Three subtypes of PPARs have been identified and cloned: PPAR α , PPAR β/δ , and PPAR γ . These subtypes are distinguished by their tissue distribution, and to a lesser degree, by their ligand specificity. PPAR α has been implicated in hepatocellular carcinoma in rodents, whereas activation of PPAR β/δ promotes human lung carcinoma cell proliferation through phosphatidylinositol 3-kinase/Akt activation^[7-9]. Of the three PPARs identified to date, PPAR γ represents the most promising target in view of the many reports implicating this molecule in lung carcinoma cell growth. As a tumor growth modifier, PPAR γ is involved in the regulation of cancer cell apoptosis, proliferation, and differentiation, and through

its actions on the tumor cell environment, it affects angiogenesis, inflammation, and immune cell functions^[10]. Hence, many studies are underway to test the impact of targeting this receptor for therapeutic purposes. This review focuses on PPAR γ , its role in lung carcinogenesis, and the potential therapeutic role of PPAR γ agonists in lung cancer.

FUNCTION OF PPAR γ

PPAR γ was discovered based on its similarity to PPAR α . By utilizing three different promoters, a single PPAR γ gene encodes three isoforms namely PPAR γ 1, PPAR γ 2 and PPAR γ 3^[11]. Analysis of PPAR γ 1 and γ 3 transcripts revealed that they both translate into the same PPAR γ 1 protein^[12]. PPAR γ 2 protein contains an additional 30 amino acids at its N-terminus compared to PPAR γ 1 (Figure 1A). Like all nuclear receptors, PPAR γ shares a similar structure with functional domains called A/B (ligand-independent domain), C (DNA binding domain), D (hinge domain) and E-F (ligand binding domain) (Figure 1B). PPAR γ is highly expressed in adipose tissue and it is a master regulator of adipocyte differentiation^[13,14]. In addition to its role in adipogenesis, PPAR γ serves as an important transcriptional regulator of glucose and lipid metabolism, and it has been implicated in the regulation of insulin sensitivity, atherosclerosis, and inflammation^[15,16]. PPAR γ is also expressed in multiple tissues such as breast, colon, lung, ovary, prostate, and thyroid where it was demonstrated to regulate cellular proliferation, differentiation, and apoptosis^[17,18]. Several leukocyte populations, including monocytes/macrophages, lymphocytes, and dendritic cells, have also been shown to express PPAR γ suggesting a role for this molecule in the regulation of immune responses^[19]. In that regard, PPAR γ appears to be a negative regulator of macrophage function since its activation suppresses the production of inflammatory cytokines, chemokines, metalloproteases, and nitric oxide^[20,21]. These PPAR γ -mediated anti-inflammatory effects are not restricted to monocytes, as treatment with PPAR γ agonists results in the inhibition of cytokine/chemokine production in several epithelial and stromal cells^[22].

Several natural and synthetic compounds have been identified as activators of PPAR γ . The insulin sensitizing anti-diabetic drugs known as *thiazolidinediones* (TZDs) were the first compounds identified as PPAR γ agonists^[23]. The TZDs, rosiglitazone and pioglitazone, are currently in clinical use for the treatment of type-II diabetes, while troglitazone was withdrawn from clinical use because it was linked to idiosyncratic liver toxicity^[24]. Other non-TZD synthetic ligands include certain non-steroidal anti-inflammatory drugs such as isoxzolidinedione JTT-501^[25], tyrosine-based GW7845^[26] and DH9, a newly synthesized PPAR γ agonist^[27]. Naturally occurring compounds that activate PPAR γ include long chain polyunsaturated fatty acids which are found in fish oil (e.g. n-3-PUFA, n-6-PUFA), eicosanoids




Figure 1 Structure of human peroxisome proliferator-activated receptor (PPAR) γ protein isoforms and domains. A: AR proteins. The three subtypes of mRNAs give rise to two different PPAR γ proteins. Transcription of the PPAR γ 1 and 3 promoters result in the same protein of 477 amino acids (aa). The PPAR γ 2 protein of 507 amino acids is produced by transcription from the promoter γ 2 area; B: main structure of PPAR γ . PPARs contain the following functional regions: an N-terminal A/B domain (ligand-independent domain), a C-domain (DNA-binding domain), a D-domain (hinge domain), and a C-terminal domain (E-F ligand-dependent domain).

[e.g. 15-deoxy- $\Delta^{12,14}$ prostaglandin J₂ (15d-PGJ₂)], lipid hydroperoxides [e.g. 9(s)-HODE and 13(s)-HODE], as well as 15d-PGJ₂ and 12/15 lipoxygenase products 15-hydroxyeicosatetraenoic acid (15-HETE) and 13-hydroxyoctadecadienoic acid (13-HETE)^[28-30]. Compounds from several medicinal plants such as Saurufuran A from *Saururus chinensis*^[31], flavonoids such as chrysin and kampferol^[32], phenolic compounds from *Glycyrrhiza uralensis*^[33], and curcumin from *Curcumin longa*^[34,35] have also been shown to activate PPAR γ . Recently, several new compounds such as (S)-3-{4-[3-(5-methyl-2-phenyl-oxazol-4-yl)-propyl]-phenyl}-2-1,2,3-triazol-2-yl-propionic acid (17), a new series of 2-aryloxy-3-phenyl-propanoic acids and *aleglitazar* were identified as potent human PPAR α/γ dual agonists with demonstrated oral bioavailability and certain encouraging responses^[36-38].

The synthetic ligands described above and some natural ligands have been used to elucidate the role of PPAR γ in cellular functions both *in vitro* and *in vivo*. However, several caveats should be taken into consideration when interpreting such studies. First, the natural ligands that regulate PPARs *in vivo* have not been completely elucidated. Second, not all PPAR γ ligands exert their effects through PPAR γ since there is strong evidence for the activation of PPAR γ independent signals, particularly with the natural ligand 15d-PGJ₂, among others^[39-41]. Third, high affinity ligands for PPAR γ (e.g. the TZDs) may exert partial agonist/antagonist activity^[42]. The latter might be due to the fact that individual TZDs induce different PPAR γ conformations that influence the recruitment of different coactivator/corepressor molecules. Thus, the activity of the PPAR γ transcriptional complex is influenced by the context of a given gene and its promoter, and by the relative availability of pertinent coactivator/corepressor molecules in the cell or tissue of interest.

PPAR γ AND PPAR γ LIGANDS IN LUNG CANCER

Among the three PPAR subtypes, the role of PPAR γ has been investigated the most in lung cancer occurrence, progression and therapy. PPAR γ is expressed in many cancers including colon, breast, and prostate, and with few exceptions, PPAR γ ligands are antiproliferative in these tumor cells. Similarly, PPAR γ is expressed in both SCLC and non-SCLC (NSCLC)^[43]. NSCLC accounts for 80% of malignant lung cancer and SCLC constitutes the remainder^[44]. Based on the cellular phenotype, NSCLC is further subdivided into squamous cell carcinoma, adenocarcinoma, and large cell carcinomas^[45]. SCLC tumors grow rapidly, and are more likely to metastasize earlier than NSCLC. PPAR γ ligands induce growth arrest and promote changes associated with differentiation as well as apoptosis in a variety of lung carcinoma cell lines, although most of the knowledge available in this area has been generated in NSCLC^[46,47]. One recent animal study demonstrated a reduction of endogenous PPAR γ ligands coinciding with increased PPAR α before the formation of lung tumors induced by treatment with 4-(methylnitrosamino)-1-(3-pyridyl)-lbutanone (NNK)^[48]. These results suggest that increased PPAR γ activity by its ligands and inhibition of PPAR α could prevent the formation of lung tumors and/or enhance the effectiveness of therapy against lung cancer^[48]. This study also suggests the possibility of using endogenous PPAR γ ligands such as 13-HETE and 15-HETE as tumor markers for lung cancer.

The exact mechanisms linking modulation of PPAR γ to cancer growth inhibition remain unclear, but include effects on transcription factors and gene expression, among others. In addition, current evidence suggests that PPAR γ ligands affect the intracellular machinery involved in cell signaling and cell cycle control, the suppression of mitogenic factors and tumor promoters, the induction of tumor suppressors, the prevention of tumor cell recognition of extracellular mitogenic signals, the break down of nicotine and nicotinic acetylcholine receptor-induced cell survival, and the expression of angiogenic factors needed for the development of the vascular networks that supply tumor cells (Figure 2)^[49]. These mechanisms are discussed below as they relate to the actions of PPAR γ ligands in lung cancer.

PPAR γ ligands, cell cycle progression and apoptotic-signaling pathways

Several studies demonstrate that PPAR γ ligands affect apoptosis and cell cycle control in lung cancer cells. For example, PPAR γ ligands have been found to inhibit the growth of A549 adenocarcinoma cells due to G0/G1 cell cycle arrest through the upregulation of mitogen-activated protein kinases extracellular signal-regulated kinases 1 and 2 (ERK1/2) and the downregulation of G1 cyclins D and E^[22]. Troglitazone inhibits NSCLC proliferation in




Figure 2 Anti-lung cancer actions of PPAR γ ligands. Through PPAR γ -dependent and -independent signals, PPAR γ ligands activate or inactivate (mostly) kinase signaling pathways [e.g. stress-activated protein kinase/c-Jun NH2-terminal kinase (SAPK/JNK), integrin-linked kinase (ILK), phosphatidylinositol 3-kinase (PI3-K)/Akt/GSK-3 β , extracellular signal-regulated kinases 1 and 2 (ERK1/2), p38MAPK, Src, FAK, and AMP-activated protein kinase (AMPK)/tuberous sclerosis complex 2 (TSC2)/mammalian target of rapamycin (mTOR)/p70S6K]. This results in the regulation of multiple depicted downstream effectors including expression of growth factors, tumor promoters, cytokines, chemokines, cell cycle control genes, nicotinic acetylcholine receptors, apoptotic genes, expression of tumor suppressor gene through inhibition or induction of transcription factors [e.g. Sp1, AP-1, AP-2, nuclear factor- κ B (NF- κ B), CRE, *etc.*]. These effects contribute to the inhibition of cell growth and induction of apoptosis in human lung cancer cells. Note that PPAR γ signaling has also been associated with tumor promoter activity in some cancer cells such as colon and breast, and this was linked to increased β -catenin, c-Myc, cyclin D1, vascular endothelial growth factor (VEGF), Angpt4 and Wnt 5 expression. COX: Cyclooxygenase; TGF: Transforming growth factor; DR-5: Death receptor 5; MMP: Matrix metalloproteinase; C-FLIP: Cellular FLICE inhibitory protein.

part by stimulating the expression of the GADD 153 (for *growth arrest and DNA damage inducible gene-153*)^[50]. Also, troglitazone was found to induce apoptosis in NCI-H23 cells *via* a mitochondrial pathway through the activation of ERK1/2^[51]. Others have shown similar results using CRL-202 cells, and further demonstrated that troglitazone downregulated the expression of the anti-apoptotic molecules Bcl-w and Bcl-2 as well as decreased the activity of stress-activated protein kinase (SAPK)/c-Jun NH2-terminal kinase (JNK)^[52]. PPAR γ ligands also induce the expression of death receptor 5 (DR5) and increase DR5 distribution at the cell surface in addition to reducing cellular FLICE-like inhibitory protein levels in human lung cancer cells. These agents cooperated with tumor necrosis factor-related apoptosis-inducing ligand to enhance apoptosis in human lung carcinoma cells^[53]. One report found that PPAR γ ligands 1-[(trans-methylimino-N-oxy)-6-(2-morpholinoethoxy)-3-phenyl-(1H-indene-2-carboxylic acid ethyl ester (KR-62980))] and rosiglitazone induce NSCLC apoptotic cell death mainly through PPAR γ -dependent reactive oxygen species formation *via* increased expression of proline oxidase, a redox enzyme expressed in mitochondria^[46].

PPAR γ ligands and kinase signaling pathways

Reports implicate alterations in the mammalian target of rapamycin (mTOR) signaling pathway in the anti-tumor effects of PPAR γ ligands. Rosiglitazone, for example, was reported to reduce the phosphorylation of Akt, an upstream positive modulator of mTOR, and increase phosphatase and tensin homolog deleted on chromosome 10 (PTEN), a negative modulator of mTOR, in NSCLC H1792 and H1838 cells; this resulted in inhibition of cell proliferation^[54]. Although the effects of rosiglitazone on Akt and PTEN were blocked by the selective PPAR γ antagonist GW9662 and restored by transient overexpression of PPAR γ , cell growth was not entirely restored suggesting the involvement of additional PPAR γ -independent mechanisms of action. Further work revealed that rosiglitazone increased the phosphorylation of AMP-activated protein kinase α , a target of LKB1, and tuberous sclerosis complex 2 (TSC2), another potential tumor suppressor and upstream downregulator of mTOR. The latter pathway was independent of PPAR γ since GW9662 and PPAR γ siRNA did not affect it^[54,55]; others have shown similar increases in PTEN expression induced by rosiglitazone^[56].

One recent study showed that troglitazone may bind directly to EGFR, inhibit its signaling, and stimulate its internalization independent of PPAR γ in several cells including lung cancer cells^[41]. In that work, inhibition of EGF-induced Akt phosphorylation most likely accounted for the growth arrest of lung cancer cells treated with troglitazone^[41].

Tumor suppressor genes are also affected by PPAR γ ligands. For example, PGJ₂ and ciglitazone stimulated the expression of p21 mRNA and protein expression in NSCLC, and this coincided with a reduction in cyclin D1 mRNA expression^[57]. Of note, p21 antisense oligonucleotides significantly blocked lung carcinoma cell growth inhibition observed with PPAR γ ligands thereby establishing an important role for p21 in this process. These findings are consistent with those of others showing that the proliferation of A549 cells injected subcutaneously into nude mice was inhibited significantly by treatment with ciglitazone, and this coincided with increased expression of PPAR γ and p21, and with downregulation of cyclin D1^[58]. A connection between another tumor suppressor gene, p53, and PPAR γ ligands has also been demonstrated by showing that 15-deoxy-PGJ₂, together with docetaxel, stimulates apoptosis in NSCLC through inhibition of Bcl-2 and cyclin D1, and overexpression of caspases and p53^[47].

More recently, we reported that rosiglitazone and dietary compounds such as fish oil (which contain certain kinds of fatty acids such as ω 3 and ω 6 polyunsaturated fatty acids known to work as PPAR γ ligands) inhibit integrin-linked kinase (ILK) expression through PPAR γ signaling and the recruitment of a PPAR γ co-activator, PGC-1 α ^[59]. ILK is a unique intracellular adaptor and kinase that links cell-adhesion receptors, integrins, and growth factors to the actin cytoskeleton and to a range

of signaling pathways that are implicated in the regulation of anchorage-dependent tumor cell growth/survival, cell cycle progression, invasion and migration, and tumor angiogenesis^[60]. This effect was associated with activation of p38 MAPK followed by induction of the transcription factor AP-2 α and, ultimately, inhibition of NSCLC cell proliferation^[59]. Docosahexaenoic acid, a component of ω 3 polyunsaturated fatty acid, is reported to inhibit the growth of lung cancer cells mainly through the induction of pro-apoptotic signaling pathways such as ERK1/2 and p38 MAPK suggesting its chemopreventive effect in lung cancer^[61].

PPAR γ ligands and cyclooxygenase-2-related pathways

PPAR γ ligands also exert anti-tumor effects by blocking access to mitogenic agents such as prostaglandin E2 (PGE₂), a major cyclooxygenase (COX) metabolite that plays important roles in tumor biology. The functions of PGE₂ are mediated through one or more of its receptors EP1, EP2, EP3, and EP4^[62]. Human NSCLC cell lines express EP2 receptors, among other EP receptors, and the inhibition of cell growth by PPAR γ ligands like GW1929, PGJ₂, ciglitazone, troglitazone, and rosiglitazone, is associated with a significant decrease in EP2 mRNA and protein expression. Notably, the inhibitory effects of rosiglitazone and ciglitazone, but not PGJ₂, were reversed by a specific PPAR γ antagonist GW9662, suggesting the involvement of PPAR γ -dependent and -independent mechanisms^[62]. Also, ciglitazone suppressed COX-2 mRNA expression and COX-2 promoter activity, while upregulating peroxisome proliferator response element promoter activity in NSCLC cells further suggesting a negative modulator role for PPAR γ ligands on the COX-2/PGE₂ pathway in NSCLC^[63]. Of note, *in vitro* studies and xenograft models have demonstrated that elevated COX-2 expression is critical for promoting lung tumorigenesis, and that the anti-tumorigenic effects of PPAR γ ligands are mediated through suppression of COX-2 *via* increased activity of PTEN, decreased levels of phospho-Akt, and inhibition of nuclear factor- κ B (NF- κ B) activity^[64].

PPAR γ and tobacco-related cancer progression

Tobacco is the most common etiologic agent in lung cancer worldwide. Recently, attention has been focused on the role of nicotine and its derivatives in lung cancer and how PPAR γ affects this. For example, a recent case-control study of 500 incident lung cancer cases and 517 age- and sex frequency-matched cancer-free controls suggested that PPAR γ polymorphisms in Chinese smokers may contribute to the etiology of lung cancer^[65]. Also, monocytes and monocyte-derived macrophages from healthy smokers showed increased PPAR γ expression as compared to those from healthy non-smokers, which was reproduced by nicotine *in vitro*^[66]. Interestingly, concomitant administration of PPAR γ agonists can effectively attenuate the effects of nicotine on alveolar type II cells^[67]. Among the carcinogenic chemi-

cals of cigarette smoking, tobacco-specific nitrosamine 4-(N-methyl-N-nitrosoamino)-1-(3-pyridyl)-1-butanone (NNK) is the most potent. One recent report showed that troglitazone blocked NNK-induced up-regulation of Heme oxygenase-1, Bcl-2, and cellular inhibitor of apoptosis protein 2; and restored Bad activity which was suppressed by NNK through activation of PPAR γ ^[68]. These findings reveal a novel molecular pathway of PPAR γ activation against cigarette smoking-related lung cancer. Evidence to date suggests that these effects of nicotine and its derivatives are mediated by nicotinic acetylcholine receptors expressed on the surface of tumor cells, thereby contributing to tumor progression^[69-71]. We recently found that rosiglitazone reduced nicotine-induced NSCLC cell growth through downregulation of α 4 nAChR-dependent signals including ERK and p38 MAPK; this effect appeared to be PPAR γ -independent^[72]. We also found that nicotine increases PPAR β/δ gene expression through α 7 nAChR-mediated activation of PI3K/mTOR signals. This is important since activation of PPAR β/δ is associated with enhanced cancer progression. These studies unveil a novel mechanism by which nicotine promotes human lung carcinoma cell growth and the impact of PPARs^[73].

PPAR γ and tumor cell-stromal interaction

Several studies suggest that PPAR γ ligands might prevent the interaction of tumor cells with their surrounding stroma, thereby interfering with host-derived and tumor-derived factors with mitogenic and pro-survival effects. An example of this is fibronectin, a matrix glycoprotein residing in the lung stroma that is increased in most, if not all, chronic forms of lung disease^[74]. This is true for tobacco-related lung disorders and fibrotic disorders, all associated with increased incidence of lung cancer^[75]. Several studies suggest that fibronectin serves as a mitogen and survival factor for NSCLC^[76], and fibronectin was recently shown to stimulate tumor cell expression of matrix metalloproteinases, proteases implicated in metastatic disease^[77]. These observations support the idea that tumor cell interactions with fibronectin through surface integrin receptors are advantageous for tumors since they stimulate proliferation, survival, and metastases^[76]. This idea was suggested by work showing reduced proliferative and metastatic capacity in tumor cells not expressing a fibronectin receptor α 5 β 1 integrin^[78]. Interestingly, PPAR γ ligands were shown to inhibit fibronectin expression in NSCLC cells by inhibiting transcription factors involved in the regulation of fibronectin gene expression^[79]. PPAR γ ligands (rosiglitazone and GW1929, but not PGJ₂) were also recently reported to inhibit the expression of the gene encoding for the α 5 integrin subunit resulting in reduced expression of the integrin α 5 β 1^[80]. Thus, by inhibiting the expression of fibronectin and its integrin α 5 β 1, PPAR γ ligands might reduce tumor cell recognition of fibronectin with consequent changes in cell proliferation and apoptosis.

Table 1 PPAR γ -dependent signals mediate the effects of PPAR γ ligands in lung cancer cells

PPAR γ ligands inhibit cancer cell growth and induce apoptosis <i>via</i> :
↓PGE $_2$ receptors (e.g. EP2 and EP4)
↑Tumor suppressors (e.g. PTEN, p21, AP-2 α , p53)
↓Inflammatory factors (e.g. NF- κ B, MCP-1, COX-2)
↓Angiogenic factor (e.g. VEGF)
↓Survival factors (e.g. SAPK/JNK, ILK, Src, FAK, PI3-K/Akt, mTOR)
↑↓Other kinase signals (e.g. ERK, p38 MAPK)
↓Growth factor receptors (e.g. EGF-R, PDGF-R)
↓Extracellular matrices (e.g. Fibronectin, MMP-2, MMP-9)
↓Integrin receptors (e.g. α 5 β 1)
↑↓Others [e.g. cytokines (e.g. IL-13, IL-21, TGF- β 1) and chemokines (e.g. MIP-1 β)]
↓Bcl-1, c-IAP2, <i>etc.</i>
PPAR γ ligands stimulate cancer cell growth and reduce apoptosis <i>via</i> :
↑Wnt signaling and oncogenes (e.g. cyclin D1, β -catenin, c-Myc)
↑Angiogenic signaling (e.g. VEGF, Angptl4)

PPAR: Peroxisome proliferator-activated receptor; SAPK/JNK: Stress-activated protein kinase/c-Jun NH2-terminal kinase; ILK: Integrin-linked kinase; PI3-K: Phosphatidylinositol 3-kinase; ERK: Extracellular signal-regulated kinase; mTOR: Mammalian target of rapamycin; NF- κ B: Nuclear factor- κ B; VEGF: Vascular endothelial growth factor; IL: Interleukin; TGF: Transforming growth factor; COX: Cyclooxygenase; MCP-1: Monocyte chemoattractant protein-1.

PPAR γ and angiogenesis

PPAR γ might also regulate the generation of the complex vascular network that supplies tumor cells. This idea is supported by studies showing a reduction in blood vessel density in lung tumors generated by the injection of A549 cells into the flanks of SCID mice treated with PPAR γ ligands^[81]. *In vitro* studies showed that the treatment of A549 cells with troglitazone or their transient transfection with a constitutively active PPAR γ construct blocked the production of angiogenic molecules such as ELR + CXC chemokines IL-8 (CXC-8), ENA-78 (CXCL5), and Gro- α (CXCL1)^[81]. Furthermore, PPAR γ activation inhibited NF- κ B, a transcription factor known to regulate the expression of many of the pro-angiogenic factors mentioned above. Similarly, rosiglitazone was shown to inhibit mouse lung tumor cell growth and metastasis *in vivo* through direct and indirect anti-angiogenic effects^[82].

Although the above studies reveal important anti-cancer effects for PPAR γ ligands, it is important to note that PPAR γ signaling has also been associated with tumor promoter activity in some cancer cells such as colon and breast, and that this effect was linked to increased β -catenin, c-Myc, Angptl4 and Wnt 5 expression^[83-85] (Table 1). PPAR γ ligands enhanced 7,12-dimethylbenz(*a*)anthracene-induced rat mammary adenocarcinoma^[86] and promoted colonic tumor growth in *Apc*^{Min} mice fed a high-fat diet^[87]. Targeted expression of activated PPAR γ in the mammary gland also enhanced tumorigenesis induced by polyoma middle T antigen^[84]. These findings need to be confirmed and tested in other tumors. However, these data suggest that activation of specific PPAR γ -related pathways may differ depending upon the cells and tumors examined. Internal genetic

variations and other factors may be responsible for the outcomes, and these need to be explored further followed by confirmation using relevant *in vivo* models of cancer.

IMPLICATIONS FOR THERAPY AND OTHER CONSIDERATIONS

The studies mentioned above suggest that PPARs are involved in lung cancer cell biology. However, their roles remain uncertain and much needs to be learned before they are targeted for therapeutic intervention, especially when considering PPAR γ . Nevertheless, activation of PPAR γ is strongly associated with decreased lung carcinoma cell proliferation both *in vitro* and *in vivo*. Furthermore, in primary NSCLC, the expression of PPAR γ has been correlated with tumor histological type and grade, and decreased PPAR γ expression was correlated with poor prognosis^[88]. Because of this, and the fact that synthetic agonists of PPAR γ with good safety profiles are currently in use in the clinical arena, PPAR γ has emerged as a reasonable target for the development of novel anti-lung cancer therapies. Synthetic and natural PPAR γ activators might be useful as well. For example, arachidonic acid inhibits the growth of A549 cells, and this effect is blocked by the synthetic PPAR γ inhibitor GW9662^[89]. MK886, a 5-lipoxygenase activating protein-directed inhibitor, stimulates apoptosis and reduces the growth of A549 cells through activation of PPAR γ ^[90]. These and related drugs can be used alone or in combination with other drugs for synergistic effects. This was observed when using low doses of MK886 in combination with ciglitazone and 13-cis-retinoic acid on A549 and H1299 cells^[90]. Also, dramatic synergistic anticancer effects have been reported for lovastatin (an HMG-CoA reductase inhibitor) and the PPAR γ ligand troglitazone in several cell lines including lung cancer cells^[91]. An enhancement of the anti-tumor effects of gefitinib by rosiglitazone on A549 cell growth was recently noted suggesting that combination strategies using selective nuclear receptor activators in conjunction with EGFR inhibitors might be effective^[92]. More recently, one report showed that the combination of clinically achievable concentrations of troglitazone and nonselective COX inhibitor, aspirin, can produce a strong synergistic effect on the inhibition of lung cancer cell growth and induction of apoptosis^[93].

One study demonstrated that combining the PPAR γ ligand rosiglitazone with carboplatin dramatically reduced lung tumor growth *in vivo*^[94]. Another study showed that the combination of PPAR γ ligand with platinum-based drugs exerted beneficial effects in the treatment of lung cancers including those tumors resistant to chemotherapy or acquired resistance to targeted therapy^[95]. More recently, one study using selenium (antioxidant), rosiglitazone, sodium phenylbutyrate or valproic acid (histone deacetylase inhibitors) and hydralazine (cytosine-demethylating agent) to prevent the progression of lung cancer in A/J mice treated with NNK demonstrated that chronic administration of rosiglitazone significantly blocked the progression of lung cancer in

Table 2 PPAR γ -independent signals triggered by PPAR γ ligands in lung cancer cells

<p>↑Tumor suppressors (e.g. LKB1, AMPK, TSC2) ↑ROS production and ERK, SAPK/JNK, p38 MAPK activation (note that this also occurs in PPARγ-dependent pathways) ↓Effects on transcription factors (e.g. AP-1, NF-κB, Smads, Sp1, CRE) ↓Nicotine receptor signaling (e.g. α4 and α7 nAChRs) ↓Apoptosis-related signals (e.g. Bcl-2, cyclin D1, c-FLIP, DR-5) ↑Apoptosis-related signals (e.g. casease 3/7, cyclin D1, p53)</p>
--

AMPK: AMP-activated protein kinase (AMPK); TSC2: Tuberous sclerosis complex 2.

the A/J mouse model^[96]. More tantalizing data were derived from a retrospective analysis demonstrating that thiazolidinedione use was associated with reduced risk of lung cancer. This study revealed a 33% reduction in lung cancer risk among thiazolidinedione users as compared to nonusers after adjusting other variables^[97]. Interestingly, a similar risk reduction was not observed for colorectal and prostate cancers^[97]. Clearly, as described previously, TZDs have many effects other than PPAR γ activation; the elucidation of such mechanisms holds the promise of unveiling new targets for the development of new anti-cancer therapies.

Despite the above findings, enthusiasm for the use of PPAR γ ligands as anti-cancer agents should be tempered by the fact that PPAR γ ligands stimulated PPAR γ transactivation in lung adenocarcinoma cell lines, while little to no effects were noted in squamous cell or large cell carcinomas^[98]. Also, it is important that we better define PPAR γ -independent pathways triggered by PPAR γ ligands to avoid unforeseen effects and to identify new targets for intervention^[92,98] (Table 2).

Furthermore, a novel splice variant of human PPAR γ 1, which is expressed strongly in tumor tissues of primary human lung SCC, was recently identified. This splice variant exhibits dominant-negative properties in human lung tumor cells, and its overexpression renders transfected cells more resistant to chemotherapeutic drug- and chemical-induced cell death^[99]. This suggests that the decreased drug sensitivity of PPAR γ 1-expressing cells may be associated with increased tumor aggressiveness and poor clinical prognosis in patients. Thus, a better understanding of the mechanisms of action of activated PPARs in tumors (and host cells) is required since the dissection of these pathways might unveil better targets for therapy. Nevertheless, the data available to date regarding PPAR γ are promising and justify engaging in carefully designed clinical studies to determine the true role of PPAR γ ligands in lung cancer, while further work should be performed to identify more selective and effective strategies.

CONCLUSION

Although the exact role of PPAR γ in controlling lung tumor growth and apoptosis remains incompletely defined, PPAR γ has been implicated both as a tumor

suppressor (in most cases) and tumor promoter (in rare cases). Hence, targeting this receptor for therapeutic purposes while minimizing side effects represents a great challenge. Nevertheless, it is clear that selective PPAR γ modulation of desired gene sets can be achieved by targeting co-repressor interactions, separating transactivation from transrepression, and favoring specific subsets of co-activators. PPAR γ activation results in inhibition of lung tumor growth (particularly NSCLC) both *in vitro* and *in vivo*. Although the exact mechanisms mediating this effect remain incompletely elucidated, data available to date regarding this member of the PPAR family is promising and justify engaging in prospective, randomized clinical studies to determine the true role of PPAR γ ligands in lung cancer biology. Further epidemiologic studies are required in patients treated with PPAR γ ligands for possible effects on tumor development.

REFERENCES

- Gilligan D, Nicolson M, Smith I, Groen H, Dalesio O, Goldstraw P, Hatton M, Hopwood P, Manegold C, Schramel F, Smit H, van Meerbeeck J, Nankivell M, Parmar M, Pugh C, Stephens R. Preoperative chemotherapy in patients with resectable non-small cell lung cancer: results of the MRC LU22/NVALT 2/EORTC 08012 multicentre randomised trial and update of systematic review. *Lancet* 2007; **369**: 1929-1937
- Cokkinides V, Bandi P, McMahon C, Jemal A, Glynn T, Ward E. Tobacco control in the United States--recent progress and opportunities. *CA Cancer J Clin* 2009; **59**: 352-365
- Mendoza L. Targeted therapies in the treatment of advanced non-small-cell lung cancer: update. *Klin Onkol* 2009; **22**: 131-138
- Pérez-Soler R. Individualized therapy in non-small-cell lung cancer: future versus current clinical practice. *Oncogene* 2009; **28** Suppl 1: S38-S45
- Roman J. Peroxisome proliferator-activated receptor gamma and lung cancer biology: implications for therapy. *J Investig Med* 2008; **56**: 528-533
- Krishnan A, Nair SA, Pillai MR. Biology of PPAR gamma in cancer: a critical review on existing lacunae. *Curr Mol Med* 2007; **7**: 532-540
- Pedchenko TV, Gonzalez AL, Wang D, DuBois RN, Massion PP. Peroxisome proliferator-activated receptor beta/delta expression and activation in lung cancer. *Am J Respir Cell Mol Biol* 2008; **39**: 689-696
- Han S, Ritzenthaler JD, Wingerd B, Roman J. Activation of peroxisome proliferator-activated receptor beta/delta (PPARbeta/delta) increases the expression of prostaglandin E2 receptor subtype EP4. The roles of phosphatidylinositol 3-kinase and CCAAT/enhancer-binding protein beta. *J Biol Chem* 2005; **280**: 33240-33249
- Han S, Ritzenthaler JD, Zheng Y, Roman J. PPARbeta/delta agonist stimulates human lung carcinoma cell growth through inhibition of PTEN expression: the involvement of PI3K and NF-kappaB signals. *Am J Physiol Lung Cell Mol Physiol* 2008; **294**: L1238-L1249
- Michalik L, Wahli W. PPARs Mediate Lipid Signaling in Inflammation and Cancer. *PPAR Res* 2008; **2008**: 134059
- Fajas L, Auboeuf D, Raspé E, Schoonjans K, Lefebvre AM, Saladin R, Najib J, Laville M, Fruchart JC, Deeb S, Vidal-Puig A, Flier J, Briggs MR, Staels B, Vidal H, Auwerx J. The organization, promoter analysis, and expression of the human PPARgamma gene. *J Biol Chem* 1997; **272**: 18779-18789
- Fajas L, Fruchart JC, Auwerx J. PPARgamma3 mRNA: a distinct PPARgamma mRNA subtype transcribed from an

- independent promoter. *FEBS Lett* 1998; **438**: 55-60
- 13 **Spiegelman BM**. PPAR γ in monocytes: less pain, any gain? *Cell* 1998; **93**: 153-155
 - 14 **Tontonoz P**, Hu E, Spiegelman BM. Regulation of adipocyte gene expression and differentiation by peroxisome proliferator activated receptor gamma. *Curr Opin Genet Dev* 1995; **5**: 571-576
 - 15 **Lehrke M**, Lazar MA. The many faces of PPAR γ . *Cell* 2005; **123**: 993-999
 - 16 **Semple RK**, Chatterjee VK, O'Rahilly S. PPAR gamma and human metabolic disease. *J Clin Invest* 2006; **116**: 581-589
 - 17 **Zhang GY**, Ahmed N, Riley C, Oliva K, Barker G, Quinn MA, Rice GE. Enhanced expression of peroxisome proliferator-activated receptor gamma in epithelial ovarian carcinoma. *Br J Cancer* 2005; **92**: 113-119
 - 18 **Elstner E**, Müller C, Koshizuka K, Williamson EA, Park D, Asou H, Shintaku P, Said JW, Heber D, Koeffler HP. Ligands for peroxisome proliferator-activated receptor gamma and retinoic acid receptor inhibit growth and induce apoptosis of human breast cancer cells in vitro and in BXN mice. *Proc Natl Acad Sci USA* 1998; **95**: 8806-8811
 - 19 **Daynes RA**, Jones DC. Emerging roles of PPARs in inflammation and immunity. *Nat Rev Immunol* 2002; **2**: 748-759
 - 20 **Jiang C**, Ting AT, Seed B. PPAR-gamma agonists inhibit production of monocyte inflammatory cytokines. *Nature* 1998; **391**: 82-86
 - 21 **Ricote M**, Huang J, Fajas L, Li A, Welch J, Najib J, Witztum JL, Auwerx J, Palinski W, Glass CK. Expression of the peroxisome proliferator-activated receptor gamma (PPAR-gamma) in human atherosclerosis and regulation in macrophages by colony stimulating factors and oxidized low density lipoprotein. *Proc Natl Acad Sci USA* 1998; **95**: 7614-7619
 - 22 **Keshamouni VG**, Reddy RC, Arenberg DA, Joel B, Thanickal VJ, Kalemkerian GP, Standiford TJ. Peroxisome proliferator-activated receptor-gamma activation inhibits tumor progression in non-small-cell lung cancer. *Oncogene* 2004; **23**: 100-108
 - 23 **Lehmann JM**, Moore LB, Smith-Oliver TA, Wilkison WO, Willson TM, Kliewer SA. An antidiabetic thiazolidinedione is a high affinity ligand for peroxisome proliferator-activated receptor gamma (PPAR gamma). *J Biol Chem* 1995; **270**: 12953-12956
 - 24 **Watkins PB**, Whitcomb RW. Hepatic dysfunction associated with troglitazone. *N Engl J Med* 1998; **338**: 916-917
 - 25 **Shibata T**, Matsui K, Nagao K, Shinkai H, Yonemori F, Wakitani K. Pharmacological profiles of a novel oral antidiabetic agent, JTT-501, an isoxazolidinedione derivative. *Eur J Pharmacol* 1999; **364**: 211-219
 - 26 **Suh N**, Wang Y, Williams CR, Risingsong R, Gilmer T, Willson TM, Sporn MB. A new ligand for the peroxisome proliferator-activated receptor-gamma (PPAR-gamma), GW7845, inhibits rat mammary carcinogenesis. *Cancer Res* 1999; **59**: 5671-5673
 - 27 **Liu M**, Fu L, Liu C, Xiong X, Gao X, Xiao M, Cai H, Hu H, Wang X, Mei C. DH9, a novel PPARgamma agonist suppresses the proliferation of ADPKD epithelial cells: An association with an inhibition of beta-catenin signaling. *Invest New Drugs* 2009; Epub ahead of print
 - 28 **Nagy L**, Tontonoz P, Alvarez JG, Chen H, Evans RM. Oxidized LDL regulates macrophage gene expression through ligand activation of PPARgamma. *Cell* 1998; **93**: 229-240
 - 29 **Chêne G**, Dubourdeau M, Balard P, Escoubet-Lozach L, Orfila C, Berry A, Bernad J, Aries MF, Charveron M, Pipy B. n-3 and n-6 polyunsaturated fatty acids induce the expression of COX-2 via PPARgamma activation in human keratinocyte HaCaT cells. *Biochim Biophys Acta* 2007; **1771**: 576-589
 - 30 **Schopfer FJ**, Lin Y, Baker PR, Cui T, Garcia-Barrio M, Zhang J, Chen K, Chen YE, Freeman BA. Nitrolinoleic acid: an endogenous peroxisome proliferator-activated receptor gamma ligand. *Proc Natl Acad Sci USA* 2005; **102**: 2340-2345
 - 31 **Hwang BY**, Lee JH, Nam JB, Kim HS, Hong YS, Lee JJ. Two new furanoditerpenes from *Saururus chinensis* and their effects on the activation of peroxisome proliferator-activated receptor gamma. *J Nat Prod* 2002; **65**: 616-617
 - 32 **Liang YC**, Tsai SH, Tsai DC, Lin-Shiau SY, Lin JK. Suppression of inducible cyclooxygenase and nitric oxide synthase through activation of peroxisome proliferator-activated receptor-gamma by flavonoids in mouse macrophages. *FEBS Lett* 2001; **496**: 12-18
 - 33 **Kuroda M**, Mimaki Y, Sashida Y, Mae T, Kishida H, Nishiyama T, Tsukagawa M, Konishi E, Takahashi K, Kawada T, Nakagawa K, Kitahara M. Phenolics with PPAR-gamma ligand-binding activity obtained from licorice (*Glycyrrhiza uralensis* roots) and ameliorative effects of glycyrrin on genetically diabetic KK-A(y) mice. *Bioorg Med Chem Lett* 2003; **13**: 4267-4272
 - 34 **Xu J**, Fu Y, Chen A. Activation of peroxisome proliferator-activated receptor-gamma contributes to the inhibitory effects of curcumin on rat hepatic stellate cell growth. *Am J Physiol Gastrointest Liver Physiol* 2003; **285**: G20-G30
 - 35 **Zheng S**, Chen A. Activation of PPARgamma is required for curcumin to induce apoptosis and to inhibit the expression of extracellular matrix genes in hepatic stellate cells in vitro. *Biochem J* 2004; **384**: 149-157
 - 36 **Casimiro-Garcia A**, Bigge CF, Davis JA, Padelino T, Pulaski J, Ohren JF, McConnell P, Kane CD, Royer LJ, Stevens KA, Auerbach B, Collard W, McGregor C, Song K. Synthesis and evaluation of novel alpha-heteroaryl-phenylpropanoic acid derivatives as PPARalpha/gamma dual agonists. *Bioorg Med Chem* 2009; **17**: 7113-7125
 - 37 **Fracchiolla G**, Laghezza A, Piemontese L, Tortorella P, Mazza F, Montanari R, Pochetti G, Lavecchia A, Novellino E, Pierno S, Conte Camerino D, Loiodice F. New 2-aryloxy-3-phenyl-propanoic acids as peroxisome proliferator-activated receptors alpha/gamma dual agonists with improved potency and reduced adverse effects on skeletal muscle function. *J Med Chem* 2009; **52**: 6382-6393
 - 38 **Henry RR**, Lincoff AM, Mudaliar S, Rabbia M, Chognot C, Herz M. Effect of the dual peroxisome proliferator-activated receptor-alpha/gamma agonist aleglitazar on risk of cardiovascular disease in patients with type 2 diabetes (SYNCHRONY): a phase II, randomised, dose-ranging study. *Lancet* 2009; **374**: 126-135
 - 39 **Berry EB**, Keelan JA, Helliwell RJ, Gilmour RS, Mitchell MD. Nanomolar and micromolar effects of 15-deoxy-delta 12,14-prostaglandin J2 on amnion-derived WISH epithelial cells: differential roles of peroxisome proliferator-activated receptors gamma and delta and nuclear factor kappa B. *Mol Pharmacol* 2005; **68**: 169-178
 - 40 **Kasai K**, Banba N, Hishinuma A, Matsumura M, Kakishita H, Matsumura M, Motohashi S, Sato N, Hattori Y. 15-Deoxy-Delta(12,14)-prostaglandin J(2) facilitates thyroglobulin production by cultured human thyrocytes. *Am J Physiol Cell Physiol* 2000; **279**: C1859-C1869
 - 41 **Li X**, Yang X, Xu Y, Jiang X, Li X, Nan F, Tang H. Troglitazone inhibits cell proliferation by attenuation of epidermal growth factor receptor signaling independent of peroxisome proliferator-activated receptor gamma. *Cell Res* 2009; **19**: 720-732
 - 42 **Reginato MJ**, Bailey ST, Krakow SL, Minami C, Ishii S, Tanaka H, Lazar MA. A potent antidiabetic thiazolidinedione with unique peroxisome proliferator-activated receptor gamma-activating properties. *J Biol Chem* 1998; **273**: 32679-32684
 - 43 **Inoue K**, Kawahito Y, Tsubouchi Y, Yamada R, Kohno M, Hosokawa Y, Katoh D, Bishop-Bailey D, Hla T, Sano H. Expression of peroxisome proliferator-activated receptor (PPAR)-gamma in human lung cancer. *Anticancer Res* 2001; **21**: 2471-2476
 - 44 **Ettinger DS**. Overview and state of the art in the manage-

- ment of lung cancer. *Oncology* (Williston Park) 2004; **18**: 3-9
- 45 **Pirozynski M.** 100 years of lung cancer. *Respir Med* 2006; **100**: 2073-2084
- 46 **Kim KY, Ahn JH, Cheon HG.** Apoptotic action of peroxisome proliferator-activated receptor-gamma activation in human non small-cell lung cancer is mediated via proline oxidase-induced reactive oxygen species formation. *Mol Pharmacol* 2007; **72**: 674-685
- 47 **Fulzele SV, Chatterjee A, Shaik MS, Jackson T, Ichite N, Singh M.** 15-Deoxy-Delta12,14-prostaglandin J2 enhances docetaxel anti-tumor activity against A549 and H460 non-small-cell lung cancer cell lines and xenograft tumors. *Anti-cancer Drugs* 2007; **18**: 65-78
- 48 **Li MY, Yuan H, Ma LT, Kong AW, Hsin MK, Yip J, Underwood MJ, Chen GG.** Roles of PPAR{alpha} and PPAR{gamma} in the Development of Non-small Cell Lung Cancer. *Am J Respir Cell Mol Biol* 2010; Epub ahead of print
- 49 **Han SW, Roman J.** Activated PPARgamma Targets Surface and Intracellular Signals That Inhibit the Proliferation of Lung Carcinoma Cells. *PPAR Res* 2008; **2008**: 254108
- 50 **Satoh T, Toyoda M, Hoshino H, Monden T, Yamada M, Shimizu H, Miyamoto K, Mori M.** Activation of peroxisome proliferator-activated receptor-gamma stimulates the growth arrest and DNA-damage inducible 153 gene in non-small cell lung carcinoma cells. *Oncogene* 2002; **21**: 2171-2180
- 51 **Li M, Lee TW, Yim AP, Mok TS, Chen GG.** Apoptosis induced by troglitazone is both peroxisome proliferator-activated receptor-gamma- and ERK-dependent in human non-small lung cancer cells. *J Cell Physiol* 2006; **209**: 428-438
- 52 **Li M, Lee TW, Mok TS, Warner TD, Yim AP, Chen GG.** Activation of peroxisome proliferator-activated receptor-gamma by troglitazone (TGZ) inhibits human lung cell growth. *J Cell Biochem* 2005; **96**: 760-774
- 53 **Zou W, Liu X, Yue P, Khuri FR, Sun SY.** PPARgamma ligands enhance TRAIL-induced apoptosis through DR5 upregulation and c-FLIP downregulation in human lung cancer cells. *Cancer Biol Ther* 2007; **6**: 99-106
- 54 **Han S, Roman J.** Rosiglitazone suppresses human lung carcinoma cell growth through PPARgamma-dependent and PPARgamma-independent signal pathways. *Mol Cancer Ther* 2006; **5**: 430-437
- 55 **Han S, Zheng Y, Roman J.** Rosiglitazone, an Agonist of PPARgamma, Inhibits Non-Small Cell Carcinoma Cell Proliferation In Part through Activation of Tumor Sclerosis Complex-2. *PPAR Res* 2007; **2007**: 29632
- 56 **Lee SY, Hur GY, Jung KH, Jung HC, Lee SY, Kim JH, Shin C, Shim JJ, In KH, Kang KH, Yoo SH.** PPAR-gamma agonist increase gefitinib's antitumor activity through PTEN expression. *Lung Cancer* 2006; **51**: 297-301
- 57 **Han S, Sidell N, Fisher PB, Roman J.** Up-regulation of p21 gene expression by peroxisome proliferator-activated receptor gamma in human lung carcinoma cells. *Clin Cancer Res* 2004; **10**: 1911-1919
- 58 **Zhang W, Zhang H, Xing L.** Influence of ciglitazone on A549 cells growth in vitro and in vivo and mechanism. *J Huazhong Univ Sci Technolog Med Sci* 2006; **26**: 36-39
- 59 **Han S, Sun X, Ritzenthaler JD, Roman J.** Fish oil inhibits human lung carcinoma cell growth by suppressing integrin-linked kinase. *Mol Cancer Res* 2009; **7**: 108-117
- 60 **Hannigan G, Troussard AA, Dedhar S.** Integrin-linked kinase: a cancer therapeutic target unique among its ILK. *Nat Rev Cancer* 2005; **5**: 51-63
- 61 **Serini S, Trombino S, Oliva F, Piccioni E, Monego G, Resci F, Boninsegna A, Picci N, Ranelletti FO, Calviello G.** Docosahexaenoic acid induces apoptosis in lung cancer cells by increasing MKP-1 and down-regulating p-ERK1/2 and p-p38 expression. *Apoptosis* 2008; **13**: 1172-1183
- 62 **Han S, Roman J.** Suppression of prostaglandin E2 receptor subtype EP2 by PPARgamma ligands inhibits human lung carcinoma cell growth. *Biochem Biophys Res Commun* 2004; **314**: 1093-1099
- 63 **Hazra S, Dubinett SM.** Ciglitazone mediates COX-2 dependent suppression of PGE2 in human non-small cell lung cancer cells. *Prostaglandins Leukot Essent Fatty Acids* 2007; **77**: 51-58
- 64 **Bren-Mattison Y, Meyer AM, Van Putten V, Li H, Kuhn K, Stearman R, Weiser-Evans M, Winn RA, Heasley LE, Nemenoff RA.** Antitumorigenic effects of peroxisome proliferator-activated receptor-gamma in non-small-cell lung cancer cells are mediated by suppression of cyclooxygenase-2 via inhibition of nuclear factor-kappaB. *Mol Pharmacol* 2008; **73**: 709-717
- 65 **Chen D, Jin G, Wang Y, Wang H, Liu H, Liu Y, Fan W, Ma H, Miao R, Hu Z, Sun W, Qian J, Jin L, Wei Q, Shen H, Huang W, Lu D.** Genetic variants in peroxisome proliferator-activated receptor-gamma gene are associated with risk of lung cancer in a Chinese population. *Carcinogenesis* 2008; **29**: 342-350
- 66 **Amoruso A, Bardelli C, Gunella G, Fresu LG, Ferrero V, Brunelleschi S.** Quantification of PPAR-gamma protein in monocyte/macrophages from healthy smokers and non-smokers: a possible direct effect of nicotine. *Life Sci* 2007; **81**: 906-915
- 67 **Rehan VK, Wang Y, Sugano S, Santos J, Patel S, Sakurai R, Boros LG, Lee WP, Torday JS.** In utero nicotine exposure alters fetal rat lung alveolar type II cell proliferation, differentiation, and metabolism. *Am J Physiol Lung Cell Mol Physiol* 2007; **292**: L323-L333
- 68 **Li MY, Hsin MK, Yip J, Mok TS, Underwood MJ, Chen GG.** PPARgamma activation extinguishes smoking carcinogen by inhibiting NNK-mediated proliferation. *Am J Respir Cell Mol Biol* 2010; **42**: 113-122
- 69 **Dasgupta P, Chellappan SP.** Nicotine-mediated cell proliferation and angiogenesis: new twists to an old story. *Cell Cycle* 2006; **5**: 2324-2328
- 70 **Lam DC, Girard L, Ramirez R, Chau WS, Suen WS, Sheridan S, Tin VP, Chung LP, Wong MP, Shay JW, Gazdar AF, Lam WK, Minna JD.** Expression of nicotinic acetylcholine receptor subunit genes in non-small-cell lung cancer reveals differences between smokers and nonsmokers. *Cancer Res* 2007; **67**: 4638-4647
- 71 **Dasgupta P, Rastogi S, Pillai S, Ordonez-Ercan D, Morris M, Haura E, Chellappan S.** Nicotine induces cell proliferation by beta-arrestin-mediated activation of Src and Rb-Raf-1 pathways. *J Clin Invest* 2006; **116**: 2208-2217
- 72 **Sun X, Ritzenthaler JD, Zheng Y, Roman J, Han S.** Rosiglitazone inhibits alpha4 nicotinic acetylcholine receptor expression in human lung carcinoma cells through peroxisome proliferator-activated receptor gamma-independent signals. *Mol Cancer Ther* 2009; **8**: 110-118
- 73 **Sun X, Ritzenthaler JD, Zhong X, Zheng Y, Roman J, Han S.** Nicotine stimulates PPARbeta/delta expression in human lung carcinoma cells through activation of PI3K/mTOR and suppression of AP-2alpha. *Cancer Res* 2009; **69**: 6445-6453
- 74 **Limper AH, Roman J.** Fibronectin. A versatile matrix protein with roles in thoracic development, repair and infection. *Chest* 1992; **101**: 1663-1673
- 75 **Roman J, Ritzenthaler JD, Gil-Acosta A, Rivera HN, Roser-Page S.** Nicotine and fibronectin expression in lung fibroblasts: implications for tobacco-related lung tissue remodeling. *FASEB J* 2004; **18**: 1436-1438
- 76 **Han S, Khuri FR, Roman J.** Fibronectin stimulates non-small cell lung carcinoma cell growth through activation of Akt/mammalian target of rapamycin/S6 kinase and inactivation of LKB1/AMP-activated protein kinase signal pathways. *Cancer Res* 2006; **66**: 315-323
- 77 **Han S, Ritzenthaler JD, Sitaraman SV, Roman J.** Fibronectin increases matrix metalloproteinase 9 expression through activation of c-Fos via extracellular-regulated kinase and phosphatidylinositol 3-kinase pathways in human lung car-

- cinoma cells. *J Biol Chem* 2006; **281**: 29614-29624
- 78 **Roman J**, Ritzenthaler JD, Roser-Page S, Sun X, Han S. {alpha}5{beta}1 Integrin Expression is Essential for Tumor Progression in Experimental Lung Cancer. *Am J Respir Cell Mol Biol* 2010; Epub ahead of print
- 79 **Han S**, Ritzenthaler JD, Rivera HN, Roman J. Peroxisome proliferator-activated receptor-gamma ligands suppress fibronectin gene expression in human lung carcinoma cells: involvement of both CRE and Sp1. *Am J Physiol Lung Cell Mol Physiol* 2005; **289**: L419-L428
- 80 **Han S**, Rivera HN, Roman J. Peroxisome proliferator-activated receptor-gamma ligands inhibit alpha5 integrin gene transcription in non-small cell lung carcinoma cells. *Am J Respir Cell Mol Biol* 2005; **32**: 350-359
- 81 **Keshamouni VG**, Arenberg DA, Reddy RC, Newstead MJ, Anthwal S, Standiford TJ. PPAR-gamma activation inhibits angiogenesis by blocking ELR+CXC chemokine production in non-small cell lung cancer. *Neoplasia* 2005; **7**: 294-301
- 82 **Panigrahy D**, Singer S, Shen LQ, Butterfield CE, Freedman DA, Chen EJ, Moses MA, Kilroy S, Duensing S, Fletcher C, Fletcher JA, Hlatky L, Hahnfeldt P, Folkman J, Kaipainen A. PPARgamma ligands inhibit primary tumor growth and metastasis by inhibiting angiogenesis. *J Clin Invest* 2002; **110**: 923-932
- 83 **Choi IK**, Kim YH, Kim JS, Seo JH. PPAR-gamma ligand promotes the growth of APC-mutated HT-29 human colon cancer cells in vitro and in vivo. *Invest New Drugs* 2008; **26**: 283-288
- 84 **Saez E**, Rosenfeld J, Livolsi A, Olson P, Lombardo E, Nelson M, Banayo E, Cardiff RD, Izpisua-Belmonte JC, Evans RM. PPAR gamma signaling exacerbates mammary gland tumor development. *Genes Dev* 2004; **18**: 528-540
- 85 **Tian L**, Zhou J, Casimiro MC, Liang B, Ojeifo JO, Wang M, Hyslop T, Wang C, Pestell RG. Activating peroxisome proliferator-activated receptor gamma mutant promotes tumor growth in vivo by enhancing angiogenesis. *Cancer Res* 2009; **69**: 9236-9244
- 86 **Pighetti GM**, Novosad W, Nicholson C, Hitt DC, Hansens C, Hollingsworth AB, Lerner ML, Brackett D, Lightfoot SA, Gimble JM. Therapeutic treatment of DMBA-induced mammary tumors with PPAR ligands. *Anticancer Res* 2001; **21**: 825-829
- 87 **Saez E**, Tontonoz P, Nelson MC, Alvarez JG, Ming UT, Baird SM, Thomazy VA, Evans RM. Activators of the nuclear receptor PPARgamma enhance colon polyp formation. *Nat Med* 1998; **4**: 1058-1061
- 88 **Theocharis S**, Kanelli H, Politi E, Margeli A, Karkandaris C, Philippides T, Koutselinis A. Expression of peroxisome proliferator activated receptor-gamma in non-small cell lung carcinoma: correlation with histological type and grade. *Lung Cancer* 2002; **36**: 249-255
- 89 **Muzio G**, Trombetta A, Maggiora M, Martinasso G, Vasilioiu V, Lassen N, Canuto RA. Arachidonic acid suppresses growth of human lung tumor A549 cells through down-regulation of ALDH3A1 expression. *Free Radic Biol Med* 2006; **40**: 1929-1938
- 90 **Avis I**, Martínez A, Tauler J, Zudaire E, Mayburd A, Abu-Ghazaleh R, Ondrey F, Mulshine JL. Inhibitors of the arachidonic acid pathway and peroxisome proliferator-activated receptor ligands have superadditive effects on lung cancer growth inhibition. *Cancer Res* 2005; **65**: 4181-4190
- 91 **Yao CJ**, Lai GM, Chan CF, Cheng AL, Yang YY, Chuang SE. Dramatic synergistic anticancer effect of clinically achievable doses of lovastatin and troglitazone. *Int J Cancer* 2006; **118**: 773-779
- 92 **Nemenoff RA**, Winn RA. Role of nuclear receptors in lung tumorigenesis. *Eur J Cancer* 2005; **41**: 2561-2568
- 93 **Yan KH**, Yao CJ, Chang HY, Lai GM, Cheng AL, Chuang SE. The synergistic anticancer effect of troglitazone combined with aspirin causes cell cycle arrest and apoptosis in human lung cancer cells. *Mol Carcinog* 2010; **49**: 235-246
- 94 **Girnun GD**, Naseri E, Vafai SB, Qu L, Szwaya JD, Bronson R, Alberta JA, Spiegelman BM. Synergy between PPARgamma ligands and platinum-based drugs in cancer. *Cancer Cell* 2007; **11**: 395-406
- 95 **Girnun GD**, Chen L, Silvaggi J, Drapkin R, Chirieac LR, Padera RF, Upadhyay R, Vafai SB, Weissleder R, Mahmood U, Naseri E, Buckley S, Li D, Force J, McNamara K, Demetri G, Spiegelman BM, Wong KK. Regression of drug-resistant lung cancer by the combination of rosiglitazone and carboplatin. *Clin Cancer Res* 2008; **14**: 6478-6486
- 96 **Lyon CM**, Klinge DM, Do KC, Grimes MJ, Thomas CL, Damiani LA, March TH, Stidley CA, Belinsky SA. Rosiglitazone prevents the progression of preinvasive lung cancer in a murine model. *Carcinogenesis* 2009; **30**: 2095-2099
- 97 **Govindarajan R**, Ratnasinghe L, Simmons DL, Siegel ER, Midathada MV, Kim L, Kim PJ, Owens RJ, Lang NP. Thiazolidinediones and the risk of lung, prostate, and colon cancer in patients with diabetes. *J Clin Oncol* 2007; **25**: 1476-1481
- 98 **Allred CD**, Kilgore MW. Selective activation of PPARgamma in breast, colon, and lung cancer cell lines. *Mol Cell Endocrinol* 2005; **235**: 21-29
- 99 **Kim HJ**, Hwang JY, Kim HJ, Choi WS, Lee JH, Kim HJ, Chang KC, Nishinaka T, Yabe-Nishimura C, Seo HG. Expression of a peroxisome proliferator-activated receptor gamma 1 splice variant that was identified in human lung cancers suppresses cell death induced by cisplatin and oxidative stress. *Clin Cancer Res* 2007; **13**: 2577-2583

S- Editor Cheng JX L- Editor Webster JR E- Editor Zheng XM