

World Journal of *Clinical Oncology*

World J Clin Oncol 2024 April 24; 15(4): 464-575



EDITORIAL

- 464 Classificatory updates in verrucous and cuniculatum carcinomas: Insights from the 5th edition of WHO-IARC head and neck tumor classification
Silveira FM, Schuch LF, Bologna-Molina R
- 468 Understanding the role of transmembrane 9 superfamily member 1 in bladder cancer pathogenesis
Gade VKV, Yadav BS
- 472 Management of lateral pelvic lymph nodes in rectal cancer: Is it time to reach an Agreement?
Romero-Zoghbi SE, López-Campos F, Couñago F
- 478 Tumor infiltrating lymphocytes in gastric cancer: Unraveling complex interactions for precision medicine
Kapoor M, Sehwat A, Karthik J, Sundriyal D

REVIEW

- 482 Focus on current and emerging treatment options for glioma: A comprehensive review
Lucke-Wold B, Rangwala BS, Shafique MA, Siddiq MA, Mustafa MS, Danish F, Nasrullah RMU, Zainab N, Haseeb A
- 496 Immune pathway through endometriosis to ovarian cancer
Calmon MS, Lemos FFB, Silva Luz M, Rocha Pinheiro SL, de Oliveira Silva LG, Correa Santos GL, Rocha GR, Freire de Melo F

MINIREVIEWS

- 523 Britanin – a beacon of hope against gastrointestinal tumors?
Kajdanek A, Kolat D, Zhao LY, Kciuk M, Pasieka Z, Kaluzińska-Kolat Ż
- 531 Molecular targets and mechanisms of different aberrant alternative splicing in metastatic liver cancer
Geng DY, Chen QS, Chen WX, Zhou LS, Han XS, Xie QH, Guo GH, Chen XF, Chen JS, Zhong XP

ORIGINAL ARTICLE

Retrospective Cohort Study

- 540 Comparative effectiveness of immunotherapy and chemotherapy in patients with metastatic colorectal cancer stratified by microsatellite instability status
Niu CG, Zhang J, Rao AV, Joshi U, Okolo P

Retrospective Study

- 548 Elevated cardiovascular risk and acute events in hospitalized colon cancer survivors: A decade-apart study of two nationwide cohorts
Desai R, Mondal A, Patel V, Singh S, Chauhan S, Jain A

Basic Study

- 554** Regulation of *TMEM100* expression by epigenetic modification, effects on proliferation and invasion of esophageal squamous carcinoma

Xu YF, Dang Y, Kong WB, Wang HL, Chen X, Yao L, Zhao Y, Zhang RQ

CASE REPORT

- 566** Low-grade myofibrosarcoma of the maxillary sinus: Two case reports

Mydlak A, Ścibik Ł, Durzynska M, Zwoliński J, Buchajska K, Lenartowicz O, Kucharz J

ABOUT COVER

Peer Reviewer of *World Journal of Clinical Oncology*, Ramiro Manuel Fernández-Placencia, FACS, MD, Professor, Surgical Oncologist, Abdominal Surgery Department, Instituto Nacional de Enfermedades Neoplásicas (INEN), Lima Lima034, Lima, Peru. ramirofp02@gmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Clinical Oncology* (WJCO, *World J Clin Oncol*) is to provide scholars and readers from various fields of oncology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJCO mainly publishes articles reporting research results and findings obtained in the field of oncology and covering a wide range of topics including art of oncology, biology of neoplasia, breast cancer, cancer prevention and control, cancer-related complications, diagnosis in oncology, gastrointestinal cancer, genetic testing for cancer, gynecologic cancer, head and neck cancer, hematologic malignancy, lung cancer, melanoma, molecular oncology, neurooncology, palliative and supportive care, pediatric oncology, surgical oncology, translational oncology, and urologic oncology.

INDEXING/ABSTRACTING

The WJCO is now abstracted and indexed in PubMed, PubMed Central, Emerging Sources Citation Index (Web of Science), Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for WJCO as 2.8; IF without journal self cites: 2.8; 5-year IF: 3.0; Journal Citation Indicator: 0.36.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Yin-Qing Zhao*; Production Department Director: *Xu Guo*; Cover Editor: *Xu Guo*.

NAME OF JOURNAL

World Journal of Clinical Oncology

ISSN

ISSN 2218-4333 (online)

LAUNCH DATE

November 10, 2010

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Hiten RH Patel, Stephen Safe, Jian-Hua Mao, Ken H Young

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2218-4333/editorialboard.htm>

PUBLICATION DATE

April 24, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Focus on current and emerging treatment options for glioma: A comprehensive review

Brandon Lucke-Wold, Burhanuddin Sohail Rangwala, Muhammad Ashir Shafique, Mohammad Arham Siddiq, Muhammad Saqlain Mustafa, Fnu Danish, Rana Muhammad Umer Nasrullah, Noor Zainab, Abdul Haseeb

Specialty type: Oncology

Provenance and peer review:

Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0

Grade B (Very good): B

Grade C (Good): 0

Grade D (Fair): 0

Grade E (Poor): E

P-Reviewer: Chu SH, China; Shibata Y, Japan

Received: November 22, 2023

Peer-review started: November 22, 2023

First decision: January 12, 2024

Revised: January 22, 2024

Accepted: February 28, 2024

Article in press: February 28, 2024

Published online: April 24, 2024



Brandon Lucke-Wold, Department of Neurosurgery, University of Florida, Gainesville, FL 32608, United States

Burhanuddin Sohail Rangwala, Muhammad Ashir Shafique, Mohammad Arham Siddiq, Muhammad Saqlain Mustafa, Fnu Danish, Rana Muhammad Umer Nasrullah, Abdul Haseeb, Department of Neurosurgery, Jinnah Sindh Medical University, Karachi 75510, Pakistan

Noor Zainab, Department of Neurosurgery, Army Medical College, Rawalpindi 46000, Pakistan

Corresponding author: Muhammad Ashir Shafique, MBBS, Doctor, Researcher, Department of Neurosurgery, Jinnah Sindh Medical University, Haji Rafique Road Karachi Sindh, Karachi 75510, Pakistan. ashirshafique109@gmail.com

Abstract

This comprehensive review delves into the current updates and challenges associated with the management of low-grade gliomas (LGG), the predominant primary tumors in the central nervous system. With a general incidence rate of 5.81 per 100000, gliomas pose a significant global concern, necessitating advancements in treatment techniques to reduce mortality and morbidity. This review places a particular focus on immunotherapies, discussing promising agents such as Zotiraciclib and Lerapolturev. Zotiraciclib, a CDK9 inhibitor, has demonstrated efficacy in glioblastoma treatment in preclinical and clinical studies, showing its potential as a therapeutic breakthrough. Lerapolturev, a viral immunotherapy, induces inflammation in glioblastoma and displays positive outcomes in both adult and pediatric patients. Exploration of immunotherapy extends to Pembrolizumab, Nivolumab, and Entrectinib, revealing the challenges and variabilities in patient responses. Despite promising preclinical data, the monoclonal antibody Depatuxizumab has proven ineffective in glioblastoma treatment, emphasizing the critical need to understand resistance mechanisms. The review also covers the success of radiation therapy in pediatric LGG, with evolving techniques, such as proton therapy, showing potential improvements in patient quality of life. Surgical treatment is discussed in the context of achieving a balance between preserving the patient's quality of life and attaining gross total resection, with the extent of surgical resection significantly influencing the survival outcomes. In addition to advancements in cancer vaccine development, this review highlights the evolving landscape of LGG treatment, emphasizing a shift toward personalized and targeted therapies. Ongoing research is essential for refining strategies and enhancing

outcomes in the management of LGG.

Key Words: Low-grade gliomas; Monoclonal antibody; Lerapolturev; Glioblastoma; CDK9 inhibitor

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Our manuscript explores the dynamic landscape of glioma treatment, emphasizing the urgent need for innovative therapies to combat this prevalent central nervous system malignancy. We delve into the promising realm of immunotherapies, highlighting novel agents like zotiraciclib, pembrolizumab, and lerapolturev, offering insights into their mechanisms and clinical efficacy. Furthermore, we discuss the evolving role of radiation therapy, emphasizing recent advancements in reducing treatment-related toxicities while improving outcomes. Surgical strategies, including subtotal resection and intraoperative radiotherapy, are also explored, showcasing their potential to enhance survival while minimizing neurological morbidities.

Citation: Lucke-Wold B, Rangwala BS, Shafique MA, Siddiq MA, Mustafa MS, Danish F, Nasrullah RMU, Zainab N, Haseeb A. Focus on current and emerging treatment options for glioma: A comprehensive review. *World J Clin Oncol* 2024; 15(4): 482-495

URL: <https://www.wjgnet.com/2218-4333/full/v15/i4/482.htm>

DOI: <https://dx.doi.org/10.5306/wjco.v15.i4.482>

INTRODUCTION

Gliomas represent the most prevalent primary tumors in the central nervous system (CNS) across various age groups[1, 2]. Gliomas have a general incidence rate of 5.81 per 100000 people, with older individuals having a threefold higher frequency than young children. Gliomas account for 29%-35% of all central nervous system tumors in the adolescent and young adult demographic (ages 15-39 years), with an incidence of 3.41 per 100000[3-5]. Gliomas continue to be a global concern, emphasizing the vital need to improve treatment techniques for lowering both mortality and morbidity, elevating it to the top of the neuro-oncology priority list[6,7].

Clinical care, therapeutic response, and outcomes differ significantly between pediatric and adult glioma patients. Children with high-grade gliomas (HGGs) have poor prognosis, with frequently limited long-term survival ranging from months to a few years after diagnosis[8,9]. In contrast, pediatric patients with low-grade gliomas (LGG) have good overall survival (OS)[10,11], despite significant tumor- and treatment-related morbidity[12] (Table 1). The increased likelihood of malignant transformation, which is extremely rare in children) adds to a less favorable prognosis in adults with low-grade gliomas[13,14].

IMMUNOTHERAPIES

Despite advancements in surgery, radiotherapy, and chemotherapy for LGG, the disease remains incurable and often progresses to secondary malignant transformation. Immunotherapeutic strategies have demonstrated success in various cancers, including lung, skin, colon, and blood-related cancers (Figure 1). Given that low-grade gliomas, particularly in younger patients, exhibit slower growth compared to high-grade gliomas, there is a suggestion that immunotherapies may be more effective due to the healthier immune systems of younger individuals, potentially leading to better treatment responses. Immunotherapies, including Zotiraciclib and Lerapolturev, exert their effects through distinct mechanism (Table 2).

Zotiraciclib

Zotiraciclib, a potent CDK9 inhibitor, exhibits efficacy against glioblastoma by suppressing transcription and disrupting cellular energy production. Preclinical studies, both in vitro and in vivo, have revealed its synergistic effect with temozolomide. In clinical trials, Zotiraciclib demonstrated the ability to cross the blood-brain barrier and suppress CDK9 activity in tumor tissues[15]. This promising mechanism, targeting multiple glioblastoma survival pathways, positions Zotiraciclib as a potential therapeutic breakthrough[16-19].

A two-stage, two-arm, randomized phase 1 clinical trial further investigated the potential of zotiraciclib in recurrent high-grade gliomas. This study included a comprehensive evaluation of pharmacokinetics, patient-reported outcomes, and a detailed examination of rapid-onset neutropenia. Despite this observed neutropenia, a thorough analysis concluded that it did not compromise patient safety, allowing the research and development of this novel CDK9 inhibitor to progress[19].

Pembrolizumab or nivolumab

Immunotherapy has garnered significant interest as a potential treatment for glioblastoma (GBM). Nevertheless, a recent

Table 1 World Health Organization classification of gliomas

Grade	Name	Description and characteristics
I	Pilocytic astrocytoma	Well-differentiated, often cystic, slow-growing, generally benign
II	Diffuse astrocytoma	Infiltrative, moderately cellular, tends to recur, can progress to higher grades
II	Oligodendroglioma	Composed of oligodendrocyte-like cells, often associated with 1p/19q co-deletion
II	Mixed oligoastrocytoma	Combination of features of oligodendroglioma and diffuse astrocytoma
III	Anaplastic astrocytoma	Higher grade astrocytoma with increased cellularity and mitotic activity
III	Anaplastic oligodendroglioma	Higher grade oligodendroglioma with increased cellularity and atypia
III	Anaplastic oligoastrocytoma	Higher grade mixed tumor with features of both anaplastic astrocytoma and anaplastic oligodendroglioma
IV	Glioblastoma	Highly aggressive, necrosis, endothelial proliferation, molecular heterogeneity

clinical study focusing on recurrent glioblastoma and employing PD-1 immune checkpoint inhibitors revealed that a minority of patients (8%) exhibited noticeable improvements in their condition[20]. The mechanistic underpinnings of the variability in response patterns remain unclear.

Enhanced T cell infiltration in the tumor microenvironment and elevated mutational burdens in various cancer types have been associated with improved responses to anti-PD-1 therapy[21-23]. However, GBM presents a more immunosuppressive tumor microenvironment and a lower burden of somatic mutations than melanomas or non-small cell lung cancer[24]. Immunosuppression in GBM is facilitated by the expression of PD-1 ligands (PD-L1/2) in tumor cells, leading to T cell exhaustion and apoptosis. The binding of PD-1 to the surface of cytotoxic T cells hampers their ability to mount an effective anti-tumor response. PD-1 inhibitor therapy disrupts this immune checkpoint, reinforcing the immune response against tumors[23].

PD-1 inhibitors, such as pembrolizumab and nivolumab, have gained attention for glioblastoma treatment. However, recent clinical studies have revealed variable responses, necessitating deeper understanding of the underlying mechanisms. Glioblastoma's immunosuppressive microenvironment and lower mutation burden compared to other cancers pose challenges. Molecular-tailored strategies hold promise for optimizing patient selection for immunotherapy, although further testing is required to validate their efficacy[25].

Lerapolturev

Lerapolturev, a viral immunotherapy, operates *via* a unique mechanism. As a polio-rhinovirus chimera, it induces persistent type-I interferon-dominant inflammation in glioblastoma, leading to polyfunctional antitumor CD8+ T-cell responses. Clinical trials involving Lerapolturev for recurrent adult glioblastoma demonstrated a 16% survival rate of at least 36 months, with a manageable safety profile[26-29].

In pediatric high-grade gliomas, Lerapolturev showed promise, with no grade 3 or 4 toxicity observed in early trials. The safety of treatment at this dose allows for further trials, including patients as young as 9 years of age. Ongoing research is crucial to understand the immunological factors influencing the efficacy of Lerapolturev in pediatric versus adult high-grade gliomas[30]. Our group's previous research in adults gained additional support from the inclusion of patients as young as 9 years old, including one individual with WHO grade 3 glioma[30]. Moreover, pediatric high-grade gliomas typically exhibit significantly different molecular profiles compared to adult high-grade gliomas[31]. However, whether immunological factors affecting viral immunotherapies, such as Lerapolturev, vary between pediatric and adult high-grade gliomas remains uncertain[30].

Depatuxizumab

Depatuxizumab (formerly ABT-806) is a humanized monoclonal antibody developed against epidermal growth factor receptor variant III (EGFRvIII) that also binds to wild-type EGFR at elevated levels[32,33]. The antibody-drug conjugate (ADC) Depatuxizumab mafodotin (formerly ABT-414) connects the depatux to the cytotoxic payload monomethyl auristatin F (MMAF or mafodotin). Upon binding to activated EGFR, ADC is internalized, degraded in acidic compartments, and releases the toxin, causing cell death. Unlike other treatments, this direct cytotoxic effect does not rely on inhibition of EGFR signaling and avoids typical toxicities[34]. Although unconjugated depatux is ineffective against GBMs, depatux-m demonstrates efficacy in GBM cell lines and models with EGFR amplification or EGFRvIII, showing effectiveness alone and in combination with radiotherapy and temozolomide[35]. ADCs, including depatux-m, show promise in various cancers[36], surpassing unconjugated monoclonal antibodies in efficacy, with numerous ADCs under investigation under diverse conditions[37].

Despite promising preclinical and early clinical data, depatux-m has proven ineffective in treating GBM. This disappointing outcome may result from the emergence of resistant clones over time, negating any overall survival benefit [38]. Limited penetration of depatux-m into large tumors and challenges in reaching intracranial tumors[39], especially in the non-enhancing tumor region, underscore crucial lessons for future studies involving large molecules[38]. Safety concerns with depatux-m were reversible, with adverse events, such as sensitivity to light and thrombocytopenia, being the most frequently observed.

Table 2 List of Immunotherapy

Ref.	Completion year	Demographics	Study phase	Identifier	Experimental drug	Sample size	Primary endpoint/outcomes	Results for primary outcome
BRAF/MEK inhibitors								
Nicolaides <i>et al</i> [107], 2020	2023	Pediatrics	Phase 2	NCT01748149 (Ongoing Trial)	Vemurafenib	40	Safety and pharmacokinetics	Not yet reported
Hargrave <i>et al</i> [108], 2019	2020	Pediatrics	Phase 1/2a	NCT01677741	Dabrafenib	32	Objective response rates and safety	Objective response rate was 44% and 91% experienced adverse effects
Kaley <i>et al</i> [109], 2018	2016	Adults	Phase 2	NCT01524978	Vemurafenib	24	Confirmed objective response rate, PFS, OS and safety	Confirmed objective response rate was 25% and median PFS was 5.5 months
FGFR inhibitors								
Lassman <i>et al</i> [110], 2022	2018	Adults	Phase 2	NCT01975701	Infigratinib	26	6-month PFS	6-month PFS rate was 16.0%
Bahleda <i>et al</i> [111], 2019	2017	Adults	Phase 1	NCT01703481	Erdafitinib	187	Safety	Most common treatment-related adverse events were hyperphosphatemia, dry mouth, and asthenia, generally grade 1/2 severity
HDAC inhibitors								
Wood <i>et al</i> [112], 2018	2018	Pediatrics	Phase 1	ACTRN12609000978268	Panobinostat	9	Safety and pharmacokinetics	2 patients experienced Grade 3-4 thrombocytopenia, 1 experienced Grade 3 anemia, and 2 experienced Grade 3 neutropenia
Imipridone								
Arrillaga-Romany <i>et al</i> [113], 2020	2023		Phase 2	NCT02525692 (Ongoing Trial)	ONC201	89	6-month PFS	Not yet reported
PI3K/mTOR inhibitors								
Wen <i>et al</i> [114], 2022	2023	Adults	Phase 2	NCT03522298	Paxalisib	32	Safety and pharmacokinetics	Well-tolerated with adverse events consistent with other PI3K inhibitors
Wen <i>et al</i> [115], 2020	2020	Adults	Phase 1	NCT01547546	GDC-0084	47	Safety and pharmacokinetics	Well-tolerated with adverse events consistent with other PI3K inhibitors
Franz <i>et al</i> [116], 2015	2014	Adults/Pediatrics	Phase 1/2	NCT00411619	Enviroximes	28	6-month change in the volume of sub ependymal giant-cell astrocytoma	Statistically significant reduction in the volume of the primary sub ependymal giant-cell astrocytoma at 6 months
NTRK/ALK inhibitors								
NCT02637687 [117]	2026	Pediatrics	Phase	NCT02637687 (Ongoing)	Larotrectinib	155	Objective response rates	Not yet reported

			1/2	Trial)				
NCT02576431[118]	2025	Adults/Pediatrics	Phase 2	NCT02576431 (Ongoing Trial)	Larotrectinib	204	Objective response rates, PFS, OS, Safety	Not yet reported
Desai <i>et al</i> [119], 2022	2025	Adults/Pediatrics	Phase 1/2	NCT02650401 (Ongoing Trial)	Entrectinib	69	Maximum Tolerated Dose and Objective response rates	Not yet reported
IDH inhibitors								
NCT05588141[120]	2029	Adults	Phase 1/2	NCT05588141 (Ongoing Trial)	Zotiraciclib	96	12-months PFS	Not yet reported
Mellinghoff <i>et al</i> [121], 2023	2027	Adults	Phase 3	NCT04164901	Vorasidenib	340	PFS	Significantly improved PFS
Mellinghoff <i>et al</i> [122], 2019	2024	Adults	Phase 1	NCT03343197	AG-120, AG881	49	2-hydroxyglutarate concentrationin resectedtumors	decreased tumorcell proliferationand immune cellactivation
EGFR inhibitors								
Weller <i>et al</i> [123], 2017	2016	Adults	Phase 3	NCT01480479	Rindopepimut/Temozolomide	745	OS	Median OS was 20.1 months in the Rindopepimut group versus 20.0 months in the control group
Lassman <i>et al</i> [124], 2023	2022	Adults	Phase 3	NCT02573324	Depatuxizumab mafodotin	691	OS	No OS benefit for depatux-m in treating EGFR-amp newly diagnosed GBM

PFS: Progression-free survival; OS: Overall survival.

Entrectinib

Entrectinib, approved by both the United States Food and Drug Administration and European Medicines Agency for tumors containing TRK or ROS1 fusions[40], encounters a challenge in treating brain neoplasms due to the blood-brain barrier (BBB)[35]. Effective targeted therapies for leptomenigeal disseminated tumors depend on their ability to penetrate this barrier. Although entrectinib, designed to cross the BBB, has demonstrated promise with a 79% objective response rate in various solid tumors, including CNS tumors, information on its cerebrospinal fluid penetrance in brain tumor patients is currently lacking[41].

The potential therapeutic efficacy of entrectinib, a selective pan TRK inhibitor, has been explored in patients with leptomenigeal disseminated pediatric high-grade gliomas (pHGG) harboring NTRK or ROS1 fusions[42,43]. The STARTRK-NG trial reported positive radiographic responses in four pHGG patients treated with entrectinib, indicating promise for CNS tumors[44,45]. This study investigated the in vitro sensitivity of pHGG cell models to entrectinib and suggested potential combination therapies[46]. The need for further studies to understand resistance mechanisms is emphasized, along with the generally well-tolerated nature of entrectinib. The observed CNS penetrance of entrectinib in a gliosarcoma patient has been discussed, highlighting its ability to cross the blood-brain barrier[47]. The text also considers the combination of entrectinib with radiotherapy and suggests the importance of intrathecal therapy in cases of leptomenigeal dissemination[41]. This conclusion underscores the need for comprehensive investigations and prospective clinical studies to establish the role of entrectinib and potential combination therapies in pHGG with ROS1/ NTRK fusions.




Figure 1 Illustrates a flow chart with drugs according to mutation.

ONC201 and paxalisib

ONC201, an oral small-molecule imipridone anticancer therapy, has demonstrated early clinical success in patients with diffuse intrinsic pontine glioma (DIPG)[48] and recurrent H3K27M-mutant diffuse midline gliomas[49]. Investigations across various cancer types have shown that ONC201-induced apoptosis in cancer cells, independent of p53, occurs through an atypical integrated stress response involving the expression of the antitumor protein TRAIL. This mechanism has shown promise in hematological[50], colorectal[51], breast[52], uterine cancers[53], and glioblastoma[54]. A sustained positive response was observed in a patient with secondary glioblastoma carrying an H3.3K27M mutation, prompting further exploration in patients with similar mutations, including those with DIPG[54].

Studies have discussed the therapeutic benefits of combining ONC201, a dopamine receptor D2 antagonist[55], with the blood-brain barrier-penetrant PI3K/Akt inhibitor, paxalisib, for treating DIPG. Mechanistic insights indicate that ONC201, by decreasing tyrosine hydroxylase expression, exhibits global DRD2 antagonism, with ClpP identified as a crucial target that causes mitochondrial dysfunction and oxidative stress[56]. The combination of paxalisib shows promising results in preclinical and preliminary clinical trials, leading to symptom resolution and tumor regression. Challenges related to immunologically cold tumor microenvironments in DIPG have been acknowledged, but potential changes in the epigenetic landscape and metabolic plasticity following ONC201 treatment may enhance immunogenicity [57,58]. The observed link between H3K27M mutations, metabolic changes, and the immune response highlights the complexity of DIPG treatment, presenting a potential avenue for the effective administration of therapy for glioblastoma [59].

RADIATION THERAPY

Radiation therapy (RT) is a successful management method for pediatric low-grade gliomas using both initial and salvage treatment approaches. Historically, RT was the chosen initial therapy for quickly progressing or unresectable tumors, with 10-year progression-free survival (PFS) and OS rates of 70% and 80%, respectively[60-62]. Furthermore, RT has been used as an adjuvant therapy, particularly when surgery is limited to partial resection or biopsy, particularly for tumors in the optic system, hypothalamus, deep midline tissues, and brainstem[63,64]. Adjuvant RT is suggested in cases of partial resection because PFS is greatly reduced[65,66]. However, there is a lack of agreement on its use, which is attributable in part to the paucity of randomized prospective studies[67,68].

For older children who have not responded to numerous systemic medications, RT is preferred as part of the care plan. Historically, postponing RT was motivated by concerns about RT-related toxicities such as cognitive impairment[69,70], endocrine dysfunction[71], secondary malignancies[72], vascular damage[72,73], and growth abnormalities[74]. The severity of these symptoms is directly related to the location of the tumor and the patient's age, particularly in patients under the age of 10[69,72].

An institutional evaluation covering a median follow-up of 11 years found 8-year PFS and OS rates of 83% and 100%, respectively[75]. Overall neurocognitive performance did not deteriorate in the trial; however, significant cognitive impairment was noted in young children (under 7 years old) and in patients who received high doses to the left temporal

lobe or hippocampus. Higher dosages to the hypothalamus or pituitary caused endocrine disruption, and two patients developed Moya disease. The 5-year PFS and OS rates in a recently published prospective research including 174 pediatric patients with LGG who received proton treatment were 84% and 92%, respectively, with a median follow-up of 4.4 years[76]. Four patients experienced severe late toxicities, including brainstem necrosis, symptomatic vasculopathy, radiation retinopathy, and fatal secondary cancers. While acknowledging the relevance of radiation-related damage, it is vital to emphasize that recent research has yielded promising outcomes. The extended latency of toxicity should be considered in light of the rapid developments in the field[77].

Concerns about RT-related toxicity originate mostly from long-term data collected from studies conducted during the 1970s and the 1990s using 2-dimensional RT methods that did not allow for accurate radiation dose administration. Significant technical progress has been achieved in reducing the radiation dose that reaches the normal structures surrounding the tumor. This began with the use of 3-dimensional conformal external beam RT (3D-CRT) and progressed in the 2000s with the advent of intensity-modulated RT (IMRT). Significantly, the introduction of proton therapy has reduced radiation exit dosage[78,79], contributing to its growing role in pediatric patients. Several studies have suggested that proton therapy might improve both patient quality of life and the cost-effectiveness of pediatric brain tumor treatment[80,81].

SURGICAL TREATMENT

The primary objective of glioma treatment is to strike a balance between preserving the patient's quality of life and improving PFS and OS[82,83]. The choice between oncological and surgical treatment depends on factors such as tumor size, location, and individual patient characteristics, including age and comorbidities[82,84,85]. Patients aged > 40 years at diagnosis, those with incomplete resection, and those with wild-type isocitrate dehydrogenase (IDH) status are typically considered to be at increased risk. The conventional treatment approach involves cytoreductive surgery to achieve gross total resection (GTR), followed by a combination of chemotherapy and/or radiation therapy[86,87].

The prognosis for gliomas, encompassing both LGG and HGG, is significantly influenced by the extent of surgical resection (EOSR) (Table 3). In LGG, EOSR is measured by the percentage of the FLAIR signal that is excised, whereas in HGG, it is determined by the removal of the percentage of enhancing tissue and the necrotic center. Extensive research on EOSR in LGG consistently shows that achieving GTR significantly improves survival rates, particularly among younger patients, classifying them as low-risk individuals compared to those who undergo only partial resection[82-84].

Similarly, investigations into EOSR in HGG consistently demonstrated a strong correlation between the extent of resection and survival outcomes, assuming no surgery-related neurological morbidities[88-90]. In most studies, the surgical goal is unequivocally defined as achieving GTR or complete tumor removal, typically amounting to 100% resection[91].

Recent clinical investigations have explored the concept of subtotal resection for gliomas[92]. This surgical approach aims to achieve GTR while simultaneously eliminating the FLAIR signal surrounding the necrotic and enhancing tumor mass in high-grade gliomas. Low-grade gliomas involve complete removal of the FLAIR signal along with additional radiographic extraction of the normal brain tissue adjacent to the tumor. Subtotal resection in LGG surgery was confirmed by observing that the resection cavity exceeded the initial FLAIR volume on the postoperative MRI at the three-month mark. Subtotal resection is considered justifiable when minimal neurological risks are involved, with the aim of eliminating invasive cells near the radiographic boundary[93-95]. Evidence from clinical case series of glioblastoma multiforme and HGG presents a conflicting picture, as performing supra total resection may entail an increased risk of neurological function decline, despite potential improvements in PFS and OS[92,96]. Additionally, there has been increased focus on the utilization of laser interstitial thermal (LIT) treatment for brain tumors. Recent trials investigating LIT have shown that achieving a greater level of ablation, including subtotal ablation, can lead to improved progression-free survival and overall survival outcomes in patients with HGG[97,98].

Intraoperative radiotherapy with a single high radiation dose administered following tumor resection, intraoperative radiotherapy (IORT), a novel and non-conventional form of radiotherapy, can eradicate any remaining tumor cells[99]. A wide range of cancers, including breast, pancreatic, lung, and colon cancers, have been treated with IORT[100-102]. The lack of a discernible increase in survival in IORT treatment reports for primary malignant gliomas has been ascribed to angle errors, low electrons, and small electron cones, which result in inadequate coverage of the target volume[103]. A mobile IORT unit, INTRABEAM (Zeiss, Oberkochen, Germany), can deliver an equal dose of low-energy radiation in all directions within a tumor cavity, along with spherical irradiation. According to research, IORT with low-energy X-rays increases glioblastoma patients' survival rates without causing new problems[104].

Vaccine development

Cancer vaccines targeting high-grade gliomas, predating coronavirus disease 2019, are gaining momentum. Strategies include peptide-based vaccines, dendritic cells, viral vectors, and personalized neoantigen vaccines. They are also being explored for the treatment of LGG. For IDH-mutant LGG, adjuvants such as poly (I:C) and poly-ICLC enhance immune responses, collectively reflecting a determined push for glioma immunotherapy[105]. To bolster the weak immune response in LGGs, synthetic double-stranded RNA molecules, such as polyinosinic acid homopolymers annealed to a polycytidylic acid homopolymer, have demonstrated potential[106]. They mimic viral infections and promote the release of interferon type 1 and other immune-boosting substances. Safely used as adjuvants with dendritic cells or peptide vaccines, they enhance therapeutic responses[106,107].

Table 3 Supratentorial surgical treatment options for glioma, *n* (%)

Ref.	Study origin	Study design	Total number of patients	Supratotal resection sample	Male, %	Age at resection	Permanent neurological deficits	Progression-free survival	Overall survival
Gajjar <i>et al</i> [63], 1997	United States	Cohort study	142	48 (68/142)	61	7 median (0.17-19)	Not reported	70 ± 5 at 4 years	90 ± 3 at 4 years
Fisher <i>et al</i> [67], 2008	United States	Cohort study	278	19 (52/278)	58	9.1 ± 0.3	Not reported	55 ± 3 at 5 years	87 ± 2 at 5 years
Wisoff <i>et al</i> [125], 2010	United States	Prospective trial	518	64 (332/518)	54	7.9 median (0.6-20.5)	Not reported	78 ± 2 at 8 years	96 ± 0.9 at 8 years
Yordanova <i>et al</i> [93], 2011	France	Case series	15	100.00	53.3	36.4 (24-59)	2, 13.3	73.3 at 38 months	100 at study end
Youland <i>et al</i> [11], 2013	United States	Retrospective cohort	351	67 (235/351)	55	10.9 (0.05-19.6)	Not reported	75.8 at 5 years	94.9 at 5 years
Lima <i>et al</i> [126], 2015	France	Case series	21	19.0 (4/21)	28.57	35 (18-57)	0, 0	100 at study end	100 at study end
Duffau <i>et al</i> [127], 2016	France	Cohort study	16	100.00	43.75	41.3 (26-63)	0, 0	50 relapse rate (avg 70 months)	100 at study end
Lima <i>et al</i> [92], 2017	France	Two-center prospective study	19	26.3 (5/19)	42.1	31.2 (19-51)	0, 0	100 at study end	100 at study end
Rossi <i>et al</i> [86], 2019	Italy	Case series	449	32 (145/449)	53.1	37.9 (median 36.5)	1, 0.69 (SupTR group)	Not reported	Not reported
Ng <i>et al</i> [128], 2020	France	Case series	74	28 (21/74)	41.89	35.7 (18-66)	0, 0	Not reported	100 at 5 years
Ng <i>et al</i> [129], 2020	France	Case series	47	26 (12/47)	34.04	39.2 ± 11.3	0, 0	Not reported	100 at study end
Goel <i>et al</i> [130], 2021	India	Cohort study	74	34 (25/74)	62.16	33 (21-55)	0, 0	98.7 at 2 years	100 at study end
Rossi <i>et al</i> [94], 2021	Italy	Case series	319	35 (110/319)	61.1	38.9 (18-75)	6, 1.9	94 at 92 months (SupTR group)	100 at 80 months (SupTR group)
Ius <i>et al</i> [131], 2022	United States, Canada, France, and Italy	Four center retrospective review	267	9 (24/267)	41.9	39.2 (18-71)	8, 3.1	Not reported	100 at 100 months (SupTR)

CONCLUSION

In conclusion, advancements in LGG treatment span immunotherapies, targeted therapies, radiation, surgery, and vaccine strategies. Immunotherapies like Zotiraciclib and Lerapolturev show promise, while targeted therapies such as Entrectinib and ONC201/Paxalisib combination demonstrate early success. Radiation therapy, evolving with proton therapy, remains crucial, and surgical approaches aim to achieve gross total resection. Cancer vaccines including synthetic RNA adjuvants have emerged. The evolving landscape underscores a shift toward personalized and targeted therapies, with ongoing research being essential for refining strategies and improving outcomes in LGG treatment.

FOOTNOTES

Author contributions: Shafique MA and Lucke-Wold B conceptualized the study; Danish F and Mustafa MS conducted a literature search; Drafting of the manuscript and writing was done by Nasrullah RMU, Rangwala BS, Siddiq MA, Zainab N and AH performed the editing; All authors read and agreed to the final version of the manuscript; All the authors have revised the manuscript accordingly.

Conflict-of-interest statement: All authors have no conflicts of interest to disclose.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the

original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: United States

ORCID number: Brandon Lucke-Wold 0000-0001-6577-4080; Burhanuddin Sohail Rangwala 0009-0008-5812-9049; Muhammad Ashir Shafique 0000-0000-7420-1292; Mohammad Arham Siddiq 0000-0002-8750-1419; Muhammad Saqlain Mustafa 0000-0002-3067-3543; Fnu Danish 0000-0003-0595-0315; Rana Muhammad Umer Nasrullah 0009-0004-0490-6464; Noor Zainab 0009-0009-4989-8679; Abdul Haseeb 0009-0004-6875-4850.

S-Editor: Liu JH

L-Editor: A

P-Editor: Yuan YY

REFERENCES

- Braunstein S, Raleigh D, Bindra R, Mueller S, Haas-Kogan D. Pediatric high-grade glioma: current molecular landscape and therapeutic approaches. *J Neurooncol* 2017; **134**: 541-549 [PMID: 28357536 DOI: 10.1007/s11060-017-2393-0]
- Stylli SS, Luwor RB, Ware TM, Tan F, Kaye AH. Mouse models of glioma. *J Clin Neurosci* 2015; **22**: 619-626 [PMID: 25698543 DOI: 10.1016/j.jocn.2014.10.013]
- Diwanji TP, Engelman A, Snider JW, Mohindra P. Epidemiology, diagnosis, and optimal management of glioma in adolescents and young adults. *Adolesc Health Med Ther* 2017; **8**: 99-113 [PMID: 28989289 DOI: 10.2147/AHMT.S53391]
- Lim-Fat MJ, Das S. Unique molecular, clinical, and treatment aspects of gliomas in adolescents and young adults: a review. *J Neurosurg* 2023; **139**: 1619-1627 [PMID: 37243561 DOI: 10.3171/2023.4.JNS.221800]
- Mughal ZUN, Ahmad TKF, Haseeb A, Shafique MA, Ahmdon OEA, Mahgoub AMA. Dabrafenib and trametinib as a promising treatment option for pediatric population with low-grade gliomas that have BRAF V600E mutation; a breakthrough in the field of neuro-oncology. *IJS Global Health* 2024; **7**: e0395 [DOI: 10.1097/GH9.0000000000000395]
- Bush NA, Chang SM, Berger MS. Current and future strategies for treatment of glioma. *Neurosurg Rev* 2017; **40**: 1-14 [PMID: 27085859 DOI: 10.1007/s10143-016-0709-8]
- Naeem A, Aziz N, Nasir M, Rangwala HS, Fatima H, Mubarak F. Accuracy of MRI in Detecting 1p/19q Co-deletion Status of Gliomas: A Single-Center Retrospective Study. *Cureus* 2024; **16**: e51863 [PMID: 38327950 DOI: 10.7759/cureus.51863]
- Eisenstat DD, Pollack IF, Demers A, Sapp MV, Lambert P, Weisfeld-Adams JD, Burger PC, Gilles F, Davis RL, Packer R, Boyett JM, Finlay JL. Impact of tumor location and pathological discordance on survival of children with midline high-grade gliomas treated on Children's Cancer Group high-grade glioma study CCG-945. *J Neurooncol* 2015; **121**: 573-581 [PMID: 25431150 DOI: 10.1007/s11060-014-1669-x]
- Bavle A, Chintagumpala M. Pediatric high-grade glioma: a review of biology, prognosis, and treatment. *J Radiat Oncol* 2018; **7**: 7-15 [DOI: 10.1007/s13566-018-0344-9]
- Sievert AJ, Fisher MJ. Pediatric low-grade gliomas. *J Child Neurol* 2009; **24**: 1397-1408 [PMID: 19841428 DOI: 10.1177/0883073809342005]
- Youland RS, Khwaja SS, Schomas DA, Keating GF, Wetjen NM, Laack NN. Prognostic factors and survival patterns in pediatric low-grade gliomas over 4 decades. *J Pediatr Hematol Oncol* 2013; **35**: 197-205 [PMID: 22983418 DOI: 10.1097/MPH.0b013e3182678bf8]
- Ris MD, Beebe DW. Neurodevelopmental outcomes of children with low-grade gliomas. *Dev Disabil Res Rev* 2008; **14**: 196-202 [PMID: 18924158 DOI: 10.1002/dvrr.27]
- Rees JH. Low-grade gliomas in adults. *Curr Opin Neurol* 2002; **15**: 657-661 [PMID: 12447102 DOI: 10.1097/01.wco.0000044760.39452.71]
- Grier JT, Batchelor T. Low-grade gliomas in adults. *Oncologist* 2006; **11**: 681-693 [PMID: 16794247 DOI: 10.1634/theoncologist.11-6-681]
- Su YT, Chen R, Wang H, Song H, Zhang Q, Chen LY, Lappin H, Vasconcelos G, Lita A, Maric D, Li A, Celiku O, Zhang W, Meetze K, Estok T, Larion M, Abu-Asab M, Zhuang Z, Yang C, Gilbert MR, Wu J. Novel Targeting of Transcription and Metabolism in Glioblastoma. *Clin Cancer Res* 2018; **24**: 1124-1137 [PMID: 29254993 DOI: 10.1158/1078-0432.CCR-17-2032]
- Hofmeister CC, Berdeja JG, Vesole DH, Suvannasankha A, Parrott T, Abonour R. TG02, an oral CDK9-inhibitor, in combination with carfilzomib demonstrated objective responses in carfilzomib refractory multiple myeloma patients. *Blood* 2015; **126**: 3052 [DOI: 10.1182/blood.V126.23.3052.3052]
- Nechaev S, Adelman K. Pol II waiting in the starting gates: Regulating the transition from transcription initiation into productive elongation. *Biochim Biophys Acta* 2011; **1809**: 34-45 [PMID: 21081187 DOI: 10.1016/j.bbagr.2010.11.001]
- Loyer P, Trembley JH, Katona R, Kidd VJ, Lahti JM. Role of CDK/cyclin complexes in transcription and RNA splicing. *Cell Signal* 2005; **17**: 1033-1051 [PMID: 15935619 DOI: 10.1016/j.cellsig.2005.02.005]
- Wu J, Yuan Y, Long Priel DA, Fink D, Peer CJ, Sissung TM, Su YT, Pang Y, Yu G, Butler MK, Mendoza TR, Vera E, Ahmad S, Bryla C, Lindsley M, Grajkowska E, Mentges K, Boris L, Antony R, Garren N, Siegel C, Lollo N, Cordova C, Aboud O, Theeler BJ, Burton EM, Penas-Prado M, Leeper H, Gonzales J, Armstrong TS, Calvo KR, Figg WD, Kuhns DB, Gallin JJ, Gilbert MR. Phase I Study of Zotiraciclib in Combination with Temozolomide for Patients with Recurrent High-grade Astrocytomas. *Clin Cancer Res* 2021; **27**: 3298-3306 [PMID: 33785481 DOI: 10.1158/1078-0432.CCR-20-4730]
- Filley AC, Henriquez M, Dey M. Recurrent glioma clinical trial, CheckMate-143: the game is not over yet. *Oncotarget* 2017; **8**: 91779-91794 [PMID: 29207684 DOI: 10.18632/oncotarget.21586]
- Le DT, Uram JN, Wang H, Bartlett BR, Kemberling H, Eyring AD, Skora AD, Luber BS, Azad NS, Laheru D, Biedrzycki B, Donehower RC, Zaheer A, Fisher GA, Crocenzi TS, Lee JJ, Duffy SM, Goldberg RM, de la Chapelle A, Koshiji M, Bhajee F, Huebner T, Hruban RH, Wood LD, Cuka N, Pardoll DM, Papadopoulos N, Kinzler KW, Zhou S, Cornish TC, Taube JM, Anders RA, Eshleman JR, Vogelstein B, Diaz LA Jr. PD-1 Blockade in Tumors with Mismatch-Repair Deficiency. *N Engl J Med* 2015; **372**: 2509-2520 [PMID: 26028255 DOI: 10.1056/NEJMoa1500596]
- Rizvi NA, Hellmann MD, Snyder A, Kvistborg P, Makarov V, Havel JJ, Lee W, Yuan J, Wong P, Ho TS, Miller ML, Rehkman N, Moreira

- AL, Ibrahim F, Bruggeman C, Gasmi B, Zappasodi R, Maeda Y, Sander C, Garon EB, Merghoub T, Wolchok JD, Schumacher TN, Chan TA. Cancer immunology. Mutational landscape determines sensitivity to PD-1 blockade in non-small cell lung cancer. *Science* 2015; **348**: 124-128 [PMID: 25765070 DOI: 10.1126/science.aaa1348]
- 23 **Tumeh PC**, Harview CL, Yearley JH, Shintaku IP, Taylor EJ, Robert L, Chmielowski B, Spasic M, Henry G, Ciobanu V, West AN, Carmona M, Kivork C, Seja E, Cherry G, Gutierrez AJ, Grogan TR, Mateus C, Tomasic G, Glaspy JA, Emerson RO, Robins H, Pierce RH, Elashoff DA, Robert C, Ribas A. PD-1 blockade induces responses by inhibiting adaptive immune resistance. *Nature* 2014; **515**: 568-571 [PMID: 25428505 DOI: 10.1038/nature13954]
 - 24 **Alexandrov LB**, Kim J, Haradhvala NJ, Huang MN, Tian Ng AW, Wu Y, Boot A, Covington KR, Gordenin DA, Bergstrom EN, Islam SMA, Lopez-Bigas N, Klimczak LJ, McPherson JR, Morganella S, Sabarinathan R, Wheeler DA, Mustonen V; PCAWG Mutational Signatures Working Group, Getz G, Rozen SG, Stratton MR; PCAWG Consortium. The repertoire of mutational signatures in human cancer. *Nature* 2020; **578**: 94-101 [PMID: 32025018 DOI: 10.1038/s41586-020-1943-3]
 - 25 **Zhao J**, Chen AX, Gartrell RD, Silverman AM, Aparicio L, Chu T, Bordbar D, Shan D, Samanamud J, Mahajan A, Filip I, Orenbuch R, Goetz M, Yamaguchi JT, Cloney M, Horbinski C, Lukas RV, Raizer J, Rae AI, Yuan J, Canoll P, Bruce JN, Saenger YM, Sims P, Iwamoto FM, Sonabend AM, Rabadan R. Immune and genomic correlates of response to anti-PD-1 immunotherapy in glioblastoma. *Nat Med* 2019; **25**: 462-469 [PMID: 30742119 DOI: 10.1038/s41591-019-0349-y]
 - 26 **Brown MC**, Mosaheb MM, Mohme M, McKay ZP, Holl EK, Kastan JP, Yang Y, Beasley GM, Hwang ES, Ashley DM, Bigner DD, Nair SK, Gromeier M. Viral infection of cells within the tumor microenvironment mediates antitumor immunotherapy via selective TBK1-IRF3 signaling. *Nat Commun* 2021; **12**: 1858 [PMID: 33767151 DOI: 10.1038/s41467-021-22088-1]
 - 27 **Mosaheb MM**, Dobrikova EY, Brown MC, Yang Y, Cable J, Okada H, Nair SK, Bigner DD, Ashley DM, Gromeier M. Genetically stable poliovirus vectors activate dendritic cells and prime antitumor CD8 T cell immunity. *Nat Commun* 2020; **11**: 524 [PMID: 31988324 DOI: 10.1038/s41467-019-13939-z]
 - 28 **Brown MC**, Holl EK, Boczkowski D, Dobrikova E, Mosaheb M, Chandramohan V, Bigner DD, Gromeier M, Nair SK. Cancer immunotherapy with recombinant poliovirus induces IFN-dominant activation of dendritic cells and tumor antigen-specific CTLs. *Sci Transl Med* 2017; **9** [PMID: 28931654 DOI: 10.1126/scitranslmed.aan4220]
 - 29 **Gromeier M**, Alexander L, Wimmer E. Internal ribosomal entry site substitution eliminates neurovirulence in intergeneric poliovirus recombinants. *Proc Natl Acad Sci USA* 1996; **93**: 2370-2375 [PMID: 8637880 DOI: 10.1073/pnas.93.6.2370]
 - 30 **Thompson EM**, Landi D, Brown MC, Friedman HS, McLendon R, Herndon JE 2nd, Buckley E, Bolognesi DP, Lipp E, Schroeder K, Becher OJ, Friedman AH, McKay Z, Walter A, Threalt S, Jagers D, Desjardins A, Gromeier M, Bigner DD, Ashley DM. Recombinant polio-rhinovirus immunotherapy for recurrent paediatric high-grade glioma: a phase 1b trial. *Lancet Child Adolesc Health* 2023; **7**: 471-478 [PMID: 37004712 DOI: 10.1016/S2352-4642(23)00031-7]
 - 31 **Mackay A**, Burford A, Carvalho D, Izquierdo E, Fazal-Salom J, Taylor KR, Bjerke L, Clarke M, Vinci M, Nandhabalan M, Temelso S, Popov S, Molinari V, Raman P, Waanders AJ, Han HJ, Gupta S, Marshall L, Zacharoulis S, Vaidya S, Mandeville HC, Bridges LR, Martin AJ, Al-Sarraj S, Chandler C, Ng HK, Li X, Mu K, Trabelsi S, Ibrahim DH, Kisljakov AN, Konovalov DM, Moore AS, Carcaboso AM, Sunol M, de Torres C, Cruz O, Mora J, Shats LI, Stavale JN, Bidinotto LT, Reis RM, Entz-Werle N, Farrell M, Cryan J, Crimmins D, Caird J, Pears J, Monje M, Debily MA, Castel D, Grill J, Hawkins C, Nikbakht H, Jabado N, Baker SJ, Pfister SM, Jones DTW, Fouladi M, von Bueren AO, Baudis M, Resnick A, Jones C. Integrated Molecular Meta-Analysis of 1,000 Pediatric High-Grade and Diffuse Intrinsic Pontine Glioma. *Cancer Cell* 2017; **32**: 520-537.e5 [PMID: 28966033 DOI: 10.1016/j.ccell.2017.08.017]
 - 32 **Jungbluth AA**, Stockert E, Huang HJ, Collins VP, Coplan K, Iversen K, Kolb D, Johns TJ, Scott AM, Gullick WJ, Ritter G, Cohen L, Scanlan MJ, Cavenee WK, Old LJ. A monoclonal antibody recognizing human cancers with amplification/overexpression of the human epidermal growth factor receptor. *Proc Natl Acad Sci USA* 2003; **100**: 639-644 [PMID: 12515857 DOI: 10.1073/pnas.232686499]
 - 33 **Gan HK**, Burgess AW, Clayton AH, Scott AM. Targeting of a conformationally exposed, tumor-specific epitope of EGFR as a strategy for cancer therapy. *Cancer Res* 2012; **72**: 2924-2930 [PMID: 22659454 DOI: 10.1158/0008-5472.CAN-11-3898]
 - 34 **Doronina SO**, Bovee TD, Meyer DW, Miyamoto JB, Anderson ME, Morris-Tilden CA, Senter PD. Novel peptide linkers for highly potent antibody-auristatin conjugate. *Bioconjug Chem* 2008; **19**: 1960-1963 [PMID: 18803412 DOI: 10.1021/bc800289a]
 - 35 **Brastianos PK**, Ippen FM, Hafeez U, Gan HK. Emerging Gene Fusion Drivers in Primary and Metastatic Central Nervous System Malignancies: A Review of Available Evidence for Systemic Targeted Therapies. *Oncologist* 2018; **23**: 1063-1075 [PMID: 29703764 DOI: 10.1634/theoncologist.2017-0614]
 - 36 **Birrer MJ**, Moore KN, Betella I, Bates RC. Antibody-Drug Conjugate-Based Therapeutics: State of the Science. *J Natl Cancer Inst* 2019; **111**: 538-549 [PMID: 30859213 DOI: 10.1093/jnci/djz035]
 - 37 **Gan HK**, van den Bent M, Lassman AB, Reardon DA, Scott AM. Antibody-drug conjugates in glioblastoma therapy: the right drugs to the right cells. *Nat Rev Clin Oncol* 2017; **14**: 695-707 [PMID: 28675164 DOI: 10.1038/nrclinonc.2017.95]
 - 38 **Marin BM**, Porath KA, Jain S, Kim M, Conage-Pough JE, Oh JH, Miller CL, Talele S, Kitange GJ, Tian S, Burgenske DM, Mladek AC, Gupta SK, Decker PA, McMinn MH, Stopka SA, Regan MS, He L, Carlson BL, Bakken K, Burns TC, Parney IF, Giannini C, Agar NYR, Eckel-Passow JE, Cochran JR, Elmquist WF, Vaubel RA, White FM, Sarkaria JN. Heterogeneous delivery across the blood-brain barrier limits the efficacy of an EGFR-targeting antibody drug conjugate in glioblastoma. *Neuro Oncol* 2021; **23**: 2042-2053 [PMID: 34050676 DOI: 10.1093/neuonc/noab133]
 - 39 **Gan HK**, Parakh S, Lassman AB, Seow A, Lau E, Lee ST, Ameratunga M, Perchyonok Y, Cao D, Burvenich IJG, O'Keefe GJ, Rigopoulos A, Gomez E, Maag D, Scott AM. Tumor volumes as a predictor of response to the anti-EGFR antibody drug conjugate depatuxizumab mafadotin. *Neurooncol Adv* 2021; **3**: vdab102 [PMID: 34549181 DOI: 10.1093/noonjnl/vdab102]
 - 40 **Cocco E**, Scaltriti M, Drilon A. NTRK fusion-positive cancers and TRK inhibitor therapy. *Nat Rev Clin Oncol* 2018; **15**: 731-747 [PMID: 30333516 DOI: 10.1038/s41571-018-0113-0]
 - 41 **Doebele RC**, Drilon A, Paz-Ares L, Siena S, Shaw AT, Farago AF, Blakely CM, Seto T, Cho BC, Tosi D, Besse B, Chawla SP, Bazhenova L, Krauss JC, Chae YK, Barve M, Garrido-Laguna I, Liu SV, Conkling P, John T, Fakih M, Sigal D, Loong HH, Buchschacher GL Jr, Garrido P, Nieva J, Steuer C, Overbeck TR, Bowles DW, Fox E, Riehl T, Chow-Maneval E, Simmons B, Cui N, Johnson A, Eng S, Wilson TR, Demetri GD; trial investigators. Entrectinib in patients with advanced or metastatic NTRK fusion-positive solid tumours: integrated analysis of three phase 1-2 trials. *Lancet Oncol* 2020; **21**: 271-282 [PMID: 31838007 DOI: 10.1016/S1470-2045(19)30691-6]
 - 42 **Drilon A**, Siena S, Ou SI, Patel M, Ahn MJ, Lee J, Bauer TM, Farago AF, Wheler JJ, Liu SV, Doebele R, Giannetta L, Cerea G, Marrapese G, Schirru M, Amatu A, Bencardino K, Palmeri L, Sartore-Bianchi A, Vanzulli A, Cresta S, Damiani S, Duca M, Ardini E, Li G, Christiansen J, Kowalski K, Johnson AD, Patel R, Luo D, Chow-Maneval E, Hornby Z, Multani PS, Shaw AT, De Braud FG. Safety and Antitumor Activity

- of the Multitargeted Pan-TRK, ROS1, and ALK Inhibitor Entrectinib: Combined Results from Two Phase I Trials (ALKA-372-001 and STARTRK-1). *Cancer Discov* 2017; **7**: 400-409 [PMID: [28183697](#) DOI: [10.1158/2159-8290.CD-16-1237](#)]
- 43 **Farago AF**, Le LP, Zheng Z, Muzikansky A, Drilon A, Patel M, Bauer TM, Liu SV, Ou SH, Jackman D, Costa DB, Multani PS, Li GG, Hornby Z, Chow-Maneval E, Luo D, Lim JE, Iafrate AJ, Shaw AT. Durable Clinical Response to Entrectinib in NTRK1-Rearranged Non-Small Cell Lung Cancer. *J Thorac Oncol* 2015; **10**: 1670-1674 [PMID: [26565381](#) DOI: [10.1097/01.JTO.0000473485.38553.f0](#)]
- 44 **Alvarez-Breckenridge C**, Miller JJ, Nayyar N, Gill CM, Kaneb A, D'Andrea M, Le LP, Lee J, Cheng J, Zheng Z, Butler WE, Multani P, Chow Maneval E, Ha Paek S, Toyota BD, Dias-Santagata D, Santagata S, Romero J, Shaw AT, Farago AF, Yip S, Cahill DP, Batchelor TT, Iafrate AJ, Brastianos PK. Clinical and radiographic response following targeting of BCAN-NTRK1 fusion in glioneuronal tumor. *NPJ Precis Oncol* 2017; **1**: 5 [PMID: [29872694](#) DOI: [10.1038/s41698-017-0009-y](#)]
- 45 **Walter AW**, Kandula VVR, Shah N. Larotrectinib imaging response in low-grade glioma. *Pediatr Blood Cancer* 2020; **67**: e28002 [PMID: [31544356](#) DOI: [10.1002/pbc.28002](#)]
- 46 **Mayr L**, Guntner AS, Madlener S, Schmook MT, Peyrl A, Azizi AA, Dieckmann K, Reisinger D, Stepien NM, Schramm K, Laemmerer A, Jones DTW, Ecker J, Sahm F, Milde T, Pajtlér KW, Blattner-Johnson M, Strbac M, Dorfer C, Czech T, Kirchhofer D, Gabler L, Berger W, Haberler C, Müllauer L, Buchberger W, Slave I, Lötsch-Gojo D, Gojo J. Cerebrospinal Fluid Penetration and Combination Therapy of Entrectinib for Disseminated ROS1/NTRK-Fusion Positive Pediatric High-Grade Glioma. *J Pers Med* 2020; **10** [PMID: [33353026](#) DOI: [10.3390/jpm10040290](#)]
- 47 **Fischer H**, Ullah M, de la Cruz CC, Hunsaker T, Senn C, Wirz T, Wagner B, Draganov D, Vazvaei F, Donzelli M, Paehler A, Merchant M, Yu L. Entrectinib, a TRK/ROS1 inhibitor with anti-CNS tumor activity: differentiation from other inhibitors in its class due to weak interaction with P-glycoprotein. *Neuro Oncol* 2020; **22**: 819-829 [PMID: [32383735](#) DOI: [10.1093/neuonc/noaa052](#)]
- 48 **Duchatel RJ**, Mannan A, Woldu AS, Hawtrey T, Hindley PA, Douglas AM, Jackson ER, Findlay IJ, Germon ZP, Staudt D, Kearney PS, Smith ND, Hindley KE, Cain JE, André N, La Madrid AM, Nixon B, De Iuliis GN, Nazarian J, Irish K, Alvaro F, Eisenstat DD, Beck A, Vitanza NA, Mueller S, Morris JC, Dun MD. Preclinical and clinical evaluation of German-sourced ONC201 for the treatment of H3K27M-mutant diffuse intrinsic pontine glioma. *Neurooncol Adv* 2021; **3**: vdab169 [PMID: [34988452](#) DOI: [10.1093/oaajnl/vdab169](#)]
- 49 **Venneti S**, Kawakibi AR, Ji S, Waszak SM, Sweha SR, Mota M, Pun M, Deogharkar A, Chung C, Tarapore RS, Ramage S, Chi A, Wen PY, Arrillaga-Romany I, Batchelor TT, Butowski NA, Sumrall A, Shonka N, Harrison RA, de Groot J, Mehta M, Hall MD, Daghistani D, Cloughesy TF, Ellingson BM, Beccaria K, Varlet P, Kim MM, Umemura Y, Garton H, Franson A, Schwartz J, Jain R, Kachman M, Baum H, Burant CF, Mottl SL, Cartaxo RT, John V, Messinger D, Qin T, Peterson E, Sajjakulnukit P, Ravi K, Waugh A, Walling D, Ding Y, Xia Z, Schwendeman A, Hawes D, Yang F, Judkins AR, Wahl D, Lyssiotis CA, de la Nava D, Alonso MM, Eze A, Spitzer J, Schmidt SV, Duchatel RJ, Dun MD, Cain JE, Jiang L, Stopka SA, Baquer G, Regan MS, Filbin MG, Agar NYR, Zhao L, Kumar-Sinha C, Mody R, Chinnaiyan A, Kurokawa R, Pratt D, Yadav VN, Grill J, Kline C, Mueller S, Resnick A, Nazarian J, Allen JE, Odia Y, Gardner SL, Koschmann C. Clinical Efficacy of ONC201 in H3K27M-Mutant Diffuse Midline Gliomas Is Driven by Disruption of Integrated Metabolic and Epigenetic Pathways. *Cancer Discov* 2023; **13**: 2370-2393 [PMID: [37584601](#) DOI: [10.1158/2159-8290.CD-23-0131](#)]
- 50 **Ishizawa J**, Kojima K, Chachad D, Ruvolo P, Ruvolo V, Jacamo RO, Borthakur G, Mu H, Zeng Z, Tabe Y, Allen JE, Wang Z, Ma W, Lee HC, Orlowski R, Sarbassov dos D, Lorenzi PL, Huang X, Neelapu SS, McDonnell T, Miranda RN, Wang M, Kantarjian H, Konopleva M, Davis RE, Andreoff M. ATF4 induction through an atypical integrated stress response to ONC201 triggers p53-independent apoptosis in hematological malignancies. *Sci Signal* 2016; **9**: ra17 [PMID: [26884599](#) DOI: [10.1126/scisignal.aac4380](#)]
- 51 **Wagner J**, Kline CL, Zhou L, Khazak V, El-Deiry WS. Anti-tumor effects of ONC201 in combination with VEGF-inhibitors significantly impacts colorectal cancer growth and survival in vivo through complementary non-overlapping mechanisms. *J Exp Clin Cancer Res* 2018; **37**: 11 [PMID: [29357916](#) DOI: [10.1186/s13046-018-0671-0](#)]
- 52 **Greer YE**, Porat-Shliom N, Nagashima K, Stuelten C, Crooks D, Koparde VN, Gilbert SF, Islam C, Ubaldini A, Ji Y, Gattinoni L, Soheilian F, Wang X, Hafner M, Shetty J, Tran B, Jailwala P, Cam M, Lang M, Voeller D, Reinhold WC, Rajapakse V, Pommier Y, Weigert R, Linehan WM, Lipkowitz S. ONC201 kills breast cancer cells in vitro by targeting mitochondria. *Oncotarget* 2018; **9**: 18454-18479 [PMID: [29719618](#) DOI: [10.18632/oncotarget.24862](#)]
- 53 **Fang Z**, Wang J, Clark LH, Sun W, Yin Y, Kong W, Pierce SR, West L, Sullivan SA, Tran AQ, Prabhu VV, Zhou C, Bae-Jump V. ONC201 demonstrates anti-tumorigenic and anti-metastatic activity in uterine serous carcinoma in vitro. *Am J Cancer Res* 2018; **8**: 1551-1563 [PMID: [30210923](#)]
- 54 **Arrillaga-Romany I**, Chi AS, Allen JE, Oster W, Wen PY, Batchelor TT. A phase 2 study of the first imipridone ONC201, a selective DRD2 antagonist for oncology, administered every three weeks in recurrent glioblastoma. *Oncotarget* 2017; **8**: 79298-79304 [PMID: [29108308](#) DOI: [10.18632/oncotarget.17837](#)]
- 55 **Madhukar NS**, Khade PK, Huang L, Gayvert K, Galletti G, Stogniew M, Allen JE, Giannakakou P, Elemento O. A Bayesian machine learning approach for drug target identification using diverse data types. *Nat Commun* 2019; **10**: 5221 [PMID: [31745082](#) DOI: [10.1038/s41467-019-12928-6](#)]
- 56 **Pruss M**, Dwucet A, Tanriover M, Hlavac M, Kast RE, Debatin KM, Wirtz CR, Halatsch ME, Siegelin MD, Westhoff MA, Karpel-Massler G. Dual metabolic reprogramming by ONC201/TIC10 and 2-Deoxyglucose induces energy depletion and synergistic anti-cancer activity in glioblastoma. *Br J Cancer* 2020; **122**: 1146-1157 [PMID: [32115576](#) DOI: [10.1038/s41416-020-0759-0](#)]
- 57 **Chung C**, Sweha SR, Pratt D, Tamrazi B, Panwalkar P, Banda A, Bayliss J, Hawes D, Yang F, Lee HJ, Shan M, Cieslik M, Qin T, Werner CK, Wahl DR, Lyssiotis CA, Bian Z, Shotwell JB, Yadav VN, Koschmann C, Chinnaiyan AM, Blüml S, Judkins AR, Venneti S. Integrated Metabolic and Epigenomic Reprogramming by H3K27M Mutations in Diffuse Intrinsic Pontine Gliomas. *Cancer Cell* 2020; **38**: 334-349.e9 [PMID: [32795401](#) DOI: [10.1016/j.ccell.2020.07.008](#)]
- 58 **Allen JE**, Kringsfeld G, Mayes PA, Patel L, Dicker DT, Patel AS, Dolloff NG, Messaris E, Scata KA, Wang W, Zhou JY, Wu GS, El-Deiry WS. Dual inactivation of Akt and ERK by TIC10 signals Foxo3a nuclear translocation, TRAIL gene induction, and potent antitumor effects. *Sci Transl Med* 2013; **5**: 171ra17 [PMID: [23390247](#) DOI: [10.1126/scitranslmed.3004828](#)]
- 59 **Harutyunyan AS**, Krug B, Chen H, Papillon-Cavanagh S, Zeinieh M, De Jay N, Deshmukh S, Chen CCL, Belle J, Mikael LG, Marchione DM, Li R, Nikbakht H, Hu B, Cagnone G, Cheung WA, Mohammadnia A, Bechet D, Faury D, McConechy MK, Pathania M, Jain SU, Ellezam B, Weil AG, Montpetit A, Salomoni P, Pastinen T, Lu C, Lewis PW, Garcia BA, Kleinman CL, Jabado N, Majewski J. H3K27M induces defective chromatin spread of PRC2-mediated repressive H3K27me2/me3 and is essential for glioma tumorigenesis. *Nat Commun* 2019; **10**: 1262 [PMID: [30890717](#) DOI: [10.1038/s41467-019-09140-x](#)]
- 60 **Taveras JM**, Mount LA, Wood EH. The value of radiation therapy in the management of glioma of the optic nerves and chiasm. *Radiology* 1956; **66**: 518-528 [PMID: [13323290](#) DOI: [10.1148/66.4.518](#)]

- 61 **Erkal HS**, Serin M, Cakmak A. Management of optic pathway and chiasmatic-hypothalamic gliomas in children with radiation therapy. *Radiother Oncol* 1997; **45**: 11-15 [PMID: [9364626](#) DOI: [10.1016/s0167-8140\(97\)00102-3](#)]
- 62 **Cappelli C**, Grill J, Raquin M, Pierre-Kahn A, Lellouch-Tubiana A, Terrier-Lacombe MJ, Habrand JL, Couanet D, Brauner R, Rodriguez D, Hartmann O, Kalifa C. Long-term follow up of 69 patients treated for optic pathway tumours before the chemotherapy era. *Arch Dis Child* 1998; **79**: 334-338 [PMID: [9875044](#) DOI: [10.1136/adc.79.4.334](#)]
- 63 **Gajjar A**, Sanford RA, Heideman R, Jenkins JJ, Walter A, Li Y, Langston JW, Muhlbauer M, Boyett JM, Kun LE. Low-grade astrocytoma: a decade of experience at St. Jude Children's Research Hospital. *J Clin Oncol* 1997; **15**: 2792-2799 [PMID: [9256121](#) DOI: [10.1200/jco.1997.15.8.2792](#)]
- 64 **Armstrong GT**, Conklin HM, Huang S, Srivastava D, Sanford R, Ellison DW, Merchant TE, Hudson MM, Hoehn ME, Robison LL, Gajjar A, Morris EB. Survival and long-term health and cognitive outcomes after low-grade glioma. *Neuro Oncol* 2011; **13**: 223-234 [PMID: [21177781](#) DOI: [10.1093/neuonc/nuq178](#)]
- 65 **Oh KS**, Hung J, Robertson PL, Garton HJ, Muraszko KM, Sandler HM, Hamstra DA. Outcomes of multidisciplinary management in pediatric low-grade gliomas. *Int J Radiat Oncol Biol Phys* 2011; **81**: e481-e488 [PMID: [21470783](#) DOI: [10.1016/j.ijrobp.2011.01.019](#)]
- 66 **Mishra KK**, Puri DR, Missett BT, Lamborn KR, Prados MD, Berger MS, Banerjee A, Gupta N, Wara WM, Haas-Kogan DA. The role of up-front radiation therapy for incompletely resected pediatric WHO grade II low-grade gliomas. *Neuro Oncol* 2006; **8**: 166-174 [PMID: [16495375](#) DOI: [10.1215/15228517-2005-011](#)]
- 67 **Fisher PG**, Tihan T, Goldthwaite PT, Wharam MD, Carson BS, Weingart JD, Repka MX, Cohen KJ, Burger PC. Outcome analysis of childhood low-grade astrocytomas. *Pediatr Blood Cancer* 2008; **51**: 245-250 [PMID: [18386785](#) DOI: [10.1002/pbc.21563](#)]
- 68 **Packer RJ**, Sutton LN, Atkins TE, Radcliffe J, Bunin GR, D'Angio G, Siegel KR, Schut L. A prospective study of cognitive function in children receiving whole-brain radiotherapy and chemotherapy: 2-year results. *J Neurosurg* 1989; **70**: 707-713 [PMID: [2709111](#) DOI: [10.3171/jns.1989.70.5.0707](#)]
- 69 **Merchant TE**, Conklin HM, Wu S, Lustig RH, Xiong X. Late effects of conformal radiation therapy for pediatric patients with low-grade glioma: prospective evaluation of cognitive, endocrine, and hearing deficits. *J Clin Oncol* 2009; **27**: 3691-3697 [PMID: [19581535](#) DOI: [10.1200/JCO.2008.21.2738](#)]
- 70 **Brauner R**, Malandry F, Rappaport R, Zucker JM, Kalifa C, Pierre-Kahn A, Bataini P, Dufier JL. Growth and endocrine disorders in optic glioma. *Eur J Pediatr* 1990; **149**: 825-828 [PMID: [2226566](#) DOI: [10.1007/bf02072067](#)]
- 71 **Tsang DS**, Murphy ES, Merchant TE. Radiation Therapy for Optic Pathway and Hypothalamic Low-Grade Gliomas in Children. *Int J Radiat Oncol Biol Phys* 2017; **99**: 642-651 [PMID: [29280458](#) DOI: [10.1016/j.ijrobp.2017.07.023](#)]
- 72 **Bowers DC**, Mulne AF, Reisch JS, Elterman RD, Munoz L, Booth T, Shapiro K, Dosey DL. Nonperioperative strokes in children with central nervous system tumors. *Cancer* 2002; **94**: 1094-1101 [PMID: [11920480](#) DOI: [10.1002/cncr.10353.abs](#)]
- 73 **Armstrong GT**, Liu Q, Yasui Y, Huang S, Ness KK, Leisenring W, Hudson MM, Donaldson SS, King AA, Stovall M, Krull KR, Robison LL, Packer RJ. Long-term outcomes among adult survivors of childhood central nervous system malignancies in the Childhood Cancer Survivor Study. *J Natl Cancer Inst* 2009; **101**: 946-958 [PMID: [19535780](#) DOI: [10.1093/jnci/djp148](#)]
- 74 **Greenberger BA**, Pulsifer MB, Ebb DH, MacDonald SM, Jones RM, Butler WE, Huang MS, Marcus KJ, Oberg JA, Tarbell NJ, Yock TI. Clinical outcomes and late endocrine, neurocognitive, and visual profiles of proton radiation for pediatric low-grade gliomas. *Int J Radiat Oncol Biol Phys* 2014; **89**: 1060-1068 [PMID: [25035209](#) DOI: [10.1016/j.ijrobp.2014.04.053](#)]
- 75 **Indelicato DJ**, Rotondo RL, Uezono H, Sandler ES, Aldana PR, Ranalli NJ, Beier AD, Morris CG, Bradley JA. Outcomes Following Proton Therapy for Pediatric Low-Grade Glioma. *Int J Radiat Oncol Biol Phys* 2019; **104**: 149-156 [PMID: [30684665](#) DOI: [10.1016/j.ijrobp.2019.01.078](#)]
- 76 **Bitterman DS**, MacDonald SM, Yock TI, Tarbell NJ, Wright KD, Chi SN, Marcus KJ, Haas-Kogan DA. Revisiting the Role of Radiation Therapy for Pediatric Low-Grade Glioma. *J Clin Oncol* 2019; **37**: 3335-3339 [PMID: [31498029](#) DOI: [10.1200/JCO.19.01270](#)]
- 77 **Harrabi SB**, Bougatf N, Mohr A, Haberer T, Herfarth K, Combs SE, Debus J, Adeberg S. Dosimetric advantages of proton therapy over conventional radiotherapy with photons in young patients and adults with low-grade glioma. *Strahlenther Onkol* 2016; **192**: 759-769 [PMID: [27363701](#) DOI: [10.1007/s00066-016-1005-9](#)]
- 78 **Takizawa D**, Mizumoto M, Yamamoto T, Oshiro Y, Fukushima H, Fukushima T, Terunuma T, Okumura T, Tsuboi K, Sakurai H. A comparative study of dose distribution of PBT, 3D-CRT and IMRT for pediatric brain tumors. *Radiat Oncol* 2017; **12**: 40 [PMID: [28228150](#) DOI: [10.1186/s13014-017-0775-2](#)]
- 79 **Yock TI**, Bhat S, Szymonifka J, Yeap BY, Delahaye J, Donaldson SS, MacDonald SM, Pulsifer MB, Hill KS, DeLaney TF, Ebb D, Huang M, Tarbell NJ, Fisher PG, Kuhlthau KA. Quality of life outcomes in proton and photon treated pediatric brain tumor survivors. *Radiother Oncol* 2014; **113**: 89-94 [PMID: [25304720](#) DOI: [10.1016/j.radonc.2014.08.017](#)]
- 80 **Verma V**, Mishra MV, Mehta MP. A systematic review of the cost and cost-effectiveness studies of proton radiotherapy. *Cancer* 2016; **122**: 1483-1501 [PMID: [26828647](#) DOI: [10.1002/cncr.29882](#)]
- 81 **Jakola AS**, Skjalsvik AJ, Myrnes KS, Sjøvik K, Unsgård G, Torp SH, Aaberg K, Berg T, Dai HY, Johnsen K, Kloster R, Solheim O. Surgical resection versus watchful waiting in low-grade gliomas. *Ann Oncol* 2017; **28**: 1942-1948 [PMID: [28475680](#) DOI: [10.1093/annonc/mdx230](#)]
- 82 **Bogdańska MU**, Bodnar M, Piotrowska MJ, Murek M, Schucht P, Beck J, Martínez-González A, Pérez-García VM. A mathematical model describes the malignant transformation of low grade gliomas: Prognostic implications. *PLoS One* 2017; **12**: e0179999 [PMID: [28763450](#) DOI: [10.1371/journal.pone.0179999](#)]
- 83 **Aghi MK**, Nahed BV, Sloan AE, Ryken TC, Kalkanis SN, Olson JJ. The role of surgery in the management of patients with diffuse low grade glioma: A systematic review and evidence-based clinical practice guideline. *J Neurooncol* 2015; **125**: 503-530 [PMID: [26530265](#) DOI: [10.1007/s11060-015-1867-1](#)]
- 84 **Opoku-Darko M**, Lang ST, Artindale J, Cairncross JG, Seivick RJ, Kelly JJP. Surgical management of incidentally discovered diffusely infiltrating low-grade glioma. *J Neurosurg* 2018; **129**: 19-26 [PMID: [28984519](#) DOI: [10.3171/2017.3.JNS17159](#)]
- 85 **Kurdi M**, Moshref RH, Katib Y, Faizo E, Najjar AA, Bahakeem B, Bamaga AK. Simple approach for the histomolecular diagnosis of central nervous system gliomas based on 2021 World Health Organization Classification. *World J Clin Oncol* 2022; **13**: 567-576 [PMID: [36157161](#) DOI: [10.5306/wjco.v13.i7.567](#)]
- 86 **Rossi M**, Ambrogi F, Gay L, Gallucci M, Conti Nibali M, Leonetti A, Puglisi G, Sciortino T, Howells H, Riva M, Pessina F, Navarria P, Franzese C, Simonelli M, Rudà R, Bello L. Is supratotal resection achievable in low-grade gliomas? Feasibility, putative factors, safety, and functional outcome. *J Neurosurg* 2019; **132**: 1692-1705 [PMID: [31100730](#) DOI: [10.3171/2019.2.JNS183408](#)]
- 87 **Hervey-Jumper SL**, Berger MS. Maximizing safe resection of low- and high-grade glioma. *J Neurooncol* 2016; **130**: 269-282 [PMID: [27363701](#) DOI: [10.1007/s00066-016-1005-9](#)]

- 27174197 DOI: [10.1007/s11060-016-2110-4](https://doi.org/10.1007/s11060-016-2110-4)]
- 88 **Brown TJ**, Brennan MC, Li M, Church EW, Brandmeir NJ, Rakszawski KL, Patel AS, Rizk EB, Suki D, Sawaya R, Glantz M. Association of the Extent of Resection With Survival in Glioblastoma: A Systematic Review and Meta-analysis. *JAMA Oncol* 2016; **2**: 1460-1469 [PMID: [27310651](https://pubmed.ncbi.nlm.nih.gov/27310651/) DOI: [10.1001/jamaoncol.2016.1373](https://doi.org/10.1001/jamaoncol.2016.1373)]
- 89 **Barnett GH**, Voigt JD, Alhuwalia MS. A Systematic Review and Meta-Analysis of Studies Examining the Use of Brain Laser Interstitial Thermal Therapy versus Craniotomy for the Treatment of High-Grade Tumors in or near Areas of Eloquence: An Examination of the Extent of Resection and Major Complication Rates Associated with Each Type of Surgery. *Stereotact Funct Neurosurg* 2016; **94**: 164-173 [PMID: [27322392](https://pubmed.ncbi.nlm.nih.gov/27322392/) DOI: [10.1159/000446247](https://doi.org/10.1159/000446247)]
- 90 **Kreatsoulas D**, Damante M, Gruber M, Duru O, Elder JB. Supratotal Surgical Resection for Low-Grade Glioma: A Systematic Review. *Cancers (Basel)* 2023; **15** [PMID: [37173957](https://pubmed.ncbi.nlm.nih.gov/37173957/) DOI: [10.3390/cancers15092493](https://doi.org/10.3390/cancers15092493)]
- 91 **de Leeuw CN**, Vogelbaum MA. Supratotal resection in glioma: a systematic review. *Neuro Oncol* 2019; **21**: 179-188 [PMID: [30321384](https://pubmed.ncbi.nlm.nih.gov/30321384/) DOI: [10.1093/neuonc/noy166](https://doi.org/10.1093/neuonc/noy166)]
- 92 **Lima GLO**, Dezamis E, Corns R, Rigaux-Viode O, Moritz-Gasser S, Roux A, Duffau H, Pallud J. Surgical resection of incidental diffuse gliomas involving eloquent brain areas. Rationale, functional, epileptological and oncological outcomes. *Neurochirurgie* 2017; **63**: 250-258 [PMID: [28161011](https://pubmed.ncbi.nlm.nih.gov/28161011/) DOI: [10.1016/j.neuchi.2016.08.007](https://doi.org/10.1016/j.neuchi.2016.08.007)]
- 93 **Yordanova YN**, Duffau H. Supratotal resection of diffuse gliomas - an overview of its multifaceted implications. *Neurochirurgie* 2017; **63**: 243-249 [PMID: [28185647](https://pubmed.ncbi.nlm.nih.gov/28185647/) DOI: [10.1016/j.neuchi.2016.09.006](https://doi.org/10.1016/j.neuchi.2016.09.006)]
- 94 **Rossi M**, Gay L, Ambroggi F, Conti Nibali M, Sciortino T, Puglisi G, Leonetti A, Mocellini C, Caroli M, Cordera S, Simonelli M, Pessina F, Navarria P, Pace A, Soffietti R, Rudà R, Riva M, Bello L. Association of supratotal resection with progression-free survival, malignant transformation, and overall survival in lower-grade gliomas. *Neuro Oncol* 2021; **23**: 812-826 [PMID: [33049063](https://pubmed.ncbi.nlm.nih.gov/33049063/) DOI: [10.1093/neuonc/noaa225](https://doi.org/10.1093/neuonc/noaa225)]
- 95 **Incekara F**, Koene S, Vincent AJPE, van den Bent MJ, Smits M. Association Between Supratotal Glioblastoma Resection and Patient Survival: A Systematic Review and Meta-Analysis. *World Neurosurg* 2019; **127**: 617-624.e2 [PMID: [31004858](https://pubmed.ncbi.nlm.nih.gov/31004858/) DOI: [10.1016/j.wneu.2019.04.092](https://doi.org/10.1016/j.wneu.2019.04.092)]
- 96 **Kaisman-Elbaz T**, Xiao T, Grabowski MM, Barnett GH, Mohammadi AM. The Impact of Extent of Ablation on Survival of Patients With Newly Diagnosed Glioblastoma Treated With Laser Interstitial Thermal Therapy: A Large Single-Institutional Cohort. *Neurosurgery* 2023; **93**: 427-435 [PMID: [36861990](https://pubmed.ncbi.nlm.nih.gov/36861990/) DOI: [10.1227/neu.0000000000002430](https://doi.org/10.1227/neu.0000000000002430)]
- 97 **Shah AH**, Semonche A, Eichberg DG, Borowy V, Luther E, Sarkiss CA, Morell A, Mahavadi AK, Ivan ME, Komotar RJ. The Role of Laser Interstitial Thermal Therapy in Surgical Neuro-Oncology: Series of 100 Consecutive Patients. *Neurosurgery* 2020; **87**: 266-275 [PMID: [31742351](https://pubmed.ncbi.nlm.nih.gov/31742351/) DOI: [10.1093/neuros/nyz424](https://doi.org/10.1093/neuros/nyz424)]
- 98 **Schueler P**, Micke O, Palkovic S, Schroeder J, Moustakis C, Bruns F, Schuck A, Wassmann H, Willich N. 12 years' experience with intraoperative radiotherapy (IORT) of malignant gliomas. *Strahlenther Onkol* 2005; **181**: 500-506 [PMID: [16044217](https://pubmed.ncbi.nlm.nih.gov/16044217/) DOI: [10.1007/s00066-005-1354-2](https://doi.org/10.1007/s00066-005-1354-2)]
- 99 **Vaidya JS**, Joseph DJ, Tobias JS, Bulsara M, Wenz F, Saunders C, Alvarado M, Flyger HL, Massarut S, Eiermann W, Keshtgar M, Dewar J, Kraus-Tiefenbacher U, Sütterlin M, Esserman L, Holtveg HM, Roncadin M, Pigorsch S, Metaxas M, Falzon M, Matthews A, Corica T, Williams NR, Baum M. Targeted intraoperative radiotherapy versus whole breast radiotherapy for breast cancer (TARGIT-A trial): an international, prospective, randomised, non-inferiority phase 3 trial. *Lancet* 2010; **376**: 91-102 [PMID: [20570343](https://pubmed.ncbi.nlm.nih.gov/20570343/) DOI: [10.1016/S0140-6736\(10\)60837-9](https://doi.org/10.1016/S0140-6736(10)60837-9)]
- 100 **Nishioka A**, Ogawa Y, Miyatake K, Tadokoro M, Nogami M, Hamada N, Kubota K, Kariya S, Kohsaki T, Saibara T, Okabayashi T, Hanazaki K. Safety and efficacy of image-guided enzyme-targeting radiosensitization and intraoperative radiotherapy for locally advanced unresectable pancreatic cancer. *Oncol Lett* 2014; **8**: 404-408 [PMID: [24959285](https://pubmed.ncbi.nlm.nih.gov/24959285/) DOI: [10.3892/ol.2014.2101](https://doi.org/10.3892/ol.2014.2101)]
- 101 **Zhou GX**, Zeng TW, Wang LY, Ma L. Analysis of the long-term effect of intraoperative radiotherapy (IORT) for non-small cell lung carcinoma (NSCLC). *Zhongguo Zhongliu Linchuang* 2007; **4**: 65-70 [DOI: [10.1007/s11805-007-0065-0](https://doi.org/10.1007/s11805-007-0065-0)]
- 102 **Nemoto K**, Ogawa Y, Matsushita H, Takeda K, Takai Y, Yamada S, Kumabe T. Intraoperative radiation therapy (IORT) for previously untreated malignant gliomas. *BMC Cancer* 2002; **2**: 1 [PMID: [11818027](https://pubmed.ncbi.nlm.nih.gov/11818027/) DOI: [10.1186/1471-2407-2-1](https://doi.org/10.1186/1471-2407-2-1)]
- 103 **Fangusaro J**, Onar-Thomas A, Poussaint TY, Wu S, Ligon AH, Lindeman N, Banerjee A, Packer RJ, Kilburn LB, Goldman S. LGG-02. A phase II Prospective Trial of Selumetinib In Children With Recurrent/Progressive pediatric low-grade glioma (pLGG) with a focus upon optic pathway/hypothalamic tumors and visual acuity outcomes: a pediatric brain tumor consortium (PBTC) study, PBTC-029B. *Neuro-Oncology* 2019; **21**: ii98-ii99 [DOI: [10.1093/neuonc/noz036.145](https://doi.org/10.1093/neuonc/noz036.145)]
- 104 **De Waele J**, Verhezen T, van der Heijden S, Berneman ZN, Peeters M, Lardon F, Wouters A, Smits ELJM. A systematic review on poly(I:C) and poly-ICLC in glioblastoma: adjuvants coordinating the unlocking of immunotherapy. *J Exp Clin Cancer Res* 2021; **40**: 213 [PMID: [34172082](https://pubmed.ncbi.nlm.nih.gov/34172082/) DOI: [10.1186/s13046-021-02017-2](https://doi.org/10.1186/s13046-021-02017-2)]
- 105 **Trippett T**, Toledano H, Campbell Hewson Q, Verschuur A, Langevin AM, Aerts I, Howell L, Gallego S, Rossig C, Smith A, Patel D, Pereira LR, Cheeti S, Musib L, Hutchinson KE, Devlin C, Bernardi R, Geoerger B. Cobimetinib in Pediatric and Young Adult Patients with Relapsed or Refractory Solid Tumors (iMATRIX-cobi): A Multicenter, Phase I/II Study. *Target Oncol* 2022; **17**: 283-293 [PMID: [35715627](https://pubmed.ncbi.nlm.nih.gov/35715627/) DOI: [10.1007/s11523-022-00888-9](https://doi.org/10.1007/s11523-022-00888-9)]
- 106 **Cantwell-Dorris ER**, O'Leary JJ, Sheils OM. BRAFV600E: implications for carcinogenesis and molecular therapy. *Mol Cancer Ther* 2011; **10**: 385-394 [PMID: [21388974](https://pubmed.ncbi.nlm.nih.gov/21388974/) DOI: [10.1158/1535-7163.MCT-10-0799](https://doi.org/10.1158/1535-7163.MCT-10-0799)]
- 107 **Nicolaidis T**, Nazemi KJ, Crawford J, Kilburn L, Minturn J, Gajjar A, Gauvain K, Leary S, Dhall G, Aboian M, Robinson G, Long-Boyle J, Wang H, Molinaro AM, Mueller S, Prados M. Phase I study of vemurafenib in children with recurrent or progressive BRAF(V600E) mutant brain tumors: Pacific Pediatric Neuro-Oncology Consortium study (PNOC-002). *Oncotarget* 2020; **11**: 1942-1952 [PMID: [32523649](https://pubmed.ncbi.nlm.nih.gov/32523649/) DOI: [10.18632/oncotarget.27600](https://doi.org/10.18632/oncotarget.27600)]
- 108 **Hargrave DR**, Bouffet E, Tabori U, Broniscer A, Cohen KJ, Hansford JR, Geoerger B, Hingorani P, Dunkel IJ, Russo MW, Tseng L, Dasgupta K, Gasal E, Whitlock JA, Kieran MW. Efficacy and Safety of Dabrafenib in Pediatric Patients with BRAF V600 Mutation-Positive Relapsed or Refractory Low-Grade Glioma: Results from a Phase I/IIa Study. *Clin Cancer Res* 2019; **25**: 7303-7311 [PMID: [31811016](https://pubmed.ncbi.nlm.nih.gov/31811016/) DOI: [10.1158/1078-0432.CCR-19-2177](https://doi.org/10.1158/1078-0432.CCR-19-2177)]
- 109 **Kaley T**, Touat M, Subbiah V, Hollebecque A, Rodon J, Lockhart AC, Keedy V, Bielle F, Hofheinz RD, Joly F, Blay JY, Chau I, Puzanov I, Raju NS, Wolf J, DeAngelis LM, Makrutzki M, Riehl T, Pitcher B, Baselga J, Hyman DM. BRAF Inhibition in BRAF(V600)-Mutant Gliomas: Results From the VE-BASKET Study. *J Clin Oncol* 2018; **36**: 3477-3484 [PMID: [30351999](https://pubmed.ncbi.nlm.nih.gov/30351999/) DOI: [10.1200/JCO.2018.78.9990](https://doi.org/10.1200/JCO.2018.78.9990)]

- 110 **Lassman AB**, Sepúlveda-Sánchez JM, Cloughesy TF, Gil-Gil MJ, Puduvalli VK, Raizer JJ, De Vos FYF, Wen PY, Butowski NA, Clement PMJ, Groves MD, Belda-Iniesta C, Giglio P, Soifer HS, Rowsey S, Xu C, Avogadri F, Wei G, Moran S, Roth P. Infigratinib in Patients with Recurrent Gliomas and FGFR Alterations: A Multicenter Phase II Study. *Clin Cancer Res* 2022; **28**: 2270-2277 [PMID: [35344029](#) DOI: [10.1158/1078-0432.CCR-21-2664](#)]
- 111 **Bahleda R**, Italiano A, Hierro C, Mita A, Cervantes A, Chan N, Awad M, Calvo E, Moreno V, Govindan R, Spira A, Gonzalez M, Zhong B, Santiago-Walker A, Poggessi I, Parekh T, Xie H, Infante J, Tabernero J. Multicenter Phase I Study of Erdafitinib (JNJ-42756493), Oral Pan-Fibroblast Growth Factor Receptor Inhibitor, in Patients with Advanced or Refractory Solid Tumors. *Clin Cancer Res* 2019; **25**: 4888-4897 [PMID: [31088831](#) DOI: [10.1158/1078-0432.CCR-18-3334](#)]
- 112 **Wood PJ**, Strong R, McArthur GA, Michael M, Algar E, Muscat A, Rigby L, Ferguson M, Ashley DM. A phase I study of panobinostat in pediatric patients with refractory solid tumors, including CNS tumors. *Cancer Chemother Pharmacol* 2018; **82**: 493-503 [PMID: [29987369](#) DOI: [10.1007/s00280-018-3634-4](#)]
- 113 **Arrillaga-Romany I**, Kurz SC, Tarapore R, Sumrall A, Butowski NA, Harrison RA, De Groot JF, Chi AS, Shonka NA, Umemura Y, Odia Y. Single-agent ONC201 in recurrent H3 K27M-mutant diffuse midline glioma. *J Clin Oncol* 2020; **38** [DOI: [10.1200/JCO.2020.38.15_suppl.3615](#)]
- 114 **Wen PY**, De Groot JF, Battiste J, Goldlust SA, Garner JS, Friend J, Simpson JA, Damek D, Olivero A, Cloughesy TF. Paxalisib in patients with newly diagnosed glioblastoma with unmethylated MGMT promoter status: Final phase 2 study results. *J Clin Oncol* 2022 [DOI: [10.1200/jco.2022.40.16_suppl.2047?af=r](#)]
- 115 **Wen PY**, Cloughesy TF, Olivero AG, Morrissey KM, Wilson TR, Lu X, Mueller LU, Coimbra AF, Ellingson BM, Gerstner E, Lee EQ, Rodon J. First-in-Human Phase I Study to Evaluate the Brain-Penetrant PI3K/mTOR Inhibitor GDC-0084 in Patients with Progressive or Recurrent High-Grade Glioma. *Clin Cancer Res* 2020; **26**: 1820-1828 [PMID: [31937616](#) DOI: [10.1158/1078-0432.CCR-19-2808](#)]
- 116 **Franz DN**, Agricola K, Mays M, Tudor C, Care MM, Holland-Bouley K, Berkowitz N, Miao S, Peyrard S, Krueger DA. Everolimus for subependymal giant cell astrocytoma: 5-year final analysis. *Ann Neurol* 2015; **78**: 929-938 [PMID: [26381530](#) DOI: [10.1002/ana.24523](#)]
- 117 **National Library of Medicine (U.S.)**. (2024, February) A Study to Test the Safety and Efficacy of the Drug Larotrectinib for the Treatment of Tumors With NTRK-fusion in Children (SCOUT) [Identifier: NCT02637687] Available from: <https://clinicaltrials.gov/study/NCT02637687>
- 118 **National Library of Medicine (U.S.)**. (2024, February) A Study to Test the Effect of the Drug Larotrectinib in Adults and Children With NTRK-fusion Positive Solid Tumors (NAVIGATE) [Identifier: NCT02576431] Available from: <https://clinicaltrials.gov/study/NCT02576431>
- 119 **Desai AV**, Robinson GW, Gauvain K, Basu EM, Macy ME, Maese L, Whipple NS, Sabnis AJ, Foster JH, Shusterman S, Yoon J, Weiss BD, Abdelbaki MS, Armstrong AE, Cash T, Pratilas CA, Corradini N, Marshall LV, Farid-Kapadia M, Chohan S, Devlin C, Meneses-Lorente G, Cardenas A, Hutchinson KE, Berghold G, Caron H, Chow Maneval E, Gajjar A, Fox E. Entrectinib in children and young adults with solid or primary CNS tumors harboring NTRK, ROS1, or ALK aberrations (STARTRK-NG). *Neuro Oncol* 2022; **24**: 1776-1789 [PMID: [35395680](#) DOI: [10.1093/neuonc/noac087](#)]
- 120 **National Library of Medicine (U.S.)**. (2023, November) Study of Zotiraciclib for Recurrent High-Grade Gliomas with Isocitrate Dehydrogenase 1 or 2 (IDH1 or IDH2) Mutations [Identifier: NCT05588141] Available from: <https://clinicaltrials.gov/study/NCT05588141>
- 121 **Mellinghoff IK**, van den Bent MJ, Blumenthal DT, Touat M, Peters KB, Clarke J, Mendez J, Yust-Katz S, Welsh L, Mason WP, Ducray F, Umemura Y, Nabors B, Holdhoff M, Hottinger AF, Arakawa Y, Sepulveda JM, Wick W, Soffietti R, Perry JR, Giglio P, de la Fuente M, Maher EA, Schoenfeld S, Zhao D, Pandya SS, Steelman L, Hassan I, Wen PY, Cloughesy TF; INDIGO Trial Investigators. Vorasidenib in IDH1- or IDH2-Mutant Low-Grade Glioma. *N Engl J Med* 2023; **389**: 589-601 [PMID: [37272516](#) DOI: [10.1056/NEJMoa2304194](#)]
- 122 **Mellinghoff IK**, Cloughesy TF, Wen PY, Taylor JW, Maher EA, Arrillaga I, Peters KB, Choi C, Ellingson BM, Lin AP, Thakur SB. A phase I, open label, perioperative study of AG-120 and AG-881 in recurrent IDH1 mutant, low-grade glioma: Results from cohort 1. *J Clin Oncol* 2019; **37** [DOI: [10.1200/JCO.2019.37.15_suppl.2003](#)]
- 123 **Weller M**, Butowski N, Tran DD, Recht LD, Lim M, Hirte H, Ashby L, Mechtler L, Goldlust SA, Iwamoto F, Drappatz J, O'Rourke DM, Wong M, Hamilton MG, Finocchiaro G, Perry J, Wick W, Green J, He Y, Turner CD, Yellin MJ, Keler T, Davis TA, Stupp R, Sampson JH; ACT IV trial investigators. Rindopepimut with temozolomide for patients with newly diagnosed, EGFRvIII-expressing glioblastoma (ACT IV): a randomised, double-blind, international phase 3 trial. *Lancet Oncol* 2017; **18**: 1373-1385 [PMID: [28844499](#) DOI: [10.1016/S1470-2045\(17\)30517-X](#)]
- 124 **Lassman AB**, Pugh SL, Wang TJC, Aldape K, Gan HK, Preusser M, Vogelbaum MA, Sulman EP, Won M, Zhang P, Moazami G, Macsai MS, Gilbert MR, Bain EE, Blot V, Ansell PJ, Samanta S, Kundu MG, Armstrong TS, Wefel JS, Seidel C, de Vos FY, Hsu S, Cardona AF, Lombardi G, Bentsion D, Peterson RA, Gedye C, Bourg V, Wick A, Curran WJ, Mehta MP. Depatuxizumab mafodotin in EGFR-amplified newly diagnosed glioblastoma: A phase III randomized clinical trial. *Neuro Oncol* 2023; **25**: 339-350 [PMID: [35849035](#) DOI: [10.1093/neuonc/noac173](#)]
- 125 **Wisoff JH**, Sanford RA, Heier LA, Sposto R, Burger PC, Yates AJ, Holmes EJ, Kun LE. Primary neurosurgery for pediatric low-grade gliomas: a prospective multi-institutional study from the Children's Oncology Group. *Neurosurgery* 2011; **68**: 1548-54; discussion 1554 [PMID: [21368693](#) DOI: [10.1227/NEU.0b013e318214a66e](#)]
- 126 **Lima GL**, Duffau H. Is there a risk of seizures in "preventive" awake surgery for incidental diffuse low-grade gliomas? *J Neurosurg* 2015; **122**: 1397-1405 [PMID: [25723301](#) DOI: [10.3171/2014.9.JNS141396](#)]
- 127 **Duffau H**. Long-term outcomes after supratotal resection of diffuse low-grade gliomas: a consecutive series with 11-year follow-up. *Acta Neurochir (Wien)* 2016; **158**: 51-58 [PMID: [26530708](#) DOI: [10.1007/s00701-015-2621-3](#)]
- 128 **Ng S**, Herbet G, Moritz-Gasser S, Duffau H. Return to Work Following Surgery for Incidental Diffuse Low-Grade Glioma: A Prospective Series With 74 Patients. *Neurosurgery* 2020; **87**: 720-729 [PMID: [31813972](#) DOI: [10.1093/neuros/nyz513](#)]
- 129 **Ng S**, Herbet G, Lemaitre AL, Cochereau J, Moritz-Gasser S, Duffau H. Neuropsychological assessments before and after awake surgery for incidental low-grade gliomas. *J Neurosurg* 2020; **135**: 871-880 [PMID: [33276332](#) DOI: [10.3171/2020.7.JNS201507](#)]
- 130 **Goel A**, Shah A, Vutha R, Dandpat S, Hawaldar A. Is "En Masse" Tumor Resection a Safe Surgical Strategy for Low-Grade Gliomas? Feasibility Report on 74 Patients Treated Over Four Years. *Neurol India* 2021; **69**: 406-413 [PMID: [33904464](#) DOI: [10.4103/0028-3886.314527](#)]
- 131 **Ius T**, Ng S, Young JS, Tomasino B, Polano M, Ben-Israel D, Kelly JJP, Skrap M, Duffau H, Berger MS. The benefit of early surgery on overall survival in incidental low-grade glioma patients: A multicenter study. *Neuro Oncol* 2022; **24**: 624-638 [PMID: [34498069](#) DOI: [10.1093/neuonc/noab210](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: office@baishideng.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

