

## Intersecting pathways in inflammation and cancer: Hepatocellular carcinoma as a paradigm

Philip Y Wai, Paul C Kuo

Philip Y Wai, Paul C Kuo, Department of Surgery, Loyola University Chicago, Stritch School of Medicine, Maywood, IL 60153, United States

**Author contributions:** Both Wai PY and Kuo PC contributed to the writing of this manuscript.

**Correspondence to:** Paul C Kuo, MD, MS, MBA, Department of Surgery, Loyola University Medical Center, 2160 S. First Avenue, EMS Building, Maywood, IL 60153, United States. [pkuo@lumc.edu](mailto:pkuo@lumc.edu)

Telephone: +708-3272705 Fax: +708-3272582

Received: November 2, 2011 Revised: January 29, 2012

Accepted: February 6, 2012

Published online: February 10, 2012

### Abstract

Viral infection and chemical carcinogens trigger somatic changes resulting in activation of oncogenes during tumor initiation in the development of cancer. However, a critical interaction resides in the synergism between these somatic changes and an inflamed tumor microenvironment where myeloid and hematopoietic cells are subverted to enhance tumor progression. The causative molecular mechanisms leading to the development of hepatocellular cancer remain incompletely understood but appear to result from multiple factors related to direct hepatocyte injury and the ensuing inflammatory changes mediated by the host response to tissue injury, DNA damage, repair of cellular damage, and chronic, repetitive injury. In this review, the molecular and cellular changes that regulate inflammation and tissue repair will be compared to the activated local tumor microenvironment. Cell-cell signaling within this microenvironment that enhances tumor progression and inhibits anti-tumor immunity will be discussed

© 2012 Baishideng. All rights reserved.

**Key words:** Hepatocellular cancer; Inflammation; Tumor microenvironment

**Peer reviewers:** Paul J Higgins, PhD, Professor and Director, Center for Cell Biology and Cancer Research, Albany Medical College, Mail Code 165, Room MS-341, Albany Medical College, 47 New Scotland Avenue, Albany, NY 12208, United States; Yong-Song Guan, Professor, Oncology and Radiology, State Key Laboratory of Biotherapy, West China Hospital of Sichuan University, Chengdu, 610041, Sichuan Province, China; Naoya Sakamoto, MD, PhD, Associate Professor, Department of Gastroenterology and Hepatology, Tokyo Medical and Dental University, 1-5-45 Yushima, Bunkyo-ku, Tokyo 113-8519, Japan; Shahriar Behboudi, Senior Lecturer, Faculty of Biomedical Sciences, The Institute of Hepatology, 69-75 Chenies Mews, London WC1E 6HX, United Kingdom

Wai PY, Kuo PC. Intersecting pathways in inflammation and cancer: Hepatocellular carcinoma as a paradigm. *World J Clin Oncol* 2012; 3(2): 15-23 Available from: URL: <http://www.wjgnet.com/2218-4333/full/v3/i2/15.htm> DOI: <http://dx.doi.org/10.5306/wjco.v3.i2.15>

### INTRODUCTION

Cellular injury resulting in inflammation, tissue repair, and fibrosis has been hypothesized to play a functional role in cancer development since Virchow's original observations<sup>[1]</sup>. Key somatic changes caused by viral or chemical carcinogens establishing a "subthreshold neoplastic state" (initiation) is often supplemented by a subsequent stimulus from the local niche or microenvironment, such as chronic inflammation (promotion). These synergistic pathways complement each other to augment tumor invasiveness<sup>[2,3]</sup>. In certain types of cancers, inflammatory conditions are present before a malignant change occurs and investigators have sought to define a causal relationship between inflammation, innate immunity and cancer development. Certainly, epidemiological studies have shown that chronic inflammation predisposes individuals to various types of cancer including bladder, cervical, gastric, intestinal, esophageal, ovarian, prostate, liver, and thyroid tumors. Perhaps the most

convincing evidence is the fact that adoptive transfer of inflammatory cells from the local tumor environment or overexpression of inflammatory cytokines promotes tumor development<sup>[4]</sup>. Despite these observations, the fundamental mechanisms by which inflammation leads to cancer or supports the progression of cancer remain unclear.

In this editorial, we will discuss the critical components of the inflammatory milieu or tumor microenvironment (TME) that promote a neoplastic and/or metastatic phenotype. The canonical physiology of inflammation will be examined within the context of cancer development highlighting similarities and differences between the two analogous processes. Finally, hepatitis and chronic liver cirrhosis will be discussed as a model of fibrogenesis and hepatic carcinogenesis with potential therapeutic targets.


## INFLAMMATION, TISSUE REPAIR, TUMOR PROGRESSION AND METASTASIS

Mantovani and colleagues defined the complementary oncogenic and inflammatory processes in cancer development as the “intrinsic” and “extrinsic” pathways<sup>[5]</sup>. The intrinsic pathway activates oncogenes and inhibits tumor suppressors by mutation, chromosomal rearrangement or amplification and drives transformation within targeted cells. Tumor cells generated in this fashion subsequently produce cytokines that recruit and populate the inflammatory TME. The extrinsic pathway derives from inflammatory or infectious conditions that amplify the cancer risk (e.g., inflammatory bowel disease, hepatitis, *Helicobacter pylori*). These two mutually dependent pathways eventually converge, appropriating-necessary components and signals from the other while also supplying reciprocally useful building blocks to fuel transformation and metastasis in a cooperated fashion. Oncogenes that can produce signals for inflammatory cell recruitment include the RET proto-oncogene in papillary thyroid cancer, K-RAS in pancreatic cancer, and BRAF-MAPK in melanoma<sup>[5]</sup>. It is no coincidence that inflammation and wound healing physiology parallels the tissue remodeling processes that occur in cancer progression. Dvorak<sup>[6]</sup> recognized that the composition of the tumor stroma strongly resembles the granulation tissue of healing skin wounds. These important, essential inflammatory cascades promote cell proliferation, migration, invasion through the extracellular matrix, angiogenesis, and ultimately provide the necessary components for host tissue repair and survival. In many types of cancer, these attributes can be subverted by nascent tumor cells as tools for cancer progression and metastasis.

The steps of the inflammatory cascade associated with tissue repair are well characterized. Tissue injury created by toxins, infection, or a chronic inflammatory stimulus results in a host response focused on recruiting

cells that initiate healing (Figure 1). Critical members of this microenvironment include neutrophils, monocytes, macrophages, mast cells, dendritic cells, fibroblasts and endothelial cells. The wound healing process often involves partially overlapping phases: blood clotting, inflammation, new tissue formation, and tissue remodeling<sup>[7]</sup> with key cell types recruited to the niche during specific phases. Important pro-inflammatory signals produced during this cascade include IL-1 $\beta$ , IL-6, IL-23, TNF- $\alpha$ , and TGF- $\beta$ 1. Activation of the selectin family of adhesion molecules (L-, P-, and E-selectin) facilitates leukocyte “rolling” along the injured vascular endothelium, activating integrin binding and immobilization ( $\alpha$ 4 $\beta$ 1 and  $\alpha$ 4 $\beta$ 7 binding to VCAM-1 and MadCAM-1), and ultimately transmigration through the endothelium into the site of injury<sup>[7]</sup>. Release of cytokines, chemokines, and prostaglandins to recruit additional inflammatory cells, the production of reactive oxygen species (ROS) to destroy infectious vectors, the generation of pro-angiogenic factors, and modulation of apoptosis represent other essential, activated functions.

Physiological inflammation is often self-limiting through downstream release of anti-inflammatory regulators (IL-10, IL-11, IL-13) which temper the pro-inflammatory cascade. However, cancer-associated inflammation can often persist, or be driven without regulation, to elicit pathologically persistent signals for cellular proliferation, migration, basement membrane invasion and angiogenesis. In this context, tumors have been comparatively described as “wounds that do not heal”<sup>[6]</sup>. The key cells residing in the TME are the same constituents that facilitate wound healing and inflammation as described above. However, the tumor-associated cells recruited often display altered functions that lend themselves to cancer development. This alteration in function derives from the upregulated expression of pro-tumor cytokines. For example, dendritic cells in neoplastic infiltrates are regulated by tumor-derived granulocyte-macrophage colony-stimulating factor (GM-CSF) and interleukin (IL)-4 and are frequently immature, less effective at capturing antigens, and defective in T-cell stimulatory capacity<sup>[2]</sup>. IL-10 released into the TME is a potent inhibitor of dendritic cell activation and differentiation allowing evasion of host adaptive immunity<sup>[5]</sup>. Similar processes occur in tumor-associated macrophages (TAMs) which produce a number of potent angiogenic and lymphangiogenic growth factors, cytokines and proteases that mediate neoplastic progression. For example, in human cervical carcinogenesis, TAMs express VEGF-C, VEGF-D, and VEGF receptor-3, to facilitate angiogenesis<sup>[8]</sup>. In a murine mammary cancer metastasis model, CSF-1 regulates tumor growth by supporting and cultivating the TME. In CSF-1<sup>-/-</sup> mice, advanced mammary tumors and pulmonary metastases fail to develop due to decreased TAMs recruitment into the neoplastic tissue<sup>[9]</sup>. CSF-1 has been shown to promote progression of mammary tumors to malignancy as replacement of transgenic CSF-1 into mammary epitheli-



**Figure 1** The intrinsic and extrinsic pathways combine to create a local microenvironment around the injured and transformed hepatocyte to augment tumor promoting mechanisms. ROS: Reactive oxygen species; HIF1 $\alpha$ : Hypoxia inducible factor 1 alpha; NK cell: Natural killer cell; MDSC: Myeloid derived suppressor cell; IL: Interleukin; TGF- $\beta$ : Transforming growth factor Beta; HGF: Hepatocyte growth factor; NF- $\kappa$ B: Nuclear factor kappa-light-chain-enhancer of activated B cells; CXCL: Chemokine ligand; STAT3: Signal transducer and activator of transcription 3.

um restores macrophage recruitment, primary tumor development, and metastatic potential<sup>[9]</sup>. A powerful stimulus for tumor progression within the TME includes the ROS derived from infiltrating leukocytes. In the presence of chronic inflammation and repetitive injury, leukocytes and other phagocytic cells induce DNA damage in proliferating cells through the generation of reactive oxygen and nitrogen species such as peroxynitrite. Irreversible DNA mutations generated by these reactive species can provide the critical trigger for neoplastic transformation. In addition to these mechanisms, the inhibition of tumor-suppressor pathways represent yet another strategy for promoting tumor growth. Macrophage migration inhibitory factor (MIF) released from TAMs and T lymphocytes is a potent cytokine that suppresses *p53* transcriptional activity. MIF released into the TME creates a niche with a deficient response to DNA damage<sup>[10]</sup>. TAMs will be diverted into the M2 phenotype in human tumors so that macrophage functions will be focused on promoting tumor growth, remodeling tissues, promoting angiogenesis, and suppressing adaptive immunity<sup>[11,12]</sup>. Another class of cells that are recruited to the TME include the myeloid-derived suppressor cells (MDSCs). These cells are abundant in tumors and strongly inhibit anti-tumor immunity<sup>[7]</sup>. MDSCs represent an immature population of myeloid cells that inhibit both innate and adaptive immunity and are present in cancer patients and in experimental animals with sizable tumor burden<sup>[13]</sup>. Although no definitive molecular characterization ex-

ists, many investigators have found human MDSCs to express CD33, CD11b and CD15 cell surface markers<sup>[13]</sup>. MDSC inhibition of anti-tumor immunity is mediated by suppression of CD4<sup>+</sup> T-cells<sup>[14]</sup>, inducing T regulatory cells<sup>[15]</sup>, by down-regulating macrophage production of the type 1 cytokine, IL-12<sup>[16]</sup>, and potentially suppressing natural killer cell cytotoxicity<sup>[17]</sup>. In hepatocellular carcinoma models, trafficking and accumulation of MDSCs appears to be gp130 dependent and downregulation of NK cell cytokine production to be NKp30 dependent<sup>[18]</sup>. Recent studies have also focused on the myofibroblast as another cell type that is commonly found in wounds and in the TME and has been implicated in tumor progression. The presence of large numbers of fibroblasts and myofibroblasts is a hallmark of cancer with many tumors producing a desmoplastic response<sup>[7]</sup>. Although tumor fibroblasts can be derived from the stroma surrounding tumors, there is evidence to suggest that cells recruited from the bone-marrow also 'home in' on the TME<sup>[19]</sup>. Auto- and paracrine PDGF and TGF- $\beta$  dependent signaling centered on the myofibroblast is considered fundamental to tumor progression, the development of epithelial-mesenchymal transition (EMT), and generation of cancer stem cells (CSCs). CSCs exhibit a CD44<sup>high</sup>/CD24<sup>low</sup> antigenic phenotype, demonstrate upregulation of the mesenchymal markers and the transcription factors, N-cadherin, fibronectin, vimentin, FOXC2, SIP1, Hedgehog (Hh), Snail, and Twist, and possess self-renewal capability enabling CSCs to exit tissue reservoirs,

enter and survive in the circulation, and exit into secondary tissue sites (“stemness”)<sup>[20]</sup>. In the liver, cancer-associated fibroblasts are important contributors to the TME<sup>[21]</sup> and their precise origin continue to be unclear with a variety of hepatic cells able to generate stem-cell characteristics including hepatocytes, oval cells/hepatic progenitor cells, and bone marrow-derived cells<sup>[22]</sup>.

## HEPATIC FIBROGENESIS AND TUMOR PROGRESSION

Constituents of the cancer-associated inflammatory stroma vary between different tumor types suggesting that TMEs can be disease-specific. Distinguishing mechanistic pathways that are conserved versus pathways that are tissue- and tumor- specific is important. Friedman and colleagues described the difference between “core” and “non-core” pathways leading to end-stage inflammation or fibrosis with “core” pathways maintaining a dominant role through different organ systems, processes, and species<sup>[23]</sup>. Non-core pathways may regulate core pathways, but are not necessarily fundamental to the fibrotic process, and may be specific to one model system without correlation to other disease states. Discovery of universal anti-fibrotic cures may require specific attention to core pathways. However, such integral proteins and signals (e.g., TGF- $\beta$ ) may be required in normal tissue function and may not be appropriate targets for inhibition. In this context, regulatory signaling pathways that are tissue-specific or even disease/pathology-specific may provide realistic targets that can be candidates for therapeutic intervention.

Hepatic fibrosis is a reversible wound-healing response to liver injury and is characterized by inflammation, accumulation of extracellular matrix (ECM), and ultimately scarring, as described above. If the injury is self-limiting, the inflammatory changes are transient and the liver tissue is restored to its normal configuration. However, when the injury or the resultant inflammatory response is persistent, the liver architecture is irreversibly transformed leading to progressive fibrosis and cirrhosis. Agents that injure the liver in such a way include toxins (CCL4, alcohol, or bile from biliary stasis), chronic infections (hepatitis B, hepatitis C), or remodeling processes (adipose tissue in non-alcoholic fatty liver disease). Chemical toxicity, viral infection, and metabolic derangements damage hepatocytes and this injury triggers a cascade designed to contain the injury by removing or repairing damaged cells, defense against further infection or injury, tissue regeneration and repair. Chronic inflammation due to repetitive injury (toxin) or inability to remove the offending agent (viral infection) results in a deranged, decompensated response (Figure 1). The liver is composed of constituent cells capable of mounting a robust inflammatory response. The space of Disse is a subendothelial space that separates the hepatic sinusoids from the hepatocytes. It contains a low density basal membrane-like matrix that is essential for


maintaining the differentiated function of parenchymal cells but can become porous enough to enable metabolic exchange between the bloodstream and hepatocytes<sup>[24]</sup>. Inflammatory cells that can be activated to secrete cytokines and regulate ECM in this space include the hepatic macrophages (Kupffer cells), hepatic stellate cells (HSCs) (that can be activated to become fibroblasts), dendritic cells, natural killer cells, hepatocytes and cholangiocytes. Progression to cirrhosis leads to the formation of nodules of regenerative parenchyma surrounded by sheets of fibrotic septae. During chronic liver injury, extracellular matrix (ECM) deposition is upregulated. Interactions between the ECM and its surrounding cells then become mutually dependent with decorin and biglycan binding of TGF- $\beta$ ; fibronectin and laminin binding of TNF- $\alpha$ ; and collagen binding to PDGF, HGF, and IL-2<sup>[24]</sup>. HSCs become activated from a quiescent vitamin A-rich cell to a contractile, highly fibrogenic, myofibroblast cell-type. Activated HSCs leads to upregulation of integrin receptors  $\alpha 2\beta 1$ ,  $\alpha 6\beta 4$ ,  $\alpha v\beta 8$ ,  $\alpha v\beta 6$ , and  $\alpha 5\beta 1$ , regulation of TGF- $\beta 1$ , PDGF, and Hedgehog (Hh) signaling pathways, expression of ADAMTS-13, ADAMTS-1, and secretion of collagenase-1 and -3, stromelysin-1 and -2, gelatinases, metalloelastases (MT-MMP1) and tissue inhibitor of metalloproteinase-1 (TIMP-1)<sup>[24]</sup> (Figure 2).

Accumulating evidence suggests that hepatocellular epithelial mesenchymal transition (EMT) plays a pivotal role in the dissemination of malignant hepatocytes during HCC progression<sup>[25]</sup>. Complete EMT often occurs in cancer cells as they lose cell-cell contacts, acquire a fibroblast-like morphology, and express mesenchymal marker proteins such as alpha smooth muscle actin ( $\alpha$ SMA) and fibroblast-specific protein 1 (FSP1)<sup>[7]</sup>, while shedding the epithelial cell markers cytokeratin and E-cadherin. This process often allows a firmly attached epithelial cell with appropriate apical-basal polarity to migrate into the interstitium and acquire characteristics of mesenchymal cells (highly motile, invasive, resistant to apoptosis, produce ECM, transmigrate through basement membrane, and intravasate into endothelium). Hepatic cancer cells that undergo EMT appear to acquire the properties of stem cells<sup>[20]</sup>. After liver injury occurs, the major trigger of EMT is the release of chemokines, MMPs, and other growth factors with PDGF, TGF- $\beta$  and Hh occupying principal roles in this process (Figure 2).

### TGF- $\beta$

TGF- $\beta$  is secreted by a variety of cell types and exists as the following isoforms: TGF- $\beta 1$ , TGF- $\beta 2$ , and TGF- $\beta 3$ . The predominant isoform in inflammation and fibrosis is TGF- $\beta 1$  and signaling is mediated through the type II receptor. PDGF receptors transduce their signals through the PI3K/Akt pathways while TGF- $\beta$  mediates signaling through the Smad proteins. Inhibition of PDGF signaling decreases migration *in vitro* and efficient tumor suppression *in vivo* indicating that TGF- $\beta$  mediated EMT of neoplastic hepatocytes is PDGF-dependent<sup>[25,26]</sup>. Activation of the type II receptor results in di-





**Figure 2** The complex cellular network in the tumor microenvironment mediated by chemokines, cytokines, and cellular transcription factors. NK cell: Natural killer cell; MDSC: Myeloid derived suppressor cell; IL: Interleukin; TGF-β: Transforming growth factor Beta; αSMA: Alpha smooth actin; FSP-1: Fibroblast specific protein; PDGF: Platelet derived growth factor; Hh: Hedgehog; NF-κB: Nuclear factor kappa-light-chain-enhancer of activated B cells; CXCL: Chemokine ligand; STAT3: Signal transducer and activator of transcription 3; Treg: T regulatory cells; MT-MMP: Membrane type matrix metalloproteinase; HNF-4: Hepatocyte nuclear factor-4; EMT: Epithelial-mesenchymal-transition; MIF: migration inhibitory factor.

merization with the type I receptor and promotes binding to Smad2 and Smad3. The phosphorylated Smad2/Smad3 complex associates with Smad4 and activates transcription. TGF-β is a tumor suppressor protein that is frequently involved in tumor progression of human cancer and also in tumor-promoting inflammation. Many liver cell types, including HSCs, hepatocytes, and liver sinusoidal endothelial cells are regulated by TGF-β<sup>[27]</sup>. Interestingly, as in other cancer cell types, TGF-β has been shown to have a dual role in HCC with TGF-β displaying antitumor effects initially. For example, loss-of-function of TGF-β type II receptor results in enhanced susceptibility to HCC, suggesting that TGF-β retains some tumor-suppressor functions<sup>[28]</sup>. Alternatively, transgenic mice with upregulated Smad7 expression restricted to hepatocytes demonstrate significantly diminished liver damage and fibrosis, demonstrating that TGF-β signaling in hepatocytes is required for fibrogenesis progression<sup>[29]</sup>. The significance of the dual nature of these effects is unclear but are reflected across other cancer cell types and suggest that the effects of TGF-β may be time-, and context-dependent. For example, inactivation of type II TGF-β receptor in an animal model of breast carcinoma increases CXCL5- and CXCL12-mediated recruitment of myeloid-derived suppressor cells (MDSCs) which are potent suppressors of the adaptive immune response to tumors<sup>[5]</sup>. Smad7 activation or RNA interference

against Smad4 decreases TGF-β signaling and attenuates the expression of pro-fibrotic genes<sup>[29,30]</sup>. However, hepatocytes isolated from livers exposed to high TGF-β *in vivo* demonstrate elongated, fibroblastoid hepatocytes expressing vimentin and collagen I in comparison to healthy mouse livers<sup>[31]</sup>. Other evidence for pro-tumor TGF-β-mediated downstream effects include studies by Morris *et al.*<sup>[32]</sup> that show loss of TGF-β receptor type II in the context of loss of *p53* decreased the incidence of HCC in a murine model of liver cancer. Cumulative evidence from the Fabregat group demonstrates that TGF-β signaling regulates seemingly contradictory processes in normal liver cells and in HCC. TGF-β-mediated growth inhibition and apoptosis (tumor-suppressor characteristics) occur in non-transformed human fetal hepatocytes while transdifferentiation into a mesenchymal-stem cell-like phenotype with increased expression of Snail, decreased E-cadherin expression, increased Vimentin and N-cadherin expression (pro-tumor) is also TGF-β-mediated<sup>[33]</sup>. Indeed, parallel experiments using siRNA-mediated down regulation of Snail showed that hepatocytes became sensitized to TGF-β mediated apoptosis and that Snail and induction of the EMT phenotype impairs TGF-β apoptosis in HCC cells<sup>[34]</sup>.

### Nuclear factor-kappa B

Nuclear factor-kappa B (NF-κB) is an important regula-

tor of innate immunity, inflammation and also of tumor progression. NF- $\kappa$ B is activated downstream from Toll-like receptor (TLR)-MyD88 signaling and by signaling through the TNF- $\alpha$  and IL-1 $\beta$  pathways. NF- $\kappa$ B can also be activated as a result of genetic alterations (amplification, mutations, or deletions) in tumor cells. NF- $\kappa$ B activation is controlled by multiple factors and its transcriptional activity is linked to various inflammatory or pro-tumor states. Deficiency of TIR8, an inhibitor of Toll/IL-1 (TIR)-mediated NF- $\kappa$ B signaling, results in increased NF- $\kappa$ B activation with increased susceptibility to intestinal inflammation and carcinogenesis<sup>[35,36]</sup>. The causative role of NF- $\kappa$ B in inflammation and cancer is further supported by studies in TAMs where the p50 homodimers confer a pro-tumor phenotype through constitutively activated NF- $\kappa$ B<sup>[37]</sup>. Inhibition of IKK- $\alpha$  releases inhibition of maspin, a tumor suppressor, and reduces metastatic spread in malignant prostate epithelial cells<sup>[38]</sup>. Evidence also suggests that NF- $\kappa$ B determines the balance between pro- and anti-tumor effects. When NF- $\kappa$ B signaling is inhibited specifically in TAMs, they switch to a "classically" activated M1 phenotype (IL-12<sup>high</sup>; IL-10<sup>low</sup>) and TAMs become cytotoxic to tumor cells again<sup>[39]</sup>. Clearly, the mechanisms involved in NF- $\kappa$ B activation are complex and contradictory functions occur in liver physiology and hepatic carcinogenesis. In Mdr2-knockout mouse (a murine model of chronic inflammation induced-HCC), NF- $\kappa$ B inhibition with inducible I $\kappa$ B super-repressor resulted in decreased tumor progression<sup>[40]</sup>. In contrast, Maeda and colleagues demonstrated that IKK- $\beta$  knockout in hepatocytes during an acute liver injury model with diethylnitrosamine (DEN) resulted in mice with an absence of chronic inflammation but increased hepatocarcinogenesis<sup>[41]</sup>. The timing and context of NF- $\kappa$ B activation or inhibition determines the associated phenotype.

### Signal transducer and activator of transcription

The signal transducer and activator of transcription (STAT) family of transcription factors also play a critical role in tumor and immune cells. STAT1 and STAT3 play a key role in liver fibrosis, antiviral defense, liver inflammation, and liver regeneration. STAT1 confers a protective effect and functions in down-regulating pro-fibrotic mechanisms in the liver by inhibiting HSC proliferation, suppression of  $\beta$ -PDGF receptor expression, inhibition of TGF- $\beta$ /Smad3 signaling, and stimulation of NK cell cytotoxicity<sup>[42]</sup>. In these loss of function studies, STAT1<sup>-/-</sup> mice demonstrated accelerated CCL4-induced liver fibrosis and HSC proliferation.

STAT3 is one of the main signaling proteins activated by HGF and EGF receptors, and is involved in oncogenesis, inhibition of apoptosis, inhibition of dendritic cells, and increased evasion of the immune system<sup>[43]</sup>. Mice that lack STAT3 in the epidermis suffer from reduced wound re-epithelization, and are resistant to carcinogen-induced skin cancer development. Conversely, mice that overexpress a constitutively active form of STAT3 develop skin

cancers with a shorter latency period<sup>[44]</sup>.

### Chemokines

Tumor cells can regulate their chemokine expression profile to recruit inflammatory cells, but also use these factors to enhance tumor growth and expression<sup>[2]</sup>. The chemokines CXCL1, CXCL2, CXCL3, and CXCL8 have been extensively studied in melanoma and they play a role in the regulation of tumor growth. Inhibition of the CXCR2 receptor attenuates melanoma cell proliferation<sup>[45]</sup>, whereas overexpression of CXCL1, CXCL2, CXCL3 enhances tumor cell colony-forming activity and tumorigenicity in nude mice<sup>[46,47]</sup>. Macrophage pro-inflammatory chemokine 3 $\alpha$  (CCL20), is a CC chemokine where the two cysteine (C) residues lie adjacent to each other, is overexpressed in pancreatic carcinoma cells and infiltrating macrophages adjacent to tumor cells, and enhances migration of TAMs<sup>[48]</sup>. Angiogenesis is associated with chronic inflammatory states including arthritis, infections, tumor growth and metastasis. The Glutamic acid-Leucine-Arginine (ELR) motif upstream of the CXC domain in the chemokine family can stimulate endothelial cell chemotaxis and retains pro-angiogenic function. Metastatic potential also appears in part to be governed by chemokine receptors and chemokine signaling. CXCL12 [stromal-cell derived factor-1 (SDF-1)] is a product of resting cells in multiple organs and also myofibroblasts. Binding to its receptor CXCR4 appears to play a critical role in cancer metastasis. Use of CXCR4 antibodies can limit CXCL12-mediated chemotaxis of mammary cells to distant organs during metastasis<sup>[49]</sup>. Indeed the amount of CXCR4 expression in primary human tumors correlates with the extent to which metastasis to the lymph nodes occurs in colorectal, breast, liver and esophageal cancer<sup>[50-52]</sup>. The critical threshold of pro-inflammatory cytokines reached by the myriad of contributing cells within the local tumor environment may augment the invasive capacity of transforming cells in a para- and auto-crine fashion. For example, autocrine TNF- $\alpha$  upregulates the expression of CXCR4 in ovarian cancer<sup>[53]</sup>.

### IAP

Recently, the inhibitor of apoptosis (IAP) family of proteins have been shown to play a significant role in cancer-related inflammation and metastasis<sup>[54]</sup>. Alterations in IAPs are found in many types of human cancer including HCC and are associated with chemoresistance, disease progression, and poor prognosis<sup>[55]</sup>. IAPs function by regulating caspases (cysteine proteases that are involved in apoptosis) and through ubiquitin (Ub)-dependent activation of NF- $\kappa$ B transcription factors. IAP1 (cIAP1; encoded by *BIRC2*), cIAP2 (encoded by *BIRC3*) and XIAP (encoded by *BIRC4*) have particular roles that have been demonstrated in tumor maintenance and progression and deserve particular attention. Using a model of breast cancer metastasis with MDA-MB-231 cells and also MCF-7 cells stably transfected with survivin, Mehrotra and colleagues demonstrated that the

survivin-XIAP complex activates NF- $\kappa$ B and increases metastasis in a splenic injection model of hepatic metastasis<sup>[56]</sup>. Amplification of chromosome 11q22 containing the *BIRC2* and *BIRC3* exons, occurs at high frequency in hepatocellular carcinoma with cIAP1 and cIAP2 functioning as key mediators of TNF- $\alpha$ -induced activation of NF- $\kappa$ B<sup>[55]</sup>. TNF-related apoptosis-inducing ligand (TRAIL) induces apoptosis in hepatocellular cancer in a cIAP1-dependent fashion<sup>[54]</sup>.

### Other signals

Hepatic EMT defined as loss of cell polarity and decrease in cell-cell adhesion was shown to be associated with reduced hepatocyte nuclear factor (HNF)-4 expression while restoration of HNF-4 $\alpha$ 1 expression in aggressive HCC cells re-programmed the cells into resting hepatocytes<sup>[57]</sup>. Analysis of Wnt/ $\beta$ -catenin signaling demonstrated that nuclear accumulation of  $\beta$ -catenin results in loss of epithelial markers and increases expression of hepatic mesenchymal markers such as M2-pyruvate kinase (M2-PK) and cytokeratin (CK19)<sup>[58]</sup>. *In vivo* loss of E-cadherin, upregulation of Twist (a negative inhibitor of E-cadherin transcription) and reduction of cytosolic  $\beta$ -catenin correlate with more invasive HCC phenotypes, increases metastasis and reduces patient survival<sup>[59-61]</sup>. Recent studies have also shown that HCV core proteins lower Smad3 expression and decrease E-cadherin expression promoting EMT in human HCC cells<sup>[62]</sup>, while hepatitis B virus encoded HBX upregulates STAT5b in HCC cell lines to augment EMT and cell invasion by repressing E-cadherin<sup>[63]</sup>.

### CONCLUSION

Chronic inflammation and fibrosis in the liver can generate a local environment or niche that supports oncogenesis, tumor progression and metastasis, and in this regard the generation of hepatocellular carcinoma parallels the key inflammatory physiology that is observed in wound healing and EMT. The current increased interest in cancer-associated inflammation may identify targets that modulate the more advanced or metastatic phenotypes in cancer and may yield novel therapeutic options.

### REFERENCES

- 1 Balkwill F, Mantovani A. Inflammation and cancer: back to Virchow? *Lancet* 2001; **357**: 539-545
- 2 Coussens LM, Werb Z. Inflammation and cancer. *Nature* 2002; **420**: 860-867
- 3 Rous P, Kidd JG. CONDITIONAL NEOPLASMS AND SUBTHRESHOLD NEOPLASTIC STATES: A STUDY OF THE TAR TUMORS OF RABBITS. *J Exp Med* 1941; **73**: 365-390
- 4 Jinushi M, Chiba S, Yoshiyama H, Masutomi K, Kinoshita I, Dosaka-Akita H, Yagita H, Takaoka A, Tahara H. Tumor-associated macrophages regulate tumorigenicity and anti-cancer drug responses of cancer stem/initiating cells. *Proc Natl Acad Sci U S A* 2011; **108**: 12425-12430
- 5 Mantovani A, Allavena P, Sica A, Balkwill F. Cancer-related inflammation. *Nature* 2008; **454**: 436-444
- 6 Dvorak HF. Tumors: wounds that do not heal. Similarities between tumor stroma generation and wound healing. *N Engl J Med* 1986; **315**: 1650-1659
- 7 Schäfer M, Werner S. Cancer as an overhealing wound: an old hypothesis revisited. *Nat Rev Mol Cell Biol* 2008; **9**: 628-638
- 8 Schoppmann SF, Birner P, Stöckl J, Kalt R, Ullrich R, Caucig C, Kriehuber E, Nagy K, Alitalo K, Kerjaschki D. Tumor-associated macrophages express lymphatic endothelial growth factors and are related to peritumoral lymphangiogenesis. *Am J Pathol* 2002; **161**: 947-956
- 9 Lin EY, Nguyen AV, Russell RG, Pollard JW. Colony-stimulating factor 1 promotes progression of mammary tumors to malignancy. *J Exp Med* 2001; **193**: 727-740
- 10 Hudson JD, Shoaibi MA, Maestro R, Carnero A, Hannon GJ, Beach DH. A proinflammatory cytokine inhibits p53 tumor suppressor activity. *J Exp Med* 1999; **190**: 1375-1382
- 11 De Palma M, Venneri MA, Galli R, Sergi L, Politi LS, Sampaioles M, Naldini L. Tie2 identifies a hematopoietic lineage of proangiogenic monocytes required for tumor vessel formation and a mesenchymal population of pericyte progenitors. *Cancer Cell* 2005; **8**: 211-226
- 12 Mantovani A, Sozzani S, Locati M, Allavena P, Sica A. Macrophage polarization: tumor-associated macrophages as a paradigm for polarized M2 mononuclear phagocytes. *Trends Immunol* 2002; **23**: 549-555
- 13 Ostrand-Rosenberg S, Sinha P. Myeloid-derived suppressor cells: linking inflammation and cancer. *J Immunol* 2009; **182**: 4499-4506
- 14 Sinha P, Clements VK, Ostrand-Rosenberg S. Reduction of myeloid-derived suppressor cells and induction of M1 macrophages facilitate the rejection of established metastatic disease. *J Immunol* 2005; **174**: 636-645
- 15 Huang B, Pan PY, Li Q, Sato AI, Levy DE, Bromberg J, Divino CM, Chen SH. Gr-1+CD115+ immature myeloid suppressor cells mediate the development of tumor-induced T regulatory cells and T-cell anergy in tumor-bearing host. *Cancer Res* 2006; **66**: 1123-1131
- 16 Sinha P, Clements VK, Bunt SK, Albelda SM, Ostrand-Rosenberg S. Cross-talk between myeloid-derived suppressor cells and macrophages subverts tumor immunity toward a type 2 response. *J Immunol* 2007; **179**: 977-983
- 17 Li H, Han Y, Guo Q, Zhang M, Cao X. Cancer-expanded myeloid-derived suppressor cells induce anergy of NK cells through membrane-bound TGF- $\beta$  1. *J Immunol* 2009; **182**: 240-249
- 18 Hoechst B, Voigtlaender T, Ormandy L, Gamrekelashvili J, Zhao F, Wedemeyer H, Lehner F, Manns MP, Greten TF, Korangy F. Myeloid derived suppressor cells inhibit natural killer cells in patients with hepatocellular carcinoma via the NKp30 receptor. *Hepatology* 2009; **50**: 799-807
- 19 Direkze NC, Hodiola-Dilke K, Jeffery R, Hunt T, Poulson R, Oukrif D, Alison MR, Wright NA. Bone marrow contribution to tumor-associated myofibroblasts and fibroblasts. *Cancer Res* 2004; **64**: 8492-8495
- 20 Mani SA, Guo W, Liao MJ, Eaton EN, Ayyanan A, Zhou AY, Brooks M, Reinhard F, Zhang CC, Shipitsin M, Campbell LL, Polyak K, Briskin C, Yang J, Weinberg RA. The epithelial-mesenchymal transition generates cells with properties of stem cells. *Cell* 2008; **133**: 704-715
- 21 Berdiel-Acer M, Bohem ME, López-Doriga A, Vidal A, Salazar R, Martínez-Iniesta M, Santos C, Sanjuan X, Villanueva A, Molleví DG. Hepatic carcinoma-associated fibroblasts promote an adaptive response in colorectal cancer cells that inhibit proliferation and apoptosis: nonresistant cells die by nonapoptotic cell death. *Neoplasia* 2011; **13**: 931-946
- 22 Alison MR, Choong C, Lim S. Application of liver stem cells for cell therapy. *Semin Cell Dev Biol* 2007; **18**: 819-826
- 23 Mehal WZ, Iredale J, Friedman SL. Scraping fibrosis: expressway to the core of fibrosis. *Nat Med* 2011; **17**: 552-553
- 24 Hernandez-Gea V, Friedman SL. Pathogenesis of liver fi-



- brosis. *Annu Rev Pathol* 2011; **6**: 425-456
- 25 **van Zijl F**, Zulehner G, Petz M, Schneller D, Kornauth C, Hau M, Machat G, Grubinger M, Huber H, Mikulits W. Epithelial-mesenchymal transition in hepatocellular carcinoma. *Future Oncol* 2009; **5**: 1169-1179
  - 26 **Gotzmann J**, Fischer AN, Zojer M, Mikula M, Proell V, Huber H, Jechlinger M, Waerner T, Weith A, Beug H, Mikulits W. A crucial function of PDGF in TGF-beta-mediated cancer progression of hepatocytes. *Oncogene* 2006; **25**: 3170-3185
  - 27 **Dooley S**, Ten Dijke P. TGF- $\beta$  in progression of liver disease. *Cell Tissue Res* 2012; **347**: 245-256
  - 28 **Kanzler S**, Meyer E, Lohse AW, Schirmacher P, Henninger J, Galle PR, Blessing M. Hepatocellular expression of a dominant-negative mutant TGF-beta type II receptor accelerates chemically induced hepatocarcinogenesis. *Oncogene* 2001; **20**: 5015-5024
  - 29 **Dooley S**, Hamzavi J, Ciucan L, Godoy P, Ilkavets I, Ehnert S, Ueberham E, Gebhardt R, Kanzler S, Geier A, Breitkopf K, Weng H, Mertens PR. Hepatocyte-specific Smad7 expression attenuates TGF-beta-mediated fibrogenesis and protects against liver damage. *Gastroenterology* 2008; **135**: 642-659
  - 30 **Kaimori A**, Potter J, Kaimori JY, Wang C, Mezey E, Koteish A. Transforming growth factor-beta1 induces an epithelial-to-mesenchymal transition state in mouse hepatocytes in vitro. *J Biol Chem* 2007; **282**: 22089-22101
  - 31 **Nitta T**, Kim JS, Mohucz D, Behrns KE. Murine cirrhosis induces hepatocyte epithelial mesenchymal transition and alterations in survival signaling pathways. *Hepatology* 2008; **48**: 909-919
  - 32 **Morris SM**, Baek JY, Koszarek A, Kannurn S, Knoblaugh SE, Grady WM. Transforming growth factor-beta signaling promotes hepatocarcinogenesis induced by p53 loss. *Hepatology* 2012; **55**: 121-131
  - 33 **Caja L**, Bertran E, Campbell J, Fausto N, Fabregat I. The transforming growth factor-beta (TGF- $\beta$ ) mediates acquisition of a mesenchymal stem cell-like phenotype in human liver cells. *J Cell Physiol* 2011; **226**: 1214-1223
  - 34 **Franco DL**, Mainez J, Vega S, Sancho P, Murillo MM, de Frutos CA, Del Castillo G, López-Blau C, Fabregat I, Nieto MA. Snail1 suppresses TGF-beta-induced apoptosis and is sufficient to trigger EMT in hepatocytes. *J Cell Sci* 2010; **123**: 3467-3477
  - 35 **Garlanda C**, Riva F, Veliz T, Polentarutti N, Pasqualini F, Radaelli E, Sironi M, Nebuloni M, Zorini EO, Scanziani E, Mantovani A. Increased susceptibility to colitis-associated cancer of mice lacking TIR8, an inhibitory member of the interleukin-1 receptor family. *Cancer Res* 2007; **67**: 6017-6021
  - 36 **Xiao H**, Gulen MF, Qin J, Yao J, Bulek K, Kish D, Altuntas CZ, Wald D, Ma C, Zhou H, Tuohy VK, Fairchild RL, de la Motte C, Cua D, Vallance BA, Li X. The Toll-interleukin-1 receptor member SIGIRR regulates colonic epithelial homeostasis, inflammation, and tumorigenesis. *Immunity* 2007; **26**: 461-475
  - 37 **Saccani A**, Schioppa T, Porta C, Biswas SK, Nebuloni M, Vago L, Bottazzi B, Colombo MP, Mantovani A, Sica A. p50 nuclear factor-kappaB overexpression in tumor-associated macrophages inhibits M1 inflammatory responses and anti-tumor resistance. *Cancer Res* 2006; **66**: 11432-11440
  - 38 **Luo JL**, Tan W, Ricono JM, Korchynskyi O, Zhang M, Gonias SL, Cheres DA, Karin M. Nuclear cytokine-activated IKKalpha controls prostate cancer metastasis by repressing Maspin. *Nature* 2007; **446**: 690-694
  - 39 **Hagemann T**, Lawrence T, McNeish I, Charles KA, Kulbe H, Thompson RG, Robinson SC, Balkwill FR. "Re-educating" tumor-associated macrophages by targeting NF-kappaB. *J Exp Med* 2008; **205**: 1261-1268
  - 40 **Mauad TH**, van Nieuwkerk CM, Dingemans KP, Smit JJ, Schinkel AH, Notenboom RG, van den Bergh Weerman MA, Verkruijsen RP, Groen AK, Oude Elferink RP. Mice with homozygous disruption of the *mdr2* P-glycoprotein gene. A novel animal model for studies of nonsuppurative inflammatory cholangitis and hepatocarcinogenesis. *Am J Pathol* 1994; **145**: 1237-1245
  - 41 **Maeda S**, Kamata H, Luo JL, Leffert H, Karin M. IKKbeta couples hepatocyte death to cytokine-driven compensatory proliferation that promotes chemical hepatocarcinogenesis. *Cell* 2005; **121**: 977-990
  - 42 **Jeong WI**, Park O, Radaeva S, Gao B. STAT1 inhibits liver fibrosis in mice by inhibiting stellate cell proliferation and stimulating NK cell cytotoxicity. *Hepatology* 2006; **44**: 1441-1451
  - 43 **Kortylewski M**, Kujawski M, Wang T, Wei S, Zhang S, Pilon-Thomas S, Niu G, Kay H, Mulé J, Kerr WG, Jove R, Pardoll D, Yu H. Inhibiting Stat3 signaling in the hematopoietic system elicits multicomponent antitumor immunity. *Nat Med* 2005; **11**: 1314-1321
  - 44 **Sano S**, Chan KS, DiGiovanni J. Impact of Stat3 activation upon skin biology: a dichotomy of its role between homeostasis and diseases. *J Dermatol Sci* 2008; **50**: 1-14
  - 45 **Norgauer J**, Metzner B, Schraufstatter I. Expression and growth-promoting function of the IL-8 receptor beta in human melanoma cells. *J Immunol* 1996; **156**: 1132-1137
  - 46 **Balentien E**, Mufson BE, Shattuck RL, Derynck R, Richmond A. Effects of MGSA/GRO alpha on melanocyte transformation. *Oncogene* 1991; **6**: 1115-1124
  - 47 **Owen JD**, Strieter R, Burdick M, Haghnegahdar H, Nannery L, Shattuck-Brandt R, Richmond A. Enhanced tumor-forming capacity for immortalized melanocytes expressing melanoma growth stimulatory activity/growth-regulated cytokine beta and gamma proteins. *Int J Cancer* 1997; **73**: 94-103
  - 48 **Kleeff J**, Kusama T, Rossi DL, Ishiwata T, Maruyama H, Friess H, Büchler MW, Zlotnik A, Korc M. Detection and localization of Mip-3alpha/LARC/Exodus, a macrophage proinflammatory chemokine, and its CCR6 receptor in human pancreatic cancer. *Int J Cancer* 1999; **81**: 650-657
  - 49 **Müller A**, Homey B, Soto H, Ge N, Catron D, Buchanan ME, McClanahan T, Murphy E, Yuan W, Wagner SN, Barrera JL, Mohar A, Verástegui E, Zlotnik A. Involvement of chemokine receptors in breast cancer metastasis. *Nature* 2001; **410**: 50-56
  - 50 **Kaifi JT**, Yekebas EF, Schurr P, Obonyo D, Wachowiak R, Busch P, Heinecke A, Pantel K, Izbicki JR. Tumor-cell homing to lymph nodes and bone marrow and CXCR4 expression in esophageal cancer. *J Natl Cancer Inst* 2005; **97**: 1840-1847
  - 51 **Salvucci O**, Bouchard A, Baccarelli A, Deschênes J, Sauter G, Simon R, Bianchi R, Basik M. The role of CXCR4 receptor expression in breast cancer: a large tissue microarray study. *Breast Cancer Res Treat* 2006; **97**: 275-283
  - 52 **Kim J**, Takeuchi H, Lam ST, Turner RR, Wang HJ, Kuo C, Foshag L, Bilchik AJ, Hoon DS. Chemokine receptor CXCR4 expression in colorectal cancer patients increases the risk for recurrence and for poor survival. *J Clin Oncol* 2005; **23**: 2744-2753
  - 53 **Kulbe H**, Hagemann T, Szlosarek PW, Balkwill FR, Wilson JL. The inflammatory cytokine tumor necrosis factor-alpha regulates chemokine receptor expression on ovarian cancer cells. *Cancer Res* 2005; **65**: 10355-10362
  - 54 **Guicciardi ME**, Mott JL, Bronk SF, Kurita S, Fingas CD, Gores GJ. Cellular inhibitor of apoptosis 1 (cIAP-1) degradation by caspase 8 during TNF-related apoptosis-inducing ligand (TRAIL)-induced apoptosis. *Exp Cell Res* 2011; **317**: 107-116
  - 55 **Gyrd-Hansen M**, Meier P. IAPs: from caspase inhibitors to modulators of NF-kappaB, inflammation and cancer. *Nat Rev Cancer* 2010; **10**: 561-574
  - 56 **Mehrotra S**, Languino LR, Raskett CM, Mercurio AM, Dohi T, Altieri DC. IAP regulation of metastasis. *Cancer Cell* 2010;



- 17: 53-64
- 57 **Lazarevich NL**, Cheremnova OA, Varga EV, Ovchinnikov DA, Kudrjavitseva EI, Morozova OV, Fleishman DI, Engelhardt NV, Duncan SA. Progression of HCC in mice is associated with a downregulation in the expression of hepatocyte nuclear factors. *Hepatology* 2004; **39**: 1038-1047
  - 58 **Zulehner G**, Mikula M, Schneller D, van Zijl F, Huber H, Sieghart W, Grasl-Kraupp B, Waldhör T, Peck-Radosavljevic M, Beug H, Mikulits W. Nuclear beta-catenin induces an early liver progenitor phenotype in hepatocellular carcinoma and promotes tumor recurrence. *Am J Pathol* 2010; **176**: 472-481
  - 59 **Zhai B**, Yan HX, Liu SQ, Chen L, Wu MC, Wang HY. Reduced expression of E-cadherin/catenin complex in hepatocellular carcinomas. *World J Gastroenterol* 2008; **14**: 5665-5673
  - 60 **Fransvea E**, Angelotti U, Antonaci S, Giannelli G. Blocking transforming growth factor-beta up-regulates E-cadherin and reduces migration and invasion of hepatocellular carcinoma cells. *Hepatology* 2008; **47**: 1557-1566
  - 61 **Lee TK**, Poon RT, Yuen AP, Ling MT, Kwok WK, Wang XH, Wong YC, Guan XY, Man K, Chau KL, Fan ST. Twist overexpression correlates with hepatocellular carcinoma metastasis through induction of epithelial-mesenchymal transition. *Clin Cancer Res* 2006; **12**: 5369-5376
  - 62 **Battaglia S**, Benzoubir N, Nobilet S, Charneau P, Samuel D, Zignego AL, Atfi A, Bréchot C, Bourgeade MF. Liver cancer-derived hepatitis C virus core proteins shift TGF-beta responses from tumor suppression to epithelial-mesenchymal transition. *PLoS One* 2009; **4**: e4355
  - 63 **Lee TK**, Man K, Poon RT, Lo CM, Yuen AP, Ng IO, Ng KT, Leonard W, Fan ST. Signal transducers and activators of transcription 5b activation enhances hepatocellular carcinoma aggressiveness through induction of epithelial-mesenchymal transition. *Cancer Res* 2006; **66**: 9948-9956

S- Editor Yang XC L- Editor Webster JR E- Editor Yang XC