


World Journal of *Orthopedics*

World J Orthop 2020 January 18; 11(1): 1-75





MINIREVIEWS

- 1 Patents and intellectual property in orthopaedics and arthroplasty
Uzoigwe CE, Shoaib A

ORIGINAL ARTICLE

Basic Study

- 10 Double-row repair of rotator cuff tears: Comparing tendon contact area between techniques
Ng SHA, Tan CHJ

Case Control Study

- 18 National trends in total hip arthroplasty for traumatic hip fractures: An analysis of a nationwide all-payer database
Boniello AJ, Lieber AM, Denehy K, Cavanaugh P, Kerbel YE, Star A

Retrospective Cohort Study

- 27 Corrections in alpha angle following two different operative approaches for CAM-type femoral acetabular impingement - Ganz surgical hip dislocation *vs* anterior mini-open
Haug EC, Novicoff WM, Cui Q
- 36 Good accuracy of the alpha-defensin lateral flow test for hip periprosthetic joint infection: A pilot study in a retrospective cohort of 52 patients
Kuiper JW, Pander P, Vos SJ

Randomized Controlled Trial

- 47 Effect of deep transverse friction massage *vs* stretching on football players' performance
Fakhro MA, Chahine H, Srouf H, Hijazi K

META-ANALYSIS

- 57 Use of three-dimensional printing in preoperative planning in orthopaedic trauma surgery: A systematic review and meta-analysis
Morgan C, Khatri C, Hanna SA, Ashrafian H, Sarraf KM

CASE REPORT

- 68 Rapid spontaneous resolution of lumbar ganglion cysts: A case report
Chiarella V, Ramieri A, Giugliano M, Domenicucci M

ABOUT COVER

Editorial Board Member of *World Journal of Orthopedics*, Charles L Nelson, MD, Associate Professor, Chief Doctor, Surgeon, Department of Orthopaedic Surgery, University of Pennsylvania Perelman School of Medicine, Philadelphia, PA 19104, United States

AIMS AND SCOPE

The primary aim of *World Journal of Orthopedics* (*WJO*, *World J Orthop*) is to provide scholars and readers from various fields of orthopedics with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJO mainly publishes articles reporting research results and findings obtained in the field of orthopedics and covering a wide range of topics including arthroscopy, bone trauma, bone tumors, hand and foot surgery, joint surgery, orthopedic trauma, osteoarthritis, osteoporosis, pediatric orthopedics, spinal diseases, spine surgery, and sports medicine.

INDEXING/ABSTRACTING

The *WJO* is now abstracted and indexed in PubMed, PubMed Central, Emerging Sources Citation Index (Web of Science), Scopus, China National Knowledge Infrastructure (CNKI), China Science and Technology Journal Database (CSTJ), and Superstar Journals Database.

RESPONSIBLE EDITORS FOR THIS ISSUE

Responsible Electronic Editor: *Mei-Yi Liu*

Proofing Production Department Director: *Xiang Li*

NAME OF JOURNAL

World Journal of Orthopedics

ISSN

ISSN 2218-5836 (online)

LAUNCH DATE

November 18, 2010

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Bao-Gan Peng

EDITORIAL BOARD MEMBERS

<http://www.wjgnet.com/2218-5836/editorialboard.htm>

EDITORIAL OFFICE

Ruo-Yu Ma, Director

PUBLICATION DATE

January 18, 2020

COPYRIGHT

© 2020 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Use of three-dimensional printing in preoperative planning in orthopaedic trauma surgery: A systematic review and meta-analysis

Catrin Morgan, Chetan Khatri, Sammy A Hanna, Hutan Ashrafian, Khaled M Sarraf

ORCID number: Catrin Morgan (0000-0003-1273-8312); Chetan Khatri (0000-0002-3961-1314); Sammy A Hanna (0000-0001-7381-701X); Hutan Ashrafian (0000-0003-1668-0672); Khaled Sarraf (0000-0003-1428-9688).

Author contributions: Morgan C, Khatri C, Ashrafian H and Sarraf K designed research; Morgan C, Khatri C and Ashrafian H performed research; Ashrafian H and Hanna SA contributed new reagents or analytic tools; Morgan C, Khatri C and Ashrafian H analyzed data; Morgan C, Khatri C, Hanna SA, Ashrafian H and Sarraf K wrote the paper.

Conflict-of-interest statement: All authors declare that they have no conflicts of interest.

PRISMA 2009 Checklist statement: Manuscript meets requirements of PRISMA statement

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited

Catrin Morgan, Chetan Khatri, Khaled M Sarraf, Trauma and Orthopaedics, Imperial College NHS Trust, London W2 1NY, United Kingdom

Sammy A Hanna, The Royal London Hospital, Barts Health NHS Trust, London E1 1BB, United Kingdom

Hutan Ashrafian, Department of Surgery and Cancer, Imperial College London, London W2 1NY, United Kingdom

Corresponding author: Catrin Morgan, BSc, MBBS, MRCS, Trauma and Orthopaedics, Imperial College NHS Trust, Praed Street, London W21NY, United Kingdom. catrin.morgan@nhs.net

Abstract

BACKGROUND

With the increasing complexity of surgical interventions performed in orthopaedic trauma surgery and the improving technologies used in three-dimensional (3D) printing, there has been an increased interest in the concept. It has been shown that 3D models allow surgeons to better visualise anatomy, aid in planning and performing complex surgery. It is however not clear how best to utilise the technique and whether this results in better outcomes.

AIM

To evaluate the effect of 3D printing used in pre-operative planning in orthopaedic trauma surgery on clinical outcomes.

METHODS

We performed a comprehensive systematic review of the literature and a meta-analysis. Medline, Ovid and Embase were searched from inception to February 8, 2018. Randomised controlled trials, case-control studies, cohort studies and case series of five patients or more were included across any area of orthopaedic trauma. The primary outcomes were operation time, intra-operative blood loss and fluoroscopy used.

RESULTS

Seventeen studies (922 patients) met our inclusion criteria and were reviewed. The use of 3D printing across all specialties in orthopaedic trauma surgery demonstrated an overall reduction in operation time of 19.85% [95% confidence intervals (CI): (-22.99, -16.71)], intra-operative blood loss of 25.73% [95%CI: (-31.07, -20.40)], and number of times fluoroscopy was used by 23.80% [95%CI: (-38.49, -9.10)].

manuscript

Received: June 11, 2019

Peer-review started: June 11, 2019

First decision: September 21, 2019

Revised: September 25, 2019

Accepted: November 25, 2019

Article in press: November 25, 2019

Published online: January 18, 2020

P-Reviewer: Huang AHC, Khajehei M, Lin JA, Xavier-Elsas P, Zhang ZH

S-Editor: Tang JZ

L-Editor: A

E-Editor: Liu MY


CONCLUSION

Our results suggest that the use of 3D printing in pre-operative planning in orthopaedic trauma reduces operative time, intraoperative blood loss and the number of times fluoroscopy is used.

Key words: Orthopaedic surgery; Trauma; Three-dimensional printing

©The Author(s) 2020. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: The use of three-dimensional (3D) printing in preoperative planning in orthopaedic trauma surgery is a relatively novel field. It has been shown that 3D models allow surgeons to better visualise anatomy, aid in planning and performing complex surgery. It is not clear how to utilise this technique and whether this results in better outcomes. This is the first systematic review and meta-analysis in this field to date. Our results suggest that the use of 3D printing in pre-operative planning in orthopaedic trauma reduces operative time, intraoperative blood loss and the number of times fluoroscopy is used.

Citation: Morgan C, Khatri C, Hanna SA, Ashrafian H, Sarraf KM. Use of three-dimensional printing in preoperative planning in orthopaedic trauma surgery: A systematic review and meta-analysis. *World J Orthop* 2020; 11(1): 57-67

URL: <https://www.wjgnet.com/2218-5836/full/v11/i1/57.htm>

DOI: <https://dx.doi.org/10.5312/wjo.v11.i1.57>

INTRODUCTION

With the increasing complexity of operations and surgical decision-making, three-dimensional (3D) printing is a novel modality with the potential to make a huge impact in the surgical field. In 1984, the first patent for a 3D printer was filled in the United States by Charles Hall titled, "Apparatus for production of 3D objects by stereolithography", which was, in effect, the world's first 3D printer (Patent number: US4575330A). It has been used in a variety of different surgical specialties including plastic surgery, neurosurgery, cardiothoracic surgery, oral and maxillofacial surgery^[1,2]. The first reported use in orthopaedics was in 1999 as an aid to pre-operative planning in complex spinal surgery^[3].

3D printing also known as "rapid prototyping" or "additive manufacturing" creates a 3D model through different techniques. As opposed to traditional manufacturing techniques that are a "subtractive" process (removing excess material), 3D printing is an "additive" process. This involves creating 3D objects by adding material layer-by-layer. Firstly, a high-resolution computed tomography (CT) scan of the deformity, fracture or defect is required. Computer-aided design (CAD) software then creates a digitalised representation of an object, which is then converted most commonly into a stereolithography (STL) file^[4]. STL files "cut up" the digitalised model created by the CAD software, allowing the 3D printer to print the object layer by layer^[5].

In orthopaedics, the use of 3D printing can be broadly split into three categories. This includes: (1) The use of 3D printing in pre-operative planning; (2) 3D implants; and (3) 3D patient-specific instrumentation (PSI). In pre-operative planning, 3D printed models of the fracture configuration or pathology can allow surgeons to visualise relevant anatomy and help aid executing complex operations, for example in pelvic trauma surgery^[6]. 3D printed implants can be used for direct replacement of a large defect after tumour resection and to aid reconstruction in limb-salvage surgery^[7]. PSIs have a wide application across orthopaedics and are largely used for more accurate implant placement, especially in the presence of abnormal anatomy and deformities, as well as developing templates for deformity correction and tumour resection^[8].

Despite 3D printing being described as the third industrial revolution, the necessary evidence to justify the expanding investment in 3D printing in surgery remains ambiguous. Whilst the concept of 3D printing is favoured, particularly in those countries with advanced economies, its widespread application to daily clinical practice is relatively unknown. As we approach the end of the second decade

following the first use of 3D printing in orthopaedics, an overview of this innovation may allow us better understanding of its potential application in every day practice in orthopaedic trauma.

Studies comparing the use of 3D printing with conventional approaches have suggested an improvement in orthopaedic operative outcomes, as measured by blood loss, use of fluoroscopy and operative time^[9-11]. However, the systematic assessment and meta-analysis of 3D printing in orthopaedic trauma as a single entity has not been performed. The aim of this comprehensive systematic review and meta-analysis was to draw evidence from all studies across all areas of orthopaedic trauma, irrespective of age and gender, to assess the overall role of 3D printing in orthopaedic preoperative planning and core surgical outcomes. The primary outcome measures in this review were (1) Operation time; (2) Intra-operative blood loss; and (3) Fluoroscopy used.

MATERIALS AND METHODS

This study was performed in accordance to PRISMA guidelines for the reporting of systematic reviews^[12]. The study protocol was pre-defined and registered on PROSPERO and can be accessed at http://www.crd.york.ac.uk/PROSPERO/display_record.php?ID=CRD42018078429.

Inclusion criteria

This paper included: (1) Full text papers in English language; (2) Humans of any age undergoing orthopaedic trauma procedures; (3) Randomised controlled trials (RCTs), case-control studies, cohort studies and case series of greater than 4 patients in orthopaedic trauma; and (4) Studies reporting the use of 3D printing in preoperative planning in orthopaedic trauma surgery.

Exclusion criteria

Studies were excluded if they were (1) Case reports or case series with less than five patients; (2) Abstract only publication; (3) Involved 3D imaging but not 3D printing during the operative procedure; (4) Involved 3D printing but not preoperative planning; (5) Duplicate data set; (6) Studies assessing the use of 3D printing in manufacturing custom implants and PSI; and (7) Non-trauma (elective) orthopaedic surgery.

Search strategy and study selection

We performed a search of Medline, Ovid and Embase from inception to February, 8 2018 and imported citations into EndNote X7 (New York, United States) reference manager software. Duplicate citations were removed and the remaining citations were screened using title and abstract to match the eligibility criteria. Two authors (Morgan C and Khatri C) independently assessed each paper with discrepancies being resolved with discussion with senior authors (Ashrafian H, Hanna SA, Sarraf K). Additional papers that were appropriate *via* a manual search were added. The full search strategy can be found in the supplementary materials.

Outcome measures

The primary outcome measures in this review were (1) Operation time; (2) Intra-operative blood loss; and (3) Fluoroscopy used.

Data extraction

Data was extracted and entered to Microsoft Excel (Microsoft Cop. Redmond, WA, United States) by two authors (Morgan C and Khatri C). Demographic data was extracted from each paper including year of publication, country of publication and type of 3D printing technology used. Specifically for the participants, the number of participants, age and gender ratio was determined.

Quality assessment

Two authors (Morgan C and Khatri C) independently assessed the quality of included papers. For randomised controlled trials, the Jadad score was used^[13]. This is a well-validated score that assesses the methodological quality of clinical trials by assessing the randomisation process and blinding the study has used. The Jadad score uses five questions, with a point scale of zero to five. We classified a score of ≥ 3 as "higher quality" and < 3 as "lower quality". For all other study types, the Newcastle-Ottawa scale was used^[14]. This score assesses studies based on three categories: (1) The selection of the study groups; (2) The comparability of the study groups; and (3) The ascertainment of the outcome measure in cohort studies. We classified studies with \geq

7 stars as “higher quality” and < 7 stars as “lower quality”.

Statistical analysis

Meta-analysis was performed through the inverse-variance, random-effects model of DerSimonian and Laird for both continuous and categorical outcomes. Continuous outcomes were analysed by calculating the ratio of means for each study, with expression of uncertainty of each result represented by the 95% confidence intervals (CI). We substituted median for mean in studies where only the median was reported^[15]. We performed meta-analysis if two or more separate studies were available. This was accomplished using Stata 15 (StataCorp., College Station, TX, United States). The I² statistic was used to estimate the degree of heterogeneity between studies, where larger values indicate increasing heterogeneity.

RESULTS

Seventeen studies were found that fulfilled the inclusion criteria, producing a data set of 922 patients (Figure 1). Six of these studies were RCTs, two cohort studies (1 prospective, 1 retrospective) and nine case series (8 prospective, 1 retrospective). The area of orthopaedics and type of pathology varied and are shown in Table 1. The overall mean follow up was 18.16 mo. Data from studies with comparison groups that specifically looked at the primary outcomes were included in the meta-analysis.

Operation time

Operation time was measured in eight of these studies; six were RCTs^[6,9-11,16-19]. Meta-analysis of these studies confirmed a significant reduction in operation time of 19.85% in the 3D printing group [95%CI: (-22.99, -16.71)] in comparison to the conventional group (see Figure 2). The heterogeneity was high ($I^2 = 100.0\%$).

Intra-operative blood loss

Intra-operative blood loss was investigated in seven studies; six were RCTs^[6,9-11,16-18]. Meta-analysis revealed a significant reduction in blood loss of 25.73% (see Figure 3) with the use of 3D printing [95%CI: (-31.07, -20.40)]. The heterogeneity was high ($I^2 = 100.0\%$).

Fluoroscopy used

The number of times fluoroscopy was used was measured in four RCTs^[6,9-11]. The use of 3D printing led to a significant reduction of 23.80% (see Figure 4) in the number of times fluoroscopy was used [95%CI: (-38.49, -9.10)]. The heterogeneity was high ($I^2 = 100.0\%$).

DISCUSSION

Our analysis demonstrates that the use of 3D printing in preoperative planning in orthopaedic trauma, can lead to a reduction in operation time, intra-operative blood loss and fluoroscopy used. Our systematic review represents the largest and most comprehensive in this area and is the first meta-analysis to date. Although research in the field consists of small-sized studies, findings confirm the current evidence in these studies of the positive effect of 3D printing in preoperative planning.

Operation time

A shortened operation time of 19.85% with the use of 3D printing confers significant benefits to the patient and costs associated. Increased operative time has been found to be associated with an increased risk of developing post-operative complications, such as wound infections and deep vein thrombosis^[29,30]. The reasons behind this are multifactorial and include: (1) the surgeon having a stronger appreciation of pathological anatomy through geometric characterisation by the 3D printed model; as such it allows the surgeon to physically review complex cases. For example, in pelvic surgery it allows the surgeon greater understanding of acetabular fractures and reduces the degree of inter-observer variability in the classification of these injuries^[28]; (2) It facilitates pre-operative instrumentation decisions; and (3) Facilitates the planning of the surgical approach^[10]; for example in trimalleolar fractures, the surgeon is able to determine the size of the internal fixation plate, clarify the placement position of the plate and screws and prebend the plate according to the morphology of the bone^[16]; and (4) The 3D model also enables simulation of the surgery *in vitro* allowing the surgeon better understanding of fracture reduction, hence leading to a reduction in the operation time (in comparison to the control groups)^[11]. Together,

Table 1 Results summary

Ref.	Yr	Pathology	Area of orthopaedics	Country of study	Design	Newcastle ottawa Score	Jadad score (RCT)	3DP technology	Total participants	Mean age	% Male
Bagaria <i>et al</i> ^[20]	2017	All	Mix	India	Retrospective case series	1		FDM	50		
Belien <i>et al</i> ^[21]	2017	Os acromiale and acrominal fractures	Upper limb	Belgium	Prospective case series	3		STL	5		
Chen <i>et al</i> ^[6]	2017	Die-Punch fractures	Upper limb	China	Randomised control trial		0	STL	107	28	60.7
Zheng <i>et al</i> ^[18]	2017	Intertrochanteric fracture	Hip	China	Retrospective cohort study	7		STL	39	66	56.4
Zheng <i>et al</i> ^[10]	2017	Calcaneal fractures	Lower limb	China	Randomised control trial		1	STL	75	45	58.7
Zheng <i>et al</i> ^[9]	2018	Humeral Intercondylar fractures	Upper limb	China	Randomised control trial		1	STL	91	44.6	53.84
Yang <i>et al</i> ^[17]	2017	Elbow fractures	Upper limb	China	Randomised control trial		2	STL	40	38.6	70
Zhang <i>et al</i> ^[22]	2017	Lower limb fractures	Lower limb	China	Prospective case series	1		STL	78	56	60.3
Bizzotto <i>et al</i> ^[23]	2016	Articular fractures	Upper and lower limb	Italy	Prospective case series	1		STL	102		44.1
Yang <i>et al</i> ^[16]	2016	Trimalleolar fractures	Lower limb	China	Randomised control trial		1	STL	30	36.5	53.3
Yang <i>et al</i> ^[24]	2016	Lateral tibial plateau fractures	Lower limb	China	Prospective case series	2		FDM	7	44	42.9
You <i>et al</i> ^[11]	2016	Proximal humeral fractures	Upper limb	China	Randomised control trial		2		66	66	40.9
Zeng <i>et al</i> ^[25]	2016	Acetabular fracture	Pelvis	China	Prospective case series	3		STL	10		50
Li <i>et al</i> ^[19]	2016	Tile C pelvic fracture	Pelvis	China	Retrospective cohort study	5			157	33	67.5
Kim <i>et al</i> ^[26]	2015	Midshaft clavicle fractures	Upper limb	China	Prospective case series	0		STL	7		
Zeng <i>et al</i> ^[27]	2015	Pelvic fracture	Pelvis	China	Prospective case series	1			38	32	65.8
Hurson <i>et al</i> ^[28]	2007	Acetabular fractures	Pelvis	Ireland	Prospective case series	1		SLS	20		

RCT: Randomised controlled trial; FDM: Fused deposition modeling; STL: Stereolithography; SLS: Selective laser sintering.

these elements of 3D printing that lead to a decreased operative time can translate into cost savings, specifically through decreased staff and operating room overheads. There is evidence to suggest that a decrease in operative time confers cost savings with a 45-min reduction in operation time leading to a saving of almost \$2700 per case for a paediatric population^[31].

Intra-operative blood loss

Reducing blood loss and the need for blood transfusion in orthopaedic surgery remains a concern among surgeons and physicians in the perioperative period^[32].




Figure 1 Search strategy.

Many different methods have been developed including; controlled hypotensive anaesthesia, cell salvage techniques^[33], and pharmacological interventions (tranexamic acid)^[34]. Our results show a significant reduction in intra-operative blood loss (25.73%) with the use of 3D printing which carries powerful implications for pre- and peri-operative planning and practice especially in the context of trauma. A parallel can be drawn with a reduction in blood loss found with the use of robotic surgery^[15]. Reasons studies attributed to the reduction of intra-operative blood loss are similar to those for reduction in operation time (better understanding of pathology and ability to plan the operation). The use of 3D printing in pelvic ring fractures demonstrated a reduced volume of fluid infused intra-operatively, with patients less likely to require a surgical drain or blood transfusion post-operatively^[19]. Similar findings were reported with the use of 3D models in elbow fractures, it allowed the plate to be chosen and contoured accurately to ensure optimal compactness with the bone surface^[17]. This minimised dissection of the surrounding tissue and protected the blood supply surrounding the fracture fragments, hence reducing intraoperative blood loss.

Fluoroscopy used

A small number of studies looked at the number of times intraoperative fluoroscopy was used and overall they reported a reduction in fluoroscopy of 23.80% with the use of 3D printing. This included calcaneal fractures, distal radius fractures and humeral fractures^[6,9-11]. A reduction in fluoroscopy carries safety benefits through minimising patient radiation exposure. Radiation exposure remains a significant occupational hazard to the orthopaedic surgeon, radiographers and theatre staff throughout their careers^[35] and also to patients subjected to this radiation. Although there is limited




Figure 2 Forest plot of operation time with the use of three-dimensional printing.

literature available on the appropriate amount of intraoperative fluoroscopy use in orthopaedics, a recent paper has reported reference values for common trauma operations^[36]. Patient exposure to intraoperative radiation is an important factor to consider especially in the paediatric trauma setting as children are up to ten times more sensitive to radiation damage than adults^[37].

Cost and production

An obstacle that exists for the transition of 3D printed models from small studies to large scale clinical practice is cost and production. Creating a 3D printed model, is a complex process requiring advanced computer software and a 3D printer. This process does not come without cost and the need for an experienced team that are familiar with the software. Clinically, not all fractures may require a CT scan, a plain radiograph may suffice, therefore the use of 3D printing may be most useful for complex fractures, for example in pelvic trauma. The cost of printing the model seemed to vary greatly from \$2-3 per model to \$330^[17,38]. There is also the cost of the printer to consider with studies reporting the cost to be \$1000-2200^[9,31]. In the studies reviewed, the time taken to produce the model ranged from 5 h to 72 h, which is a potential limitation to the use of 3D printing in an acute trauma setting^[6,18]. Nevertheless, an important factor to consider is that 3D printing in orthopaedics is a novel field. Therefore, as 3D printing technology progresses both the cost and production time is likely to reduce. Today, the most basic 3D printer for home use can be purchased for less than \$200 compared to when 3D printers were first used in the 1980s costing around \$300000 (at that time, corresponding to over \$750000 in the current era)^[39].

Patient understanding and surgical training

A number of studies in this review conducted patient and surgeon questionnaires on 3D printing^[6,9,10,17]. It was found that the 3D models improved patient understanding of the fracture and communication with the doctor, leading to higher compliance with post-operative rehabilitation. Patients felt that conventional medical images such as plain radiograph or CT are too complicated to understand, however 3D models allow better visualisation of the fracture^[17]. This technology therefore represents a powerful modality to enhance patient empowerment.

The reduction in the total hours worked by surgical trainees has led to an increase in the use of simulation-based training in orthopaedics^[40,41], this has been used in fields such as arthroplasty, arthroscopy and trauma surgery^[42,43]. With the increase in operative management for complex deformities, using 3D models enables these deformities to be recreated allowing the trainees to learn how to manage these challenging cases through simulation and apply this knowledge to the operating theatre. The advantages of 3D printing therefore have the potential to decrease surgical learning-curves and possibly improve surgical outcomes.




Figure 3 Forest plot of blood loss with the use of three-dimensional printing.

Strengths and limitations

This is the first comprehensive systematic review and meta-analysis conducted on the use of 3D printing in preoperative planning across all types of orthopaedic trauma surgery on core surgical outcomes. It included 17 studies in which six were randomised control trials. However, the results presented here should be considered in the context of a number of limitations. Where meta-analysis was possible, the heterogeneity was high. However, this was not unexpected given the wide area of orthopaedics (where different specialties are highly distinct), type of pathology and patient demographics. It was not possible to carry out a subgroup analysis due to the limited number of studies in the different subspecialties of orthopaedic trauma. A high level of heterogeneity in the absence of any other clear evidence highlights that this scientific field is still in its infancy where its methodology and practices are highly variable, nevertheless the results of this integrated data does indicate the results of clinical effect in the absence of more robust evidence (which is clearly necessary in future studies).

Our focus on operation time, blood loss and number of times fluoroscopy was used was based primarily on the fact that these were the most commonly reported outcomes in the literature. However, these parameters may not fully demonstrate the true value of 3D printing in orthopaedics, parameters such as post-operative function or complications were not determined.

Case series were included to address the paucity of RCTs inevitably introducing other biases associated with this study design and contributing to the heterogeneity. Of note twenty-three of these studies were carried out in China comprising of 81% (745) of the total number of patients. Apart from the randomised control trials, overall the quality scoring of the studies were low, with all 11 studies scoring six or less in the Newcastle-Ottawa Scale. All six of the randomised control trials were of low quality as determined by the Jadad scale. Reasons for low scores include lack of surgeon blinding, which is unlikely to be possible in clinical trials of 3D printing. In many trials, however, the risk of bias relating to sequence generation, adequate follow up, study dropout rate and assessment of outcome was unclear as sufficient information was not available due to poor reporting. The risk of publication bias should also be considered. The number of participants in these studies were small, with the largest being 157, therefore the possibility of “small study effect” cannot be ignored.

CONCLUSION

In summary, our meta-analysis and systematic review on the use of 3D printing in preoperative planning in orthopaedic trauma suggests that 3D printing reduces operative time, intraoperative blood loss and the number of times fluoroscopy is used. 3D printing is a rapidly evolving field and it allows surgeons to gain better




Figure 4 Forest plot of fluoroscopy used with the use of three-dimensional printing.

understanding of complex trauma cases and aid in surgical planning. The increased application of this technology has the potential to revolutionise orthopaedic practice and enhance clinical outcomes. There is however the need for an in-depth cost-analysis for the use of 3D printing in surgery, taking into account the production cost *vs* potential savings made by improved intra-operative outcomes. Further studies, in particular more randomised control trials in similar areas of orthopaedics are required to further enhance our knowledge of the role of 3D printing in orthopaedics and its application to daily clinical practice.

ARTICLE HIGHLIGHTS

Research background

With the increasing complexity of operations and surgical decision-making, three-dimensional (3D) printing is a novel modality with the potential to make a huge impact in the surgical field. In orthopaedics, the use of 3D printing can be broadly split into three categories. This includes: (1) The use of 3D printing in pre-operative planning; (2) 3D implants; and (3) 3D patient-specific instrumentation. In pre-operative planning, 3D printed models of the fracture configuration or pathology can allow surgeons to visualise relevant anatomy and help aid executing complex operations. It is however not clear how best to utilise the technique and whether this results in better outcomes.

Research motivation

The focus of this study is the use of 3D printing in preoperative planning in orthopaedic trauma surgery. Studies comparing the use of 3D printing with conventional approaches have suggested an improvement in orthopaedic operative outcomes, as measured by blood loss, use of fluoroscopy and operative time. However, the systematic assessment and meta-analysis of 3D printing in orthopaedic trauma as a single entity has not been performed. The increased application of this technology has the potential to revolutionize orthopaedic practice and enhance clinical outcomes.

Research objectives

The aim of this research was to draw evidence from all studies across all areas of orthopaedic trauma, irrespective of age and gender, to assess the overall role of 3D printing in orthopaedic preoperative planning and core surgical outcomes. The primary outcome measures in this review were (1) Operation time; (2) Intra-operative blood loss; and (3) Fluoroscopy used.

Research methods

This study was performed in accordance to PRISMA guidelines for the reporting of systematic reviews. The study protocol was pre-defined and registered on PROSPERO. A search of Medline, Ovid and Embase from inception to February 8, 2018 was carried out and citations were imported into EndNote X7 (New York, United States) reference manager software. Two authors independently assessed the quality of included papers. For randomised controlled trials, the Jadad score was used and for all other study types, the Newcastle-Ottawa scale was used. Meta-analysis was performed through the inverse-variance, random-effects model of DerSimonian and Laird for both continuous and categorical outcomes.

Research results

Seventeen studies (922 patients) met our inclusion criteria and were reviewed. The use of 3D printing across all specialties in orthopaedic trauma surgery demonstrated an overall reduction in operation time of 19.85% [95% confidence interval (CI): -22.99, -16.71], intra-operative blood loss of 25.73% (95%CI: -31.07, -20.40) and number of times fluoroscopy was used by 23.80% (95%CI: -38.49, -9.10). Overall, the quality of the studies was low and it was not possible to carry out a subgroup analysis due to the limited number of studies in the different subspecialties of orthopaedic trauma.

Research conclusions

This meta-analysis and systematic review on the use of 3D printing in preoperative planning in orthopaedic trauma suggests that 3D printing reduces operative time, intraoperative blood loss and the number of times fluoroscopy is used. 3D printing is a rapidly evolving field and it allows surgeons to gain better understanding of complex trauma cases and aid in surgical planning. 3D printing should be considered as an adjunct to improve patient care by minimising operative insult in orthopaedic trauma surgery.

Research perspectives

The study highlights the potential impact 3D printing can have in orthopaedic trauma surgery. Further studies, in particular more randomised control trials in similar areas of orthopaedics are required to further enhance our knowledge of the role of 3D printing in orthopaedics and its application to daily clinical practice. There is also the need for an in-depth cost-analysis for the use of 3D printing in surgery, taking into account the production cost *vs* potential savings made by improved intra-operative outcomes.

REFERENCES

- 1 **Giannopoulos AA**, Steigner ML, George E, Barile M, Hunsaker AR, Rybicki FJ, Mitsouras D. Cardiothoracic Applications of 3-dimensional Printing. *J Thorac Imaging* 2016; **31**: 253-272 [PMID: 27149367 DOI: 10.1097/RTI.0000000000000217]
- 2 **Chae MP**, Rozen WM, McMenamin PG, Findlay MW, Spychal RT, Hunter-Smith DJ. Emerging Applications of Bedside 3D Printing in Plastic Surgery. *Front Surg* 2015; **2**: 25 [PMID: 26137465 DOI: 10.3389/fsurg.2015.00025]
- 3 **D'Urso PS**, Askin G, Earwaker JS, Merry GS, Thompson RG, Barker TM, Effeney DJ. Spinal bi modeling. *Spine (Phila Pa 1976)* 1999; **24**: 1247-1251 [PMID: 10382253 DOI: 10.1097/00007632-199906150-00013]
- 4 **Tack P**, Victor J, Gemmel P, Annemans L. 3D-printing techniques in a medical setting: a systematic literature review. *Biomed Eng Online* 2016; **15**: 115 [PMID: 27769304 DOI: 10.1186/s12938-016-0236-4]
- 5 **Heller M**, Bauer HK, Goetze E, Gielisch M, Roth KE, Drees P, Maier GS, Dorweiler B, Ghazy A, Neufurth M, Müller WE, Schröder HC, Wang X, Vahl CF, Al-Nawas B. Applications of patient-specific 3D printing in medicine. *Int J Comput Dent* 2016; **19**: 323-339 [PMID: 28008429]
- 6 **Chen C**, Cai L, Zhang C, Wang J, Guo X, Zhou Y. Treatment of Die-Punch Fractures with 3D Printing Technology. *J Invest Surg* 2018; **31**: 385-392 [PMID: 28722489 DOI: 10.1080/08941939.2017.1339150]
- 7 **Ma L**, Zhou Y, Zhu Y, Lin Z, Chen L, Zhang Y, Xia H, Mao C. 3D printed personalized titanium plates improve clinical outcome in microwave ablation of bone tumors around the knee. *Sci Rep* 2017; **7**: 7626 [PMID: 28790331 DOI: 10.1038/s41598-017-07243-3]
- 8 **Ma L**, Zhou Y, Zhu Y, Lin Z, Wang Y, Zhang Y, Xia H, Mao C. 3D-printed guiding templates for improved osteosarcoma resection. *Sci Rep* 2016; **6**: 23335 [PMID: 26997197 DOI: 10.1038/srep23335]
- 9 **Zheng W**, Su J, Cai L, Lou Y, Wang J, Guo X, Tang J, Chen H. Application of 3D-printing technology in the treatment of humeral intercondylar fractures. *Orthop Traumatol Surg Res* 2018; **104**: 83-88 [PMID: 29248764 DOI: 10.1016/j.otsr.2017.11.012]
- 10 **Zheng W**, Tao Z, Lou Y, Feng Z, Li H, Cheng L, Zhang H, Wang J, Guo X, Chen H. Comparison of the Conventional Surgery and the Surgery Assisted by 3d Printing Technology in the Treatment of Calcaneal Fractures. *J Invest Surg* 2018; **31**: 557-567 [PMID: 28925760 DOI: 10.1080/08941939.2017.1363833]
- 11 **You W**, Liu LJ, Chen HX, Xiong JY, Wang DM, Huang JH, Ding JL, Wang DP. Application of 3D printing technology on the treatment of complex proximal humeral fractures (Neer3-part and 4-part) in old people. *Orthop Traumatol Surg Res* 2016; **102**: 897-903 [PMID: 27521179 DOI: 10.1016/j.otsr.2016.06.009]
- 12 **Moher D**, Liberati A, Tetzlaff J, Altman DG; PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ* 2009; **339**: b2535 [PMID: 19622551 DOI: 10.1136/bmj.b2535]
- 13 **Jadad AR**, Moore RA, Carroll D, Jenkinson C, Reynolds DJ, Gavaghan DJ, McQuay HJ. Assessing the quality of reports of randomized clinical trials: is blinding necessary? *Control Clin Trials* 1996; **17**: 1-12 [PMID: 8721797 DOI: 10.1016/0197-2456(95)00134-4]
- 14 **Stang A**. Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses. *Eur J Epidemiol* 2010; **25**: 603-605 [PMID: 20652370 DOI: 10.1007/s10654-010-9491-z]
- 15 **Tan A**, Ashrafi H, Scott AJ, Mason SE, Harling L, Athanasiou T, Darzi A. Robotic surgery: disruptive innovation or unfulfilled promise? A systematic review and meta-analysis of the first 30 years. *Surg Endosc* 2016; **30**: 4330-4352 [PMID: 26895896 DOI: 10.1007/s00464-016-4752-x]
- 16 **Yang L**, Shang XW, Fan JN, He ZX, Wang JJ, Liu M, Zhuang Y, Ye C. Application of 3D Printing in the Surgical Planning of Trimalleolar Fracture and Doctor-Patient Communication. *Biomed Res Int* 2016; **2016**: 2482086 [PMID: 27446944 DOI: 10.1155/2016/2482086]
- 17 **Yang L**, Grottkau B, He Z, Ye C. Three dimensional printing technology and materials for treatment of elbow fractures. *Int Orthop* 2017; **41**: 2381-2387 [PMID: 28856399 DOI: 10.1007/s00264-017-3627-7]
- 18 **Zheng SN**, Yao QQ, Mao FY, Zheng PF, Tian SC, Li JY, Yu YF, Liu S, Zhou J, Hu J, Xu Y, Tang K, Lou Y, Wang LM. Application of 3D printing rapid prototyping-assisted percutaneous fixation in the treatment

- of intertrochanteric fracture. *Exp Ther Med* 2017; **14**: 3644-3650 [PMID: 29042960 DOI: 10.3892/etm.2017.4991]
- 19 **Li B**, Chen B, Zhang Y, Wang X, Wang F, Xia H, Yin Q. Comparative use of the computer-aided angiography and rapid prototyping technology versus conventional imaging in the management of the Tile C pelvic fractures. *Int Orthop* 2016; **40**: 161-166 [PMID: 25994426 DOI: 10.1007/s00264-015-2800-0]
 - 20 **Bagaria V**, Chaudhary K. A paradigm shift in surgical planning and simulation using 3Dgraphy: Experience of first 50 surgeries done using 3D-printed biomodels. *Injury* 2017; **48**: 2501-2508 [PMID: 28882373 DOI: 10.1016/j.injury.2017.08.058]
 - 21 **Belien H**, Biesmans H, Steenwerckx A, Bijmens E, Dierickx C. Prebending of osteosynthesis plate using 3D printed models to treat symptomatic os acromiale and acromial fracture. *J Exp Orthop* 2017; **4**: 34 [PMID: 29067535 DOI: 10.1186/s40634-017-0111-7]
 - 22 **Zhang W**, Ji Y, Wang X, Liu J, Li D. Can the recovery of lower limb fractures be achieved by use of 3D printing mirror model? *Injury* 2017; **48**: 2485-2495 [PMID: 28899557 DOI: 10.1016/j.injury.2017.09.003]
 - 23 **Bizzotto N**, Tami I, Santucci A, Adani R, Poggi P, Romani D, Carpeggiani G, Ferraro F, Festa S, Magnan B. 3D Printed replica of articular fractures for surgical planning and patient consent: a two years multi-centric experience. *3D Print Med* 2015; **2**: 2 [PMID: 30050974 DOI: 10.1186/s41205-016-0006-8]
 - 24 **Yang P**, Du D, Zhou Z, Lu N, Fu Q, Ma J, Zhao L, Chen A. 3D printing-assisted osteotomy treatment for the malunion of lateral tibial plateau fracture. *Injury* 2016; **47**: 2816-2821 [PMID: 27702464 DOI: 10.1016/j.injury.2016.09.025]
 - 25 **Zeng C**, Xing W, Wu Z, Huang H, Huang W. A combination of three-dimensional printing and computer-assisted virtual surgical procedure for preoperative planning of acetabular fracture reduction. *Injury* 2016; **47**: 2223-2227 [PMID: 27372187 DOI: 10.1016/j.injury.2016.03.015]
 - 26 **Kim HN**, Liu XN, Noh KC. Use of a real-size 3D-printed model as a preoperative and intraoperative tool for minimally invasive plating of comminuted midshaft clavicle fractures. *J Orthop Surg Res* 2015; **10**: 91 [PMID: 26054648 DOI: 10.1186/s13018-015-0233-5]
 - 27 **Zeng C**, Xiao J, Wu Z, Huang W. Evaluation of three-dimensional printing for internal fixation of unstable pelvic fracture from minimal invasive para-rectus abdominis approach: a preliminary report. *Int J Clin Exp Med* 2015; **8**: 13039-13044 [PMID: 26550226]
 - 28 **Hurson C**, Tansey A, O'Donnchadha B, Nicholson P, Rice J, McElwain J. Rapid prototyping in the assessment, classification and preoperative planning of acetabular fractures. *Injury* 2007; **38**: 1158-1162 [PMID: 17884058 DOI: 10.1016/j.injury.2007.05.020]
 - 29 **Daley BJ**, Cecil W, Clarke PC, Cofer JB, Guillaumondegui OD. How slow is too slow? Correlation of operative time to complications: an analysis from the Tennessee Surgical Quality Collaborative. *J Am Coll Surg* 2015; **220**: 550-558 [PMID: 25728140 DOI: 10.1016/j.jamcollsurg.2014.12.040]
 - 30 **Kim JY**, Khavanin N, Rambachan A, McCarthy RJ, Mlodinow AS, De Oliveria GS, Stock MC, Gust MJ, Mahvi DM. Surgical duration and risk of venous thromboembolism. *JAMA Surg* 2015; **150**: 110-117 [PMID: 25472485 DOI: 10.1001/jamasurg.2014.1841]
 - 31 **Cherkasskiy L**, Caffrey JP, Szweczyk AF, Cory E, Bomar JD, Farnsworth CL, Jeffords M, Wenger DR, Sah RL, Upasani VV. Patient-specific 3D models aid planning for triplane proximal femoral osteotomy in slipped capital femoral epiphysis. *J Child Orthop* 2017; **11**: 147-153 [PMID: 28529664 DOI: 10.1302/1863-2548-11-170277]
 - 32 **Ponnusamy KE**, Kim TJ, Khanuja HS. Perioperative blood transfusions in orthopaedic surgery. *J Bone Joint Surg Am* 2014; **96**: 1836-1844 [PMID: 25378512 DOI: 10.2106/JBJS.N.00128]
 - 33 **Sharrock NE**, Mineo R, Urquhart B, Salvati EA. The effect of two levels of hypotension on intraoperative blood loss during total hip arthroplasty performed under lumbar epidural anesthesia. *Anesth Analg* 1993; **76**: 580-584 [PMID: 8452271 DOI: 10.1213/00000539-199303000-00023]
 - 34 **Poeran J**, Rasul R, Suzuki S, Danninger T, Mazumdar M, Opperer M, Boettner F, Memtsoudis SG. Tranexamic acid use and postoperative outcomes in patients undergoing total hip or knee arthroplasty in the United States: retrospective analysis of effectiveness and safety. *BMJ* 2014; **349**: g4829 [PMID: 25116268 DOI: 10.1136/bmj.g4829]
 - 35 **Giordano BD**, Grauer JN, Miller CP, Morgan TL, Rechline GR 2nd. Radiation exposure issues in orthopaedics. *J Bone Joint Surg Am* 2011; **93**: e69(1-e610) [PMID: 21498490 DOI: 10.2106/JBJS.J.01328]
 - 36 **Rashid MS**, Aziz S, Haydar S, Fleming SS, Datta A. Intra-operative fluoroscopic radiation exposure in orthopaedic trauma theatre. *Eur J Orthop Surg Traumatol* 2018; **28**: 9-14 [PMID: 28798994 DOI: 10.1007/s00590-017-2020-y]
 - 37 **Furlow B**. Radiation protection in pediatric imaging. *Radiol Technol* 2011; **82**: 421-439 [PMID: 21572064 DOI: 10.1016/j.anbehav.2011.05.013]
 - 38 **Yang JC**, Ma XY, Xia H, Wu ZH, Ai FZ, Zhang K, Yin QS. Clinical application of computer-aided design-rapid prototyping in C1-C2 operation techniques for complex atlantoaxial instability. *J Spinal Disord Tech* 2014; **27**: E143-E150 [PMID: 24866908 DOI: 10.1097/01.bsd.0000450173.95940.ed]
 - 39 **Hoang D**, Perrault D, Stevanovic M, Ghiassi A. Surgical applications of three-dimensional printing: a review of the current literature & how to get started. *Ann Transl Med* 2016; **4**: 456 [PMID: 28090512 DOI: 10.21037/atm.2016.12.18]
 - 40 **Palan J**, Roberts V, Bloch B, Kulkarni A, Bhowal B, Dias J. The use of a virtual learning environment in promoting virtual journal clubs and case-based discussions in trauma and orthopaedic postgraduate medical education: the Leicester experience. *J Bone Joint Surg Br* 2012; **94**: 1170-1175 [PMID: 22933486 DOI: 10.1302/0301-620X.94B9.28780]
 - 41 **Sonnadara RR**, Van Vliet A, Safir O, Alman B, Ferguson P, Kraemer W, Reznick R. Orthopedic boot camp: examining the effectiveness of an intensive surgical skills course. *Surgery* 2011; **149**: 745-749 [PMID: 21236456 DOI: 10.1016/j.surg.2010.11.011]
 - 42 **Coughlin RP**, Pauyo T, Sutton JC, Coughlin LP, Bergeron SG. A Validated Orthopaedic Surgical Simulation Model for Training and Evaluation of Basic Arthroscopic Skills. *J Bone Joint Surg Am* 2015; **97**: 1465-1471 [PMID: 26333743 DOI: 10.2106/JBJS.N.01140]
 - 43 **Sugand K**, Mawkin M, Gupte C. Validating Touch Surgery™: A cognitive task simulation and rehearsal app for intramedullary femoral nailing. *Injury* 2015; **46**: 2212-2216 [PMID: 26094504 DOI: 10.1016/j.injury.2015.05.013]



Published By Baishideng Publishing Group Inc
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-2238242
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

