

World Journal of *Orthopedics*

World J Orthop 2022 September 18; 13(9): 777-875



EXPERT CONSENSUS

- 777 Consensus Delphi study on guidelines for the assessment of anterior cruciate ligament injuries in children
Campón Chekroun A, Velázquez-Saornil J, Guillén Vicente I, Sánchez Milá Z, Rodríguez-Sanz D, Romero-Morales C, Fernandez-Jaén T, Garrido González JI, Sánchez-Garrido MÁ, Guillén García P

ORIGINAL ARTICLE**Case Control Study**

- 791 Histological difference in ligament flavum between degenerative lumbar canal stenosis and non-stenotic group: A prospective, comparative study
Jain M, Sable M, Tirpude AP, Sahu RN, Samanta SK, Das G

Clinical Trials Study

- 802 Short arm cast is as effective as long arm cast in maintaining distal radius fracture reduction: Results of the SLA-VER noninferiority trial
Dib G, Maluta T, Cengarle M, Bernasconi A, Marconato G, Corain M, Magnan B

Observational Study

- 812 Revision anterior cruciate ligament reconstruction: Return to sports at a minimum 5-year follow-up
Ortiz E, Zicaro JP, Garcia Mansilla I, Yacuzzi C, Costa-Paz M

SYSTEMATIC REVIEWS

- 825 Fragility of statistically significant findings from randomized clinical trials of surgical treatment of humeral shaft fractures: A systematic review
Morris SC, Gowd AK, Agarwalla A, Phipatanakul WP, Amin NH, Liu JN
- 837 Return to work following shoulder arthroplasty: A systematic review
Lalehzarian SP, Agarwalla A, Liu JN

SCIENTOMETRICS

- 853 Evolution of evidence in spinal surgery – past, present and future Scientometric analysis of randomized controlled trials in spinal surgery
Muthu S, Jeyaraman M, Jeyaraman N

CASE REPORT

- 870 Calcium pyrophosphate dihydrate crystals in a 9-year-old with osteomyelitis of the knee: A case report
Pavlis W, Constantinescu DS, Murgai R, Barnhill S, Black B

ABOUT COVER

Editorial Board Member of *World Journal of Orthopedics*, Andrei Fernandes Fernandes Joaquim, MD, PhD, Associate Professor, Neurosurgeon, Department of Neurology, State University of Campinas, Campinas 13083-970, Brazil. andjoaquim@yahoo.com

AIMS AND SCOPE

The primary aim of *World Journal of Orthopedics* (WJO, *World J Orthop*) is to provide scholars and readers from various fields of orthopedics with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJO mainly publishes articles reporting research results and findings obtained in the field of orthopedics and covering a wide range of topics including arthroscopy, bone trauma, bone tumors, hand and foot surgery, joint surgery, orthopedic trauma, osteoarthropathy, osteoporosis, pediatric orthopedics, spinal diseases, spine surgery, and sports medicine.

INDEXING/ABSTRACTING

WJO is now abstracted and indexed in PubMed, PubMed Central, Emerging Sources Citation Index (Web of Science), Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 edition of Journal Citation Reports® cites the 2021 Journal Citation Indicator (JCI) for WJO as 0.62. The WJO's CiteScore for 2021 is 2.4 and Scopus CiteScore rank 2021: Orthopedics and Sports Medicine is 139/284.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Ying-Yi Yuan*, Production Department Director: *Xiang Li*, Editorial Office Director: *Jin-Lai Wang*.

NAME OF JOURNAL

World Journal of Orthopedics

ISSN

ISSN 2218-5836 (online)

LAUNCH DATE

November 18, 2010

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Massimiliano Leigheb

EDITORIAL BOARD MEMBERS

<http://www.wjgnet.com/2218-5836/editorialboard.htm>

PUBLICATION DATE

September 18, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Return to work following shoulder arthroplasty: A systematic review

Simon P Lalehzarian, Avinesh Agarwalla, Joseph N Liu

Specialty type: Orthopedics

Provenance and peer review:

Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B, B
Grade C (Good): C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Luo ZW, China;
Mastrantonakis K, Greece

Received: February 1, 2022

Peer-review started: February 1, 2022

First decision: May 31, 2022

Revised: June 13, 2022

Accepted: August 22, 2022

Article in press: August 22, 2022

Published online: September 18, 2022



Simon P Lalehzarian, The Chicago Medical School, Rosalind Franklin University of Medicine and Science, North Chicago, IL 60064, United States

Avinesh Agarwalla, Department of Orthopedic Surgery, Westchester Medical Center, Valhalla, NY 10595, United States

Joseph N Liu, USC Epstein Family Center for Sports Medicine, Keck Medicine for USC, Los Angeles, CA 90033, United States

Corresponding author: Joseph N Liu, MD, Assistant Professor, USC Epstein Family Center for Sports Medicine, Keck Medicine for USC, 1520 San Pablo St #2000, Los Angeles, CA 90033, United States. joseph.liu@med.usc.edu

Abstract

BACKGROUND

Many patients prioritize the ability to return to work (RTW) after shoulder replacement surgeries such as total shoulder arthroplasty (TSA), reverse TSA (rTSA), and shoulder hemiarthroplasty (HA). Due to satisfactory clinical and functional long-term outcomes, the number of shoulder replacements performed will continue to rise into this next decade. With younger individuals who compose a significant amount of the workforce receiving shoulder replacements, patients will begin to place a higher priority on their ability to RTW following shoulder arthroplasty.

AIM

To summarize RTW outcomes following TSA, rTSA, and HA, and analyze the effects of workers' compensation status on RTW rates and ability.

METHODS

This systematic review and analysis was performed in accordance with Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines. A literature search regarding RTW following shoulder arthroplasty was performed using four databases (PubMed, Scopus, Embase, and Cochrane Library), and the *Reference Citation Analysis* (<https://www.referencecitationanalysis.com/>). All studies in English relevant to shoulder arthroplasty and RTW through January 2021 that had a level of evidence I to IV were included. Nonclinical studies, literature reviews, case reports, and those not reporting on RTW after shoulder arthroplasty were excluded.

RESULTS

The majority of patients undergoing TSA, rTSA, or HA were able to RTW between

one to four months, depending on work demand stratification. While sedentary or light demand jobs generally have higher rates of RTW, moderate or heavy demand jobs tend to have poorer rates of return. The rates of RTW following TSA (71%-93%) were consistently higher than those reported for HA (69%-82%) and rTSA (56%-65%). Furthermore, workers' compensation status negatively influenced clinical outcomes following shoulder arthroplasty. Through a pooled means analysis, we proposed guidelines for the average time to RTW after TSA, rTSA, and HA. For TSA, rTSA, and HA, the average time to RTW regardless of work demand stratification was 1.93 ± 3.74 mo, 2.3 ± 2.4 mo, and 2.29 ± 3.66 mo, respectively.

CONCLUSION

The majority of patients are able to RTW following shoulder arthroplasty. Understanding outcomes for rates of RTW following shoulder arthroplasty would assist in managing expectations in clinical practice.

Key Words: Shoulder replacement; Total shoulder arthroplasty; Reverse total shoulder arthroplasty; Hemiarthroplasty; Return to work

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Many patients prioritize the ability to return to work after shoulder replacement surgeries such as total shoulder arthroplasty, reverse total shoulder arthroplasty, and shoulder hemiarthroplasty. While rates of return to work have been studied in the literature following shoulder arthroplasty, a consensus on which is the most effective treatment is still controversial. Information about the ability to return to work following any type of shoulder arthroplasty would assist patients and surgeons in managing expectations and put into place evidence-based guidelines. This systematic review examines how return to work following shoulder arthroplasty has been studied and reported in the literature.

Citation: Lalehzarian SP, Agarwalla A, Liu JN. Return to work following shoulder arthroplasty: A systematic review. *World J Orthop* 2022; 13(9): 837-852

URL: <https://www.wjgnet.com/2218-5836/full/v13/i9/837.htm>

DOI: <https://dx.doi.org/10.5312/wjo.v13.i9.837>

INTRODUCTION

Over the last two decades, the number of shoulder arthroplasties, including total shoulder arthroplasty (TSA), reverse TSA (rTSA), and shoulder hemiarthroplasty (HA), has increased at exponential rates[1-4]. TSA has typically been indicated for end-stage shoulder conditions in individuals with intact rotator cuff and sufficient glenoid bone stock to allow for stable glenoid component implantation[1-4]. The TSA procedure involves replacing the humeral head and glenoid with similarly shaped prosthetic components. rTSA, on the other hand, was historically indicated for patients with massive rotator cuff tears and involves using a convex glenoid hemispheric ball and a concave humerus articulating cup to reconstruct the glenohumeral joint. HA has traditionally been indicated in patients with glenohumeral arthritis where the glenoid bone stock is inadequate for TSA[1-4]. This procedure involves removing the humeral articular surface and replacing it with a stemmed humeral component.

Due to satisfactory clinical and functional long-term outcomes, the number of shoulder replacements performed will continue to rise into this next decade, with models predicting between 174810 and 350558 procedures by 2025[2,5,6]. Historically, shoulder replacements have been performed in elderly patients for degenerative shoulder conditions; however, these procedures are becoming more prevalent in younger and more active populations[5-8]. Furthermore, individuals born between 1981 and 1996 make up the largest generation of workers in the U.S. Labor Force[9]. With younger individuals who compose a significant amount of the workforce receiving shoulder replacements, patients will begin to place a higher priority on their ability to return to work (RTW) following shoulder arthroplasty.

Prior studies have shown varying levels of RTW after shoulder arthroplasty based on arthroplasty type, diagnosis, and work intensity[10-12]. While informative, a compilation comparing various demographics, arthroplasty types, diagnoses, and work intensities has not been performed in recent years. The purpose of this systematic literature review and analysis is to summarize outcomes of RTW following TSA, rTSA, and HA as well as analyze the effects of workers' compensation (WC) status on rates and ability to RTW.

MATERIALS AND METHODS

In accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines, a systematic review and analysis was performed[13,14]. The PubMed, Scopus, Embase, and Cochrane Library databases was queried using the search terms “shoulder arthroplasty”, “shoulder replacement”, “shoulder hemiarthroplasty”, or “humeral resurfacing” combined with “return to work”. The *Reference Citation Analysis* (<https://www.referencecitationanalysis.com/>) software was also used to identify any additional studies. The final search was performed on January 8, 2021. Additionally, the references of each study were manually assessed as well for potential inclusion in this investigation. The flow diagram summarizes the progression of the literature review with 12 total references meeting the inclusion criteria (Figure 1).

Clinical studies were evaluated and included if they were in English, had level of evidence I to IV, and reported on RTW after shoulder arthroplasty. Nonclinical studies, literature reviews, case reports, and those not reporting on RTW after shoulder arthroplasty were excluded. Title and abstract reviews were performed by two of the study authors (Lalehzarian SP and Liu JN). The full texts of articles meeting inclusion criteria based on title and abstract were then reviewed by two of the study authors (Lalehzarian SP and Agarwalla A) for final inclusion in the study. As referenced in Figure 1, 23 references were initially identified by the keyword search terms described above. After the title review, 8 references were excluded as 7 were irrelevant to the topic of discussion and 1 was a case report. One reference was excluded after abstract review as it was a review article and two references were excluded after full text review as they did not include RTW data. Following the review process, there were 12 references left and all were included in this review.

Included studies were evaluated using the Methodological Index for Non-Randomized Studies (MINORS) checklist[15]. Studies were evaluated on 8 items to 12 items, with each scored 0 (not reported), 1 (reported but poorly or inadequately done), or 2 (reported, well done and adequate), with a maximum score of 16 and 24 for noncomparative and comparative studies, respectively. Articles were scored by one of the study authors (Lalehzarian SP) and confirmed by two of the study authors (Agarwalla A and Liu JN). An analysis of the 12 total articles is shown in Tables 1-3.

RESULTS

RTW after TSA

TSA has shown to be a highly effective treatment for degenerative shoulder disease with adequate long-term outcomes, low revision rates, and high implant survivorship[7,16]. The number of anatomic total shoulder arthroplasties has increased from 29414 in 2011 to 40750 in 2017 partly due to an increased demand from younger populations and expanded indications[2,17-20]. With this increase in demand and volume, RTW following anatomic TSA is an important metric for many employed patients.

In a study by Bühlhoff *et al*[10], 57 TSA patients were analyzed after meeting inclusion criteria. At the most recent follow-up, 22 total patients (39%) returned to work. It is important to note that 6 patients (11%) cited their inability to pursue work at the time of most recent follow-up due to shoulder problems. While the authors concluded that approximately 61% of their patients did not retire or cease their vocation because of TSA, a large number of patients who were not working at final follow-up had retired from work[10]. This major limitation could be responsible for a low rate of RTW.

Liu *et al*[12] reported on 52 patients (54 shoulders), who were 55 years or younger at the time of surgery, worked in the 3 years leading up to surgery, and were available for a minimum follow-up of 2 years. Forty-eight patients (92%) were able to RTW postoperatively at an average of 2.1 mo after surgery. In addition to calculating the rate of RTW, the authors stratified patients by intensity of work: sedentary, light, moderate, or heavy. Forty one of 41 (100%) patients who had sedentary, light, or moderate work preoperatively were able to return to the same level of work. However, only 7 of 11 (64%) patients who had heavy-intensity work preoperatively were able to RTW. Of the 4 patients who did not RTW, only one patient cited shoulder pain and limited range of motion as the reason[12]. Additionally, the intensity of work was positively correlated with time to RTW. The authors found a statistically greater time to RTW when comparing heavy intensity (4.2 mo) to sedentary, light, and moderate intensity, respectively.

Cvetanovich *et al*[21] analyzed 27 shoulders (24 patients) that underwent anatomic TSA with an inlay glenoid component and stemless ovoid humeral head component. Twenty five (93%) of 27 patients were able to RTW with an average duration of 3.7 mo following surgery. Of the 2 patients who were not able to return, one patient cited reasons unrelated to the shoulder and the other patient cited back issues. When stratified by job intensity preoperatively, the rates of RTW were as follows: 5/5 for sedentary, 2/2 for light, 9/9 for moderate, and 9/11 for heavy. Furthermore, of the 25 patients who returned to work, 19 (76%) were able to return to their preoperative occupational demands. The 6 patients who returned to work at a lower intensity held heavy intensity occupations[21]. In addition to corroborating high rates of RTW for patients undergoing TSA, the authors found that patients with heavier demand jobs were less likely to RTW at the same occupational level postoperatively than patients in the other work

Table 1 Study characteristics

Ref.	Design	No. of groups	Level of evidence	Mean follow-up (range), yr	MINORS score
Bülhoff <i>et al</i> [10], 2015	Retrospective	1	IV	6.2 (2.6-12.6)	7/16
Jawa <i>et al</i> [56], 2015	Retrospective	2	III	3.9 (2.0-5.6)	17/24
Morris <i>et al</i> [55], 2015	Retrospective	2	III	3.5 (2-8)	20/24
Garcia <i>et al</i> [28], 2016	Retrospective	1	IV	2.6 (1-4.7)	10/16
Garcia <i>et al</i> [36], 2016	Retrospective	1	IV	5.1 (1-7.5)	10/16
Hurwit <i>et al</i> [11], 2017	Retrospective	2	III	HHA: 5.3 (1.1-7.5); rTSA ¹ : 2.7 (1.0-4.9)	16/24
Liu <i>et al</i> [12], 2018	Retrospective	1	IV	5.4 (2.5-8.6)	10/16
Kurowicki <i>et al</i> [41], 2018	Retrospective	2	III	2.4 (0.5-7.6)	17/24
Gowd <i>et al</i> [48], 2019	Retrospective	2	III	Hemi RR: 5.7 (SD \pm 2.0); aTSA: 5.8 (SD \pm 2.2)	17/24
Cvetanovich <i>et al</i> [21], 2020	Retrospective	1	IV	3.4 (1.9-5.0)	9/16
Jayasekara <i>et al</i> [22], 2020 ²	Retrospective	3	IV	NR	NA
Liu <i>et al</i> [49], 2020	Retrospective	2	III	HHA: 5.2 (2.0-7.5); aTSA: 5.18 (2.0-7.49)	16/24

¹Represents duplicate data from Garcia *et al*[28]; not included in meta-analysis.

²Numbers are relevant to groups who underwent total shoulder arthroplasty, reverse total shoulder arthroplasty, and hemiarthroplasty.

MINORS: Methodological Index for Non-Randomized Studies; HHA: Humeral hemiarthroplasty; rTSA: Reverse total shoulder arthroplasty; NR: Not reported; NA: Not available; Hemi RR: Hemiarthroplasty with ream-and-run resurfacing; aTSA: Anatomic total shoulder arthroplasty.

demand classes.

In a large clinical series by Jayasekara *et al*[22], 1773 patients were examined. TSA was one of the twelve surgeries analyzed with a total number of 38 patients. At the six month follow-up, 27 (71%) patients were able to return to some type of work: 14 (37%) patients returned with full duty, 13 (34%) patients returned with lighter duty, and 11 (29%) patients were unable to RTW. Of the twelve surgeries analyzed, TSA at 71% was shown to have a lower rate of RTW compared to surgeries such as HA and rTSA which had 82% and 56%, respectively[22]. This lower rate of RTW may have been due to a higher average age of patients who underwent TSA compared to those in previous studies; therefore, the age of the patients may have negatively influenced their desire and ability to RTW[22].

In summary, the majority of studies cited a rate of RTW between 71% and 93% with an average duration of 1 mo to 4 mo following TSA[12,21]. Furthermore, most patients who undergo TSA are able to RTW at the same preoperative intensity level with the exception of those patients in heavy intensity jobs who are less likely to RTW after TSA.

RTW after rTSA

In 2003, the United States Food and Drug Administration approved the use of rTSA for rotator cuff arthropathy[1,23]. Since that time, the volume of rTSA has drastically increased, from 21916 in 2011 to 63845 in 2017, in part due to its encouraging results and expanded indications to cover proximal humerus fracture and previous failures of arthroplasty[24-27]. When comparing the number of rTSA to the total number of shoulder replacements from 2011 and 2017, the percentage has increased from 33% to 58%[2]. Due to the exponential increase in rTSA use, a clinical review outlining the rate of RTW after rTSA will assist orthopedic surgeons in treating future patients with shoulder conditions.

Garcia *et al*[28] conducted a study on 40 patients who had undergone rTSA. Of the 40 patients analyzed, 26 (65%) of them were able to RTW with an average time of 2.3 mo. From the 14 patients who did not RTW, only two of them retired due to shoulder reasons while the other 12 retired due to nonorthopedic causes. When stratified into intensity level, rates were comparable to the overall rate of RTW with 17 (68%) of 25 patients returning in the sedentary class and 9 (60%) of 15 returning in the light class. Additionally, patients with sedentary jobs returned to work more quickly than those with light work (1.4 mo *vs* 4.0 mo).

Jayasekara *et al*[22] evaluated 34 rTSA patients, with 19 (56%) of them able to return to some type of work at the 6 mo follow-up. Eight of the 19 patients who returned to work were able to RTW with full duties and the other eleven returned to work with lighter duties. From the twelve surgeries included in the study, rTSA was associated with the lowest rate of RTW at 56%. Jayasekara *et al*[22] concluded that this percentage is consistent with prior studies which cited a 65% rate of RTW[11,28].

Table 2 Characteristics of the patients

Ref.	No. of shoulders	Mean age (range), yr	Gender (M/F), n	Dominant/nondominant, n	BMI, kg/m ² (range)	WC/NWC	RTW (%)	Work intensity
Bülhoff <i>et al</i> [10], 2015	154	72 (33-88)	35/119	103/51	NR	NR	22/57 (38.6) ¹	NR
Jawa <i>et al</i> [56], 2015	13	55.9 (39-74)	13/0	NR	NR	13/0	4/13 (30.8)	1 light, 12 heavy
Morris <i>et al</i> [55], 2015	28	WC: 58.8 (49-69); NWC: 63.4 (50-72)	20/8	19/9	WC: 32.0 (SD ± 8.4); NWC: 27.1 (SD ± 5.3)	14/14	WC: 2/14 (14.3); NWC: 5/11 (45.5) ¹	WC: 8 sedentary/light, 6 heavy/strenuous; NWC: 3 retired, 7 sedentary/light, 4 heavy/strenuous
Garcia <i>et al</i> [28], 2016 (rTSA)	40	74.7 (56-82)	16/24	26/14	28.8 (14.8-46.2)	0/40	26/40 (65)	25 sedentary, 15 light
Garcia <i>et al</i> [36], 2016 (HHA)	79	69 (27.6-97.1)	24/55	62/17	28.3 (19.8-49.3)	0/79	34/49 (69.4) ¹	20 sedentary, 25 light, 4 moderate
Hurwit <i>et al</i> [11], 2017	81	HHA: 60.8 (40-88); rTSA: 68.6 (41-48)	33/48	52/29	HHA: 28.9; rTSA: 29.5	NR	55/81 (84.6)	44 sedentary, 33 light, 4 heavy
Liu <i>et al</i> [12], 2018	52	67.2 (56-96)	38/14	24/28	28.0 (18.1-52.9)	5/47	48/52 (92)	10 sedentary, 14 light, 17 moderate, 11 heavy
Kurowicki <i>et al</i> [41], 2018 ²	265	aTSA: 69; rTSA: 75	NR	NR	NR	NR	21% higher difference in ability to RTW following aTSA than rTSA	115 retired, 72 housework, 49 desk job, 16 prolonged standing, 11 yard work, 9 creative jobs, 5 requires lifting, 4 carpenter/construction, 5 cook/food prep
Gowd <i>et al</i> [48], 2019	53	Hemi RR: 52.8 ± 7.7; aTSA: 53.3 ± 9.2	48/5	28/25	Hemi RR: 28.5 ± 3.5; aTSA: 31.1 ± 5.7	4/49	50/53 (94.3)	17 sedentary, 12 light, 13 moderate, 11 heavy
Cvetanovich <i>et al</i> [21], 2020	27	52.1 ± 6 (42-63)	25/2	NR	NR	3/24	25/27 (92.6)	5 sedentary, 2 light, 9 moderate, 11 heavy
Jayasekara <i>et al</i> [22], 2020 ³	83	TSA: 65 ± 1.6 (48-86); rTSA: 72 ± 1.6 (54-91); Hemi: 72 ± 2.7 (57-84)	42/41	NR	NR	3/83	55/83 (66.3)	28 full duty, 27 lighter duty
Liu <i>et al</i> [49], 2020	49	HHA: 62.4 (42.7-87.7); aTSA: 61.7 (47.7-75.6)	22/27	30/19	HHA: 29.8 ± 7.1; aTSA: 29.2 ± 6.5	NR	36/49 (73.5)	20 sedentary, 21 light, 6 heavy

¹Excluding those who were retired preoperatively, retired due to medical concerns, or retired due to non-specified reasons.

²Only includes individuals who responded to question 10 of the ASES questionnaire in regards to work.

³Numbers are relevant to groups who underwent total shoulder arthroplasty, reverse total shoulder arthroplasty, and hemiarthroplasty.

M: Male; F: Female; BMI: Body mass index; WC: Workers' compensation; NWC: Non-workers' compensation; RTW: Return to work; NR: Not reported; aTSA: Anatomic total shoulder arthroplasty; rTSA: Reverse total shoulder arthroplasty; HHA: Humeral hemiarthroplasty.

Table 3 Diagnosis and surgical characteristics

Ref.	Diagnosis	Surgery	Mean time out of work (range), mo	Complications
Bühlhoff <i>et al</i> [10], 2015	Primary OA, 154 (100%)	aTSA	NR	NR
Jawa <i>et al</i> [56], 2015	OA, 11 (84.6%); capsulor-rhaphy arthropathy, 2 (15.4%)	aTSA	4.2 (2.9-6.0)	NR
Morris <i>et al</i> [55], 2015	CTA, 14; massive RCT, 8; post-traumatic malunion, 4; failed prior arthroplasty, 2	rTSA	NR	WC (4): postoperative anterior dislocation (2), intraoperative humeral shaft fracture, postoperative periprosthetic infection; NWC (2): postoperative anterior dislocation
Garcia <i>et al</i> [28], 2016 (rTSA)	CTA, 21 (53.5%); OA, 10 (25%); PHFx, 7 (17.5%); RA, 2 (5%)	rTSA	2.3 (0.5-11)	NR
Garcia <i>et al</i> [36], 2016 (HHA)	OA, 40 (50.6%); PHFx, 17 (21.5%); AVN, 11 (13.9%); CTA, 8 (10.1%); RA, 3 (3.8%)	HHA	1.4 (0.25-24)	8 complications: 4 revision HHA (2 for dislocation, 2 for periprosthetic fracture after fall); 3 HHA revised to TSA; 1 HHA revised to rTSA for continued pain/glenoid wear
Hurwit <i>et al</i> [11], 2017	CTA, 63 (77.8%); RA, 14 (17.2%); PHFx, 2 (2.5%)	rTSA; HHA	rTSA: 3.1; HHA: 2.3	rTSA: 20 chronic pain and stiffness/limited mobility; 1 returned to OR; HHA: 4 chronic pain and stiffness/limited mobility; 5 returned to OR
Liu <i>et al</i> [12], 2018	OA, 42 (81%); failed prior arthroplasty, 7 (13%); AVN, 2 (4%); RA, 1 (2%)	aTSA	2.1 (SD: 1.7)	22 complications: 17 postoperative stiffness, 6 chronic pain, 3 instability, 4 returned to OR
Kurowicki <i>et al</i> [41], 2018	NR	aTSA; rTSA	NR	NR
Gowd <i>et al</i> [48], 2019	End-stage glenohumeral OA, 53 (100%)	Hemi RR; aTSA	Hemi RR: 2.5 ± 4.8; aTSA: 1.98 ± 2.6	Hemi RR: 3 chronic pain, 1 felt unstable, 5 postoperative stiffness, 1 nagging soreness, 1 acute pain, 2 conversion to aTSA, 1 received arthroscopic debridement; aTSA: 1 chronic pain, 2 weakness, 6 postoperative stiffness, 1 subscapularis repair, 1 revision with glenoid explantation due to loosening
Cvetanovich <i>et al</i> [21], 2020	Glenohumeral OA, 23 (85.1%); post-traumatic OA, 4 (14.9%)	aTSA	3.7 ± 5.2	1 hematoma, 1 pulmonary embolism
Jayasekara <i>et al</i> [22], 2020 ¹	NR	aTSA; rTSA; HHA	NR	NR
Liu <i>et al</i> [49], 2020	End-stage glenohumeral OA, 49 (100%)	HHA; aTSA	HHA: 1.9 ± 2.3; aTSA: 1.3 ± 1.0	HHA: 15 chronic pain, 8 postoperative stiffness, 2 conversion to aTSA, 2 conversion to rTSA; aTSA: 3 postoperative stiffness

¹Numbers are relevant to groups who underwent total shoulder arthroplasty, reverse total shoulder arthroplasty, and hemiarthroplasty.

OA: Osteoarthritis; CTA: Cuff tear arthropathy; RCT: Rotator cuff tear; PHFx: Proximal humerus fracture; RA: Rheumatoid arthritis; AVN: Avascular necrosis; aTSA: Anatomic total shoulder arthroplasty; rTSA: Reverse total shoulder arthroplasty; HHA: Humeral hemiarthroplasty; TSA: Total shoulder arthroplasty; NR: Not reported; WC: Workers' compensation; NWC: Non-workers compensation; OR: Operating room.

The available data suggests that the majority of patients who undergo rTSA are able to RTW at rates between 56% and 65%. Despite this low percentage, the volume of rTSA continues to rise due to expanding indications[2].

RTW after HA

Traditionally, HA was considered a safer option compared to TSA or rTSA for patients who wished to remain active following surgery due to its low failure rate and utilization of an intact glenoid[29]. Despite exponential rises in TSA and RTSA, the rate of HA procedures has steadily declined from 15860 in 2011 to 6150 in 2017[2]. This is in part due to the increase in rTSA for shoulder replacement. Since the indications for rTSA have been expanded to include fractures, the rate of HA for fracture use has decreased by nearly 30%[30-32]. Additionally, recent studies have shown that clinical outcomes from HA are significantly inferior to that of TSA and that patients undergoing HA had statistically significantly worse functional scores[11,33-35]. With this steady decline over the last decade, there is much necessity for a clinical review that examines all available literature regarding the rates of RTW for HA.

Garcia *et al* [36] examined 49 patients who worked preoperatively and underwent HA. Thirty-four (69.4%) patients were able to return to previous employment at an average duration of 1.4 mo. Preoper-




Figure 1 Flow diagram illustrating systematic literature review process.

atively, 20 (41%) patients classified their jobs as sedentary, 25 (51%) patients as light physical work, and 4 (8%) patients as moderate physical work. Following HA, 15 of 20 (75%) patients returned as sedentary, 17 of 25 (68%) patients as light physical work, and 2 of 4 (50%) patients as moderate physical work. While no patients changed job demand level postoperatively, the average time to return to employment varied: 1.9 mo for sedentary, 2.6 mo for light, and 13.1 mo for moderate. As one of the first studies to analyze the rates of RTW following HA, Garcia *et al*[36] was able to quantify evidence that aided physicians in managing expectations of patients undergoing shoulder HA.

Jayasekara *et al*[22] included 11 patients who underwent shoulder HA. Nine (82%) patients were able to return to some type of work at 6 mo follow-up with 6 (55%) patients able to return to full duties, 3 (27%) able to return to lighter duties, and 2 (18%) unable to RTW. While the reason for not returning to work was not cited, it may be due to the fact that the average age of patients undergoing HA in this cohort was 72 years of age. Although a limitation of this study was a smaller size, Jayasekara *et al*[22] found higher rates of RTW despite an average age much higher than previous studies[11,36].

Recent literature has shown rates of RTW for shoulder HA between 69% and 82% compared to both TSA and rTSA[22,36]. Despite higher rates of RTW for HA compared to rTSA, the number of HA cases continues to decline with poor functional outcomes at long-term follow-up[11,37].

Comparison of RTW between TSA and rTSA

In patients with end-stage glenohumeral arthritis and an intact rotator cuff, TSA has shown to be a highly effective treatment with high rates of functional recovery[5]. While the original indication for rTSA was rotator cuff arthropathy, the indications for rTSA have expanded to include conditions such as TSA and HA implant failures, complex proximal humerus fractures, asymmetric glenoid wear, posterior humeral head subluxation in patients with intact rotator cuffs, and irreparable rotator cuff tears in the absence of arthritis[38-40]. Similarly, the indications for TSA have also expanded to now include a more diverse and active patient population[41]. As younger patients undergo shoulder replacements, many patients cite their ability to work as instrumental in their decision to have surgery. With increased indications for both surgeries, assessing the ability of patients to RTW following TSA and rTSA is imperative to educate future patients and manage expectations.

In one recent study, Kurowicki *et al*[41], evaluated 159 patients undergoing TSA (average age 69) and 106 patients undergoing rTSA (average age 75). Authors used the American Shoulder and Elbow Surgeons (ASES) Assessment Form as a way to track patients' ability to RTW. Among usually reported work, it is important to note that 43% of patients cited retirement as their work, with housework (27%) and desk jobs (18%) as the second and third most cited, respectively. Kurowicki *et al*[41] reported a 21% higher difference in overall ability to work for patients following TSA compared to those patients who underwent rTSA. In particular, statistically significant differences were found between TSA and rTSA amongst patients who cited their work as housework or gardening.

Kurowicki *et al*[41] is the only study that compares the ability of TSA patients to RTW to rTSA patients. Based on this study, authors concluded that returning to work after TSA is more favorable than rTSA in fields of work that require low-demand activities such as housework and gardening[41]. This

study was limited by its reporting bias from survey-based studies, small sample size within work subgroups, and population representation differences particularly in age. Regardless, comparisons among these groups hold importance in defining patient and surgeon expectations after surgery.

Comparison of RTW between TSA and HA

If non-operative treatment for glenohumeral osteoarthritis with intact rotator cuff integrity fails, patients are often told to consider HA or TSA. While the optimal surgical treatment remains controversial, there are benefits to both procedures. Multiple studies have shown that patients with glenohumeral arthritis who undergo TSA have improved pain relief, higher functional scores, and more range of motion compared to those who undergo HA[35,36,42-44]. However, TSA also has an increased operative time, more blood loss, more technical difficulty, and incurs the risk of glenoid loosening[45]. On the other hand, while HA has the benefits of decreased operative time, decreased blood loss, and less technical difficulty, there is some concern regarding the progression of arthritic changes especially with bone loss and the need for future revision surgeries such as conversion to TSA[45,46]. Furthermore, many patients have lifting restrictions after TSA, which may limit their ability to RTW[47].

Gowd *et al*[48] analyzed 53 total patients with glenohumeral arthritis. Twenty five patients (average age of 52.8 years) received HA with ream-and-run resurfacing and 28 patients (average age of 53.3 years) received TSA. Of the 25 patients undergoing HA, all 25 (100%) were able to RTW at an average duration of 1.98 mo. On the other hand, 25 (89%) of 28 patients receiving TSA were able to RTW with an average time of 2.5 mo following surgery. When HA patients were stratified preoperatively into work demand level, 7 patients were categorized as sedentary, 7 were light, 4 were moderate, and 7 were heavy. For TSA, 10 patients were categorized as sedentary, 5 were light, 9 were moderate, and 4 were heavy. Postoperatively, all HA patients (100%) in sedentary, light, and moderate were able to RTW. For TSA, 9 (90%) of 10 returned to sedentary work, while all (100%) light and moderate duty patients returned to work. For the heavy category, 7 (100%) of 7 HA patients were able to return compared to 2 (50%) of 4 TSA patients demonstrating that heavy duty workers undergoing HA had a significantly higher rate of RTW[48]. Of the 2 TSA heavy duty patients who were unable to RTW, only one reported permanent restriction with overhead lifting. Despite this difference, authors concluded near equivalent rates of RTW between HA and TSA.

Liu *et al*[49] evaluated 49 total patients with end-stage glenohumeral osteoarthritis. Twenty-six patients underwent HA (average age of 62.4 years) and 23 patients underwent TSA (average age 61.7). Sixteen (62%) of 26 HA patients were able to RTW at an average duration of 1.88 mo following surgery. Of the patients undergoing TSA, 20 (87%) were able to RTW at an average time of 1.29 mo following surgery. From the 10 HA patients who did not RTW, only one had retired postoperatively due to shoulder issues. The other nine patients retired preoperatively due to the shoulder, other medical reasons, or postoperatively due to non-specified reasons. Of the three TSA patients who did not RTW, zero had retired postoperatively due to the shoulder. Patients either retired preoperatively due to the shoulder, other medical concerns, or non-specified reasons. For patients who underwent TSA, 7 (100%) of 7 returned to a sedentary work demand level, 9 (82%) of 11 returned to a light work demand level, and 3 (100%) of 3 returned to a heavy work demand level (Table 4). For patients who underwent HA, 8 (62%) of 13 returned to a sedentary work demand level, 7 (70%) of 10 returned to a light work demand level, and 1 (33%) of 3 returned to a heavy work demand level (Table 4). (68%) of 25 returned to a sedentary work level and 9 (60%) of 15 returned to a light work demand level (Table 4). Liu *et al*[49] concluded that patients with osteoarthritis undergoing TSA have higher rates of RTW and function compared to those undergoing HA.

From these two studies, there is still a discrepancy in terms of ability to RTW between HA and TSA. The mixed results could potentially be due to the limitations of each study. For example, in Gowd *et al* [48], surgeons counseled their TSA patients that they would have permanent overhead lifting restrictions, whereas those who underwent HA would not receive these restrictions. Comparatively, in Liu *et al*[49], surgeons placed no postoperative work restrictions on either group. Furthermore, the average age of individuals in Gowd *et al*[48] (52.8 and 53.3 years of age) was significantly lower than the average of individuals in Liu *et al*[49] (62.4 and 61.7 years of age) possibly indicating that older patients either hold more sedentary, less demanding occupations or may benefit more in their ability to RTW following TSA compared to HA[48,49].

Comparison of RTW between HA and rTSA

When TSA is contraindicated, in cases such as rotator cuff or deltoid dysfunction, deficiencies in glenoid bone stock, or proximal humerus fractures, patients must be educated on the benefits and drawbacks of HA *vs* rTSA[50]. Many studies over the last decade have shown more predictable and superior outcomes for rTSA compared to HA[51-53]. Yet, in the younger population, especially those who want to remain employed following surgery, surgeons often feel more comfortable recommending HA given the theoretical risk of glenoid component loosening or failure in rTSA[54]. Furthermore, surgeons tend to place more activity restrictions on patients who undergo rTSA, which could significantly limit their ability to RTW.

Table 4 Return to work after total shoulder arthroplasty vs hemiarthroplasty[49]

Occupation intensity	RTW after TSA (%)	RTW after HA (%)
Sedentary	7/7 (100)	8/13 (62)
Light	9/11 (82)	7/10 (70)
Heavy	3/3 (100)	1/3 (33)
Total	20/23 (87)	16/26 (62)

RTW: Return to work; TSA: Total shoulder arthroplasty; HA: Hemiarthroplasty.

Hurwit *et al*[11] compared 40 rTSA patients (average age of 68.6 years) to 41 HA patients (average age of 60.8 years) all of whom had end-stage glenohumeral arthritis with rotator cuff dysfunction, deficiencies in glenoid bone stock that prohibited the insertion of an anatomic glenoid component, or proximal humerus fracture. Of the patients who underwent rTSA, 26 (65%) of them were able to RTW at an average duration of 2.3 mo following surgery. Only two patients who were unable to RTW cited their main reason as issues with the shoulder following surgery, while the other twelve either retired preoperatively due to medical reasons or non-specified reasons. Twenty-nine (71%) of the 41 HA patients were able to RTW at an average time of 3.1 mo after surgery. In this cohort, only one patient retired postoperatively due to shoulder issues. The other eleven had retired preoperatively due to the shoulder, medical reasons, or non-specified reasons. For patients who underwent HA, 14 (74%) of 19 were able to return to a sedentary work demand level, 13 (72%) of 18 returned to a light demand level, and 2 (50%) of 4 returned to work at a heavy work level (Table 5). For patients who underwent rTSA, 17 (68%) of 25 returned to a sedentary work level and 9 (60%) of 15 returned to a light work demand level (Table 5). Hurwit *et al*[11] concluded no significant difference between the two groups in terms of return to low- and moderate-intensity work, despite an older age for patients undergoing rTSA.

Despite a higher rate of RTW for HA patients, no significant differences were found by Hurwit *et al* [11]. A potential limitation with this study was the no significant difference in average age of each cohort (68.6 years for rTSA patients and 60.8 years for HA patients), even though this did not affect RTW rates [11]. Furthermore, this study only had sufficient sample sizes for sedentary and light duty workers. Due to the lack of heavy duty workers, especially in rTSA, it is possible to hypothesize that heavy laborers may have experienced more difficulty in returning to work.

Comparison of RTW between WC and non-WC

Work-related injuries are a common cause of disability in the United States and have significant implications for workers, employers, insurers, and physicians[18,55,56]. WC status has shown to have a detrimental effect on clinical outcomes following orthopedic surgery[57,58]. The impact of WC status on postoperative outcomes is an important consideration for patients undergoing shoulder arthroplasty.

Morris *et al*[55] compared 14 WC patients who underwent rTSA to a matched cohort of 14 patients without WC status who also underwent rTSA. From the patients with WC claims, only 2 (14%) of 14 were able to RTW. Of the 12 patients who were not able to RTW, one was unemployed and seeking employment at the time of follow-up, five were disabled, and six had retired following rTSA. In the matched cohort of non-WC patients, only 11 patients had worked prior to the surgery. From these 11 patients, 5 (46%) were able to RTW, one was disabled, and five had retired after rTSA. No patients, WC or non-WC, were able to return to heavy/strenuous work demands after rTSA. Despite significant improvement from preoperative to final follow-up outcomes, WC patients had significantly worse Constant scores, ASES scores, Western Ontario Osteoarthritis of the Shoulder Index scores, and less external rotation compared with the matched cohort group. Morris *et al*[55] reported that while WC patients had significant improvements following rTSA, they achieved significantly worse outcomes compared to non-WC patients after rTSA.

In Jawa *et al*[56], a cohort of 13 WC patients (average age of 55.9 years) who underwent TSA were compared to a control group of 63 patients (average age of 63.2 years) who also underwent TSA. While RTW rates were not cited for the control group, only 4 (31%) of the 13 WC patients were able to RTW following TSA. Of the four patients who returned, one returned to the same job with lifting restrictions and the other three changed jobs to those that require less lifting. From the nine patients who did not return, 7 did not return due to functional restrictions after the surgery and 2 had retired. Additionally, Jawa *et al*[56] found the ASES score to be significantly lower in the WC cohort compared to the control group. From this study, authors concluded no WC patients were able to return to full duty work at their current job and that WC patients receiving TSA had poorer outcomes compared to non-WC patients.

Despite the lack of difference in RTW rates following shoulder arthroplasty for patients with or without WC claims, many studies in orthopedic literature have found poorer outcomes, lower satisfaction rates, and more pain in patients with WC status after shoulder arthroplasty[55,56,59,60]. Similar findings exist in the shoulder literature outside of shoulder arthroplasty. For example, in

Table 5 Return to work after hemiarthroplasty vs reverse total shoulder arthroplasty[11]

Occupation intensity	RTW after HA (%)	RTW after rTSA (%)
Sedentary	14/19 (74)	17/25 (68)
Light	13/18 (72)	9/15 (60)
Heavy ¹	2/4 (50)	-
Total	29/41 (71)	26/40 (65)

¹No reverse total shoulder arthroplasty patients were classified into the heavy work category.

RTW: Return to work; rTSA: Reverse total shoulder arthroplasty; HA: Hemiarthroplasty.

numerous rotator cuff studies, patients with WC status have been found to be significantly less compliant with postoperative protocols and have less improvement in functional outcomes and pain after controlling for confounding factors such as age, marital status, education level, preoperative expectations, work demands, smoking, comorbidities, duration of symptoms, size of tear, and repair technique[61-64]. Furthermore, other similar results have been found in WC cohorts undergoing acromioplasty, superior labral anterior-posterior tear, and biceps tenodesis for failed superior labral anterior-posterior repair[65-68]. Regardless of procedure type, the differences in pain and outcomes persist, suggesting that WC status may play a crucial role in inferior outcomes.

DISCUSSION

RTW guidelines

While the decision to RTW depends on a variety of factors, all physicians have the goal of returning patients to maximal function in the shortest period of time with the least residual disability[69,70]. Based on the available literature, guidelines can be proposed for average time to RTW for each work demand level within each type of shoulder arthroplasty (Table 6). We determined these averages through a pooled analysis[71].

Throughout the rehabilitation process, physicians must assess patients, especially those with WC status, in terms of work restrictions and limitations. Given the little published evidence for guidelines regarding physical restrictions after shoulder arthroplasty, the work restrictions are commonly based on the physician's clinical judgment[69]. On the other hand, work limitations are easier to define as they are based on the patient's ability to perform a certain task[69].

Particularly for WC patients who undergo shoulder replacement and rehabilitation and have still failed to RTW at their desired work demand level, work conditioning or work hardening therapy regimens can be prescribed[69,72]. Work conditioning, a task simulation program lasting two to four hours per day for three to five days per week, is meant to develop a patient's ability to tolerate specific tasks they would typically encounter at work. Work hardening has the same goal in mind with a higher intensity lasting up to eight hours per day for five days per week[69].

When recovery from shoulder arthroplasty has reached a therapeutic plateau for either non-WC or WC patients, a physician must rate the residual permanent impairment and individually assess how long each injured patient should remain on this plateau before considering them at maximum medical improvement (MMI)[69]. MMI is established when no further treatment will significantly change the patient's outcome; at this point, a patient can be recovered completely without any residual impairment or have some permanent impairment[69,73]. Specifically for TSA, Cabarcas *et al*[74] established MMI at twelve months postoperatively. While Puzzitiello *et al*[75] established MMI for rTSA at twelve months following surgery, Matar *et al*[76] found patients undergoing rTSA may reach MMI as early as six months after surgery. If a patient has reached MMI, but has failed to achieve their pre-injury or prior level of work status, then a physician can utilize a functional capacity evaluation to determine the patient's ability and impose final work restrictions[69]. Although the results of the FCE are often used to set work limitations, some studies have questioned its utility as FCE does not take biopsychosocial factors into account and possibly measures a patient's tolerance to an activity as opposed to the patient's true ability[69,77,78].

After MMI has been reached, there are two outcomes: (1) The patient is able to RTW with or without permanent restrictions at the same job; or (2) The patient finds a new job because the employer cannot accommodate the patient's work limitations[69]. Using evidence-based guidelines to determine MMI for TSA, rTSA, and HA is important not only for counseling patients, but also modifying their expectations prior to surgery.

Table 6 Time to return to work (mo)

	Intensity ¹				
	Sedentary	Light	Moderate	Heavy	Overall
TSA					
Gowd <i>et al</i> [48], 2019	2.1 ± 3.8	1.3 ± 1.2	2.1 ± 2.0	3.0 ± 2.8	2.0 ± 2.6
Liu <i>et al</i> [12], 2018	1.3 ± 1.2	1.6 ± 1.3	2 ± 1.7	4.2 ± 2.0	2.1 ± 1.7
Liu <i>et al</i> [49], 2020	1.04 ± 0.87	1.06 ± 0.73	-	1.83 ± 1.04	1.29 ± 0.96
Cvetanovich <i>et al</i> [21], 2020	-	-	-	-	3.7 ± 5.2
Average ²	1.19 ± 1.24	1.25 ± 0.99	2.03 ± 1.79	2.96 ± 3.23	1.93 ± 3.74
rTSA					
Garcia <i>et al</i> [28], 2016/ Hurwit <i>et al</i> [11], 2017 ³	1.38 ± 0.93	4 ± 3.4	-	-	2.3 ± 2.4
HA					
Hurwit <i>et al</i> [11], 2017/Garcia <i>et al</i> [24], 2015 ⁴	1.96 ± 3.0	2.72 ± 2.6	-	13.13 ± 15.4	3.1 ± 4.9
Gowd <i>et al</i> [48], 2019	0.9 ± 1.1	1.0 ± 1.7	6.8 ± 11.5	3.1 ± 2.3	2.5 ± 4.8
Liu <i>et al</i> [49], 2020	1.06 ± 0.98	2.76 ± 3.27	-	2.25 ⁵	1.88 ± 2.34
Average ²	1.09 ± 1.36	2.00 ± 3.36	6.8 ± 11.5	3.16 ± 2.74	2.29 ± 3.66

¹Intensity as based on US Department of Labor[56].

²Pooled means using meta analysis[57].

³Both studies used the same reverse total shoulder arthroplasty population.

⁴Both studies used the same hemiarthroplasty population.

⁵Only one patient in the heavy group, so no standard deviation available.

TSA: Total shoulder arthroplasty; rTSA: Reverse total shoulder arthroplasty; HA: Hemiarthroplasty.

Limitations and future research directions

Our narrative systematic review and analysis has several limitations. First, identification and inclusion of references utilized for this review relied on the previously described search strategy in 4 different databases. We searched 4 different databases in order to limit the possibility of overlooking studies related to shoulder arthroplasty and RTW. Second, our data relied on the data reported in the included studies. Therefore, we are limited by the clarity of the results reported as well as the study design and level of evidence. As a result, we utilized the MINORS score to evaluate the quality of the 12 included studies and any potential publication bias. We found that the 12 studies were of acceptable quality and determined no findings suggestive of publication bias. Additionally, our data shows a high level of heterogeneity which may lead to treatment bias effect. Similarly, with regard to work intensity, our study is limited by what was reported and those studies may exclude important nuances that could have led to functional consequences. Furthermore, the heterogeneity of our data is reflective of the reality of clinical practice and often most accurately represents what orthopedic surgeons encounter in the clinical setting[79-81]. Despite these limitations, the findings in our study provide important data that help orthopedic surgeons manage patient expectations about RTW following TSA, rTSA, or HA.

In the future, systematic reviews and analyses regarding shoulder arthroplasty and RTW will hopefully have access to references that are more homogenous with higher levels of evidence. Although the reality that a high level of heterogeneity may be inevitable in the clinical research setting, additional research should be conducted that compares short- and long-term outcomes following TSA, rTSA, and HA and a patients' ability to RTW. Furthermore, revision arthroplasty and ability to RTW may be a topic worth exploring as the average age of patients undergoing shoulder replacement is decreasing.

CONCLUSION

The majority of patients are able to RTW following TSA, rTSA, and shoulder HA. The rates of RTW following TSA (71%-93%) seem to be consistently higher than those reported for HA (69%-82%) and rTSA (56%-65%), although this may reflect demographic differences such as age in patient populations. Sedentary, light demand jobs generally have higher rates of RTW than moderate or heavy demand jobs. On average, most patients who underwent TSA, rTSA, or HA were able to RTW at an average duration between 1 mo to 4 mo depending on work demand level. Furthermore, WC status negatively influenced clinical outcomes following shoulder arthroplasty.

ARTICLE HIGHLIGHTS

Research background

Over the last two decades, the number of shoulder arthroplasties, including total shoulder arthroplasty (TSA), reverse TSA (rTSA), and shoulder hemiarthroplasty (HA), has increased at exponential rates. Due to satisfactory clinical and functional long-term outcomes, the number of shoulder replacements performed will continue to rise into this next decade. Additionally, these procedures are becoming more prevalent in younger and more active populations. With younger individuals who compose a significant amount of the workforce receiving shoulder replacements, patients will begin to place a higher priority on their ability to return to work following shoulder arthroplasty.

Research motivation

Prior studies have shown varying levels of return to work after shoulder arthroplasty based on arthroplasty type, diagnosis, and work intensity. While informative, a compilation comparing various demographics, arthroplasty types, diagnoses, and work intensities has not been performed in recent years.

Research objectives

The aim of the review article was to summarize return to work outcomes following TSA, rTSA, and HA, and analyze the effects of workers' compensation status on return to work rates and ability.

Research methods

This systematic review and analysis was performed in accordance with Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines. A literature search regarding return to work following shoulder arthroplasty was performed using four databases through January 2021. All studies included in this review were analyzed by at least two authors. Included studies were then evaluated using the Methodological Index for Non-Randomized Studies checklist.

Research results

The majority of patients undergoing TSA, rTSA, or HA were able to return to work between one to four months, depending on work demand stratification. While sedentary or light demand jobs generally have higher rates of return to work, moderate or heavy demand jobs tend to have poorer rates of return. Furthermore, workers' compensation status negatively influenced clinical outcomes following shoulder arthroplasty. Through a pooled means analysis, we proposed guidelines for the average time to return to work following TSA, rTSA, and HA.

Research conclusions

The majority of patients were able to return to work following TSA, rTSA, or HA. Understanding outcomes for rates of return to work following shoulder arthroplasty should assist surgeons and patients in managing expectations in clinical practice.

Research perspectives

Further research and analyses comparing short- and long-term outcomes following TSA, rTSA, and HA and a patients' ability to return to work would provide tremendous benefit. Additionally, revision arthroplasty and ability to return to work may be a topic worth exploring as the average age of patients undergoing shoulder replacement is decreasing.

FOOTNOTES

Author contributions: All authors made significant contributions toward the preparation of this manuscript. Lalehzarian SP wrote the article, critically revised the article, and participated in the final approval of the version to be published; Agarwalla A critically revised the article and participated in the final approval of the version to be published; Liu JN designed the work, critically revised the article, and was responsible for final approval of the version to be published.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

PRISMA 2009 Checklist statement: The authors have read the PRISMA 2009 Checklist, and the manuscript was prepared and revised according to the PRISMA 2009 Checklist.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license

their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: United States

ORCID number: Simon P Lalehzarian 0000-0003-3406-3651; Avinesh Agarwalla 0000-0001-5056-6780; Joseph N Liu 0000-0002-3801-8885.

S-Editor: Gao CC

L-Editor: A

P-Editor: Gao CC

REFERENCES

- 1 **Kim SH**, Wise BL, Zhang Y, Szabo RM. Increasing incidence of shoulder arthroplasty in the United States. *J Bone Joint Surg Am* 2011; **93**: 2249-2254 [PMID: 22258770 DOI: 10.2106/JBJS.I.01994]
- 2 **Wagner ER**, Farley KX, Higgins I, Wilson JM, Daly CA, Gottschalk MB. The incidence of shoulder arthroplasty: rise and future projections compared with hip and knee arthroplasty. *J Shoulder Elbow Surg* 2020; **29**: 2601-2609 [PMID: 33190759 DOI: 10.1016/j.jse.2020.03.049]
- 3 **Jain NB**, Yamaguchi K. The contribution of reverse shoulder arthroplasty to utilization of primary shoulder arthroplasty. *J Shoulder Elbow Surg* 2014; **23**: 1905-1912 [PMID: 25304043 DOI: 10.1016/j.jse.2014.06.055]
- 4 **Schairer WW**, Nwachukwu BU, Lyman S, Craig EV, Gulotta LV. National utilization of reverse total shoulder arthroplasty in the United States. *J Shoulder Elbow Surg* 2015; **24**: 91-97 [PMID: 25440519 DOI: 10.1016/j.jse.2014.08.026]
- 5 **Fehring EV**, Kopjar B, Boorman RS, Churchill RS, Smith KL, Matsen FA 3rd. Characterizing the functional improvement after total shoulder arthroplasty for osteoarthritis. *J Bone Joint Surg Am* 2002; **84**: 1349-1353 [PMID: 12177264 DOI: 10.2106/00004623-200208000-00009]
- 6 **Haines JF**, Trail IA, Nuttall D, Birch A, Barrow A. The results of arthroplasty in osteoarthritis of the shoulder. *J Bone Joint Surg Br* 2006; **88**: 496-501 [PMID: 16567785 DOI: 10.1302/0301-620X.88B4.16604]
- 7 **Deshmukh AV**, Koris M, Zurakowski D, Thornhill TS. Total shoulder arthroplasty: long-term survivorship, functional outcome, and quality of life. *J Shoulder Elbow Surg* 2005; **14**: 471-479 [PMID: 16194737 DOI: 10.1016/j.jse.2005.02.009]
- 8 **Padegimas EM**, Maltenfort M, Lazarus MD, Ramsey ML, Williams GR, Namdari S. Future patient demand for shoulder arthroplasty by younger patients: national projections. *Clin Orthop Relat Res* 2015; **473**: 1860-1867 [PMID: 25758376 DOI: 10.1007/s11999-015-4231-z]
- 9 **Fry R**. Millennials are the largest generation in the U.S. labor force. Pew Research Center: Fact Tank. [cited 31 January 2021]. In: Pew Research Center [Internet]. Available from: <https://www.pewresearch.org/fact-tank/2018/04/11/millennials-largest-generation-us-labor-force/>
- 10 **Bühlhoff M**, Sattler P, Bruckner T, Loew M, Zeifang F, Raiss P. Do patients return to sports and work after total shoulder replacement surgery? *Am J Sports Med* 2015; **43**: 423-427 [PMID: 25406301 DOI: 10.1177/0363546514557940]
- 11 **Hurwit DJ**, Liu JN, Garcia GH, Mahony G, Wu HH, Dines DM, Warren RF, Gulotta LV. A comparative analysis of work-related outcomes after humeral hemiarthroplasty and reverse total shoulder arthroplasty. *J Shoulder Elbow Surg* 2017; **26**: 954-959 [PMID: 28089256 DOI: 10.1016/j.jse.2016.10.004]
- 12 **Liu JN**, Garcia GH, Wong AC, Sinatro A, Wu HH, Dines DM, Warren RF, Gulotta LV. Return to Work After Anatomic Total Shoulder Arthroplasty for Patients 55 Years and Younger at Average 5-Year Follow-up. *Orthopedics* 2018; **41**: e310-e315 [PMID: 29451941 DOI: 10.3928/01477447-20180213-08]
- 13 **McInnes MDF**, Moher D, Thoms BD, McGrath TA, Bossuyt PM; and the PRISMA-DTA Group, Clifford T, Cohen JF, Deeks JJ, Gatsonis C, Hooft L, Hunt HA, Hyde CJ, Korevaar DA, Leeftang MMG, Macaskill P, Reitsma JB, Rodin R, Rutjes AWS, Salameh JP, Stevens A, Takwoingi Y, Tonelli M, Weeks L, Whiting P, Willis BH. Preferred Reporting Items for a Systematic Review and Meta-analysis of Diagnostic Test Accuracy Studies: The PRISMA-DTA Statement. *JAMA* 2018; **319**: 388-396 [PMID: 29362800 DOI: 10.1001/jama.2017.19163]
- 14 **Salameh JP**, Bossuyt PM, McGrath TA, Thoms BD, Hyde CJ, Macaskill P, Deeks JJ, Leeftang M, Korevaar DA, Whiting P, Takwoingi Y, Reitsma JB, Cohen JF, Frank RA, Hunt HA, Hooft L, Rutjes AWS, Willis BH, Gatsonis C, Levis B, Moher D, McInnes MDF. Preferred reporting items for systematic review and meta-analysis of diagnostic test accuracy studies (PRISMA-DTA): explanation, elaboration, and checklist. *BMJ* 2020; **370**: m2632 [PMID: 32816740 DOI: 10.1136/bmj.m2632]
- 15 **Slim K**, Nini E, Forestier D, Kwiatkowski F, Panis Y, Chipponi J. Methodological index for non-randomized studies (minors): development and validation of a new instrument. *ANZ J Surg* 2003; **73**: 712-716 [PMID: 12956787 DOI: 10.1046/j.1445-2197.2003.02748.x]
- 16 **Roberson TA**, Bentley JC, Griscom JT, Kissenberth MJ, Tolan SJ, Hawkins RJ, Tokish JM. Outcomes of total shoulder arthroplasty in patients younger than 65 years: a systematic review. *J Shoulder Elbow Surg* 2017; **26**: 1298-1306 [PMID: 28209327 DOI: 10.1016/j.jse.2016.12.069]
- 17 **Steinhaus ME**, Gowd AK, Hurwit DJ, Lieber AC, Liu JN. Return to work after shoulder arthroplasty: a systematic review and meta-analysis. *J Shoulder Elbow Surg* 2019; **28**: 998-1008 [PMID: 30885548 DOI: 10.1016/j.jse.2018.12.011]
- 18 **Riffkin R**. Americans Settling on Older Retirement Age. Gallup. [cited 31 January 2021]. Available from: <https://news.gallup.com/poll/182939/americans-settling-older-retirement-age.aspx>
- 19 **Mattei L**, Mortera S, Arrigoni C, Castoldi F. Anatomic shoulder arthroplasty: an update on indications, technique, results

- and complication rates. *Joints* 2015; **3**: 72-77 [PMID: 26605254 DOI: 10.11138/jts/2015.3.2.072]
- 20 **Walch G**, Boileau P, Noël E. Shoulder arthroplasty: evolving techniques and indications. *Joint Bone Spine* 2010; **77**: 501-505 [PMID: 20961793 DOI: 10.1016/j.jbspin.2010.09.004]
 - 21 **Cvetanovich GL**, Naylor AJ, O'Brien MC, Waterman BR, Garcia GH, Nicholson GP. Anatomic total shoulder arthroplasty with an inlay glenoid component: clinical outcomes and return to activity. *J Shoulder Elbow Surg* 2020; **29**: 1188-1196 [PMID: 31899092 DOI: 10.1016/j.jse.2019.10.003]
 - 22 **Jayasekara M**, Lam PH, Murrell GAC. Return to Work Following Shoulder Surgery: An Analysis of 1,773 Cases. *JB JS Open Access* 2020; **5** [PMID: 32803105 DOI: 10.2106/JBJS.OA.19.00081]
 - 23 **Westermann RW**, Pugely AJ, Martin CT, Gao Y, Wolf BR, Hettrich CM. Reverse Shoulder Arthroplasty in the United States: A Comparison of National Volume, Patient Demographics, Complications, and Surgical Indications. *Iowa Orthop J* 2015; **35**: 1-7 [PMID: 26361437]
 - 24 **Garcia GH**, Taylor SA, DePalma BJ, Mahony GT, Grawe BM, Nguyen J, Dines JS, Dines DM, Warren RF, Craig EV, Gulotta LV. Patient Activity Levels After Reverse Total Shoulder Arthroplasty: What Are Patients Doing? *Am J Sports Med* 2015; **43**: 2816-2821 [PMID: 26316610 DOI: 10.1177/0363546515597673]
 - 25 **Nam D**, Kepler CK, Neviasser AS, Jones KJ, Wright TM, Craig EV, Warren RF. Reverse total shoulder arthroplasty: current concepts, results, and component wear analysis. *J Bone Joint Surg Am* 2010; **92** Suppl 2: 23-35 [PMID: 21189245 DOI: 10.2106/JBJS.J.00769]
 - 26 **Hyun YS**, Huri G, Garbis NG, McFarland EG. Uncommon indications for reverse total shoulder arthroplasty. *Clin Orthop Surg* 2013; **5**: 243-255 [PMID: 24340143 DOI: 10.4055/cios.2013.5.4.243]
 - 27 **Drake GN**, O'Connor DP, Edwards TB. Indications for reverse total shoulder arthroplasty in rotator cuff disease. *Clin Orthop Relat Res* 2010; **468**: 1526-1533 [PMID: 20049573 DOI: 10.1007/s11999-009-1188-9]
 - 28 **Garcia GH**, Taylor SA, Mahony GT, DePalma BJ, Grawe BM, Nguyen J, Dines JS, Dines DM, Warren RF, Craig EV, Gulotta LV. Reverse Total Shoulder Arthroplasty and Work-Related Outcomes. *Orthopedics* 2016; **39**: e230-e235 [PMID: 26811957 DOI: 10.3928/01477447-20160119-03]
 - 29 **Simovitch RW**, Gerard BK, Brees JA, Fullick R, Kears JC. Outcomes of reverse total shoulder arthroplasty in a senior athletic population. *J Shoulder Elbow Surg* 2015; **24**: 1481-1485 [PMID: 25958214 DOI: 10.1016/j.jse.2015.03.011]
 - 30 **Palsis JA**, Simpson KN, Matthews JH, Traven S, Eichinger JK, Friedman RJ. Current Trends in the Use of Shoulder Arthroplasty in the United States. *Orthopedics* 2018; **41**: e416-e423 [PMID: 29658976 DOI: 10.3928/01477447-20180409-05]
 - 31 **Gadea F**, Alami G, Pape G, Boileau P, Favard L. Shoulder hemiarthroplasty: outcomes and long-term survival analysis according to etiology. *Orthop Traumatol Surg Res* 2012; **98**: 659-665 [PMID: 22944393 DOI: 10.1016/j.otsr.2012.03.020]
 - 32 **Schultz BJ**, Lowe DT, Egol KA, Zuckerman JD. Shoulder Hemiarthroplasty for Proximal Humerus Fracture. *J Orthop Trauma* 2021; **35**: S3-S4 [PMID: 34227587 DOI: 10.1097/BOT.0000000000002158]
 - 33 **Garcia GH**, Liu JN, Mahony GT, Sinatro A, Wu HH, Craig EV, Warren RF, Dines DM, Gulotta LV. Hemiarthroplasty Versus Total Shoulder Arthroplasty for Shoulder Osteoarthritis: A Matched Comparison of Return to Sports. *Am J Sports Med* 2016; **44**: 1417-1422 [PMID: 26960913 DOI: 10.1177/0363546516632527]
 - 34 **Sowa B**, Thierjung H, Bühlhoff M, Loew M, Zeifang F, Bruckner T, Raiss P. Functional results of hemi- and total shoulder arthroplasty according to diagnosis and patient age at surgery. *Acta Orthop* 2017; **88**: 310-314 [PMID: 28121220 DOI: 10.1080/17453674.2017.1280656]
 - 35 **Singh JA**, Sperling J, Buchbinder R, McMaken K. Surgery for shoulder osteoarthritis: a Cochrane systematic review. *J Rheumatol* 2011; **38**: 598-605 [PMID: 21239751 DOI: 10.3899/jrheum.101008]
 - 36 **Garcia GH**, Mahony GT, Fabricant PD, Wu HH, Dines DM, Warren RF, Craig EV, Gulotta LV. Sports- and Work-Related Outcomes After Shoulder Hemiarthroplasty. *Am J Sports Med* 2016; **44**: 490-496 [PMID: 26657261 DOI: 10.1177/0363546515613077]
 - 37 **Levine WN**, Fischer CR, Nguyen D, Flatow EL, Ahmad CS, Bigliani LU. Long-term follow-up of shoulder hemiarthroplasty for glenohumeral osteoarthritis. *J Bone Joint Surg Am* 2012; **94**: e164 [PMID: 23172331 DOI: 10.2106/JBJS.K.00603]
 - 38 **Botros M**, Curry EJ, Yin J, Jawa A, Eichinger JK, Li X. Reverse shoulder arthroplasty has higher perioperative implant complications and transfusion rates than total shoulder arthroplasty. *JSES Open Access* 2019; **3**: 108-112 [PMID: 31334437 DOI: 10.1016/j.jses.2019.03.001]
 - 39 **Wierks C**, Skolasky RL, Ji JH, McFarland EG. Reverse total shoulder replacement: intraoperative and early postoperative complications. *Clin Orthop Relat Res* 2009; **467**: 225-234 [PMID: 18685908 DOI: 10.1007/s11999-008-0406-1]
 - 40 **Flurin PH**, Roche CP, Wright TW, Marczuk Y, Zuckerman JD. A Comparison and Correlation of Clinical Outcome Metrics in Anatomic and Reverse Total Shoulder Arthroplasty. *Bull Hosp Jt Dis (2013)* 2015; **73** Suppl 1: S118-S123 [PMID: 26631207]
 - 41 **Kurowicki J**, Rosas S, Law TY, Levy JC. Participation in Work and Sport Following Reverse and Total Shoulder Arthroplasty. *Am J Orthop (Belle Mead NJ)* 2018; **47** [PMID: 29883508 DOI: 10.12788/ajo.2018.0034]
 - 42 **Bryant D**, Litchfield R, Sandow M, Gartsman GM, Guyatt G, Kirkley A. A comparison of pain, strength, range of motion, and functional outcomes after hemiarthroplasty and total shoulder arthroplasty in patients with osteoarthritis of the shoulder. A systematic review and meta-analysis. *J Bone Joint Surg Am* 2005; **87**: 1947-1956 [PMID: 16140808 DOI: 10.2106/JBJS.D.02854]
 - 43 **Edwards TB**, Kadakia NR, Boulahia A, Kempf JF, Boileau P, Némaz C, Walch G. A comparison of hemiarthroplasty and total shoulder arthroplasty in the treatment of primary glenohumeral osteoarthritis: results of a multicenter study. *J Shoulder Elbow Surg* 2003; **12**: 207-213 [PMID: 12851570 DOI: 10.1016/s1058-2746(02)86804-5]
 - 44 **Radnay CS**, Setter KJ, Chambers L, Levine WN, Bigliani LU, Ahmad CS. Total shoulder replacement compared with humeral head replacement for the treatment of primary glenohumeral osteoarthritis: a systematic review. *J Shoulder Elbow Surg* 2007; **16**: 396-402 [PMID: 17582789 DOI: 10.1016/j.jse.2006.10.017]
 - 45 **van den Bekerom MP**, Geervliet PC, Somford MP, van den Borne MP, Boer R. Total shoulder arthroplasty vs hemiarthroplasty for glenohumeral arthritis: A systematic review of the literature at long-term follow-up. *Int J Shoulder*

- Surg* 2013; **7**: 110-115 [PMID: [24167403](#) DOI: [10.4103/0973-6042.118915](#)]
- 46 **Carroll RM**, Izquierdo R, Vazquez M, Blaine TA, Levine WN, Bigliani LU. Conversion of painful hemiarthroplasty to total shoulder arthroplasty: long-term results. *J Shoulder Elbow Surg* 2004; **13**: 599-603 [PMID: [15570227](#) DOI: [10.1016/j.jse.2004.03.016](#)]
 - 47 **Golant A**, Christoforou D, Zuckerman JD, Kwon YW. Return to sports after shoulder arthroplasty: a survey of surgeons' preferences. *J Shoulder Elbow Surg* 2012; **21**: 554-560 [PMID: [21393018](#) DOI: [10.1016/j.jse.2010.11.021](#)]
 - 48 **Gowd AK**, Garcia GH, Liu JN, Malaret MR, Cabarcas BC, Romeo AA. Comparative analysis of work-related outcomes in hemiarthroplasty with concentric glenoid reaming and total shoulder arthroplasty. *J Shoulder Elbow Surg* 2019; **28**: 244-251 [PMID: [30269934](#) DOI: [10.1016/j.jse.2018.07.026](#)]
 - 49 **Liu JN**, Garcia GH, Gowd AK, Mahony G, Sinatro A, Wu HH, Dines DM, Warren RF, Gulotta LV. Return to Work After Shoulder Replacement for Glenohumeral Osteoarthritis Is Similar When Hemiarthroplasty Is Compared to Total Shoulder Arthroplasty. *HSS J* 2020; **16**: 212-217 [PMID: [33088235](#) DOI: [10.1007/s11420-019-09692-0](#)]
 - 50 **Lin DJ**, Wong TT, Kazam JK. Shoulder Arthroplasty, from Indications to Complications: What the Radiologist Needs to Know. *Radiographics* 2016; **36**: 192-208 [PMID: [26761537](#) DOI: [10.1148/rg.2016150055](#)]
 - 51 **Alentorn-Geli E**, Guirro P, Santana F, Torrens C. Treatment of fracture sequelae of the proximal humerus: comparison of hemiarthroplasty and reverse total shoulder arthroplasty. *Arch Orthop Trauma Surg* 2014; **134**: 1545-1550 [PMID: [25138037](#) DOI: [10.1007/s00402-014-2074-9](#)]
 - 52 **Boyle MJ**, Youn SM, Frampton CM, Ball CM. Functional outcomes of reverse shoulder arthroplasty compared with hemiarthroplasty for acute proximal humeral fractures. *J Shoulder Elbow Surg* 2013; **22**: 32-37 [PMID: [22652065](#) DOI: [10.1016/j.jse.2012.03.006](#)]
 - 53 **Leung B**, Horodyski M, Struk AM, Wright TW. Functional outcome of hemiarthroplasty compared with reverse total shoulder arthroplasty in the treatment of rotator cuff tear arthropathy. *J Shoulder Elbow Surg* 2012; **21**: 319-323 [PMID: [21872496](#) DOI: [10.1016/j.jse.2011.05.023](#)]
 - 54 **Magnussen RA**, Mallon WJ, Willems WJ, Moorman CT 3rd. Long-term activity restrictions after shoulder arthroplasty: an international survey of experienced shoulder surgeons. *J Shoulder Elbow Surg* 2011; **20**: 281-289 [PMID: [21051242](#) DOI: [10.1016/j.jse.2010.07.021](#)]
 - 55 **Morris BJ**, Haigler RE, Laughlin MS, Elkousy HA, Gartsman GM, Edwards TB. Workers' compensation claims and outcomes after reverse shoulder arthroplasty. *J Shoulder Elbow Surg* 2015; **24**: 453-459 [PMID: [25306491](#) DOI: [10.1016/j.jse.2014.07.009](#)]
 - 56 **Jawa A**, Dasti UR, Fasulo SM, Vaickus MH, Curtis AS, Miller SL. Anatomic total shoulder arthroplasty for patients receiving workers' compensation. *J Shoulder Elbow Surg* 2015; **24**: 1694-1697 [PMID: [26159842](#) DOI: [10.1016/j.jse.2015.04.017](#)]
 - 57 **Gruson KI**, Huang K, Wanich T, Depalma AA. Workers' compensation and outcomes of upper extremity surgery. *J Am Acad Orthop Surg* 2013; **21**: 67-77 [PMID: [23378370](#) DOI: [10.5435/JAAOS-21-02-67](#)]
 - 58 **Harris I**, Mulford J, Solomon M, van Gelder JM, Young J. Association between compensation status and outcome after surgery: a meta-analysis. *JAMA* 2005; **293**: 1644-1652 [PMID: [15811984](#) DOI: [10.1001/jama.293.13.1644](#)]
 - 59 **Chen AL**, Bain EB, Horan MP, Hawkins RJ. Determinants of patient satisfaction with outcome after shoulder arthroplasty. *J Shoulder Elbow Surg* 2007; **16**: 25-30 [PMID: [17097315](#) DOI: [10.1016/j.jse.2006.04.013](#)]
 - 60 **Vajapey SP**, Cvetanovich GL, Bishop JY, Neviaser AS. Psychosocial factors affecting outcomes after shoulder arthroplasty: a systematic review. *J Shoulder Elbow Surg* 2020; **29**: e175-e184 [PMID: [31899094](#) DOI: [10.1016/j.jse.2019.09.043](#)]
 - 61 **Cuff DJ**, Pupello DR. Prospective evaluation of postoperative compliance and outcomes after rotator cuff repair in patients with and without workers' compensation claims. *J Shoulder Elbow Surg* 2012; **21**: 1728-1733 [PMID: [22652063](#) DOI: [10.1016/j.jse.2012.03.002](#)]
 - 62 **Henn RF 3rd**, Tashjian RZ, Kang L, Green A. Patients with workers' compensation claims have worse outcomes after rotator cuff repair. *J Bone Joint Surg Am* 2008; **90**: 2105-2113 [PMID: [18829907](#) DOI: [10.2106/JBJS.F.00260](#)]
 - 63 **Holtby R**, Razmjou H. Impact of work-related compensation claims on surgical outcome of patients with rotator cuff related pathologies: a matched case-control study. *J Shoulder Elbow Surg* 2010; **19**: 452-460 [PMID: [19766021](#) DOI: [10.1016/j.jse.2009.06.011](#)]
 - 64 **Razmjou H**, Henry P, Costa G, Dwyer T, Holtby R. Effect of arthroscopic rotator cuff surgery in patients with preoperative restricted range of motion. *BMC Musculoskelet Disord* 2016; **17**: 99 [PMID: [26911157](#) DOI: [10.1186/s12891-016-0956-4](#)]
 - 65 **Nicholson GP**. Arthroscopic acromioplasty: a comparison between workers' compensation and non-workers' compensation populations. *J Bone Joint Surg Am* 2003; **85**: 682-689 [PMID: [12672845](#)]
 - 66 **Denard PJ**, Lädermann A, Burkhart SS. Long-term outcome after arthroscopic repair of type II SLAP lesions: results according to age and workers' compensation status. *Arthroscopy* 2012; **28**: 451-457 [PMID: [22264832](#) DOI: [10.1016/j.arthro.2011.09.005](#)]
 - 67 **Verma NN**, Garretson R, Romeo AA. Outcome of arthroscopic repair of type II SLAP lesions in worker's compensation patients. *HSS J* 2007; **3**: 58-62 [PMID: [18751771](#) DOI: [10.1007/s11420-006-9023-2](#)]
 - 68 **Werner BC**, Pehlivian HC, Hart JM, Lyons ML, Gilmore CJ, Garrett CB, Carson EW, Diduch DR, Miller MD, Brockmeier SF. Biceps tenodesis is a viable option for salvage of failed SLAP repair. *J Shoulder Elbow Surg* 2014; **23**: e179-e184 [PMID: [24332800](#) DOI: [10.1016/j.jse.2013.11.020](#)]
 - 69 **Bible JE**, Spengler DM, Mir HR. A primer for workers' compensation. *Spine J* 2014; **14**: 1325-1331 [PMID: [24462532](#) DOI: [10.1016/j.spinee.2014.01.030](#)]
 - 70 **US Department of Labor**, Office of Administrative Law Judges. Dictionary of occupational titles, revised 4th ed. [cited 31 January 2021]. In: United States Department of Labor [Internet]. Available from: <https://www.oalj.dol.gov/PUBLIC/DOT/REFERENCES/DOTAPPC.HTM>
 - 71 **Wallace BC**, Dahabreh IJ, Trikalinos TA, Lau J, Trow P, Schmid CH. Closing the Gap between Methodologists and End-Users: R as a Computational Back-End. *J Stat Softw* 2012; **49**: 1-15 [DOI: [10.18637/jss.v049.i05](#)]
 - 72 **Voss MR**, Homa JK, Singh M, Seidl JA, Griffitt WE. Outcomes of an interdisciplinary work rehabilitation program. *Work*

- 2019; **64**: 507-514 [PMID: [31658084](#) DOI: [10.3233/WOR-193012](#)]
- 73 **Knoblauch DK**, Cassaro S. Workers Compensation. [cited 31 January 2021]. In: StatPearls [Internet]. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK448106/>
- 74 **Cabarcas BC**, Gowd AK, Liu JN, Cvetanovich GL, Erickson BJ, Romeo AA, Verma NN. Establishing maximum medical improvement following reverse total shoulder arthroplasty for rotator cuff deficiency. *J Shoulder Elbow Surg* 2018; **27**: 1721-1731 [PMID: [30030030](#) DOI: [10.1016/j.jse.2018.05.029](#)]
- 75 **Puzzitiello RN**, Agarwalla A, Liu JN, Cvetanovich GL, Romeo AA, Forsythe B, Verma NN. Establishing maximal medical improvement after anatomic total shoulder arthroplasty. *J Shoulder Elbow Surg* 2018; **27**: 1711-1720 [PMID: [29730138](#) DOI: [10.1016/j.jse.2018.03.007](#)]
- 76 **Matar RN**, Gardner TJ, Kassam F, Grawe BM. When do patients truly reach maximal medical improvement after undergoing reverse shoulder arthroplasty? *JSES Int* 2020; **4**: 675-679 [PMID: [32939505](#) DOI: [10.1016/j.jseint.2020.03.010](#)]
- 77 **Ansuategui Echeita J**, Bethge M, van Holland BJ, Gross DP, Kool J, Oesch P, Trippolini MA, Chapman E, Cheng ASK, Sellars R, Spavins M, Streibelt M, van der Wurff P, Reneman MF. Correction to: Functional Capacity Evaluation in Different Societal Contexts: Results of a Multicountry Study. *J Occup Rehabil* 2019; **29**: 237-238 [PMID: [29946814](#) DOI: [10.1007/s10926-018-9797-3](#)]
- 78 **Ansuategui Echeita J**, Bethge M, van Holland BJ, Gross DP, Kool J, Oesch P, Trippolini MA, Chapman E, Cheng ASK, Sellars R, Spavins M, Streibelt M, van der Wurff P, Reneman MF. Functional Capacity Evaluation in Different Societal Contexts: Results of a Multicountry Study. *J Occup Rehabil* 2019; **29**: 222-236 [PMID: [29802582](#) DOI: [10.1007/s10926-018-9782-x](#)]
- 79 **Borm GF**, Lemmers O, Fransen J, Donders R. The evidence provided by a single trial is less reliable than its statistical analysis suggests. *J Clin Epidemiol* 2009; **62**: 711-715.e1 [PMID: [19171462](#) DOI: [10.1016/j.jclinepi.2008.09.013](#)]
- 80 **Rücker G**, Schwarzer G, Schumacher M, Carpenter J. Are large trials less reliable than small trials? *J Clin Epidemiol* 2009; **62**: 886-7; author reply 887 [PMID: [19481419](#) DOI: [10.1016/j.jclinepi.2009.03.007](#)]
- 81 **Shrier I**, Platt RW, Steele RJ. Mega-trials vs. meta-analysis: precision vs. heterogeneity? *Contemp Clin Trials* 2007; **28**: 324-328 [PMID: [17188025](#) DOI: [10.1016/j.cct.2006.11.007](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

