World Journal of *Orthopedics*

World J Orthop 2023 February 18; 14(2): 42-89





Published by Baishideng Publishing Group Inc

World Journal of Orthopedics

Contents

Monthly Volume 14 Number 2 February 18, 2023

MINIREVIEWS

Diagnosis and treatment of chronic osteomyelitis based on nanomaterials 42 Zeng M, Xu Z, Song ZQ, Li JX, Tang ZW, Xiao S, Wen J

55 Fungal arthritis: A challenging clinical entity

Mishra A, Juneja D

ORIGINAL ARTICLE

Basic Study

Mechanism of spinal cord injury regeneration and the effect of human neural stem cells-secretome 64 treatment in rat model

Semita IN, Utomo DN, Suroto H

LETTER TO THE EDITOR

83 Effect of SARS-CoV-2 infection on trauma throughput to alternative elective care approaches Joob B. Wiwanitkit V

85 New classification for septic arthritis of the hand Lipatov KV, Asatryan A, Melkonyan G, Kazantcev AD, Solov'eva EI, Gorbacheva IV, Vorotyntsev AS, Emelyanov AY



Contents

Monthly Volume 14 Number 2 February 18, 2023

ABOUT COVER

Editorial Board Member of World Journal of Orthopedics, Li-Fang Hu, PhD, Associate Professor, School of Life Sciences, Northwestern Polytechnical University, Xi'an 710072, Shaanxi Province, China. hulifang@nwpu.edu.cn

AIMS AND SCOPE

The primary aim of World Journal of Orthopedics (WJO, World J Orthop) is to provide scholars and readers from various fields of orthopedics with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJO mainly publishes articles reporting research results and findings obtained in the field of orthopedics and covering a wide range of topics including arthroscopy, bone trauma, bone tumors, hand and foot surgery, joint surgery, orthopedic trauma, osteoarthropathy, osteoporosis, pediatric orthopedics, spinal diseases, spine surgery, and sports medicine.

INDEXING/ABSTRACTING

WJO is now abstracted and indexed in PubMed, PubMed Central, Emerging Sources Citation Index (Web of Science), Scopus, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 edition of Journal Citation Reports® cites the 2021 Journal Citation Indicator (JCI) for WJO as 0.62. The WJO's CiteScore for 2021 is 2.4 and Scopus CiteScore rank 2021: Orthopedics and Sports Medicine is 139/284.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Ying-Yi Yuan, Production Department Director: Xiang Li, Editorial Office Director: Jin-Lei Wang.

NAME OF JOURNAL World Journal of Orthopedics	INSTRUCTIONS TO AUTHORS https://www.wjgnet.com/bpg/gerinfo/204
ISSN	GUIDELINES FOR ETHICS DOCUMENTS
ISSN 2218-5836 (online)	https://www.wjgnet.com/bpg/GerInfo/287
LAUNCH DATE	GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH
November 18, 2010	https://www.wjgnet.com/bpg/gerinfo/240
FREQUENCY	PUBLICATION ETHICS
Monthly	https://www.wjgnet.com/bpg/GerInfo/288
EDITORS-IN-CHIEF	PUBLICATION MISCONDUCT
Massimiliano Leigheb	https://www.wjgnet.com/bpg/gerinfo/208
EDITORIAL BOARD MEMBERS	ARTICLE PROCESSING CHARGE
http://www.wjgnet.com/2218-5836/editorialboard.htm	https://www.wjgnet.com/bpg/gerinfo/242
PUBLICATION DATE	STEPS FOR SUBMITTING MANUSCRIPTS
February 18, 2023	https://www.wjgnet.com/bpg/GerInfo/239
COPYRIGHT	ONLINE SUBMISSION
© 2023 Baishideng Publishing Group Inc	https://www.f6publishing.com

© 2023 Baishideng Publishing Group Inc. All rights reserved. 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA E-mail: bpgoffice@wjgnet.com https://www.wjgnet.com



WJD

World Journal of **Orthopedics**

Submit a Manuscript: https://www.f6publishing.com

World J Orthop 2023 February 18; 14(2): 85-89

DOI: 10.5312/wio.v14.i2.85

ISSN 2218-5836 (online)

LETTER TO THE EDITOR

New classification for septic arthritis of the hand

Konstantin V Lipatov, Arthur Asatryan, George Melkonyan, Aleksandr D Kazantcev, Ekaterina I Solov'eva, Irina V Gorbacheva, Alexander S Vorotyntsev, Andrey Y Emelyanov

Specialty type: Orthopedics

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0 Grade B (Very good): B, B Grade C (Good): 0 Grade D (Fair): 0 Grade E (Poor): 0

P-Reviewer: Doski JO, Iraq; Mehta V. India

Received: November 16, 2022 Peer-review started: November 16, 2022

First decision: November 25, 2022 Revised: November 29, 2022 Accepted: February 7, 2023 Article in press: February 7, 2023 Published online: February 18, 2023



Konstantin V Lipatov, Aleksandr D Kazantcev, Department of General Surgery, Institute of Clinical Medicine Named After N.V. Sklifosovsky, Sechenov First Moscow State Medical University (Sechenov University), Moscow 119021, Russia

Arthur Asatryan, Department of Wound and Wound Infection Surgery, State Budgetary Institution "City Clinical Hospital named after S.S. Yudin of Moscow Healthcare Department", Moscow 115446, Russia

George Melkonyan, Department of General Surgery, Hospital for War Veterans №3, Moscow 129336. Russia

Ekaterina I Solov'eva, Irina V Gorbacheva, Alexander S Vorotyntsev, Andrey Y Emelyanov, Department of General Surgery, I.M. Sechenov First Moscow State Medical University, Moscow 119048, Russia

Corresponding author: Konstantin V Lipatov, DSc, MD, Doctor, Full Professor, Professor, Surgeon, Department of General Surgery, Institute of Clinical Medicine Named After N.V. Sklifosovsky, Sechenov First Moscow State Medical University (Sechenov University), Rossolimo Street 11-2, Moscow 119021, Russia. lipatov k v@staff.sechenov.ru

Abstract

The severity of septic arthritis of the hand and the prospects for restoration of joint function are determined by a complex of factors. Among them, the leading role belongs to local changes in tissue structures. This includes the destruction of articular cartilage and bone tissue with the development of osteomyelitis, the involvement of paraarticular soft tissues in the purulent process, and the destruction of the flexor/extensor tendons of the fingers. The currently missing specialized classification of septic arthritis could help in systematizing the diseases, determining treatment tactics, and predicting the results of treatment. The classification of septic arthritis of the hand proposed for discussion is based on the following principle: Joint–Wound–Tendon $(J_x W_x T_y)$; J_x characterizes damage to the osteochondral structures of the joint, W_x is the presence of paraarticular purulent wounds or fistulas, and T_x is destruction of the flexor/extensor tendons of the finger. The classification of the diagnosis makes it possible to assess the nature and severity of damage to the structures of the joint and may also be useful when comparing the results of treatment of septic arthritis of the hand.

Key Words: Hand; Septic arthritis; Classification; Osteomyelitis; Paraarticular wounds



WJO https://www.wjgnet.com

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: The absence to date of a specialized classification of septic arthritis of the hand determines the relevance of its development. The proposed classification is based on the principle JOINT-WOUND-TENDON and reflects in aggregate the lesion of the osteo-cartilaginous structures of the joint, paraarticular soft tissues, and tendons of the flexor/extensor of the finger.

Citation: Lipatov KV, Asatryan A, Melkonyan G, Kazantcev AD, Solov'eva EI, Gorbacheva IV, Vorotyntsev AS, Emelyanov AY. New classification for septic arthritis of the hand. *World J Orthop* 2023; 14(2): 85-89 **URL**: https://www.wjgnet.com/2218-5836/full/v14/i2/85.htm **DOI**: https://dx.doi.org/10.5312/wjo.v14.i2.85

TO THE EDITOR

Septic arthritis of the hand is a common infectious pathology of the joints and ranks second in frequency after inflammation of the knee joint[1]. Given the high functional significance of the small joints of the hand, their inflammation often leads to serious consequences, sometimes ending in limitation or even disability. The most common cause of septic arthritis of the hand is various penetrating wounds. The pathogenic microflora that has entered the joint cavity causes the development of an infectious process, which, if surgical care is not provided in time, leads to the destruction of the articular cartilage and the development of osteomyelitis. In septic arthritis of the hand, a purulent process often occurs in paraarticular tissues with the formation of wounds and fistulas, which has a significant impact on the extent of surgical intervention and the results of treatment in general. The inflammatory process in the small joints of the hand, proceeding with purulent-necrotic lesions of the surrounding soft tissues, may be accompanied by destructive changes in the area of the flexor/extensor tendons of the finger, which negatively affects the prospects for restoring movements in the joint. This fact distinguishes the course of septic arthritis of the hand from purulent arthritis of large joints, in which tendon damage is uncharacteristic[2].

Thus, the course of septic arthritis of the hand and the features of treatment and its results are determined by a complex of pathological changes in the osteochondral apparatus, paraarticular soft tissues, and flexor/extensor tendons of the finger. While analyzing the literature on septic arthritis of the hand (published in the *World Journal of Orthopedics* 2022; 13: 622-630)[3], we faced problems comparing the results of surgical treatment presented by different authors.

To obtain reliable results, it is necessary to understand the nature and extent of tissue damage in the observations that we compare.

Evaluating the response to treatment of isolated septic arthritis, septic arthritis with osteomyelitis, purulent process in the paraarticular soft tissues, and tendon destruction can hardly be justified. However, in most cases, the authors use the term "septic arthritis of the hand" when formulating the diagnosis, only occasionally highlighting cases with osteomyelitis[4].

As a consequence, comparison of the results of treatment presented by different experts may be incorrect since the severity of the disease was initially different. In this regard, the use in clinical practice and in the course of scientific analysis of the classification of the pathological process is of great importance. However, to date, there has been no specialized classification of septic arthritis of the hand. According to the most common classification of hand infection given by Brown[5], only two forms can be attributed to the characteristics of purulent arthritis: Osteomyelitis and septic arthritis. However, this is clearly not enough in view of the above arguments[5]. This can be partially compensated for by using the classification of Tan *et al*[6] for infection of the joints of the hand, which was developed to characterize the inflammatory process in large joints[6]. It presents information about the damage to the osteochondral structures of the joint and paraarticular soft tissue (*isolated septic arthritis; septic arthritis with soft-tissue extension, but no osteomyelitis; septic arthritis with contiguous osteomyelitis*). At the same time, this classification does not reflect the presence/absence of destruction of the tendon apparatus, which is rare in arthritis of large joints but is extremely important in septic arthritis of the hand. In addition, the classification of Tan *et al*[6] is quite voluminous, which makes it difficult to use it in wide clinical practice.

We have significant experience in the treatment of patients with septic arthritis of the hand and would like to propose our classification of septic arthritis of the hand.

The classification is based on the principle of Joint–Wound–Tendon ($J_xW_xT_x$) and takes into account the presence of damage to the osteochondral structures of the joint, paraarticular soft tissues, and tendon apparatus.

WJO | https://www.wjgnet.com



DOI: 10.5312/wjo.v14.i2.85 Copyright ©The Author(s) 2023.

Figure 1 New classification for septic arthritis of the hand. J_xW_xT_x: Joint-Wound-Tendon.



DOI: 10.5312/wjo.v14.i2.85 Copyright ©The Author(s) 2023.

Figure 2 Septic arthritis of the proximal interphalangeal joint of the index finger (J₁W₀T₀). A: Hand after repeated treatment; B: X-ray.

The symbol J_x characterizes the lesion of the osteochondral structures of the joint, where the index "x" determines the presence or absence of a sign: J_0 -without osteomyelitis; J_1 -with osteomyelitis.

The symbol W_x characterizes the presence of purulent wounds or fistulas in the paraarticular region: W_0 -without wounds or fistulas; W_1 -with a fistula; W_2 -with a purulent wound.

The symbol T_x characterizes the destruction of the flexor/extensor tendons: T_0 -without tendon destruction; T_1 -extensor tendon destruction; T_2 -flexor tendon destruction; T_3 -destruction of both tendons.

Schematically, this classification can be represented below (Figure 1).

Examples of the application of this classification in the formulation of a diagnosis are shown below (Figures 2-4).

A 58-year-old woman pricked the index finger of her right hand with a needle while doing embroidery 3 wk ago. A day later, she noticed swelling of the finger and aching pain, aggravated by bending the finger. She was examined by a polyclinic surgeon and received a course of antibacterial drugs with some positive effect. Despite this, finger edema and pain on flexion persisted, which served as the basis for a second visit to the doctor. Upon further diagnostic testing, X-ray showed osteomyelitis in the proximal interphalangeal joint. On magnetic resonance imaging, there was purulent inflammation in the joint without destruction of the capsule and tendon apparatus (Figure 2).

A 54-year-old woman pricked the index finger of her left hand with a plant thorn while gardening a month ago. The next day, she noted swelling of the nail phalanx and reddening of the skin. She was self-treated using topical antiseptics and oral antibiotics (amoxicillin). The inflammation subsided, but a week ago, swelling and redness of the skin and pain reappeared in the same area. After using ointment dressings, a purulent fistula opened. X-ray showed signs of osteomyelitis in the area of the distal interphalangeal joint (DIP). On examination, there was a purulent fistula in the DIP projection. Intraop-



WJO https://www.wjgnet.com



DOI: 10.5312/wjo.v14.i2.85 Copyright ©The Author(s) 2023.

Figure 3 Septic arthritis of the distal interphalangeal joint of the index finger (J₁W₁T₁). A: During hospitalization; B: X-ray.



DOI: 10.5312/wjo.v14.i2.85 Copyright ©The Author(s) 2023.

Figure 4 Septic arthritis of the 2nd metacarpophalangeal joint of the right hand (J₀W₂T₁). A: During hospitalization; B: X-ray.

erative finding included the destruction of the joint capsule and extensor tendon (Figure 3).

A 30-year-old man injured his right hand as a result of a blow with a fist in a fight 10 d ago. The patient did not consult a doctor. He treated himself independently with topical antiseptics. During the last few days, the pain increased, the swelling of the hand increased, and the discharge from the wound became purulent. Wound revision revealed damage to the extensor tendon of the finger and a violation of the integrity of the capsule of the metacarpophalangeal joint and purulent exudate in the joint cavity. Upon further diagnostic testing, X-ray showed no evidence of osteomyelitis (Figure 4).

Thus, a specialized classification of septic arthritis of the hand takes into account the main pathological changes that occur in this disease: Destruction of bone and cartilage structures, paraarticular soft tissues, and tendon apparatus. Its use can help in predicting the course of the inflammatory process, determining treatment tactics, and determining the nature and timing of the start of rehabilitation measures. The use of the classification will make it possible to objectify the comparison of the results of treatment of septic arthritis of the hand presented by various authors.

FOOTNOTES

Author contributions: Lipatov KV designed the research; Asatryan A and Melkonyan G performed the research; Kazantcev AD analyzed the data; Solov'eva EI and Gorbacheva IV wrote the letter; Vorotyntsev AS and Emelyanov AY revised the letter.

Conflict-of-interest statement: All the authors report no relevant conflicts of interest for this article.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-



WJO | https://www.wjgnet.com

NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is noncommercial. See: https://creativecommons.org/Licenses/by-nc/4.0/

Country/Territory of origin: Russia

ORCID number: Konstantin V Lipatov 0000-0002-9902-2650; Arthur Asatryan 0000-0002-8409-2605; George Melkonyan 0000-0001-7234-4185; Aleksandr D Kazantcev 0000-0003-1238-1990; Ekaterina I Solov'eva 0000-0003-4143-3593; Irina V Gorbacheva 0000-0002-1060-1163; Alexander S Vorotyntsev 0000-0002-3686-4789; Andrey Y Emelyanov 0000-0002-9688-4079.

S-Editor: Fan JR L-Editor: Wang TQ P-Editor: Fan JR

REFERENCES

- 1 Sendi P, Kaempfen A, Uçkay I, Meier R. Bone and joint infections of the hand. Clin Microbiol Infect 2020; 26: 848-856 [PMID: 31917233 DOI: 10.1016/j.cmi.2019.12.007]
- 2 McBride S, Mowbray J, Caughey W, Wong E, Luey C, Siddiqui A, Alexander Z, Playle V, Askelund T, Hopkins C, Quek N, Ross K, Orec R, Mistry D, Coomarasamy C, Holland D. Epidemiology, Management, and Outcomes of Large and Small Native Joint Septic Arthritis in Adults. Clin Infect Dis 2020; 70: 271-279 [PMID: 30941403 DOI: 10.1093/cid/ciz265]
- 3 Lipatov KV, Asatryan A, Melkonyan G, Kazantcev AD, Solov'eva EI, Cherkasov UE. Septic arthritis of the hand: Current issues of etiology, pathogenesis, diagnosis, treatment. World J Orthop 2022; 13: 622-630 [PMID: 36051375 DOI: 10.5312/wio.v13.i7.622]
- 4 Wang J, Wang L. Novel therapeutic interventions towards improved management of septic arthritis. BMC Musculoskelet Disord 2021; 22: 530 [PMID: 34107951 DOI: 10.1186/s12891-021-04383-6]
- 5 Brown H. Hand infections. Am Fam Physician 1978; 18: 79-85 [PMID: 685801]
- 6 Tan V, Pepe MD, Esterhai JL. Sepsis of the shoulder girdle. In: Disoders of the shoulder: diagnosis and management. Edited by J. Iannotti, G.R. Williams. Philadelphia, Baltimore, New York, London, Buenos Aires, Hong Kong, Sydney, Tokyo. Lippincott Williams and Wilkins, 1998.-P.951-976. In: Habermeyer P, Magosch P, Lichtenberg S editors. Classifications and Scores of the Shoulder. Springer 2006: 179-181 [DOI: 10.1007/3-540-35142-6]





Published by Baishideng Publishing Group Inc 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA Telephone: +1-925-3991568 E-mail: bpgoffice@wjgnet.com Help Desk: https://www.f6publishing.com/helpdesk https://www.wjgnet.com

