

Antimicrobial potential of *Terminalia chebula* Retz. fruit extracts against ear pathogens

Chetan Sharma, Kamal R Aneja, Ramkrashan Kasera, Ashish Aneja

Chetan Sharma, Kamal R Aneja, Department of Microbiology, Kurukshetra University, Kurukshetra 136119, Haryana, India
 Ramkrashan Kasera, Department of Biotechnology, Pune University, Pune 411007, India

Ashish Aneja, University Health Center, Kurukshetra University, Kurukshetra 136119, Haryana, India

Author contributions: Sharma C and Aneja KR performed the majority of experiments; Kasera R helped in identification of bacterial pathogens; and Aneja A helped in collection of ear samples of otitis externa patients.

Supported by Kurukshetra University, Kurukshetra, Haryana

Correspondence to: Dr. Chetan Sharma, Department of Microbiology, Kurukshetra University, Kurukshetra 136119, Haryana, India. chetanmicro147@gmail.com

Telephone: +91-98-12287101 Fax: +91-17-44238277

Received: October 25, 2011 Revised: December 14, 2011

Accepted: February 27, 2012

Published online: April 28, 2012

Abstract

AIM: To evaluate the antimicrobial potential of *Terminalia chebula* (*T. chebula*) extracts against pathogens causing otitis externa and compare it with ear drops.

METHODS: Four different extracts, methanol, ethanol, acetone and aqueous (hot and cold) extracts, from the *T. chebula* were tested for their antimicrobial activity through the agar well diffusion method and minimum inhibitory concentration (MIC)/minimum bactericidal concentration (MBC) values were determined through the macrodilution broth method against six different microorganism, five bacterial (one gram positive and four gram negative) and one yeast.

RESULTS: Organic and aqueous fruit extracts displayed activity against all five tested bacterial ear pathogens with a maximum zone of inhibition of 31.6 mm against *Staphylococcus aureus*, followed by *Acinetobacter* sp. (24.6 mm), *Pseudomonas aeruginosa* (23.6 mm), *Proteus mirabilis* (21 mm) and *Escherichia coli* (19.3

mm). Of the four solvents evaluated, acetonetic fruit extract of *T. chebula* was found to be best. The MIC values ranged between 0.78 mg/mL and 50 mg/mL for the different bacterial ear pathogens and MBC values ranged between 1.56 mg/mL and 50 mg/mL. The acetonetic fruit extract showed larger inhibition zones compared to the herbal ear drops, Kan pip with lowest MIC of 0.78 mg/mL and MBC of 1.56 mg/mL.

CONCLUSION: Acetonetic extract of *T. chebula* fruit may be used to treat otitis externa. However, more detailed studies, such as *in vivo* testing and pharmacokinetics properties, are needed to determine its therapeutic potential.

© 2012 Baishideng. All rights reserved.

Key words: Otitis externa; *Terminalia chebula*; Antimicrobial activity; Minimum inhibitory concentration; Minimum bactericidal concentration; Organic and aqueous extracts

Peer reviewers: Francesco Dispenza, MD, PhD, Department of Otorhinolaryngology, Hospital S Giovanni di Dio ASP Agrigento, Via Oreto 339, 90124 Palermo, Italy; Nuray Bayar Muluk, MD, Professor, Department of Otorhinolaryngology, Kırıkkale University, Ankara 06610, Turkey

Sharma C, Aneja KR, Kasera R, Aneja A. Antimicrobial potential of *Terminalia chebula* Retz. fruit extracts against ear pathogens. *World J Otorhinolaryngol* 2012; 2(2): 8-13 Available from: URL: <http://www.wjgnet.com/2218-6247/full/v2/i2/8.htm>
 DOI: <http://dx.doi.org/10.5319/wjo.v2.i2.8>

INTRODUCTION

Otitis externa refers to a spectrum of infections of the external auditory canal and auricle, usually associated with microbial infection of macerated skin and subcutaneous cellular tissue. It affects between 5% and 20%

of patients attending ear, nose and throat (ENT) clinics. Manifestations of otitis externa include pain, pruritus and erythema but as the disease progresses, edema, otorrhea and conductive hearing loss may also develop^[1-3]. The main causative agents of the diseases are *Staphylococcus aureus* (*S. aureus*), *Staphylococcus epidermidis* (*S. epidermidis*), *Escherichia coli* (*E. coli*), *Pseudomonas aeruginosa* (*P. aeruginosa*), *Acinetobacter calcoaceticus*, *Proteus mirabilis* (*P. mirabilis*), *Aspergillus niger*, *Aspergillus flavus*, *Aspergillus fumigatus*, *Aspergillus luchuensis*, *Aspergillus terreus*, *Penicillium* sp. and *Candida albicans* (*C. albicans*)^[4-6].

Due to indiscriminate use of commercial antimicrobial drugs and development of multiple drug resistant strains of bacteria and fungi, there is a worldwide emphasis to search for new antimicrobials from natural sources. Plant metabolites and plant based drugs appear to be one of the better alternatives as they are known to have minimal toxicity and are cost-effective in contrast to synthetic agents^[7].

India is sitting on a gold mine of well-recorded and traditional well practised knowledge of herbal medicine^[8]. *Terminalia chebula* (*T. chebula*) Retz., commonly known as Black Myrobalan and Harad, is a member of the family *Combretaceae*. It is native to the Indian subcontinent and the adjacent areas such as Pakistan, Nepal, south-west China and Sri Lanka. In Tibet, it is called the “king of medicine”. It is a medium to large deciduous tree attaining a height of up to 30 m, with widely spreading branches and a broad roundish crown. The leaves are elliptical, oblong with an acute tip, cordate at the base, entire margins, glabrous above with a yellowish pubescence below. The flowers are monoecious, dull white to yellow, with a strong unpleasant odor, borne in terminal spikes or short panicles. The fruits are glabrous, ellipsoid to ovoid drupes, yellow to orange brown in color, containing a single angle stone^[9-11].

The dried ripe fruit has traditionally been used in the treatment of asthma, sore throat, vomiting, hiccup, diarrhea, bleeding piles, gout and heart and bladder disease. In addition, the plant is commonly used for acidity, chronic lung disease, skin diseases and eye disorders^[12-14]. It has been reported to exhibit a variety of biological activities, such as anticancer, antioxidant, antidiabetic, antibacterial, antifungal, antiviral, antianaphylactic, anti-ulcerogenic and antispasmodic. It also possesses cardioprotective, hepatoprotective, radioprotective and wound healing activities^[9,15,16].

This plant is known to be an important source of secondary metabolites, of which, 33% of the total phytoconstituents are hydrolysable tannins (which may vary from 20%-50%) and are responsible for pharmacological activity. The chief constituents of tannin are chebulic acid, chebulagic acid, corilagin and gallic acid^[17-19]. Tannin of *T. chebula* is of the pyrogallol (hydrolyzable) type. Hydrolyzable tannins (gallic acid, chebulic acid, punicalagin, chebulanin, corilagin, neochebulinic acid, ellagic acid, chebulogic acid, chebulinic acid, 1,2,3,4,6- penta-O-gallo-

yl-H-D-glucose, 1,6-di-O-galloyl-D-glucose, casuarinin, 3,4,6-tri-O-galloyl-D- glucose and terchebulin) have been reported from *T. chebula* fruits^[20-23].

In view of these reported medicinal values, the present study was carried out to examine the antimicrobial potential of *T. chebula* fruits against the locally isolated microbes, bacteria and yeasts, from otitis externa patients and to compare its efficacy with locally available ear drops.

MATERIALS AND METHODS

Plant collection

The fruits of *T. chebula* were obtained from the local market in Kurukshetra, Haryana. The taxonomic identity of the plant was confirmed by Dr. Vashishta BD, plant taxonomist, Chairman of Botany Department, Kurukshetra University, Kurukshetra.

Extraction of plant material

The samples were carefully washed under running tap water followed by sterile distilled water and air dried at room temperature (35-40 °C) for 4-5 d, homogenized to a fine powder using a sterilized mixer grinder and stored in air tight bottles. Four different solvents, namely ethanol, methanol, acetone and aqueous (hot and cold), were used for extraction. Homogenized fruits, 10 g each, were separately soaked in conical flasks each containing 100 mL of acetone, ethanol, methanol (95%) and sterile distilled water. Also, an equal amount (i.e., 10 g) of homogenized fruits was immersed separately in 100 mL of hot sterile distilled water in conical flasks and allowed to stand for 30 min in a water bath (at 100 °C) with occasional shaking, followed by keeping all the flasks on rotary shaker at 200 rpm for 24 h. Each preparation was filtered through a sterilized Whatman No. 1 filter paper and finally concentrated to dryness under vacuum at 40 °C using a rota evaporator. The dried extract thus obtained was sterilized by overnight ultra violet-irradiation, checked for sterility on nutrient agar plates and stored at 4 °C in labelled sterile bottles until further use^[24].

Test microorganisms

Five bacteria, namely *S. aureus* (HM626197, nucleotide sequence of all the five bacteria have been submitted to GenBank database which provided the GenBank accession number, HM626197-HM626201), *Acinetobacter* sp. (HM626198), *Proteus mirabilis* (HM626199), *E. coli* (HM626200), *P. aeruginosa* (HM626201) and one yeast, *C. albicans*, were isolated from patients with an ear infection from the local ENT clinics in Kurukshetra^[4]. Bacterial strains were identified on the basis of gram staining, biochemical and molecular characteristics (16S rRNA sequencing)^[25] and on the basis of staining, morphological and cultural characteristics for the yeast^[26,27]. The bacterial isolates were subcultured on nutrient agar and *C. albicans* on malt yeast agar and incubated aerobically at 37 °C. The media were procured from Hi Media Laboratory Pvt. Ltd., Bombay, India.

Ear drops

Three commonly prescribed ear drops by otolaryngologists, two allopathic ciprofloxacin (antibacterial), candid (antifungal) and a herbal ear drop Kan pip (antimicrobial), were procured from the local market in Kurukshetra.

Screening for antimicrobial activity

The acetone, methanol, ethanol and hot and cold aqueous *T. chebula* fruits extracts were used for evaluation of antimicrobial activity by the agar well diffusion method. In this method, a pure isolate of each microbe was grown on agar plates at 37 °C for 24 h. One plate of each microorganism was taken and a minimum of four colonies were transferred into normal saline (0.85%) under aseptic conditions. Density of each microbial suspension was adjusted to be equal to that of 10^6 cfu/mL (standardized by 0.5McFarland standard) and used as the inoculum for performing an agar well diffusion assay. One hundred microliter (100 µL) of the inoculum of each test organism was spread onto the agar plates so as to achieve a confluent growth. The agar plates were allowed to dry and 8 mm wells were made with a sterile borer in the inoculated agar plates. The lower portion of each well was sealed with molten agar medium. The dried extracts were reconstituted to 20% in dimethylsulphoxide (DMSO) for the bioassay analysis. A 100 µL volume of each extract was propelled directly into the wells (in triplicate) of the inoculated agar plates for each test organism. The plates were allowed to stand for 1 h at room temperature (40 °C) for diffusion of the extract into agar and incubated at 37 °C for 24 h. Sterile DMSO (20%) served as the negative control and ciprofloxacin (for bacteria), candid (for fungi) and Kan pip (antimicrobial) ear drops served as the positive controls. The antimicrobial activity, indicated by an inhibition zone surrounding the well containing the extract, was recorded if the zone was greater than 8 mm. The experiments were performed in triplicate and the mean values of the diameter of inhibition zones \pm standard deviations were calculated^[24].

Determination of minimum inhibitory concentration

Minimum inhibitory concentration (MIC) for each test organism was determined by the macrodilution broth method. A twofold serial dilution of each extract was prepared by first reconstituting the dried extract (100 mg/mL) in 20% DMSO, followed by dilution in Mueller-Hinton broth (bacteria) and Malt yeast broth (yeast) to achieve a decreasing concentration range of 50 mg/mL to 0.39 mg/mL. Each dilution was seeded with 100 µL of the standardized microbial inoculum (1.5×10^6 cfu/mL). The inoculated culture tubes were incubated at 37 °C for 24 h. A set of tubes containing only broth was kept as control. Afterwards, incubation tubes were examined for changes in turbidity as an indicator of growth. The lowest concentration that did not permit any visible growth was considered as MIC^[28,29].

Determination of minimum bactericidal concentration

Minimum bactericidal concentration (MBC) is the lowest concentration of antimicrobial agent that will not allow the growth of an organism after subculturing on antibiotic free media. MBC was determined by subculturing the preparations that did not show any bacterial growth in the MIC determination. A 100 µL aliquot from the selected tube (showing MIC) was spread over the MHA plate and incubated at 37 °C for 24 h and examined for bacterial growth. The MBC, the lowest concentration of the plant extract giving 99.9% reduction of the bacterial growth of various plants parts against the bacterial pathogens, was recorded^[29].

Determination of minimum fungicidal concentration

A loopful of culture from each set of tubes that did not show any visible growth of the yeast in MIC determination was subcultured on to fresh plates of MEA and incubated at 37 °C for 24-48 h. Minimum fungicidal concentration for each plant extracts against the tested yeast was recorded as the lowest concentration that did not yield any fungal growth on the solid medium^[30,31].

Statistical analysis

The experimental results were repeated thrice in triplicate each time and expressed as mean \pm SD and results were statistically evaluated using Dennett's *T*-test. *P* value less than 0.01 was considered significant.

RESULTS

The antibacterial activity of *T. chebula* fruits extracts on the agar plates varied in different organic (methanol, ethanol and acetone) and aqueous (hot and cold) extracts. Positive controls produced significantly sized inhibition zones against the tested bacteria (ranging between 16.3 mm and 56.3 mm) and the yeast (with zone of inhibition 21.6 mm), and the negative control produced no observable inhibitory effect against any of the test organism (Tables 1 and 2). A perusal of the data in Table 1 reveals that all the tested solvent fruit extracts possessed antibacterial activity against all five tested bacterial ear pathogens. However, the bioactivity against *C. albicans* was absent in all five extracts.

The acetonic fruit extract was found most effective against *S. aureus* (31.6 mm), followed by *Acinetobacter sp.* (24.6 mm), *P. aeruginosa* (23.6 mm), *P. mirabilis* (21 mm) and *E. coli* (19.3 mm). The ethanolic and methanolic fruit extracts showed moderate activity with inhibition zones of 28.6 mm and 27.3 mm against *S. aureus*, followed by *Acinetobacter sp.* (22.3 mm and 21.6 mm), *P. aeruginosa* (22.3 mm and 20.6 mm), *P. mirabilis* (20.6 mm and 19.3 mm) and *E. coli* (16.3 mm and 15.6 mm). Of the two tested aqueous extracts, hot aqueous extract exhibited more activity than the cold aqueous extract with zone of inhibition of 26.3 mm against *S. aureus*, followed by *Acinetobacter sp.* (18.3 mm), *P. mirabilis* (17.3 mm), *P. aeru-*

Table 1 Antibacterial and antiyeast activity of *Terminalia chebula* fruits extract on ear pathogens

Solvent extracts (mg/mL)	Diameter of growth of inhibition zones (mm)					
	Sa	Pm	Pa	Ec	As	Ca
Acetone	31.6 ^{a,1} ± 1.52 ²	21.0 ± 1.0	23.6 ± 1.15	19.3 ± 1.52	24.6 ± 0.57	-
Ethanol	27.3 ± 1.15	19.3 ± 1.15	20.6 ± 1.15	15.6 ± 0.57	21.6 ± 1.15	-
Methanol	28.6 ± 0.57	20.6 ± 0.57	22.3 ± 1.52	16.3 ± 1.52	22.3 ± 1.15	-
Hot aqueous	26.3 ± 1.15	17.3 ± 0.57	16.3 ± 1.15	14.3 ± 1.15	18.3 ± 0.57	-
Cold aqueous	25.6 ± 0.57	15.6 ± 1.52	14.6 ± 1.52	13.6 ± 0.57	17.6 ± 1.52	-
Ciplox ear drop	56.3 ± 0.57	46.3 ± 1.52	34.0 ± 1.0	36.3 ± 0.57	32.6 ± 0.57	NT
Kan pip ear drop	26.3 ± 1.52	20.3 ± 0.57	18.3 ± 1.52	23.6 ± 1.15	21.6 ± 1.52	16.3 ± 1.15
Candid ear drop	NT	NT	NT	NT	NT	21.6 ± 0.57

¹Values, including diameter of the well (8 mm), are means of three replicates, ²± SD. The data were analyzed by one way analysis of variance followed by Dunnett's test. ^a*P* < 0.01 vs positive control. Sa: *Staphylococcus aureus*; Pm: *Proteus mirabilis*; Pa: *Pseudomonas aeruginosa*; Ec: *Escherichia coli*; As: *Acinetobacter sp.*; Ca: *Candida albicans*; NT: Not tested.

Table 2 Minimum inhibitory concentration and minimum bactericidal concentration of *Terminalia chebula* fruits extracts against bacterial ear pathogens

Solvent extract	mg/mL	Sa	Pm	Pa	Ec	As
Acetone	MIC	0.78	12.5	6.25	25	6.25
	MBC	1.56	6.25	6.25	25	6.25
Ethanol	MIC	3.12	25	12.5	50	12.5
	MBC	6.25	25	12.5	> 50	12.5
Methanol	MIC	3.12	12.5	12.5	50	12.5
	MBC	3.12	12.5	25	50	25
Hot aqueous	MIC	3.12	25	25	50	25
	MBC	6.12	50	50	50	25
Cold aqueous	MIC	3.12	25	25	50	25
	MBC	6.12	50	50	> 50	50

Sa: *Staphylococcus aureus*; Pm: *Proteus mirabilis*; Pa: *Pseudomonas aeruginosa*; Ec: *Escherichia coli*; As: *Acinetobacter sp.* MIC: Minimum inhibitory concentration; MBC: Minimum bactericidal concentration.

ginosa (16.3 mm) and *E. coli* (14.3 mm). The antibacterial activity of the acetonic extract was found better than the standard herbal ear drop (Kan pip) against all the tested bacterial ear pathogens (Table 1).

The MIC values ranged between 0.78 mg/mL and 50 mg/mL for the different bacterial ear pathogens and MBC values ranged between 1.56 mg/mL and 50 mg/mL. The results revealed that the MBC values were either equal or twofold higher than the MIC values against the corresponding pathogens. Of all the fruit extracts in different solvents tested, the acetonic extract was the best solvent where the lowest MIC of 0.78 mg/mL and MBC of 1.56 mg/mL was found against *S. aureus* that increased to MIC value of 6.25 mg/mL against *P. aeruginosa*, *P. mirabilis* and *Acinetobacter sp.*, and MIC was 25 mg/mL for *E. coli* (Table 2).

Kan pip ear drop showed the inhibition of all the tested pathogens with a zone of inhibition ranging between 26.3 mm and 16.3 mm but the allopathic antibacterial ear drop ciplox, containing ciprofloxacin and antifungal ear drop candid containing clotrimazole, produced a zone of inhibition ranging between 56.3 mm and 21.6 mm. The antibacterial activity of *T. chebula* acetonic extract was found to be better than the standard herbal ear drop

(Kan pip) against all the tested bacterial ear pathogens. All the obtained results were statistically significant as they showed (*P* < 0.01) compared with control (Table 1).

DISCUSSION

Medicinal plants have been considered a boon to human society to cure a number of ailments^[32]. Several works have documented the pharmacological screening of plant extracts which have been exploited as the source of innumerable therapeutic agents^[33-35]. *T. chebula* is an important medicinal plant in Indian traditional medicine and it is the most frequently used herb in Ayurveda^[36]. Therefore, in the present investigation, different organic (ethanol, methanol, acetone) and aqueous (hot and cold) fruit extracts of this plant were evaluated for their antibacterial and antifungal potential for the first time against the pathogens causing ear infection.

In our study, the organic fruit extracts of *T. chebula* were found to be the most active in inhibiting the growth of all the five tested bacterial ear pathogens compared to aqueous extracts. They showed a broad spectrum of antibacterial activity showing inhibition of gram-positive and gram-negative bacteria. Our work is supported by earlier studies on an alcoholic extract that exhibited greater activity than the aqueous extracts against bacteria^[37,38]. There are several other reports about the antibacterial activity of *T. chebula* fruit extracts against uropathogenic *E. coli*, *Helicobacter pylori* and *S. aureus*, *Salmonella typhi*, *S. epidermidis*, *Bacillus subtilis*, *Proteus vulgaris* and *Pseudomonas aeruginosa*^[7,10,36].

Among the tested bacterial ear pathogens, gram-positive bacterial strains have been found to be more susceptible than gram-negative bacterial strains. This may be attributed to the fact that the cell wall in gram-positive bacteria consists of a single layer, whereas, the gram-negative cell wall is a multilayered structure bounded by an outer cell membrane^[39]. A majority of the described antimicrobial effects of *T. chebula* extracts have been attributed to their secondary metabolites, notably tannins. The antibacterial activities of tannins are well documented and are known to inhibit the growth of many fungi, yeasts, bacteria and viruses^[10].

Of the three organic extracts of this plant screened, the acetonc extract has been found to be more active and have a better antibacterial activity than the corresponding ethanolic and methanolic extracts (Table 1). Our results confirm the finding of Nair *et al.*^[40], Cowan^[41] and El-off^[42], who rated acetone as the best solvent. Interestingly, *T. chebula* extracts have been found to be more potent against the tested ear pathogens compared to the standard herbal ear drop (Kan pip), showing a great potential to be developed as a herbal ear drop to control microbial ear infections.

CONCLUSION

The present investigation revealed that the acetonc extracts of *T. chebula* showed promising antibacterial activity against all the tested bacterial ear pathogens, which explains its use in traditional system of medicines. *T. chebula* can be employed as a source of natural antimicrobials that can serve as an alternative to conventional medicines. However, further experiments, including phytochemical analysis, are needed to identify the active constituents responsible for the observed antibacterial activity and *in vivo* studies on this plant to determine its toxicity and their pharmacokinetics properties, for therapeutic utility in treating otitis externa infections.

ACKNOWLEDGMENTS

The authors are thankful to Dr. Vashishta BD, Chairman, Department of Botany, Kurukshetra University, Kurukshetra, for rendering help in confirmation of the identification of the plant.

COMMENTS

Background

The incidence of otitis externa is mainly found in tropical areas of the world and in most of the patients visiting the ear, nose and throat clinics.

Research frontiers

Presently, most of us are consuming antibiotics for every ailment but misuse and overuse of these antibiotics leads to the development of antimicrobial resistance. So, to combat the antimicrobial resistance, the authors in this study evaluated the antimicrobial potential of a plant compared to the ear drops.

Innovations and breakthroughs

This recent study has highlighted the significance of a herbal plant [*Terminalia chebula* (*T. chebula*) fruit] extract for treating a bacterial ear infection. Fruit extract of *T. chebula* displayed a broad spectrum activity against all the tested bacterial pathogens. This is the first study showing the use of *T. chebula* fruit extracts against the pathogens causing otitis infection.

Applications

This study suggested that the fruit extract of *T. chebula* can be used for treating bacterial otitis infection. This fruit extract can be developed as a herbal ear drop or be used in any other formulation for human beings, after testing its toxicity and pharmacokinetics properties.

Terminology

Otitis externa: refers to a spectrum of infections of the external auditory canal and auricle; Otorrhea: any flowing/drainage/discharge from the ear; *Pruritus*: itching/irritation of the skin; Erythema: redness of the skin.

Peer review

This is a good descriptive study in which the authors evaluated the antibacterial

potential of herbal plants and compare it with locally available ear drops. The results are interesting, with acetonc fruit extracts of *T. chebula* displaying more antibacterial activity compared to Ayurvedic ear drops, Kan pip. Hence, this extract can be used as a therapeutic substance that could be used in the pharmaceutical industry for treating otitis and other infections.

REFERENCES

- 1 Nogueira JC, Melo Diniz Mde F, Lima EO, Lima ZN. Identification and antimicrobial susceptibility of acute external otitis microorganisms. *Braz J Otorhinolaryngol* 2008; **74**: 526-530
- 2 Osguthorpe JD, Nielsen DR. Otitis externa: Review and clinical update. *Am Fam Physician* 2006; **74**: 1510-1516
- 3 Ong YK, Chee G. Infections of the external ear. *Ann Acad Med Singapore* 2005; **34**: 330-334
- 4 Aneja KR, Sharma C, Joshi R. Fungal infection of the ear: a common problem in the north eastern part of Haryana. *Int J Pediatr Otorhinolaryngol* 2010; **74**: 604-607
- 5 Clark WB, Brook I, Bianki D, Thompson DH. Microbiology of otitis externa. *Otolaryngol Head Neck Surg* 1997; **116**: 23-25
- 6 Hawke M, Wong J, Krajden S. Clinical and microbiological features of otitis externa. *J Otolaryngol* 1984; **13**: 289-295
- 7 Sharma A, Meena S and Barman N. Efficacy of ethyl acetate and ether extract of *Terminalia chebula* Retz against some human pathogenic strains. *Inter J Pharm Tech Res* 2011; **3**: 724-727
- 8 Dubey NK, Kumar R, Tripathi P. Global promotion of herbal medicine: India's opportunity. *Curr Sci* 2004; **86**: 37-41
- 9 Chattopadhyay RR, Bhattacharyya SK. *Terminalia chebula*: An update. *Pharmacog Rev* 2007; **1**: 151-156
- 10 Chattopadhyay RR, Bhattacharyya SK, Medda C, Chanda S, Datta S, Pal NK. Antibacterial activity of black myrobalan (Fruit of *Terminalia chebula* Retz.) against uropathogen *Escherichia coli*. *Pharmacog Mag* 2007; **11**: 212-215
- 11 Kirtikar KR, Basu BD. *Terminalia chebula*. In: Indian Medicinal Plants. 2nd ed. Allahabad: Lotit Mohan Basu Publication, 1954: 456-458
- 12 Anonymous. Indian Herbal Pharmacopoeia. Mumbai: Regional Research Laboratory, Jammutwai and Indian Drug Manufacturers Association, 1999: 51
- 13 Khanna AK, Chander R, Kapoor NK, Singh C, Srivastava AK. Hypolipidemic activity of *Terminalia chebula* in rats. *Fitoterapia* 1993; **64**: 351-356
- 14 Kirtikar KR, Basu BD. Indian Materia Medica. Dehradun: International Book distributors, 1987: 333-335
- 15 Lee HS, Jung SH, Yun BS, Lee KW. Isolation of chebulic acid from *Terminalia chebula* Retz. and its antioxidant effect in isolated rat hepatocytes. *Arch Toxicol* 2007; **81**: 211-218
- 16 Tasduq SA, Singh K, Satti NK, Gupta DK, Suri KA, Johri RK. *Terminalia chebula* (fruit) prevents liver toxicity caused by sub-chronic administration of rifampicin, isoniazid and pyrazinamide in combination. *Hum Exp Toxicol* 2006; **25**: 111-118
- 17 Chevallier. Encyclopedia of Medicinal Plants. New York: D.K. Publishing, 1996: 273
- 18 Bruneton J. Pharmacognosy, Phytochemistry, Medicinal Plants. Paris: Lavoisier Publishing, 1995: 333
- 19 Ber RM. Phytosterol in some plants materials. *Indian Sopa J* 1970; **35**: 275-277
- 20 Jayaramkumar K. Effect of geographical variation on content of tannic acid, gallic acid, chebulinic acid, and ethyl galate in *Terminalia chebula* fruits. *Nat Prod* 2006; **2**: 170-175
- 21 Juang LJ, Sheu SJ, Lin TC. Determination of hydrolyzable tannins in the fruit of *Terminalia chebula* Retz. by high-performance liquid chromatography and capillary electrophoresis. *J Sep Sci* 2004; **27**: 718-724
- 22 Creencia E, Eguchi T, Nishimura T, Kakinuma K. Isolation and structure elucidation of the biologically active components of *Terminalia chebula* Retz (Combretaceae). *KIMI-*

- KA 1996; **12**: 1-10
- 23 **Asish P**, Sashi B. Triterpenoids and their glycosides from *Terminalia chebula*. *Phytochem* 1993; **32**: 999-1002
- 24 **Aneja KR**, Sharma C. Antimicrobial potential of fruit extracts of *Elettaria Cardamomum* Maton (Chhoti elaichi) against the pathogens Causing ear infection. *Pharmacologyonl* 2010; **3**: 750-756
- 25 **Lawongsa P**, Boonkerd N, Wongkaew SO, Gara F, Teamroong N. Molecular and phenotypic characterization of potential plant growth-promoting *Pseudomonas* from rice and maize rhizospheres. *World J Microbiol Biotechnol* 2008; **24**: 1877-1884
- 26 **Cappuccino JG**, Sherman N. Microbiology Lab Manual. 7th ed. San Francisco: Benjamin Cummings Publishing Company, 2008
- 27 **Aneja KR**. Experiments in Microbiology, Plant Pathology and Biotechnology. 4th ed. New Delhi: New Age International Publishers, 2003
- 28 **Das K**, Tiwari RKS, Shrivastava DK. Techniques for evaluation of medicinal plant products as antimicrobial agent: Current methods and future trends. *J Med Plants Res* 2010; **4**: 104-111
- 29 **Ncube NS**, Afolayan AJ, Okoh AI. Assessment techniques of antimicrobial properties of natural compounds of plant origin: current methods and future trends. *Afr J Biotechnol* 2008; **7**: 1797-1806
- 30 **Mann A**, Banso A, Clifford LC. An antifungal property of crude plant extracts from *Anogeissus leiocarpus* and *Terminalia avicennioides*. *Tanzan J Health Res* 2008; **10**: 34-38
- 31 **Doughari JH**, Obidah JS. In vitro antifungal activity of stem bark extracts of *Leptadenia lancifolia*. *Int J Integrative Biol* 2008; **3**: 111-117
- 32 **Murray M**. The healing power of Herbs. Rocklin: Prima Publishing, 1995: 162-171
- 33 **Natarajan E**, Senthilkumar S, Francis Xavier T, Kalaiselvi V. Antibacterial activities of leaf extracts of *Alangium salviifolium*. *J Trop Med Plants* 2003; **4**: 9-13
- 34 **Yoshikazu S**, Murata H, Nakanishi T, Inatomi Y. Inhibitory effect of plant extracts on production of Verotoxin by enterohemorrhagic *Escherichia coli* O157: H7. *J Health Sci* 2001; **47**: 473-477
- 35 **Herrera RM**, Perez M, Martin-Herrera DA, Lopez-Garcia R, Rabanal RM. Antimicrobial activity of extracts from plants endemic to the Canary Islands. *Phytother Res* 1996; **10**: 364-366
- 36 **Kannan P**, Ramadevi SR, Waheeta Hopper. Antibacterial activity of *Terminalia chebula* fruit extract. *Afr J Microbiol Res* 2009; **3**: 180-184
- 37 **Ahmad I**, Mehmood Z, Mohammad F. Screening of some Indian medicinal plants for their antimicrobial properties. *J Ethnopharmacol* 1998; **62**: 183-193
- 38 **Phadke SA**, Kulkarni SD. Screening of in vitro antibacterial activity of *Terminalia chebula*, *Eclapta alba* and *Ocimum sanctum*. *Indian J Med Sci* 1989; **43**: 113-117
- 39 **Yao J**, Moellering R. Antibacterial agents. In: Murray P, Baron E, Pfaller M, Tenover F, Tenover F, editors. Manual of clinical microbiology. Washington: ASM, 1995: 1281-1290
- 40 **Nair R**, T Kalariya, C Sumitra. Antibacterial activity of some selected Indian medicinal plants. *Tur J Biol* 2005; **29**: 41-47
- 41 **Cowan MM**. Plant products as antimicrobial agents. *Clin Microbiol Rev* 1999; **12**: 564-582
- 42 **Eloff JN**. Which extractant should be used for the screening and isolation of antimicrobial components from plants? *J Ethnopharmacol* 1998; **60**: 1-8

S- Editor Wu X L- Editor Roemmele A E- Editor Zhang DN