

World Journal of *Respirology*

World J Respirol 2023 May 26; 12(1): 1-15



MINIREVIEWS

- 1 Monoclonal antibody for COVID-19: Unveiling the recipe of a new cocktail
Bajpai J, Kant S, Verma AK, Pradhan A

CASE REPORT

- 10 Pulmonary arterial hypertension confirmed by right heart catheterization following COVID-19 pneumonia: A case report and review of literature
Henriques King M, Ogbuka IC, Bond VC

ABOUT COVER

Peer Reviewer of *World Journal of Respiriology*, Deven Juneja, DNB, FNB, EDIC, FCCM. Director, Institute of Critical Care Medicine, Max Super Speciality Hospital, Saket, New Delhi - 110092, India. deven.juneja@maxhealthcare.com

AIMS AND SCOPE

The primary aim of *World Journal of Respiriology* (WJR, *World J Respirol*) is to provide scholars and readers from various fields of respirology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online. WJR mainly publishes articles reporting research results and findings obtained in the field of respirology and covering a wide range of topics including bronchial diseases, ciliary motility disorders, respiratory tract granuloma, laryngeal diseases, lung diseases, nose diseases, pleural diseases, respiration disorders, respiratory center abnormalities, neuromuscular disorders, respiratory hypersensitivity, respiratory system abnormalities, pulmonary vascular diseases, respiratory tract fistula, respiratory tract infections, respiratory tract neoplasms, thoracic diseases, and tracheal diseases.

INDEXING/ABSTRACTING

The WJR is now abstracted and indexed in Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Si Zhao; Production Department Director: Xiang Li; Editorial Office Director: Yu-Jie Ma.

NAME OF JOURNAL

World Journal of Respiriology

ISSN

ISSN 2218-6255 (online)

LAUNCH DATE

December 30, 2011

FREQUENCY

Continuous Publication

EDITORS-IN-CHIEF

Kazuhiro Yamaguchi

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2218-6255/editorialboard.htm>

PUBLICATION DATE

May 26, 2023

COPYRIGHT

© 2023 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Monoclonal antibody for COVID-19: Unveiling the recipe of a new cocktail

Jyoti Bajpai, Surya Kant, Ajay Kumar Verma, Akshyaya Pradhan

Specialty type: Infectious diseases

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0

Grade B (Very good): 0

Grade C (Good): C

Grade D (Fair): D

Grade E (Poor): 0

P-Reviewer: Arteaga-Livias K, Peru; Roton JC, Italy

Received: February 3, 2023

Peer-review started: February 3, 2023

First decision: February 21, 2023

Revised: March 3, 2023

Accepted: May 8, 2023

Article in press: May 8, 2023

Published online: May 26, 2023



Jyoti Bajpai, Surya Kant, Ajay Kumar Verma, Department of Respiratory Medicine, King George's Medical University, Lucknow 226003, India

Akshyaya Pradhan, Department of Cardiology, King George's Medical University, Lucknow 226003, India

Corresponding author: Jyoti Bajpai, MD, Assistant Professor, Doctor, Department of Respiratory Medicine, King George's Medical University, E 1204 Shalimar Gallant Mahanagar, Lucknow 226003, India. jyotibajpai33@gmail.com

Abstract

The coronavirus disease 2019 (COVID-19) pandemic has had a tremendous adverse impact on the global health system, public sector, and social aspects. It is unarguably the worst pandemic of the century. However, COVID-19 management is a mystery in front of us, and an authentic treatment is urgently needed. Various repurposed drugs, like ivermectin, remdesivir, tocilizumab, baricitinib, etc., have been used to treat COVID-19, but none are promising. Antibody therapy and their combinations are emerging modalities for treating moderate COVID-19, and they have shown the potential to reduce hospitalisations. One antibody monotherapy, bamlanivimab, and two cocktails, casirivimab/imdevimab and bamlanivimab/esterivimab, have received authorization for emergency use by the United States Food and Drug Administration for the treatment of mild COVID-19 in high risk individuals. The European Emergency has made similar recommendations for use of the drug in COVID-19 patients without oxygen therapy. This brief review will focus on monoclonal antibodies and their combination cocktail therapy in managing COVID-19 infection.

Key Words: SARS-CoV-2; Mild COVID-19; Antibodies; Risk factors

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: The coronavirus disease 2019 (COVID-19) pandemic is a severe public health emergency that necessitated the rapid development of novel medicines and viral detection technologies. Monoclonal antibodies against the receptor-binding domain of the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) spike protein have become an important target for the creation of therapeutic antibodies. The use of antibody cocktails is anticipated to be a key component of an efficient COVID-19 treatment plan because SARS-CoV-2 has a high mutation rate, particularly when subjected to the selection pressure of aggressively applied preventive vaccinations and neutralising antibodies.

Citation: Bajpai J, Kant S, Verma AK, Pradhan A. Monoclonal antibody for COVID-19: Unveiling the recipe of a new cocktail. *World J Respir* 2023; 12(1): 1-9

URL: <https://www.wjgnet.com/2218-6255/full/v12/i1/1.htm>

DOI: <https://dx.doi.org/10.5320/wjr.v12.i1.1>

INTRODUCTION

The coronavirus disease 2019 (COVID-19) pandemic has placed a high burden on healthcare systems globally[1]. The first case of COVID-19 was reported on January 30, 2020. As of July 20, 2021, India had the highest number of COVID-19 instances, with more than 30 million[2]. The second wave of COVID-19 was more severe than the first. There was a reported shortage of drugs, oxygen, hospital beds, and vaccines. Some patients with COVID-19 will develop acute disease and multiorgan complications, but there are currently no proven therapeutics to prevent or reduce COVID-19 related hospitalisations, complications, or mortality.

Various drugs are approved for patients hospitalised with severe COVID-19 infection, but only a few drugs for mild COVID-19 patients who are not sick enough to be hospitalised are available[3]. Monoclonal antibodies (mAbs) are a new treatment for mild COVID-19 outpatients with a high risk. Recently, cocktail therapy has been approved for mild to moderate COVID-19 patients. Immunity to viral infection is a multipronged response, comprising the innate response-which restricts viral replication and creates an antiviral state in the local tissue environment, the adaptive response- in which virus-specific CD4+ T cells, CD8+ T cells, and the antibodies produced by B cells to control and clear the infection and generate immune memory. COVID-19 appears to evade or delay the innate immune response. If adaptive immunity is delayed for too long (because of efficient viral evasion, diminished innate immunity in the patient, or both), then the inability to control infection puts patients at an increased risk for severe or even fatal COVID-19 disease[4].

Despite recent studies demonstrating immune responses to COVID-19 as far as up to 8 mo after symptom onset, much remains to be learned about post-infection immunity to COVID-19[5,6]. Immunity to seasonal human coronaviruses is usually of short duration, and reinfection has been documented in patients who have already been infected with COVID-19[7]. Moreover, some individuals might not benefit from vaccination, as the vaccine trials published to date have not shown 100% efficacy, and real-world experience has demonstrated breakthrough events[8,9]. Furthermore, large parts of the population are still not vaccinated primarily because of supply issues and in part because of vaccine hesitancy. Thus, there is a pressing need for alternative therapies for COVID-19 patients. This article reviews the currently approved cocktail therapy(-ies) in the management of COVID-19 disease.

WHAT ARE MABS?

An antibody is a protein molecule naturally developed by the immune response to infection. Antibodies are an essential factor in immunity against most viral diseases. Monoclonal antibodies (mAbs) are a single isotype with a defined specificity targeting with high potency a particular antigen *via* the antigen-binding fragment. As such, mAbs against COVID-19 have been derived from plasma donated by patients who recovered from COVID-19[10]. Polyclonal antibodies are usually defined as a mixture of diverse antibodies with mixed affinities for their targets. However, in the world of COVID-19 therapeutics, the term “polyclonal antibodies” is more descriptive of convalescent plasma with several antibody components[11].

mAbs are designed in a laboratory and mimic the natural immune system in response to infection. They are created for a specific target of infectious particles. A mAb is produced by exposing white blood cells to a particular viral protein cloned to produce antibodies to treat several infections and cancers [12]. mAbs that bind to the spike (S) protein of the COVID-19 virus stop the virus from binding the angiotensin-converting enzyme II (ACE2) receptor of human cells and prevent its invasion and replication[13]. mAbs have been effective against new COVID-19 variants B.1.1.7.

Even though more than 75 mAbs have been licensed by the United States Food and Drug Administration (FDA), only three are used to treat or prevent infectious diseases like anthrax, respiratory syncytial virus, and *Clostridium difficile*, and two are used to treat Ebola virus diseases. mAbs are intended for patients recently diagnosed with COVID-19 who are not very sick and have risk factors for severe infection[14-16].

This article focuses on mAbs with neutralising activity against severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), which work by targeting the receptor binding domain (RBD) of the viral S protein, thereby preventing viral attachment to the ACE2 receptor and preventing a critical step in viral entry and infection. Bamlanivimab is a recombinant, neutralising human immunoglobulin G1 (IgG1) mAb effective against the S protein of COVID-19. Etesevimab is a recombinant, fully human monoclonal neutralising antibody that binds to the surface S protein receptor-binding domain with high affinity and blocks viral binding to the ACE2 receptor of the host cell surface. Imdevimab and Casirivimab are IgG1 that act against SARS-CoV-2 S protein. Thus, the antibody cocktail thwarts the attachment of the virus and its entry into the human cell.

MABS THERAPY: SCIENTIFIC EVIDENCE

Casirivimab (REGN10933) and imdevimab (REGN 10987) were developed by Regeneron F and Hoffmann-La Roche Ltd pharmaceuticals, respectively, and bamlanivimab and the cocktail of bamlanivimab and etesevimab were developed by Eli Lilly and AbCellera, respectively. The following studies or their scientific data reveal how these mAbs are helpful in treating COVID-19 (Table 1).

BLAZE-1

The BLAZE 1 trial was a phase 2/3 trial that enrolled 452 ambulatory COVID-19 patients and was given bamlanivimab as one of three doses [bamlanivimab (LYCoV555) -700 mg, 2800 mg, or 7000 mg in intravenous (IV) infusion] or placebo. The quantitative virologic endpoints and clinical outcomes were assessed[17]. The immediate result was the change in the viral load by day 11. For patients who received a 2800 mg (middle) antibody dose, viral load decreased by a factor of 3.4. The patients who received the 700 mg (lower) dose or the 7000 mg (higher) amount showed a more negligible difference from the placebo in the viral load change from baseline. In addition, bamlanivimab antibody therapy resulted in fewer hospitalisations and/or emergency room visits (1.9% in 2800 mg treatment group compared to 6.3% in the placebo group).

ACTIVE-3

The ACTIVE -3 trial enrolled 314 (163 drug group and 151 placebo group) hospitalised COVID-19 patients without end organ failure[18]. All the patients were also on supportive care as background therapy, including an antiviral drug and when indicated, supplemental oxygen and glucocorticoids. Bamlanivimab at a 7000 mg or placebo dose was administered as a single IV infusion over 1 h. The results revealed that mAbs when administered with remdesivir did not show efficacy among hospitalised COVID-19 patients without end organ failure.

BLAZE-2

This randomised phase 3 clinical trial enrolled 966 residents and staff at a United States nursing facility with at least one confirmed COVID-19 index case and who were negative at baseline for COVID-19 infection and serology. The incidence of COVID-19 disease among those treated with the antibody bamlanivimab *vs* placebo (8.5% *vs* 15.2% respectively) was lower[19]. Bamlanivimab monotherapy compared with a placebo reduced the risk of COVID-19 in residents and staff of nursing facilities.

COCKTAIL THERAPY

Bamlanivimab and etesevimab

The BLAZE-1 phase 3 trial showed that the cocktail of bamlanivimab and etesevimab was associated with a significant reduction in viral load compared to the placebo. In contrast, bamlanivimab monotherapy did not result in a substantial reduction. The cocktail was also shown to reduce the number of hospitalisations. The trial included 518 patients in the treatment arm who received a single infusion of bamlanivimab 2800 mg and etesevimab 2800 mg together, and 517 patients received a placebo[20]. The primary endpoint was COVID-19 related hospitalisations or death by any cause during the 29-d follow-up. Hospitalisation or death occurred in 36 (7%) patients who received a placebo compared to 11 (2%) patients treated with bamlanivimab and etesevimab together (a 70% reduction). Ten deaths (2%) deaths occurred in the placebo group. Thus, all-cause death was significantly lower in the bamlanivimab and etesevimab group compared to the placebo group. The United States FDA granted Emergency Use Authorisation (EUA) for the 700 mg dose of bamlanivimab for ambulatory

Table 1 Properties of different studies of antibodies therapy in coronavirus disease 2019

Ref.	Study type	Dose and duration	Primary outcome	Secondary outcomes	Primary result	Additional characteristics	Adverse effects
Chen <i>et al</i> [17], BLAZE-1	Randomised, double-blind, placebo-controlled, single-dose trial	Total 452 patients; 309 in the bamlanivimab (LY-CoV555) group and 143 in the placebo group. mAb at doses of 700 mg, 2800 mg, and 7000 mg and placebo administered within 3 d after positive SARS-CoV-2 results	The change from baseline to day 11 (4 d) in SARS-CoV-2 viral load	COVID-19 related inpatient hospitalisation, a visit to the emergency department, death, safety, symptom severity, and time points for viral clearance	The viral load at day 11 was lower in patients who received 2800 mg drug compared to the placebo group	High-risk subgroups (an age of ≥ 65 yr or a BMI of ≥ 35), the percentage of hospitalisation was 4.2% in the LY-CoV555 group and 14.6% in the placebo group	Serious adverse events occurred in none of the patient treatment groups, diarrhoea was reported in 3.2% of the patients
Weinreich <i>et al</i> [23], REGN-COV2	Double-blind, phase 1-3 trial, 275 (1:1:1) non-hospitalised patients with COVID-19	REGN-COV2 is a combination of casirivimab (REGN10933) and imdevimab (REGN10987). Among the 275 patients, 90 were assigned to receive high-dose (8.0 g), 92 to receive low-dose (2.4 g), and 93 to receive placebo	The time-weighted average change in viral load from baseline (day 1) through day 7	The percentage of patients with at least one COVID-19 related medically attended visit through day 29	REGN-COV2 enhanced clearance of virus, particularly in patients in whom an endogenous immune response had not yet been initiated	The median age was 44.0 yr, 49% were male, 13% identified as Black or African American, and 56% as Hispanic or Latino	In this interim analysis, both REGN-COV2 doses (2.4 g and 8.0 g) were associated with few and low-grade toxic effects (1%) in the combined REGN-COV2 dose groups
Gottlieb <i>et al</i> [20], BLAZE 1	Multipart, 49 United States centres including phase 2/3, randomised, double-blind, placebo-controlled, single-infusion study (BLAZE-1) ambulatory patients ($n = 613$) and had one or more mild to moderate symptom	Patients were randomised to receive a single infusion of bamlanivimab [700 mg ($n = 101$), 2800 mg ($n = 107$), or 7000 mg ($n = 101$)], the combination treatment (2800 mg of bamlanivimab and 2800 mg of etesevimab ($n = 112$), or placebo ($n = 156$)	Change in SARS-CoV-2 log viral load at day 11 (± 4 d)	A total of nine prespecified secondary outcome measures were evaluated. Three focused on viral load (time to viral clearance; the proportion of patients with viral support at days 7, 11, 15, and 22; time to symptom improvement; time to symptom resolution; and the balance of patients showing symptom improvement or resolution at days 7, 11, 15, and 22), and 1 focused on clinical outcomes (the proportion of patients with a COVID-19 related hospitalisation, emergency department visit, or death) at day 29	Among the 577 patients who were randomised and received an infusion, 533 (92.4%) completed the efficacy evaluation period (day 29). The change in log viral load from baseline at day 11 was -3.72 for 700 mg, -4.08 for 2800 mg, -3.49 for 7000 mg, -4.37 for combination treatment, and -3.80 for placebo	The mean age of patients was 44.7 ± 15.7 yr. A total of 315 patients (54.6%) were female, 245 patients (42.5%) identified as Hispanic, and 387 patients (67.1%) had at least one risk factor for severe COVID-19 (aged ≥ 55 yr, BMI ≥ 30 , or ≥ 1 relevant comorbidity such as hypertension)	Immediate hypersensitivity reactions were reported in 9 patients (6 bamlanivimab, 2 combination treatment, and 1 placebo). No deaths occurred during the study treatment
BLAZE-2 [21]	Randomised, double-blind, single-dose, phase 3 placebo-controlled trial, 966 participants (300 residents and 666 staff) who tested negative for SARS-CoV-2 at baseline	Bamlanivimab 4200 mg or placebo only if a nursing home recorded at least one confirmed case of SARS-CoV-2 infection among residents or facility staff from a sample collected within the last 7 d	To find incidence of COVID-19, defined as the detection of SARS-CoV-2 by reverse transcriptase-PCR and mild or worse disease severity within 21 d of detection, within 8 wk of randomisation	To find incidence of moderate or worse COVID-19 severity and incidence of SARS-CoV-2 infection	Bamlanivimab significantly reduced the incidence of COVID-19 in the prevention population compared with placebo [8.5% vs 15.2%; odds ratio: 0.43 (95%CI: 0.28-0.68); $P < 0.001$]; absolute risk difference, -6.6 (95%CI: -10.7 to -2.6 percentage points)	Significantly reduced the incidence of moderate or worse COVID-19 compared with placebo (8.3% vs 14.1%)	The rate of participants with adverse events was 20.1% in the bamlanivimab group and 18.9% in the placebo group. The most common adverse events were urinary tract infection (2.0%) in bamlanivimab and (2.4%) placebo and hypertension (1.2%) in bamlanivimab and (1.7%) placebo
Lundgren <i>et al</i> [18],	Randomised, double-blind, placebo-	Hospitalised COVID-19 patients ($n = 314$) without	A sustained recovery, as assessed in a time-to-	Death from any cause	Hospitalised patients with COVID-19 who received mAb	The majority of patients had hypoxemia and tested the	Serious adverse events (19%) in the LY-CoV555 group and

ACTIVE-3	controlled trial	end organ failure, single infusion of the neutralising mAb antibody LY-CoV555 (at a dose of 7000 mg)	event analysis, through day 90 as well as two ordinal outcomes that were measured at day 5		did not have better clinical outcomes at day 5 than those who received placebo	effect of LY-CoV555 on a background of remdesivir and substantial glucocorticoid therapy	(14%) in the placebo
REGN-COV2067 ²⁴	Phase (I-III) adaptive randomised placebo control double-blind	COVID-19 in infected non-hospitalised patients (<i>n</i> = 4567; REGN-COV2067)	1200 mg cocktail (<i>n</i> = 736), placebo (<i>n</i> = 748), and another group cocktail dose 2400 mg IV (<i>n</i> = 1355), placebo (<i>n</i> = 1341)	Clinically significant effect on risk of COVID-19 hospitalisation or all-cause death in high-risk non-hospitalised patients and confirm safety	Cocktail of casirivimab and imdevimab significantly reduced the risk of hospitalisation or death by 70% (1200 mg IV) and 71% (2400 mg IV) compared to placebo	Cocktail therapy reduced symptom duration from 14 d to 10 d (median numbers)	Not mentioned

BMI: Body mass index; COVID-19: Coronavirus disease 2019; IV: intravenous; mAb: Monoclonal antibody; SARS-CoV-2: Severe acute respiratory syndrome coronavirus 2.

COVID-19 patients at high risk[21,22].

The EUA advised the population on the benefits of monotherapy despite uncertainties[21,22]. The authorised dosage of 700 mg bamlanivimab and 1400 mg etesevimab administered together was based on analyses of available preclinical, clinical, and virologic data as well as pharmacokinetic and pharmacodynamic modelling, which supported that the authorised dosage was expected to have a similar clinical and virologic effect to 2800 mg bamlanivimab and 2800 mg etesevimab administered together.

Casirivimab and imdevimab

REGN-COV-2: The mAbs casirivimab and imdevimab bind to the non-overlapping portion of the RBD. A phase 1/2/3 trial (NCT04425629) is taking place across several countries. The phase 3 trial results have been reported. The trial enrolled 4576 patients with one risk factor for severe COVID-19, and an IV infusion of 1200 mg or 2400 mg casirivimab/imdevimab *vs* placebo was given. The trial reached its primary outcome and depicted that the casirivimab and imdevimab cocktail significantly reduced the risk of hospitalisation or death by 70% in the 1200 mg dose arm and 71% in the 2400 mg dose arm; both were significant compared with placebo[23]. In addition, benefits in the secondary outcomes were also found, including a 4-d reduction in the median duration of symptoms *vs* placebo.

Interim data from the first 275 patients (phase 1/2 portion) revealed that the cocktail showed virological efficacy resulting in an overall reduction in viral load of 0.25 log₁₀ RNA copies/mL (95%CI: 0.60, 0.10) for the 2400 mg dose and a reduction of 0.56 log₁₀ RNA copies/mL (95%CI: 0.91, 0.21) for an 8000 mg dose (combined dose reduction was 0.41 log₁₀ RNA copies/mL, 95%CI: 0.71, 0.10) *vs* placebo at day 7.

No data on infectious virus titres or time to the cessation of viral shedding endpoints have been reported, similar to the situation with bamlanivimab or other mAb studies. An ongoing dose-ranging phase 2 companion trial in low-risk symptomatic or asymptomatic non-hospitalised patients with COVID-19 (NCT04666441) showed significant and comparable viral load reductions in casirivimab/imdevimab doses ranging from 300 mg to 2400 mg delivered *via* IV or subcutaneously (SC). The casirivimab/imdevimab cocktail has received EUA by the United States FDA for the treatment of ambulatory patients with mild to moderate COVID-19 and a high risk of hospitalisation, and the EUA

has similarly recommended casirivimab/imdevimab for use in COVID-19 patients who do not require supplemental oxygen and who are at increased risk of progressing to severe COVID-19 (Figure 1).

A trial showed that SC injection of the antibody cocktail casirivimab and imdevimab reduced the risk of symptomatic COVID-19 infection by 81% in household contact with an infected person without COVID-19 antibodies. The trial was conducted by the National Institute of Allergy and Infectious Disease. The individuals treated with the cocktail therapy with symptoms were relieved in 1 wk compared to 3 wk in placebo. The FDA has given EUA for the use of casirivimab and imdevimab antibody combination for the treatment of mild to moderate COVID-19 in adults and children over 12 years and weigh more than 40 kg who are at high risk for progressing to severe disease/hospitalisation. The trial showed that the cocktail antibodies casirivimab and imdevimab were more effective when given as early as possible[24]. The Indian drug regulatory body, the Central Drugs Standards Control Organisation, has recently approved the cocktail regimen for use in the country in the fight against COVID-19. The drug, marketed by Cipla Inc. in India, is currently in vogue for clinical use[25].

Another contemporary mAb has been evaluated and is primarily targeted against the COVID-19 S protein. Sotrovimab, a mAb, also blocks the attachment and viral entry into the host cell. A phase 1/2/3 double-blind placebo-controlled trial enrolled 583 non-hospitalised mild to moderate COVID-19 adult patients. Of these, 291 received sotrovimab, and the rest received a placebo within 5 d of symptoms. The primary endpoint was hospitalisation or death through day 29. The result showed 21 (7%) patients were hospitalised or died in the placebo arm compared to 3 (1%) patients in the sotrovimab group. An 85% reduction in hospitalisation or death in the treatment group was observed[26]. Sotrovimab showed activity against the current variants reported in the United Kingdom, South Africa, Brazil, California, New York, and India. The EUA recommends a 500-mg single IV dose of sotrovimab for non-hospitalised mild to moderate COVID-19 patients[27].

WHICH GROUPS ARE SUITABLE FOR MAB?

mAb therapies have shown promise for treating non-hospitalised patients with mild to moderate COVID-19. The EUA recommends that mAb treatment be given within 10 d of symptom onset or as early as 72 h of positive COVID-19 result. However, treatment should begin as early as possible to mitigate viral proliferation. In the REGN-COV2 study, the effect of REGN-COV2 on viral load was most pronounced among patients with a negative serum antibody test result at baseline. Furthermore, most trials administered mAb treatment within 3 d of a positive COVID-19 test result and a median of 3-4 d after symptom onset. Altogether, these studies suggest that early mAb treatment is more efficacious than the later treatment for COVID-19 patients. Indeed, by the time a patient reaches the lung injury phase of infection, it is too late for mAb treatment to be effective, as suggested by the results from the ACTIV-3 study (Figure 2).

Route, dose, and cost of mAbs

A 600 mg of each or a combined 1200 mg of the casirivimab and imdevimab cocktail has been approved for administration. This can be given either IV or SC. The administration of a total dose of cocktail antibody takes around 30 min. The patient should be kept on observation for 1 h to check for any adverse effects. The price for a dose of 1200 mg cocktail (600 mg of casirivimab and 600 mg of imdevimab) is INR 59750 (700 USD approx). This drug should be stored at 2-8 °C.

Efficacy and safety

The clinical efficacy and safety profiles do not differ between mAb monotherapy and cocktails. Yet, monotherapy *vs* combination therapy is particularly relevant given the emergence of variant strains from the United Kingdom, South Africa, Brazil, California, New York, and India. The results from one study suggested that a mAb cocktail, particularly one combining antibodies that bind distinct and non-overlapping regions, can minimise mutational escape[28]. More importantly, viral mutations can reduce the effectiveness of mAb monotherapy. A recent preprint publication reported that bamlanivimab and casirivimab are effective against the South African variant. Several variants have been labelled by the Centers for Disease Control and Prevention as “variants of concern” because the mutations they carry increase transmission, increase disease severity, and reduce the efficacy of mAb therapy and vaccinations.

CONCLUSION

There is growing evidence that mAb treatment is effective, safe, and well-tolerated. Patients should know that mAb treatment is available to all patients at a high cost in India and that mAb treatment should be started within 72 h of a positive COVID-19 test result to affect the clinical course of COVID-19. Further studies on mAb efficacy and safety in different patient populations (*e.g.*, young children and




Figure 1 Contraindications of monoclonal antibody therapy. COVID-19: Coronavirus disease 2019.

> 65 years of age	> 55 years of age any of the following risk factor	> 12 years of age any of the following risk factor	12-17 years of age with any of the following risk factors
<ul style="list-style-type: none"> No additional risk factors required 	<ul style="list-style-type: none"> CVD HTN COPD/other 	<ul style="list-style-type: none"> BMI > 40 Kg/M² CKD Diabetes Immunosuppressive disorder and drugs 	<ul style="list-style-type: none"> Sickle cell disorder Neurodevelopmental disorder Congenital heart disease Asthma and chronic respiratory diseases BMI > 85th percentile for their age and gender

DOI: 10.5320/wjr.v12.i1.1 Copyright ©The Author(s) 2023.

Figure 2 High-risk groups indicated for treatment with monoclonal antibodies. CVD: Cardiovascular disease; HTN: Hypertension; COPD: Chronic obstructive pulmonary disease; BMI: Body mass index; CKD: Chronic kidney disease.

pregnant women) are needed.

FOOTNOTES

Author contributions: Bajpai J conceptualised the article design; Bajpai J, Pradhan A, and Verma AK searched the literature; Bajpai J and Pradhan A drafted the manuscript; Critical revision was conducted by Kant S, Pradhan A, and Bajpai J.

Conflict-of-interest statement: The authors declare that they have no conflicts of interest.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: India

ORCID number: Jyoti Bajpai 0000-0001-6337-856X; Surya Kant 0000-0001-7520-5404; Ajay Kumar Verma 0000-0002-2973-1793; Akshyaya Pradhan 0000-0002-2360-7580.

S-Editor: Zhang H

L-Editor: Filipodia

P-Editor: Zhao S

REFERENCES

- 1 **Bezemer GFG**, Garssen J. TLR9 and COVID-19: A Multidisciplinary Theory of a Multifaceted Therapeutic Target. *Front Pharmacol* 2020; **11**: 601685 [PMID: 33519463 DOI: 10.3389/fphar.2020.601685]
- 2 Corona virus update (Live). [cited 21 March 2021]. In: Worldometer [Internet]. Available from <http://www.worldometer.info/coronavirus/>
- 3 **Rotondo JC**, Martini F, Maritati M, Mazziotta C, Di Mauro G, Lanzillotti C, Barp N, Gallerani A, Tognon M, Contini C. SARS-CoV-2 Infection: New Molecular, Phylogenetic, and Pathogenetic Insights. Efficacy of Current Vaccines and the Potential Risk of Variants. *Viruses* 2021; **13** [PMID: 34578269 DOI: 10.3390/v13091687]
- 4 **Sette A**, Crotty S. Adaptive immunity to SARS-CoV-2 and COVID-19. *Cell* 2021; **184**: 861-880 [PMID: 33497610 DOI: 10.1016/j.cell.2021.01.007]
- 5 **Dan JM**, Mateus J, Kato Y, Hastie KM, Yu ED, Faliti CE, Grifoni A, Ramirez SI, Haupt S, Frazier A, Nakao C, Rayaprolu V, Rawlings SA, Peters B, Krammer F, Simon V, Saphire EO, Smith DM, Weiskopf D, Sette A, Crotty S. Immunological memory to SARS-CoV-2 assessed for up to 8 months after infection. *Science* 2021; **371** [PMID: 33408181 DOI: 10.1126/science.abf4063]
- 6 **Rotondo JC**, Martini F, Maritati M, Caselli E, Gallenga CE, Guarino M, De Giorgio R, Mazziotta C, Tramarin ML, Badiale G, Tognon M, Contini C. Advanced Molecular and Immunological Diagnostic Methods to Detect SARS-CoV-2 Infection. *Microorganisms* 2022; **10** [PMID: 35744711 DOI: 10.3390/microorganisms10061193]
- 7 **Poland GA**, Ovsyannikova IG, Kennedy RB. SARS-CoV-2 immunity: review and applications to phase 3 vaccine candidates. *Lancet* 2020; **396**: 1595-1606 [PMID: 33065034 DOI: 10.1016/S0140-6736(20)32137-1]
- 8 **Polack FP**, Thomas SJ, Kitchin N, Absalon J, Gurtman A, Lockhart S, Perez JL, Pérez Marc G, Moreira ED, Zerbini C, Bailey R, Swanson KA, Roychoudhury S, Koury K, Li P, Kalina WV, Cooper D, Frenck RW Jr, Hammitt LL, Türeci Ö, Nell H, Schaefer A, Ünal S, Tresnan DB, Mather S, Dormitzer PR, Şahin U, Jansen KU, Gruber WC; C4591001 Clinical Trial Group. Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *N Engl J Med* 2020; **383**: 2603-2615 [PMID: 33301246 DOI: 10.1056/NEJMoa2034577]
- 9 **Lin DY**, Baden LR, El Sahly HM, Essink B, Neuzil KM, Corey L, Miller J; COVE Study Group. Durability of Protection Against Symptomatic COVID-19 Among Participants of the mRNA-1273 SARS-CoV-2 Vaccine Trial. *JAMA Netw Open* 2022; **5**: e2215984 [PMID: 35675078 DOI: 10.1001/jamanetworkopen.2022.15984]
- 10 **Amit S**, Regev-Yochay G, Afek A, Kreiss Y, Leshem E. Early rate reductions of SARS-CoV-2 infection and COVID-19 in BNT162b2 vaccine recipients. *Lancet* 2021; **397**: 875-877 [PMID: 33610193 DOI: 10.1016/S0140-6736(21)00448-7]
- 11 **Marovich M**, Mascola JR, Cohen MS. Monoclonal Antibodies for Prevention and Treatment of COVID-19. *JAMA* 2020; **324**: 131-132 [PMID: 32539093 DOI: 10.1001/jama.2020.10245]
- 12 **Pantaleo G**, Correia B, Fenwick C, Joo VS, Perez L. Antibodies to combat viral infections: development strategies and progress. *Nat Rev Drug Discov* 2022; **21**: 676-696 [PMID: 35725925 DOI: 10.1038/s41573-022-00495-3]
- 13 **Ali MG**, Zhang Z, Gao Q, Pan M, Rowan EG, Zhang J. Recent advances in therapeutic applications of neutralizing antibodies for virus infections: an overview. *Immunol Res* 2020; **68**: 325-339 [PMID: 33161557 DOI: 10.1007/s12026-020-09159-z]
- 14 **Romero JR**. Palivizumab prophylaxis of respiratory syncytial virus disease from 1998 to 2002: results from four years of palivizumab usage. *Pediatr Infect Dis J* 2003; **22**: S46-S54 [PMID: 12671452 DOI: 10.1097/01.inf.0000053885.34703.84]
- 15 FDA Approves Treatment for Ebola Virus. 2020. [cited 30 March 2021]. In: US Food and Drug Administration [Internet]. Available from: <https://www.fda.gov/drugs/drug-safety-and-availability/fda-approves-treatment-ebola-virus>
- 16 **Mulangu S**, Dodd LE, Davey RT Jr, Tshiani Mbaya O, Proschan M, Mukadi D, Lusakibanza Manzo M, Nzolo D, Tshomba Oloma A, Ibanda A, Ali R, Coulibaly S, Levine AC, Grais R, Diaz J, Lane HC, Muyembe-Tamfum JJ; PALM Writing Group, Sivahera B, Camara M, Kojan R, Walker R, Digheero-Kemp B, Cao H, Mukumbayi P, Mbala-Kingebeni P, Ahuka S, Albert S, Bonnett T, Crozier I, Duvenhage M, Proffitt C, Teitelbaum M, Moench T, Aboulhab J, Barrett K, Cahill K, Cone K, Eckes R, Hensley L, Herpin B, Higgs E, Ledgerwood J, Pierson J, Smolskis M, Sow Y, Tierney J, Sivapalasingam S, Holman W, Gettlinger N, Vallée D, Nordwall J; PALM Consortium Study Team. A Randomized, Controlled Trial of Ebola Virus Disease Therapeutics. *N Engl J Med* 2019; **381**: 2293-2303 [PMID: 31774950 DOI: 10.1056/NEJMoa1910993]
- 17 **Chen P**, Nirula A, Heller B, Gottlieb RL, Boscia J, Morris J, Huhn G, Cardona J, Mocherla B, Stosor V, Shawa I, Adams AC, Van Naarden J, Custer KL, Shen L, Durante M, Oakley G, Schade AE, Sabo J, Patel DR, Klekotka P, Skovronsky DM; BLAZE-1 Investigators. SARS-CoV-2 Neutralizing Antibody LY-CoV555 in Outpatients with Covid-19. *N Engl J Med* 2021; **384**: 229-237 [PMID: 33113295 DOI: 10.1056/NEJMoa2029849]
- 18 **ACTIV-3/TICO LY-CoV555 Study Group**, Lundgren JD, Grund B, Barkauskas CE, Holland TL, Gottlieb RL, Sandkovsky U, Brown SM, Knowlton KU, Self WH, Files DC, Jain MK, Benfield T, Bowdish ME, Leshnower BG, Baker JV, Jensen JU, Gardner EM, Ginde AA, Harris ES, Johansen IS, Markowitz N, Matthay MA, Østergaard L, Chang CC, Davey VJ, Goodman A, Higgs ES, Murray DD, Murray TA, Paredes R, Parmar MKB, Phillips AN, Reilly C, Sharma S, Dewar RL, Teitelbaum M, Wentworth D, Cao H, Klekotka P, Babiker AG, Gelijns AC, Kan VL, Polizzotto MN, Thompson BT, Lane HC, Neaton JD. A Neutralizing Monoclonal Antibody for Hospitalized Patients with Covid-19. *N Engl J Med* 2021; **384**: 905-914 [PMID: 33356051 DOI: 10.1056/NEJMoa2033130]
- 19 **Cohen MS**, Nirula A, Mulligan MJ, Novak RM, Marovich M, Yen C, Stemmer M, Mayer SM, Wohl D, Brengle B, Montague BT, Frank I, McCulloh RJ, Fichtenbaum CJ, Lipson B, Gabra N, Ramirez JA, Thai C, Chege W, Gomez

- Lorenzo MM, Sista N, Farrior J, Clement ME, Brown ER, Custer KL, Van Naarden J, Adams AC, Schade AE, Dabora MC, Knorr J, Price KL, Sabo J, Tuttle JL, Klekotka P, Shen L, Skovronsky DM; BLAZE-2 Investigators. Effect of Bamlanivimab vs Placebo on Incidence of COVID-19 Among Residents and Staff of Skilled Nursing and Assisted Living Facilities: A Randomized Clinical Trial. *JAMA* 2021; **326**: 46-55 [PMID: 34081073 DOI: 10.1001/jama.2021.8828]
- 20 **Gottlieb RL**, Nirula A, Chen P, Boscia J, Heller B, Morris J, Huhn G, Cardona J, Mocherla B, Stosor V, Shawa I, Kumar P, Adams AC, Van Naarden J, Custer KL, Durante M, Oakley G, Schade AE, Holzer TR, Ebert PJ, Higgs RE, Kallewaard NL, Sabo J, Patel DR, Klekotka P, Shen L, Skovronsky DM. Effect of Bamlanivimab as Monotherapy or in Combination With Etesevimab on Viral Load in Patients With Mild to Moderate COVID-19: A Randomized Clinical Trial. *JAMA* 2021; **325**: 632-644 [PMID: 33475701 DOI: 10.1001/jama.2021.0202]
- 21 Fact sheet for health care providers: Emergency Use Authorization (EUA) of bamlanivimab. [cited 21 March 2021]. In: US Food and Drug Administration [Internet]. Available from: <https://www.fda.gov/media/143603/download>
- 22 Fact sheet for health care providers: Emergency Use Authorization (EUA) of bamlanivimab and etesevimab. [cited 21 March 2021]. In: US Food and Drug Administration [Internet]. Available from: <https://www.fda.gov/media/145802/download>
- 23 **Weinreich DM**, Sivapalasingam S, Norton T, Ali S, Gao H, Bhore R, Musser BJ, Soo Y, Rofail D, Im J, Perry C, Pan C, Hosain R, Mahmood A, Davis JD, Turner KC, Hooper AT, Hamilton JD, Baum A, Kyratsous CA, Kim Y, Cook A, Kampman W, Kohli A, Sachdeva Y, Graber X, Kowal B, DiCioccio T, Stahl N, Lipsich L, Braunstein N, Herman G, Yancopoulos GD; Trial Investigators. REGN-COV2, a Neutralizing Antibody Cocktail, in Outpatients with Covid-19. *N Engl J Med* 2021; **384**: 238-251 [PMID: 33332778 DOI: 10.1056/NEJMoa2035002]
- 24 Fact sheet for health care providers: Emergency Use Authorization (EUA) of REGEN-COV™ (casirivimab with imdevimab). [cited 21 March 2021]. In: US Food and Drug Administration [Internet]. Available from: <https://www.fda.gov/media/145611/download>
- 25 CDSCO approves antibody cocktail drug for restricted emergency use to treat mild Covid-19 cases. [cited 21 March 2021]. In: India Today [Internet]. Available from: <https://www.indiatoday.in/coronavirus-outbreak/story/cdco-approves-antibody-cocktail-drug-for-restricted-emergency-use-to-treat-mild-covid-19-cases-1804859-2021-05-20>
- 26 **Gupta A**, Gonzalez-Rojas Y, Juarez E, Crespo Casal M, Moya J, Falci DR, Sarkis E, Solis J, Zheng H, Scott N, Cathcart AL, Hebner CM, Sager J, Mogalian E, Tipple C, Peppercorn A, Alexander E, Pang PS, Free A, Brinson C, Aldinger M, Shapiro AE; COMET-ICE Investigators. Early Treatment for Covid-19 with SARS-CoV-2 Neutralizing Antibody Sotrovimab. *N Engl J Med*. 2021; 385: 1941-1950 [PMID: 34706189 DOI: 10.1056/NEJMoa2107934]
- 27 Fact sheet for healthcare providers: emergency use authorization (EUA) of sotrovimab. [cited 21 March 2021]. In: US Food and Drug Administration [Internet]. Available from: <https://www.fda.gov/media/149534/download>
- 28 **Baum A**, Fulton BO, Wloga E, Copin R, Pascal KE, Russo V, Giordano S, Lanza K, Negron N, Ni M, Wei Y, Atwal GS, Murphy AJ, Stahl N, Yancopoulos GD, Kyratsous CA. Antibody cocktail to SARS-CoV-2 spike protein prevents rapid mutational escape seen with individual antibodies. *Science* 2020; **369**: 1014-1018 [PMID: 32540904 DOI: 10.1126/science.abd0831]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

