World Journal of *Critical Care Medicine*

World J Crit Care Med 2021 November 9; 10(6): 310-400





Published by Baishideng Publishing Group Inc

World Journal of C C M Critical Care Medicine



Bimonthly Volume 10 Number 6 November 9, 2021

OPINION REVIEW

310 Point of care venous Doppler ultrasound: Exploring the missing piece of bedside hemodynamic assessment

Galindo P, Gasca C, Argaiz ER, Koratala A

MINIREVIEWS

Extracorporeal membrane oxygenation and inhaled sedation in coronavirus disease 2019-related acute 323 respiratory distress syndrome

Bellgardt M, Özcelik D, Breuer-Kaiser AFC, Steinfort C, Breuer TGK, Weber TP, Herzog-Niescery J

334 Role of bronchoscopy in critically ill patients managed in intermediate care units - indications and complications: A narrative review

Menditto VG, Mei F, Fabrizzi B, Bonifazi M

345 Timing of tracheostomy in mechanically ventilated COVID-19 patients

Amadi N. Trivedi R. Ahmed N

ORIGINAL ARTICLE

Retrospective Study

355 Sequential organ failure assessment score is superior to other prognostic indices in acute pancreatitis Teng TZJ, Tan JKT, Baey S, Gunasekaran SK, Junnarkar SP, Low JK, Huey CWT, Shelat VG

Observational Study

369 Intensive care unit hospitalizations and outcomes in patients with severe COVID-19 during summer and fall surges in Georgia

Olanipekun T, Abe TA, Effoe VS, Musonge-Effoe JE, Chuks A, Kwara E, Caldwell A, Obeng S, Bakinde N, Westney G, Snvder R

SYSTEMATIC REVIEWS

377 Management of genitourinary trauma – current evaluation from the Sub-Saharan region: A systematic review

Cassell III AK, Manobah B

META-ANALYSIS

Recovery after acute kidney injury requiring kidney replacement therapy in patients with left ventricular 390 assist device: A meta-analysis

Kovvuru K, Kanduri SR, Thongprayoon C, Bathini T, Vallabhajosyula S, Kaewput W, Mao MA, Cheungpasitporn W, Kashani KB



Contents

Bimonthly Volume 10 Number 6 November 9, 2021

ABOUT COVER

Peer Reviewer of World Journal of Critical Care Medicine, Muthukumaran Rangarajan, FRSC, Professor, MS, FRCS, FACS, Department of Surgeon, Kovai Medical Center, Coimbatore 608001, India. rangy68@gmail.com

AIMS AND SCOPE

The primary aim of the World Journal of Critical Care Medicine (WJCCM, World J Crit Care Med) is to provide scholars and readers from various fields of critical care medicine with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJCCM mainly publishes articles reporting research results and findings obtained in the field of critical care medicine and covering a wide range of topics including acute kidney failure, acute respiratory distress syndrome and mechanical ventilation, application of bronchofiberscopy in critically ill patients, cardiopulmonary cerebral resuscitation, coagulant dysfunction, continuous renal replacement therapy, fluid resuscitation and tissue perfusion, hemodynamic monitoring and circulatory support, ICU management and treatment control, sedation and analgesia, severe infection, etc.

INDEXING/ABSTRACTING

The WJCCM is now indexed in PubMed, PubMed Central, China National Knowledge Infrastructure (CNKI), China Science and Technology Journal Database (CSTJ), and Superstar Journals Database.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Lin-YuTong Wang; Production Department Director: Xiang Li; Editorial Office Director: Li-Li Wang,

NAME OF JOURNAL	INSTRUCTIONS TO AUTHORS		
World Journal of Critical Care Medicine	https://www.wjgnet.com/bpg/gerinfo/204		
ISSN	GUIDELINES FOR ETHICS DOCUMENTS		
ISSN 2220-3141 (online)	https://www.wjgnet.com/bpg/GerInfo/287		
LAUNCH DATE	GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH		
February 4, 2012	https://www.wjgnet.com/bpg/gerinfo/240		
FREQUENCY	PUBLICATION ETHICS		
Bimonthly	https://www.wjgnet.com/bpg/GerInfo/288		
EDITORS-IN-CHIEF	PUBLICATION MISCONDUCT		
Kam-Lun Ellis Hon	https://www.wjgnet.com/bpg/gerinfo/208		
EDITORIAL BOARD MEMBERS	ARTICLE PROCESSING CHARGE		
https://www.wjgnet.com/2220-3141/editorialboard.htm	https://www.wjgnet.com/bpg/gerinfo/242		
PUBLICATION DATE November 9, 2021	STEPS FOR SUBMITTING MANUSCRIPTS https://www.wjgnet.com/bpg/GerInfo/239		
COPYRIGHT	ONLINE SUBMISSION		
© 2021 Baishideng Publishing Group Inc	https://www.f6publishing.com		

© 2021 Baishideng Publishing Group Inc. All rights reserved. 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA E-mail: bpgoffice@wjgnet.com https://www.wjgnet.com



W T C C M World Journal of Critical Care Medicine

Submit a Manuscript: https://www.f6publishing.com

World J Crit Care Med 2021 November 9; 10(6): 369-376

DOI: 10.5492/wjccm.v10.i6.369

ISSN 2220-3141 (online)

ORIGINAL ARTICLE

Observational Study

Intensive care unit hospitalizations and outcomes in patients with severe COVID-19 during summer and fall surges in Georgia

Titilope Olanipekun, Temidayo Ayotomiwa Abe, Valery Sammah Effoe, Joffi E Musonge-Effoe, Agusiegbe Chuks, Esther Kwara, Alexandra Caldwell, Samed Obeng, Nicolas Bakinde, Gloria Westney, Richard Snyder

ORCID number: Titilope Olanipekun 0000-0002-8472-6970; Temidayo Ayotomiwa Abe 0000-0001-8371-2000; Valery Sammah Effoe 0000-0002-7297-1151; Joffi E Musonge-Effoe 0000-0003-0513-4473; Agusiegbe Chuks 0000-0003-3585-5351; Esther Kwara 0000-0003-4352-6881; Alexandra Caldwell 0000-0001-8075-0167; Samed Obeng 0000-0002-8964-195X; Nicolas Bakinde 0000-0003-1229-1311; Gloria Westney 0000-0001-9665-0750; Richard Snyder 0000-0001-6669-1818.

Author contributions: Olanipekun T and Abe T contributed to study conception and manuscript writing; Effoe VS, and Musonge-Effoe J contributed to study conception and critical review; Chuks A, Kwara E, Caldwell A and Obeng S, contributed to data extraction and critical review; Bakinde N, Westney G and Snyder R contributed to study conception and critical review; All authors have read and approved the final manuscript.

Institutional review board

statement: Data from this study used de-identified data from the National Inpatient Sample Database. A publicly available allpayer inpatient care database in the United States. Institutional Review Board Approval Form or

Titilope Olanipekun, Department of Hospital Medicine, Covenant Health System, Knoxville, TN, Covenant Health System, Knoxville, TX 37922, United States

Temidayo Ayotomiwa Abe, Agusiegbe Chuks, Esther Kwara, Alexandra Caldwell, Samed Obeng, Nicolas Bakinde, Richard Snyder, Department of Medicine, Morehouse School of Medicine, Atlanta, GA 30310, United States

Valery Sammah Effoe, Department of Cardiology, Morehouse School of Medicine, Atlanta, GA 30310, United States

Joffi E Musonge-Effoe, Department of Clinical Research, Morehouse School of Medicine, Atlanta, GA 30310, United States

Gloria Westney, Department of Medicine, Pulmonary and Critical Care Section, Morehouse School of Medicine, Atlanta, GA 30310, United States

Corresponding author: Titilope Olanipekun, MD, Academic Research, Department of Hospital Medicine, Covenant Health System, Knoxville, TN, Covenant Health System, 100 Fort Sanders W Blvd, Knoxville100 Fort Sanders W Blvd, Knoxville, Knoxville, TX 37922, United States. toolanipekun@gmail.com

Abstract

BACKGROUND

There is limited data on the difference in the clinical characteristics and outcomes of patients with severe coronavirus disease 2019 (COVID-19) infection in the summer compared to the fall surge.

AIM

To compare the sociodemographic, clinical characteristics, and outcomes among mechanically ventilated patients with severe COVID-19 infection admitted to the intensive care unit (ICU) during the summer and fall surges in the year 2020.

METHODS

We included patients admitted to the ICU and treated with invasive mechanical ventilation for COVID-19 associated respiratory failure between April 1 and December 31, 2020. Patients were categorized into summer surge for ICU admissions between June 15, 2020, and August 15, 2020, and fall surge between



Document is not required.

Informed consent statement: Data from this study used de-identified data from the National Inpatient Sample Database. A publicly available all-payer inpatient care database in the United States. Informed patient consent is not required.

Conflict-of-interest statement: The authors declare no conflict of interests for this article.

Data sharing statement: Data that support the findings of this study are publicly available at https://www.hcupus.ahrq.gov/db/nation/nis/nisdb

documentation.jsp.

STROBE statement: The authors

have read the STROBE Statement, and the manuscript was prepared and revised according to the STROBE Statement.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: htt p://creativecommons.org/License s/by-nc/4.0/

Specialty type: Critical care medicine

Country/Territory of origin: United States

Peer-review report's scientific quality classification

Grade A (Excellent): 0 Grade B (Very good): B Grade C (Good): 0 Grade D (Fair): 0 Grade E (Poor): 0

Received: June 24, 2021 Peer-review started: June 24, 2021 First decision: July 27, 2021

October 15, 2020, and December 31, 2020. We compared patients' characteristics and outcomes using descriptive and inferential statistics.

RESULTS

A total of 220 patients were admitted to the Grady Memorial Hospital ICU and mechanically ventilated for COVID-19 associated hypoxemic respiratory failure during the period considered (125 during the summer surge and 95 during the fall surge). More women were admitted in the fall compared to summer (41.1% vs 36.8%, difference, 4.3%; 95% CI: 1.2, 7.5). Patients admitted in the fall had fewer comorbidities (chronic obstructive pulmonary disease, stroke, diabetes mellitus, obstructive sleep apnea and body mass index \geq 35 kg/m²). Overall, patients in the fall had a lower ICU mortality rate (27.4% vs 38.4%, difference, -11.0; 95%CI: -6.4, -18.2), shorter length of stay on the mechanical ventilator (7 d vs 11 d, difference, 4 d; 95%CI: 2.1, 6.6) and shorter ICU length of stay (9 d vs 14 d, difference, 5 d; 95%CI: 2.7, 9.4).

CONCLUSION

Patients admitted with severe COVID-19 infection requiring mechanical ventilation had better outcomes in the fall than summer. This difference observed is likely attributable to a better understanding of the condition and advances in treatment strategies.

Key Words: COVID-19; COVID-19 surge; Georgia; Intensive care unit; Mechanical ventilation

©The Author(s) 2021. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: In this observational study, we compared the sociodemographic, clinical characteristics, and outcomes among mechanically ventilated patients with coronavirus disease 2019 (COVID-19) infection admitted to the intensive care unit (ICU) during the summer and fall surges in the year 2020. Compared to patients admitted with severe COVID-19 in the summer, those in the fall had better outcomes including decreased mortality and low length of stay in the ICU. This is likely due to the improved understanding of COVID-19 and the advances in treatment strategies.

Citation: Olanipekun T, Abe TA, Effoe VS, Musonge-Effoe JE, Chuks A, Kwara E, Caldwell A, Obeng S, Bakinde N, Westney G, Snyder R. Intensive care unit hospitalizations and outcomes in patients with severe COVID-19 during summer and fall surges in Georgia. World J Crit Care Med 2021; 10(6): 369-376

URL: https://www.wjgnet.com/2220-3141/full/v10/i6/369.htm DOI: https://dx.doi.org/10.5492/wjccm.v10.i6.369

INTRODUCTION

Coronavirus disease 2019 (COVID-19) was first reported in the United States on January 20, 2020. The World Health Organization declared the novel viral infection a pandemic on March 11, 2020[1]. Within 1 year of the pandemic, more than 31 million cases and 500000 deaths have been recorded in the United States^[2].

In the United States the pandemic has been characterized by waves of case surges attributed to holiday gatherings, relaxation of social distancing guidelines and removal of COVID-19 restriction during reopening after lockdown in different states[3,4]. Surges in COVID-19 cases are associated with increased hospitalizations including the intensive care units (ICU) placing significant strains on hospital resources[3]. The state of Georgia experienced resurgence of cases during the summer and fall seasons of 2020 with increased rates of hospitalizations^[5].

We describe the differences in the sociodemographic, clinical characteristics, and outcomes in mechanically ventilated patients with severe COVID-19 infection admitted to the ICU of Grady Memorial Hospital (GMH) in Atlanta, Georgia across the surges in the summer and fall of year 2020.



Revised: August 5, 2021 Accepted: September 16, 2021 Article in press: September 16, 2021 Published online: November 9, 2021

P-Reviewer: Braga MB S-Editor: Ma YJ L-Editor: A P-Editor: Wang LYT



MATERIALS AND METHODS

We identified patients with positive reverse transcriptase-polymerase chain reaction nasopharyngeal swab test results for severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) admitted to the GMH ICU (Morehouse school of medicine and Emory university ICU service) for hypoxemic respiratory failure and treated with invasive mechanical ventilation between April 1 and December 31, 2020. GMH is a level 1 trauma hospital in Atlanta, Georgia with more than 900 beds and one of the largest hospitals in Georgia. GMH played a significant role in the Georgia COVID-19 response by providing care to many COVID-19 patients in Georgia[6].

We extracted information on the age, sex, race/ethnicity, comorbidities, medication use, length of stay (LOS) on the mechanical ventilator, LOS in the ICU and ICU mortality outcomes from the electronic health record system. We obtained data from the Georgia State Department of Health to evaluate the trend of new COVID-19 cases between April 1 and December 31, 2020. Patients were categorized into summer surge for ICU admissions between June 15, 2020 and August 15, 2020 and fall surge between October 15, 2020 and December 31, 2020 based on the resurgence of COVID-19 cases in Georgia during the summer and fall periods. The study was approved by the Morehouse School of Medicine's Institutional Review Board (IRB). The IRB issued a waiver of HIPAA authorization to access electronic medical records.

We report summary statistics as means or medians and proportions for sociodemographic, clinical, and outcome variables of severe COVID-19 patients hospitalized in the ICU. We computed bivariable comparisons across summer and fall surges and reported the proportional differences with corresponding 95% CIs. All analyses were performed with version 3.5.2 of the R programming language (R Project for Statistical Computing; R Foundation). *P* values were 2-sided, with statistical significance set at *P* < 0.05.

RESULTS

A total of 220 patients were admitted to the GMH ICU and mechanically ventilated for COVID-19 associated hypoxemic respiratory failure during the period considered (125 during the summer surge and 95 during the fall surge). Table 1 describes the differences between the socio-demographic, clinical characteristics, and the outcomes of these patients across the two surge periods. Proportion of females was higher during the fall surge compared to the summer surge (41.1% *vs* 36.8%, difference, 4.3%; 95%CI: 1.2, 7.5). More patients had private insurance during the fall surge (36.8%*vs* 30.4%, difference, 6.4%; 95%CI: 1.5, 13.3) while fewer patients were uninsured during the summer (18.9% *vs* 28%, difference, -9.1%: 95%CI: -6.4, -12.5).

Patients admitted to the ICU during the fall surge had significantly higher burden of chronic kidney disease 4 and above, human immunodeficiency virus/acquired immune deficiency syndrome, hypertension, class 1 obesity (body mass index, BMI \geq 30 kg/m² to < 35 kg/m²) and tobacco use disorder, while fewer patients had chronic obstructive pulmonary disease (COPD), cerebrovascular accidents, diabetes mellitus (DM), obstructive sleep apnea and class 2 or greater obesity (BMI \geq 35 kg/m²). A significantly higher proportion of patients during the fall had no comorbidities at baseline compared to those in the summer (16.8% *vs* 9.6%, difference 7.2%; 95%CI: 2.8, 13.9). A greater proportion of patients in the fall surge were treated with remdesivir and dexamethasone.

The ICU mortality rate (27.4% *vs* 38.4%, difference, -11.0; 95% CI: -6.4, -18.2) was lower in the fall compared to summer. Similarly, patients in the fall had a shorter LOS on the mechanical ventilator (7 d *vs* 11 d, difference, 4 d; 95% CI: 2.1, 6.6) and shorter LOS in the ICU (9 d *vs* 14 d, difference, 5 d; 95% CI: 2.7, 9.4).

Figures 1 and 2 present trends of new COVID-19 cases and ICU hospitalizations in Georgia and GMH between April 1 – December 31, 2020. There was an increase in COVID-19 cases and ICU hospitalizations at GMH correlating with the summer and fall surges in Georgia. Dates corresponding to specific US national holidays are highlighted in Figure 1. Notably, the memorial holiday (May 25, 2020) preceded the surge of cases in the summer while the Labor Day (September 7, 2020) and Thanks-giving (November 26, 2020) holidays preceded the Fall surge.

Zaishidena® WJCCM | https://www.wjgnet.com

Table 1 Sociodemographic, comorbidity, clinical, and outcome differences between surge 1 and surge 2 of Intensive care unit hospitalizations for coronavirus disease-19 Respiratory Failure at Grady Memorial Hospital, Atlanta, Georgia

Variables	Surge 1 (Summer 2020)	Surge 2 (Fall 2020)	Difference (95%CI)	P value
Total – <i>n</i> (%)	125	95		
Age, median (IQR)	61.5 (51-69)	61 (51.5-71)	0.5	0.34
< 55 yr	41 (32.8)	30 (31.6)	-1.2 (-3.8, 6.5)	0.39
55-64 yr	38 (30.4)	25 (26.3)	-4.1 (-6.2, -2.3)	0.04
65-74 yr	27 (21.6)	27 (28.4)	6.8 (4.5, 10.3)	0.04
> 75 yr	19 (15.2)	13 (13.7)	-1.5 (-5.5, 2.5)	0.19
Race				
Non-Hispanic Black	86 (68.8)	61 (64.2)	-4.6 (-10.7, 1.1)	0.1
Non-Hispanic White	22 (17.6)	17 (17.9)	0.3 (-1.7, 2.5)	0.18
Hispanic	10 (8)	8 (8.4)	0.4 (-1.6, 2.4)	0.26
Others ¹	7 (5.6)	8 (8.4)	2.8 (0.9, 4.9)	0.09
Gender				
Female	46 (36.8)	39 (41.1)	4.3 (1.2, 7.5)	< 0.01
Male	79 (63.2)	56 (58.9)	-4.3 (-2.4, -6.2)	< 0.01
Health insurance				
Medicaid only	7 (5.6)	5 (5.3)	-0.3 (-2.3, 1.5)	0.41
Medicare only	20 (16)	15 (15.8)	-0.2 (-1.8, 1.3)	0.12
Medicaid/Medicare	25 (20)	22 (23.2)	3.2 (1.0, 5.8)	0.05
Private insurance/Self pay	38 (30.4)	35 (36.8)	6.4 (1.5, 13.3)	0.02
Uninsured	35 (28)	18 (18.9)	-9.1 (- 6.4, -12.5)	0.04
Comorbid diseases				
Asthma	13 (10.4)	9 (9.5)	-0.9 (-4.8, 2.8)	0.22
Coronary artery disease	20 (16)	12 (12.6)	-3.4 (-8.5, 1.7)	0.11
Cancer (solid organ tumors)	12 (9.6)	6 (6.3)	-3.3 (- 8.2, 1.9)	0.18
Congestive heart failure	29 (23.2)	20 (21.1)	-2.1 (- 6.6, 2.1)	0.4
Chronic kidney disease 3 and above	17 (13.6)	16 (16.8)	3.2 (1.6, 5.1)	0.02
Chronic liver disease	10 (8)	9 (9.5)	1.5 (0.8, 2.47)	0.13
Chronic obstructive pulmonary disease	22 (17.6)	12 (12.6)	-5.0 (-10.1, -5.6)	0.04
Cerebrovascular accident	21 (16.8)	11 (11.6)	-5.2 (-10.3, -5.8)	0.02
Diabetes mellitus	62 (49.6)	41 (43.2)	-6.4 (-3.1, -9.8)	< 0.01
HIV/AIDS	4 (3.2)	8 (8.4)	5.2 (2.7, 8.2)	< 0.01
Hypertension	85 (68)	67 (70.5)	2.5 (1.3, 4.9)	< 0.01
Obstructive sleep apnea	14 (11.2)	6 (6.3)	-4.9 (-1.9, 8.8)	< 0.01
Body mass index				
$< 30 \text{ kg/m}^2$	37 (29.6)	29 (30.5)	0.9 (0.4, 1.4)	0.078
$\ge 30 \text{ kg/m}^2 \text{ to} \le 35 \text{ kg/m}^2$	48 (38.4)	40 (42.1)	3.7 (1.6, 5.9)	< 0.01
\geq 35 kg/m ²	40 (32)	26 (27.4)	-4.6 (-1.3, 8.7)	< 0.01
Tobacco use (current smoker)	51 (40.8)	45 (47.4)	6.6 (3.2, 11.8)	< 0.01
No of comorbidities				
None	12 (9.6)	16 (16.8)	7.2 (2.8, 13.9)	< 0.01



Olanipekun T et al. Outcomes of severe COVID-19 during summer compared to fall

1	8 (6.4)	5 (5.3)	-1.1 (-3.4, 1.1)	0.46			
>1	105 (84)	78 (82.1)	-1.9 (-5.1, 1.3)	0.37			
Treatment received in the ICU							
Hydroxychloroquine	6 (4.8)	0 (0)	-4.8 (- 2.5, -7.6)	0.03			
Azithromycin	25 (20)	14 (14.7)	-5.3 (- 3.1, -7.9)	0.04			
Hydroxychloroquine + Azithromycin	36 (28.8)	0 (0)	-28.8 (-19.4, 41.1)	< 0.01			
Remdesivir	44 (35.2)	42 (44.2)	9.0 (3.4, 10.2)	< 0.01			
Dexamethasone	0	22 (23.2)	23.2 (18.3, 29.1)	< 0.01			
Length of stay, median (IQR) (range), d							

¹American Indian or Alaska Native, Asian, and Native Hawaiian or other Pacific Islander. Differences calculated as surge 2 - surge 1. Negative values represent decrease in surge 2. Null value = 0. HIV: Human immunodeficiency virus; AIDS: Acquired immunodeficiency syndrome; COVID-19: Coronavirus disease 2019; ICU: Intensive care unit; IQR: Interquartile range.



Figure 1 New coronavirus disease 2019 cases in Georgia State (April 1 - December 15, 2020). A: Memorial Day Holiday May 25, 2020; B: Labor Day Holiday September 7, 2020; C: Thanksgiving Holiday November 26, 2020.Cited: Georgia State Department of Health COVID-19 Daily Status Report. https://dph.georgia.gov/covid-19-daily-status-report. Accessed 12/28/2020. Figure reproduced with permission.

DISCUSSION

We observed increased ICU hospitalizations for COVID-19 associated respiratory failure requiring invasive mechanical ventilation corresponding to the surges of cases during the summer and fall of 2020 in Georgia. ICU hospitalizations during the reporting period were consistently guided by severity of symptoms, comorbidities, clinical and diagnostic findings, respiratory status, and indications for mechanical ventilation. Therefore, the higher ICU census noted during the surges likely reflects increased rates of COVID-19 prevalence in the community.

ICU mortality was 11% lower among patients in the fall cohort than those in the summer. Also, the length of time a patient spent on the mechanical ventilator and in the ICU were shorter during the fall when compared to the summer. These observed differences could be explained by a number of factors. First, patients in the fall cohort had a higher proportion of patients with no comorbidities at baseline and fewer patients relative to the summer cohort, with specific chronic medical problems such as DM, COPD and class 2 obesity or greater, which have been associated with severe COVID-19 course and poorer outcomes. Second, more patients in the fall surge received Food and Drug Administration (FDA)-approved and Centers for Disease Control and Prevention (CDC) recommended treatments, including dexamethasone and remdesivir, which have been shown to improve outcomes among COVID-19 patients. The Randomized Evaluation of Covid-19 Therapy (RECOVERY) trial demonstrated a lower 28 d mortality among COVID-19 patients treated with



WJCCM https://www.wjgnet.com



Figure 2 Daily hospital census of coronavirus disease-19 infected patients at Grady Memorial Hospital (April 1- December 15, 2020). A: Hospitalizations during the surge of new cases in the summer; B: Hospitalizations during the surge of new cases in the fall. ICU: Intensive care unit; MICU: Medical intensive care unit

dexamethasone compared to placebo[7]. There is also evidence that remdesivir, when compared to placebo is associated with a shorter duration of mechanical ventilation or extracorporeal membrane oxygenation (ECMO) for COVID-19 respiratory failure[8]. Third, the improved outcomes likely mirror a combination of better understanding of COVID-19 pathophysiology, availability of novel therapies and better medical management.

It is not surprising that the summer and fall surges were preceded by major holidays as there have been numerous similar reports globally. Attending events that involve large gatherings typically seen during holidays increases the chances of COVID-19 infection[3]. For instance, the Lunar New Year holiday coincided with the start of the pandemic when millions of people left the city of Wuhan in China to visit relatives in other parts of the country and the world [9]. In the United Kingdom, the early COVID-19 epidemic followed a one-week school holiday break from February 17 to February 21, 2020 when thousands of people came back infected with SARS-CoV-2 virus from tourist activities in northern Italy and Spain^[10]. Also, Canada reported its highest numbers of COVID-19 infection cases in the two weeks following the Thanksgiving holiday on October 12, 2020[2].

Our study has some limitations. First, this was a single center study with unique institutional practices and findings that may not be generalizable. Second, there are socio-behavioral and political circumstances that may have contributed to the surge of COVID-19 cases during the fall and summer seasons of 2020 that we could not measure in this study.

CONCLUSION

In this single-center study, we found significant differences in the sociodemographic, clinical characteristics, and outcomes among mechanically ventilated COVID-19 patients in the ICU during the 2020 summer surge compared to the fall surge. ICU mortality, LOS on mechanical ventilator, and LOS in the ICU were all significantly lower in the fall than summer. This finding is likely a result of our improved understanding of COVID-19 and advancement in management strategies.

ARTICLE HIGHLIGHTS

Research background

There is limited data on the difference in the clinical characteristics and outcomes of patients with severe coronavirus disease 2019 (COVID-19) infection in the summer compared to the fall surge.



Research motivation

Surges in COVID-19 cases are associated with increased hospitalizations including the intensive care units (ICU) placing significant strains on hospital resources. Knowledge about the differences in the clinical characteristics and outcomes between each surge will provide useful information on how to decrease related morbidity and mortality.

Research objectives

To compare the sociodemographic, clinical characteristics, and outcomes among mechanically ventilated patients with severe COVID-19 infection admitted to the (ICU) during the summer and fall surges in the year 2020.

Research methods

The authors included mechanically ventilated COVID-19 patients managed at Grady Memorial Hospital (GMH) from April 1 and December 31, 2020. Patients were categorized into two groups, those admitted in the summer (June 15, 2020 - August 15, 2020) and fall (October 15, 2020 - December 31, 2020). We compared patients' characteristics and outcomes using descriptive and inferential statistics.

Research results

A total of 220 patients were admitted to the GMH ICU and mechanically ventilated for COVID-19 (125 during the summer surge and 95 during the fall surge). Patients admitted in the fall had fewer comorbidities, lower mortality rate, shorter length of stay on the mechanical ventilator and shorter ICU length of stay.

Research conclusions

Patients admitted with severe COVID-19 infection requiring mechanical ventilation had better outcomes in the fall than in summer.

Research perspectives

Further studies are needed to replicate these findings.

REFERENCES

- Holshue ML, DeBolt C, Lindquist S, Lofy KH, Wiesman J, Bruce H, Spitters C, Ericson K, 1 Wilkerson S, Tural A, Diaz G, Cohn A, Fox L, Patel A, Gerber SI, Kim L, Tong S, Lu X, Lindstrom S, Pallansch MA, Weldon WC, Biggs HM, Uyeki TM, Pillai SK; Washington State 2019-nCoV Case Investigation Team. First Case of 2019 Novel Coronavirus in the United States. N Engl J Med 2020; 382: 929-936 [PMID: 32004427 DOI: 10.1056/NEJMoa2001191]
- 2 JHU CSSE. Dataset: GitHub CSSEGISandData/COVID-19: Novel Coronavirus (COVID-19) Cases. Dataset COVID-19, 2020. Available from: https://github.com/CSSEGISandData/COVID-19
- 3 Duque D, Morton DP, Singh B, Du Z, Pasco R, Meyers LA. Timing social distancing to avert unmanageable COVID-19 hospital surges. Proc Natl Acad Sci USA 2020; 117: 19873-19878 [PMID: 32727898 DOI: 10.1073/pnas.2009033117]
- 4 Vahidy FS, Drews AL, Masud FN, Schwartz RL, Askary BB, Boom ML, Phillips RA. Characteristics and Outcomes of COVID-19 Patients During Initial Peak and Resurgence in the Houston Metropolitan Area. JAMA 2020; 324: 998-1000 [PMID: 32789492 DOI: 10.1001/jama.2020.15301]
- Moore JX, Langston ME, George V, Coughlin SS. Epidemiology of the 2020 pandemic of COVID-5 19 in the state of Georgia: Inadequate critical care resources and impact after 7 weeks of community spread. J Am Coll Emerg Physicians Open 2020 [PMID: 32838368 DOI: 10.1002/emp2.12127]
- Jackson BR, Gold JAW, Natarajan P, Rossow J, Neblett Fanfair R, da Silva J, Wong KK, Browning SD, Bamrah Morris S, Rogers-Brown J, Hernandez-Romieu AC, Szablewski CM, Oosmanally N, Tobin-D'Angelo M, Drenzek C, Murphy DJ, Hollberg J, Blum JM, Jansen R, Wright DW, SeweSll WM, Owens JD, Lefkove B, Brown FW, Burton DC, Uyeki TM, Bialek SR, Patel PR, Bruce BB. Predictors at admission of mechanical ventilation and death in an observational cohort of adults hospitalized with COVID-19. Clin Infect Dis 2020 [PMID: 32971532 DOI: 10.1093/cid/ciaa1459]
- RECOVERY Collaborative Group, Horby P, Lim WS, Emberson JR, Mafham M, Bell JL, Linsell 7 L, Staplin N, Brightling C, Ustianowski A, Elmahi E, Prudon B, Green C, Felton T, Chadwick D, Rege K, Fegan C, Chappell LC, Faust SN, Jaki T, Jeffery K, Montgomery A, Rowan K, Juszczak E, Baillie JK, Haynes R, Landray MJ. Dexamethasone in Hospitalized Patients with Covid-19. N Engl J Med 2021; 384: 693-704 [PMID: 32678530 DOI: 10.1056/NEJMoa2021436]
- Beigel JH, Tomashek KM, Dodd LE, Mehta AK, Zingman BS, Kalil AC, Hohmann E, Chu HY, 8 Luetkemeyer A, Kline S, Lopez de Castilla D, Finberg RW, Dierberg K, Tapson V, Hsieh L, Patterson TF, Paredes R, Sweeney DA, Short WR, Touloumi G, Lye DC, Ohmagari N, Oh MD, Ruiz-Palacios GM, Benfield T, Fätkenheuer G, Kortepeter MG, Atmar RL, Creech CB, Lundgren J,



Babiker AG, Pett S, Neaton JD, Burgess TH, Bonnett T, Green M, Makowski M, Osinusi A, Nayak S, Lane HC; ACTT-1 Study Group Members. Remdesivir for the Treatment of Covid-19 - Final Report. N Engl J Med 2020; 383: 1813-1826 [PMID: 32445440 DOI: 10.1056/NEJMoa2007764]

- 9 Chen S, Yang J, Yang W, Wang C, Bärnighausen T. COVID-19 control in China during mass population movements at New Year. *Lancet* 2020; **395**: 764-766 [PMID: 32105609 DOI: 10.1016/S0140-6736(20)30421-9]
- du Plessis L, McCrone JT, Zarebski AE, Hill V, Ruis C, Gutierrez B, Raghwani J, Ashworth J, 10 Colquhoun R, Connor TR, Faria NR, Jackson B, Loman NJ, O'Toole Á, Nicholls SM, Parag KV, Scher E, Vasylyeva TI, Volz EM, Watts A, Bogoch II, Khan K; COVID-19 Genomics UK (COG-UK) Consortium, Aanensen DM, Kraemer MUG, Rambaut A, Pybus OG. Establishment and lineage dynamics of the SARS-CoV-2 epidemic in the UK. Science 2021; 371: 708-712 [PMID: 33419936 DOI: 10.1126/science.abf2946]





Published by Baishideng Publishing Group Inc 7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA Telephone: +1-925-3991568 E-mail: bpgoffice@wjgnet.com Help Desk: https://www.f6publishing.com/helpdesk https://www.wjgnet.com

