

# World Journal of *Psychiatry*

*World J Psychiatry* 2023 June 19; 13(6): 262-401



**OPINION REVIEW**

- 262 Pharmacotherapy in autism spectrum disorders, including promising older drugs warranting trials  
*Hellings J*

**REVIEW**

- 278 Operational definitions and measurement of externalizing behavior problems: An integrative review including research models and clinical diagnostic systems  
*Torres-Rosado L, Lozano OM, Sanchez-Garcia M, Fernández-Calderón F, Diaz-Batanero C*
- 298 Psychiatrists' occupational stigma conceptualization, measurement, and intervention: A literature review  
*Shi XL, Li LY, Fan ZG*

**MINIREVIEWS**

- 319 Epigenetics in psychiatry: Beyond DNA methylation  
*Kouter K, Šalamon Arčan I, Videtič Paska A*
- 331 Psychological trauma, posttraumatic stress disorder and trauma-related depression: A mini-review  
*Wang SK, Feng M, Fang Y, Lv L, Sun GL, Yang SL, Guo P, Cheng SF, Qian MC, Chen HX*

**ORIGINAL ARTICLE****Basic Study**

- 340 Acupuncture at Back-Shu point improves insomnia by reducing inflammation and inhibiting the ERK/NF- $\kappa$ B signaling pathway  
*Zhang MM, Zhao JW, Li ZQ, Shao J, Gao XY*

**Observational Study**

- 351 Psychological impact of cancer scale: Turkish validity and reliability study  
*Bahçecioglu Turan G, Karaman S, Aksoy M*
- 361 Relationship between depression, smartphone addiction, and sleep among Chinese engineering students during the COVID-19 pandemic  
*Gao WJ, Hu Y, Ji JL, Liu XQ*
- 376 Prevalence of posttraumatic stress disorder following acute coronary syndrome and clinical characteristics of patients referred to cardiac rehabilitation  
*Sopek Merkaš I, Lakušić N, Sonicki Z, Koret B, Vuk Pisk S, Filipčić I*
- 386 Abnormal volumetric brain morphometry and cerebral blood flow in adolescents with depression  
*Fu YJ, Liu X, Wang XY, Li X, Dai LQ, Ren WY, Zeng YM, Li ZL, Yu RQ*

**LETTER TO THE EDITOR**

- 397** Digital interventions empowering mental health reconstruction among students after the COVID-19 pandemic

*Liu XQ, Guo YX, Zhang XR, Zhang LX, Zhang YF*

**ABOUT COVER**

Editorial Board Member of *World Journal of Psychiatry*, Lenin Pavón, PhD, Professor, Laboratory of Psychoimmunology, National Institute of Psychiatry, “Ramón de la Fuente”, Mexico City 14070, Mexico.  
lkuriaki@gmail.com

**AIMS AND SCOPE**

The primary aim of *World Journal of Psychiatry* (WJP, *World J Psychiatry*) is to provide scholars and readers from various fields of psychiatry with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJP mainly publishes articles reporting research results and findings obtained in the field of psychiatry and covering a wide range of topics including adolescent psychiatry, biological psychiatry, child psychiatry, community psychiatry, ethnopsychology, psychoanalysis, psychosomatic medicine, etc.

**INDEXING/ABSTRACTING**

The WJP is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports/Science Edition, PubMed, PubMed Central, Reference Citation Analysis, China National Knowledge Infrastructure, China Science and Technology Journal Database, and Superstar Journals Database. The 2022 Edition of Journal Citation Reports® cites the 2021 impact factor (IF) for WJP as 3.500; IF without journal self cites: 3.313; 5-year IF: 7.380; Journal Citation Indicator: 0.62; Ranking: 89 among 155 journals in psychiatry; and Quartile category: Q3.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: *Yu-Xi Chen*; Production Department Director: *Xu Guo*; Editorial Office Director: *Jia-Ping Yan*.

**NAME OF JOURNAL**

*World Journal of Psychiatry*

**ISSN**

ISSN 2220-3206 (online)

**LAUNCH DATE**

December 31, 2011

**FREQUENCY**

Monthly

**EDITORS-IN-CHIEF**

Rajesh R Tampi, Ting-Shao Zhu, Panteleimon Giannakopoulos

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/2220-3206/editorialboard.htm>

**PUBLICATION DATE**

June 19, 2023

**COPYRIGHT**

© 2023 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>



## Psychiatrists' occupational stigma conceptualization, measurement, and intervention: A literature review

Xiao-Li Shi, Lu-Yao Li, Zhi-Guang Fan

**Specialty type:** Psychiatry

**Provenance and peer review:**

Invited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): 0  
Grade B (Very good): B  
Grade C (Good): C  
Grade D (Fair): 0  
Grade E (Poor): 0

**P-Reviewer:** Gazdag G, Hungary;  
Masaru T, Hungary

**Received:** January 28, 2023

**Peer-review started:** January 28, 2023

**First decision:** March 24, 2023

**Revised:** April 5, 2023

**Accepted:** May 11, 2023

**Article in press:** May 11, 2023

**Published online:** June 19, 2023



**Xiao-Li Shi, Lu-Yao Li, Zhi-Guang Fan**, School of Education, Jilin International Studies University, Changchun 130000, Jilin Province, China

**Corresponding author:** Zhi-Guang Fan, PhD, Associate Professor, School of Education, Jilin International Studies University, 3658 Jingyue Street, Changchun 130000, Jilin Province 130117, China. [fanyqm@126.com](mailto:fanyqm@126.com)

### Abstract

Psychiatrists require frequent contact with and treatment of patients with mental illnesses. Due to the influence of associative stigma, psychiatrists may also be targets of stigma. Occupational stigma warrants special consideration because it significantly affects psychiatrists' career advancement, well-being, and their patients' health. Given that there is no complete summary of this issue, this study reviewed the existing literature on psychiatrists' occupational stigma to clearly synthesize its concepts, measurement tools, and intervention strategies. Herein, we emphasize that psychiatrists' occupational stigma is a multifaceted concept that simultaneously encompasses physically, socially, and morally tainted aspects. Currently, standardized methods to specifically measure psychiatrists' occupational stigma are lacking. Interventions for psychiatrists' occupational stigma may consider the use of protest, contact, education, comprehensive and systematic methods, as well as the use of psychotherapeutic approaches. This review provides a theoretical basis for the development of relevant measurement tools and intervention practices. Overall, this review seeks to raise public awareness of psychiatrists' occupational stigma, thereby promoting psychiatric professionalism and reducing its stigma.

**Key Words:** Psychiatrists; Occupational stigma; Conceptualization; Measurement; Intervention; Associative stigma

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.



**Core Tip:** Psychiatrists' occupational stigma, an area that is little noticed, deserves more attention from the public and professionals considering its negative effects. This study aimed to elucidate its concept, explore the potential measurement tools, and focus on effective interventions by comprehensively reviewing related literature. It is expected to encourage more studies in this field.

**Citation:** Shi XL, Li LY, Fan ZG. Psychiatrists' occupational stigma conceptualization, measurement, and intervention: A literature review. *World J Psychiatry* 2023; 13(6): 298-318

**URL:** <https://www.wjgnet.com/2220-3206/full/v13/i6/298.htm>

**DOI:** <https://dx.doi.org/10.5498/wjp.v13.i6.298>

## INTRODUCTION

Stigma is a form of social classification. Occupational stigma results when people make derogatory, insulting, or negative comments about others practicing their occupation[1]. Occupational stigma is the negative labeling of a profession that the general public views as having dishonorable, humiliating, and shameful features, leading to negative consequences of social exclusion, status loss, demeaning, discrimination, and pessimistic rejection[2-4]. Although stigma arises from dirty work in common sense, any occupation can face various levels and types of stigma[3]. Physicians, a typically prestigious profession [5], also suffer from stigma that seriously jeopardizes health care career development[6,7]. Furthermore, psychiatrists may experience intense stigma due to the nature of their profession and the population they serve[8]. A study of trainee psychiatrists found that 75% of individuals heard denigrating or humiliating remarks about the psychiatric profession[9].

Occupational stigma exerts negative impact on psychiatrists' career advancement, well-being, and patients' health. More specifically, the stigma may cause burnout, job dissatisfaction, and low professional value[10-17], and can act as a strong predictor of health and well-being[18,19]. Numerous studies have demonstrated that occupational stigma contributes to practitioners' withdrawal behaviors and the propensity for resignation[14,20-22]. At present, the shortage of psychiatrists cannot meet the demands of mental health workforce development[23]. Alarmingly, a study suggested that just about half of psychiatry college graduates moved on to work in related disciplines[24]. It has shown that occupational stigma plays a major role in that kind of phenomenon[25-28]. In addition, as a consequence of stigma, medical students rarely consider psychiatry a future career option, with the perception of taking psychiatrists as having low professional prestige and respect[29-31]. The professional shortage in psychiatry has been existing considering the insufficient medical graduates attracted by psychiatry[32, 33].

Moreover, it is noteworthy that stigma can straightly influence the establishment of the well physician-patient relationship and the treatment process. Stigma is classified as either public stigma or self-stigma, in which patients' stigma toward psychiatrists falls under the former[34]. That public stigma can reduce patient compliance and help-seeking behavior, thereby impeding treatment[32,35]. For another, psychiatrists' self-stigma may result in an increase in defensive medical behavior[36], such as little communication with patients, unnecessary testing, excessive medication, refusal, or referral[37-39], leading to not only health-care costs rise and poor medical relationship but also the potential negative impact on patient health and treatment effect[40].

In conclusion, psychiatrists influenced by associative stigma are perceived as a heavily stigmatized profession. Despite numerous studies showing that occupational stigma exerts a significant negative impact on the development of psychiatry, there is little literature reviewing psychiatrists' occupational stigma. To that end, the purpose of this literature review is to provide a comprehensive overview of the concept, measurement, and intervention of psychiatrists' occupational stigma by reviewing the extant literature, with the prospect of providing a theoretical reference for future research.

## LITERATURE SEARCH

The authors sequentially searched the PubMed, Web of Science, and Reference Citation Analysis (<https://www.referencecitationanalysis.com/>) databases for articles containing a cross combination of the following topical keywords: "Psychiatrist," "stigma," "occupational stigma," "stress," "negative affect," "career satisfaction," "dirty work," "healthcare workers," "associative stigma," "psychiatry," "mental health professionals," "self-stigma," "mental illness," "intervention," "measurement," and "anti-stigma." March 2023 was the deadline for the keyword search, which yielded an initial total of 21098 papers. Literature selection criteria, as decided between the professor and students, were as follows: First, include a study relevant to and representative of the topic; second, any such study should be published in English or French; third, exclude duplicates. After review 195 papers met the selection

criteria and 20903 papers were excluded.

## CONCEPTUALIZATION OF PSYCHIATRISTS' OCCUPATIONAL STIGMA

Research on occupational stigma can be traced back to Hughes[2] and his exploration of dirty work[2]. Impressive findings have been attained in the following studies that explore occupational stigma across practically all professions[3,41,42]. In his research on dirty work, Hughes[2] classified occupational stigma into three categories: physically, socially, and morally tainted in the aspect of work content[2]. Based on this, Ashforth and Kreiner[43] provided a precise definition of the three different forms of stigma[43]. Particularly, the term "physically tainted" refers to jobs that involve direct contact with trash, death, or filth[44,45] such as cleaner and mortician, or directly working in dangerous and harmful environment[46,47] such as firefighter and miner. The term "socially tainted" describes jobs like prison guards and infectious disease doctors that require regular contact with stigmatized groups as part of their duties[48,49], as well as those like nannies and tour guides that include a servant-subordinate relationship[50,51]. The term "morally tainted" refers to occupations that are viewed as being sinful and unethical[52,53] such as doctor who perform abortions or sex worker, as well as occupations with leading and deceptive traits[54,55] such as anchors who lead viewers to spend money and poker players who deceive their opponents. It is critical to note that one type of stigma may predominate in a given occupation, or two or even three types of stigma may exist concurrently[3,43]. Stenger *et al*[56] extended the three-dimensional classification, arguing that occupational stigma is a negative stereotype formed by the public of certain occupations' work images, social relations, or ethics[56].

In a follow-up study, Kreiner *et al*[3] further proposed the concepts of "breadth" and "depth" of taint applied to work tasks undertaken[3]. Breadth refers to the centrality of stigma in occupational identity, and the frequency of stigma-related behaviors occurring. Depth refers to the degree to which a practitioner is directly exposed to dirt. Accordingly, occupational stigma was further divided into pervasive stigma, compartmentalized stigma, diluted stigma and idiosyncratic stigma. At the same time, Ashforth and Kreiner[42] took occupational reputation as an important dimension of occupational stigma division, and then divided occupational stigma into high/low reputation physically tainted, high/low reputation socially tainted, and high/low reputation morally tainted[42]. In a recent study, Zhang *et al* [57] analyzed the four-level stigma literature of individual, occupational, organizational, and industry, and divided the sources of stigma into six types (physical, tribal, moral, servile, emotional, associational), which extended the three-dimensional classification of occupational stigma. The sources of occupational stigma are considered to include these six types. At the same time, five characteristics of stigma (concealability, controllability, centrality, disruptiveness, malleability) are further proposed. Scholars believe that the types of stigma source and the characteristics of stigma under different social conditions will jointly influence the formation of stigma[57].

At present, different scholars do not consistently agree on the concept of occupational stigma. The above-mentioned concepts of stigma proposed by Hughes[2], Ashforth and Kreine[41] are most commonly used in related studies. To this end, this study also defines the concept of occupational stigma toward psychiatrists in the aspect of physical, social and moral, and explores the specific causes of occupational stigma.

People with mental illnesses are frequently labeled as "violent," "offensive," "dangerous," and "aggressive"[58]. Compared to doctors in other clinical departments, the public attempts to consider that psychiatrists are more likely to encounter with violence and operate in a riskier setting[59,60], which were confirmed in a survey of psychiatric healthcare professionals[61]. In a study based on a sample in China, it was found that 78% of psychiatrists reported having experienced verbal abuse, compared to more than 30% who had experienced physical abuse[62]. Similar findings about psychiatrists' susceptibility to being hurt in medical injury incidents were discovered in surveys conducted in Germany[63], Ghana[64], Turkey[65], and Kuwait[66]. Psychiatrists are considered physically tainted because of the hazardous work environment.

For psychiatrists, associative stigma serves as a major source of occupational stigma[67,68]. Due to their frequent interactions with and treatment of patients with mental illness, psychiatrists may experience associative stigma that their clients may attach to them[9,69]. Patients with mental illness are a highly stigmatized group[70,71]. Their image is portrayed as negative, dangerous, and dishonest, and their social value is severely diminished[72]. To a large extent, the formation and development of stigma toward mental illness result from both religion and culture[73]. People with mental illness were thought to be possessed by evil spirits in Christianity[74] and Islam[75]. Expulsions and floggings, the confinement of mentally ill persons in jails and insane asylums, or even burning as a form of torture were all common practices in medieval Europe including Switzerland, Germany, and France[72]. Moreover, the thought of saving face is strongly ingrained in Chinese Confucian culture[76]. Concealing or avoiding the information that a family member suffers from a mental illness is typically done to preserve the family's reputation[68]. Even today, many people who suffer from mental illness and their family members still refrain from disclosing their situations to others for fear of prejudice and rejection. This ubiquitous phenomenon exists in nations where collectivist principles are valued[77,78]. The low

use of mental health services is partly caused by the stigma around mental illness[79]. Being socially tainted is, therefore, the most typical trait of psychiatrists' occupational stigma.

The possibility of being morally tainted among psychiatrists should not be overlooked. In treating people with mental illness, coercive measures such as seclusion, restraint, and forced medication are widely used[80]. Although patients perceive coercion as harmful to the treatment, healthcare professionals take it as an effective method of caring for, protecting, and treating patients[81]. Coercive measures have been the subject of heated debate in medical ethics, as well as legal scrutiny. Then, alternative options are sought and implemented by psychiatrists actively[82]. On the one hand, psychiatrists frequently consider physical and chemical restraints necessary, but on the other hand, they also recognize their potential to undermine patient rights and negatively impact the therapeutic relationship [83]. Therefore, it is understandable that psychiatrists may experience criticism, moral condemnation, and perceived intense moral pressure due to the dangers that coercive tactics may cause[84]. Now, an increasing number of researchers are advocating for improvements in the clinical practice of mental health medicine in order to reduce the use of coercive tactics[85]. In addition, electroconvulsive therapy (ECT) is frequently utilized to treat a variety of psychiatric diseases, including depression[86] and schizophrenia[87], with positive effects. However, ECT has been stigmatized and used as "proof" of psychiatrists' violence and harm to patients due to the media and the antipsychiatry movement[88]. As a result, psychiatrists also experience being morally tainted.

As illustrated above, the psychiatrists' occupational stigma is a complex idea that calls for multidimensional interpretations. Psychiatrists are seen as simultaneously being physically tainted, socially tainted, and morally tainted considering the potentially dangerous work environment, their exposure to and treatment of highly stigmatized populations, and the use of controversial and aggressive treatment methods. Occupational stigma could result in detrimental consequences for psychiatrists including isolation, discrimination, and loss of status and then made itself an indispensable factor to impede the advancement of medical and health services in mental health.

## THE MEASUREMENT OF PSYCHIATRISTS' OCCUPATIONAL STIGMA

To date, no occupational stigma scales have been developed for psychiatrists. In such a pertinent quantitative investigation, this can be accomplished by adapting other occupational stigma instruments (see Table 1). There are four different categories of scales that can be used to gauge public stigma toward psychiatrists. One type is that one dimension or more of the scale's items indicates occupational stigma. Richmond *et al*[89] created the psychometrically reliable Trust in Doctors in General (T-DiG) and Trust in the Health Care Team scales (THCT) to assess public trust in medical professionals and healthcare teams. Both the T-DiG and THCT include 29 items, which are from seven dimensions that refer to stigma-based discrimination, communication skills, system trust, loyalty, confidentiality, fairness, and general trust[89]. The stigma-based discrimination dimension contains three items that doctors or people who work in health care would unfairly treat patients with a history of mental illness, HIV, or drug abuse. Although T-DiG and THCT can reflect the public occupational stigma against physicians to some extent, the richness of occupational stigma is not accessible by measuring a single dimension of the scale. Meanwhile, occupational stigma, the complex concept, must be examined from multiple perspectives. The T-DiG and THCT contain only one item, physicians' unfair treatment of patients with a history of mental illness, and can only assess specific aspects of occupational stigma.

The second is to measure patients' occupational stigma against physicians. Fan *et al*[90] developed the Patient Toward Physician Occupational Stigma Scale (PPOSS), which consists of 19 items divided into three dimensions, namely stereotype, prejudice, and discrimination. The cognitive, affective, and behavioral components of occupational stigma are measured respectively[90]. Although the PPOSS was created specifically to assess physicians' occupational stigma, there are significant differences between physicians in different departments regarding the source, type, and extent of occupational stigma[49,52,91,92]. Psychiatric patients were not chosen as the study subjects during the development of the PPOSS, and items reflecting the psychiatrists' professional characteristics were lacking. As a result, more evidence is needed to determine whether the PPOSS is suitable for assessing patients' occupational stigma toward psychiatrists.

The third is to measure the public's social distance from physicians to examine their stigmatizing attitudes. A key element of psychological distance is social distance, which measures the degree of intimacy between various groups and between people within the same cohort[93]. The Social Distance Scale (SDS) has been utilized as a measurement tool in a study to examine public stigma and public behavioral inclinations toward physicians[94]. However, the SDS was originally designated not to measure occupational stigma but to measure the public distance perception between various nations, ethnic groupings, races, or groups[95]. The concepts of social isolation and occupational stigma are not interchangeable because they have obvious distinctions. Currently, the SDS is employed in research on occupational stigma less frequently, particularly when assessing public stigma toward physicians. Only a small number of studies used the scale[96].



**Table 1 Measurement tool of psychiatrists' occupational stigma**

Scale title	Target	Content	Item	Dimension
[89]THCT	Public	Measure public trust in medical professionals and health care team	29	7 dimensions: Stigma-based discrimination; communication skills; system trust; loyalty; confidentiality; fairness; general trust
[90]PPOSS	Patients	Measure the cognitive, affective, and behavioral components of occupational stigma	19	3 dimensions: Stereotype (6 items); prejudice (7 items); discrimination (6 items)
[94]SDS	Public	Measure stigmatizing public perceptions of and behavioral inclinations toward the physicians	8	Single dimension
[97]Public attitudes towards psychiatrists' questionnaire	Public	Measure the public's unfavorable views and impressions of psychiatrists' professionalism, mental health, occupational authority and ethical standards	8	Single dimension
[98]OSCS	Practitioners	Measure the stigma awareness of practitioners in the service industry	6	Single dimension
[100]E16-COVID19-S	Practitioners	Measure specific aspects of physician occupational stigma	16	3 dimensions: Personalized stigma (8 items); concerns of disclosure and public attitudes (5 items); negative experiences (3 items)
[101]PSCP	Physicians	Measure perception of COVID-19-induced stigma in healthcare workers	10	2 dimensions: Environmental stigmatization; the perception of personal stigmatization
[103]Self-designed stigma questionnaire	Physicians	Measure mental health stigma	12	Single dimension
[104]DAQ	Medical students	Measure stigmatization of physicians suffering from mental disorders	3	Single dimension
[105]SOSS-D	Doctors	Measure the stigma of occupational stress and burnout among physicians	11	3 dimensions: Perceived structural stigma (5 items); perceived individual stigma (3 items); perceived other related stigma (3 items)
[106]MHPSS	Mental Health Professionals	Measure occupational stress and burnout stigma in mental health professionals	13	4 dimensions: Perceived other stigma (3 items); perceived structural stigma (4 items); personal stigma (3 items); self-stigma (3 items)
[107]PIOSS	Physicians	Measure physicians' identification with negative labels, perceptions of devaluation and discrimination, as well as the denial and disapproval of their own profession	19	3 dimensions: Label identification (5 items); status loss (8 items); career denial (6 items)
[108]FSS	Forensic	Measure internalized occupational stigma	12	2 dimensions: Dangerousness/unpredictability (7 items); responsibility/blame (5 items)
[13]Mental health professionals' questionnaire	Mental health professionals	Measure the perceived associative stigma	4	Single dimension
[110]Measuring Psychiatrist occupational stigma scale (the scale a)	Psychiatrists	Measure perceived stigma in terms of the perception of societal stereotypes	16	Single dimension
[110]Measuring Psychiatrist occupational stigma scale (the scale b)	Psychiatrists	Measure self-stigma in terms of stereotype agreement	16	Single dimension
[110]Measuring Psychiatrist occupational stigma scale (the scale c)	Psychiatrists	Measure perceived stigma in terms of structural discrimination	5	Single dimension
[110]Measuring Psychiatrist occupational stigma scale (the scale d)	Psychiatrists	Measure discrimination experiences	13	Single dimension
[110]Measuring Psychiatrist occupational stigma scale (the scale e)	Psychiatrists	Measure stigma outcomes	5	Single dimension

COVID-19: Coronavirus disease 2019; DAQ: Discriminative attitude questionnaire; FSS: Forensic Stigma Scale; OSCS: Occupational Stigma Consciousness Scale; PIOSS: Physician Internalized Occupational Stigma Scale; PPOSS: Patient Toward Physician Occupational Stigma Scale; PSCP: Perception of Stigma due to COVID-19 in Physicians; SDS: Social Distance Scale; SOSS-D: Stigma of Occupational Stress Scale for Doctors; THCT: Trust in doctors in general and Trust in the health care team scales.

Fourth, a non-standard assessment method was used to examine the public's negative perceptions of psychiatrists. Ta *et al*[97] used a self-designed Public attitudes towards psychiatrists questionnaire with eight items to assess public perceptions of psychiatrists. The questionnaire mainly measured the public's negative attitude and views of psychiatrists' professionalism, mental health, occupational authority and ethical standards[97]. Nevertheless, it is hard to state the reliability and validity of the research results because the questionnaire's reliability and validity were not assessed.

The scales that can be used to measure the self-stigma of psychiatrists could be divided into four categories. The first is the scale used to gauge psychiatrists' sensitivity to stigma. The Occupational Stigma Consciousness Scale (OSCS) is the most frequently used tool in studies on physicians' occupational stigma. The OSCS comprises six one-dimensional items that assess practitioners' perceptions of public stigmatization of their work. However, the OSCS was originally developed based on call center service workers and was primarily used to assess the stigma awareness of practitioners in the service industry[98]. According to the needs of researchers' studies, they appropriately adapted the scale for use in quantitative studies of various occupational stigma in subsequent studies[99]. It should not be ignored that psychiatrists are part of a high-prestige profession that differs significantly from dirty work regarding stigma manifestations and negative consequences.

The second is the scale for measuring specific aspects of physician occupational stigma. Based on the severe acute respiratory syndrome stigma scale, Mostafa *et al*[100] developed the new coronavirus disease 2019 (COVID-19) Stigma Scale (E16-COVID19-S)[100]. Similar to this, Oktar *et al*[101] developed the Perception of Stigma due to COVID-19 in Physicians (PSCP) with 10 items, including two dimensions that are environmental stigma and individual stigma perception[101]. These two scales measure specific components of physicians' occupational stigma, but fail to reflect its full spectrum of connotations.

The stigmatization of occupational stress and burnout among physicians has been the research focus in this field. For instance, Riley *et al*[102] discovered that physicians experience high stigma in mental health, work stress and burnout, manifested as inability to admit vulnerability and insistence on working, even if unwell[102]. A study by Wijeratne *et al*[103] on physicians' mental health stigma found that they tend to conceal their mental health conditions from colleagues and are less likely to seek help because there is a belief that physicians suffering from depression or anxiety disorders are perceived as untrustworthy[103]. This study applied a self-designed 12-item stigma questionnaire as a survey tool, which was not strictly tested for reliability nor validity, but only reported internal consistency coefficient values. Zarzycki *et al*[104] adopted a self-designed Discriminative Attitude Questionnaire (DAQ) to examine medical students' stigmatization of physicians with mental disorders[104]. The DAQ includes only three non-standardized items and is only applicable for assessing stigma regarding mental disorders. Furthermore, Clough *et al*[105] developed the 11-item Stigma of Occupational Stress Scale for Doctors (SOSS-D). There are three dimensions extracted in the SOSS-D including perceived structural stigma, perceived individual stigma, and perceived other related stigma[105]. To measure occupational stress and burnout stigma in mental health professionals, Clough *et al*[106] created the Mental Health Professional Stigma Scale (MHPSS)[106]. There are 17 items total in the MHPSS, which are broken down into four dimensions: Perceived other stigma, perceived structural stigma, personal stigma, and self-stigma. Stigmatizing attitudes, stress and burnout among psychiatrists can pose serious threats to their professional development. However, scales for measuring the stigma of occupational stress and burnout specifically among psychiatrists are lacking and should be developed in future research.

The third part include scales for assessing physicians' internalized occupational stigma. Fan *et al*[107] created a 19-item Physician Internalized Occupational Stigma Scale (PIOSS) divided into three dimensions including label identification, status loss, and career denial. The PIOSS scale is primarily used to evaluate physicians' identification with negative labels, perceptions of devaluation and discrimination, and denial and disapproval of their profession[107]. Besides, Healey *et al*[108] developed the Forensic Stigma Scale (FSS) with 12 items falling into two dimensions that refer to danger/unpredictability and blame/responsibility[108]. Although PIOSS and FSS have strong validity and reliability, the study subject did not include psychiatrists.

Fourth, there is a non-standard tool for assessing the perception of stigma among psychiatrists. A self-designed mental health professional's questionnaire was used in the study by Verhaeghe *et al*[109]. Four items comprise the questionnaire, which mainly measures the perceived associative stigma among psychiatrists and other mental health professionals. However, Cronbach's alpha coefficient for the questionnaire was just 0.51 since it had not undergone a rigorous reliability test[109].

Fifth, there is a scale specifically designed to measure the occupational stigma of psychiatrists. The World Psychiatric Association (WPA) has developed specific action plans to reduce the stigma toward psychiatry and psychiatrists. One of the essential tasks is to develop standardized questionnaires[110]. The basic version comprises five scales: (1) Perceived stigma in terms of the perception of societal stereotypes; (2) self-stigma in terms of stereotype agreement; (3) perceived stigma in terms of structural discrimination; (4) discrimination experiences; and (5) stigma outcomes. The scales (1) and (2) consist of 16 items measuring the professionals' competence, professional conduct, and personality, as well as the stigma of psychiatry as a medical specialty and its treatment methods. Scale (3) and scale (5), consisting of five items, assesses the social aspects of stigma and the negative consequences of stigma. Scale (4)

consists of 13 items, assessing experiences made in contact with doctors from other disciplines and experiences made in personal life. The scale developed by the WPA is a valid tool for assessing the internalization of occupational stigma among psychiatrists.

## THE INTERVENTION OF PSYCHIATRISTS' OCCUPATIONAL STIGMA

The study of stigma intervention strategies has been the main research subject in the stigma field. To effectively intervene with various demographics and stigma, the approaches used can be cross-referenced[111,112]. In a systematic evaluation of stigma intervention strategies, which included research findings from various countries (low-, middle-, and high-income), stigmatized populations (such as those with AIDS, mental health disorders, and leprosy), intervention targets (such as medical personnel, family members, and community members), and intervention strategies (such as contact, education, and training), the analysis discovered some similarities in the strategies, measures, and intervention outcomes achieved when intervening with different types of stigma[113]. A review of stigma intervention strategies can serve as a theoretical foundation for psychiatrists' occupational stigma intervention practice. Stigma intervention strategies can be roughly classified into six categories.

First, the protest approach is a strategy for the stigmatized community to voice their opinions and express their disapproval through public declarations, media exposure, and stigma-related commercials to minimize stigma[114]. However, protest tactics risk escalating stigma, leading to public conflict and rebellion[115]. Therefore, they are used relatively infrequently. Furthermore, compared to more conventional rallies and demonstrations, the usage of social media platforms on the internet is growing. For instance, rebuttals to stigmatizing attitudes about mental illness are shared on Twitter during Mental Health Awareness Week, an annual effort in May[116]. Similarly, Depression Awareness Week is coordinated on Twitter with the primary objective of lowering the stigma attached to mental illness [117].

Second, educational strategies are applied to alter erroneous beliefs and reduce stigma by delivering accurate information to the intervention target[118]. A few commonly employed techniques are class lectures, anti-stigma training, special lectures, workshops, role plays, case studies, watching instructional videos, reading professional publications, and creating self-reflection reports (see Table 2). In the pertinent literature, more attention is paid to the stigmatization of medical professionals and the general public toward people with mental illnesses. In contrast, relatively few research studies have been conducted on psychiatrists' occupational stigma. Educational strategies effectively change healthcare professionals' attitudes toward patients with mental illnesses and increase contact willingness and frequency[119-122]. Besides that, it is conducive to reducing self-stigma in patients with mental illnesses through education[123]. Moreover, interventions can be implemented in stand-alone educational formats, such as workshops[124] and educational videos[125], or in a combination of formats to achieve more significant results. For example, in Education Not Discrimination[126] and Mental Health First Aid[127], a combination of presentations, videos, action plan ideas, case discussions, and role plays is used.

Third, the contact strategy aims to strengthen relationships with stigmatized people to alleviate adverse stereotypes. In stigma interventions, the contact strategy has been applied most frequently and has produced promising benefits[128]. It is possible to boost intergroup connections and minimize prejudice by increasing the frequency of public contact with stigmatized individuals, especially high-quality contact[129]. According to Corrigan *et al*[130], contact approaches are approximately three times more effective than educational strategies with more prominent and persistent impact[130]. Even short-term exposure has the potential to improve attitudes and understanding about stigma[131] and lower stigma levels[132]. The question of how to accomplish the desired intervention outcomes and guarantee the effectiveness of the contact method has always been the focus of scholars[131]. Corrigan *et al*[133] proposed that the most critical factors influencing the effectiveness of interventions are design, target, staff, message, evaluation and follow-up in five areas[133]. Specifically, the intervention can be delivered face-to-face, with a person with extensive life experience serving as a speaker, designing a program that matches the target audience's characteristics and delivering the intervention through storytelling. After the intervention's completion, evaluation and follow-up of the intervention effects are also required[134].

In terms of forms of intervention, direct and indirect contact are included[135]. Direct contact as a promising anti-stigma strategy can improve communication and cooperation between patients with mental illnesses and other patients, as well as between patients and family members and the general public[136]. By increasing contact between health science students and people with mental illnesses in the Co-Production with Dialogue Program for Reducing Stigma, it is possible to significantly reduce public stigma and improve mutual understanding between patients and students[137]. Psychoeducational materials, face-to-face workshops, and interventions based on cognitive behavioral therapy were generally well received in the workplace-based multi-country intervention tackling depression, anxiety, and mental illness-related stigma study[138]. In the "Honest, Open, Proud" project, self-exposure and community-based participatory research were used to reduce self-stigma among individuals with

Table 2 Intervention of education

Strategy	Target	Content	Objective
[119]Education training	Family caregivers of patients with schizophrenia	Teaching related knowledge and skills (90 min/8 sessions)	Reduce stigma toward patients with schizophrenia
[120]Anti-stigma training	Care assistant workers	Financial assistance policy of care assistant workers + mental health knowledge (2 h) + stigma related to mental illness (1 h)	Reduce stigma toward patients with mental illness
[121]Anti-stigma training	Community mental health staff	Two parts: Knowledge introduction + stigma related to mental illness (1 d)	Increase mental health knowledge; Reduce stigma toward patients with mental illness and social distance
[122]Anti-stigma training	Probation officers	Six modules in PPT about overview of severe and persistent mental illnesses; information about key diagnoses and the medications; services associated with those diagnoses	Increase mental health knowledge; Reduce stigma toward mental illness and behavior tendency
[123]NECT	Persons with severe mental illness	20 sessions (1 h): (1) Introduction (1 wk); (2) Psychoeducation (3 wk); (3) Cognitive restructuring (8 wk); and (4) Narrative enhancement (8 wk)	Reduce stigma toward mental illness
[124]Educational workshop	High School Students	Classroom-based workshop: (1) Introduction: Mental illness and public stigma; (2) Education; and (3) Treatment of mental illness and available community resources (1 h)	Reduce stigma toward mental illness
[125]Educational videos	High school students and college students	Educational film (10 min): (1) BMM about mental illness; or (2) RCM advocated by experts	Reduce more stigma in BMM than RCM
[126]END	Medical students	Short lecture; testimonies about the experiences and stigma; role-plays	Increase mental health knowledge; Reduce stigma toward mental illness and behavior tendency
[127]Mental health first aid training	Pharmacy and non-pharmacy students	Identify, understand, and respond to signs of mental illnesses and substance use disorders (8 h); Methods: Videos, role play sessions, action plan creation, discussion, and other interactive activities	Reduce stigma toward mental illness

BMM: Biomedical messages; END: Education not discrimination project; NECT: Narrative enhancement cognitive therapy; PPT: Power point; RCM: Recommended messages.

mental illness[139].

The typical indirect contact method is video contact. Researchers prefer video contact because of its low cost, broad audience reach, reusability, low resource possession, and ease of dissemination[130]. Short video interventions are an essential and effective intervention in studies of depression stigma and help-seeking attitude stigma[140], mental health-related stigma[141], and mental illness stigma[142].

Fourth, comprehensive strategies combine two or all three of the protest, education, and contact strategies. Education and contact strategies are most frequently employed (see Table 3). For instance, Tan *et al*[143] discovered that increasing interaction (direct contact) and attending lectures (education) improved people's attitudes and levels of acceptance toward those who suffer from depression[143]. Furthermore, a three-stage intervention paradigm was subsequently developed by Ahuja *et al*[144]. Specifically, participants in the intervention were required to watch a dance drama to learn about common misconceptions and correct perceptions of mental illness. Then, individuals were asked to listen to an informational lecture and directly communicate with people suffering from mental illness. Hawke *et al*[145] used an indirect intervention strategy, showing the recorded stage play to healthcare providers, college students, people with bipolar disorder and their friends and family members, and the general public. The findings revealed that education and exposure *via* video approach similarly reduced the stigma of bipolar disorder among intervention subjects and had sound delayed effects[145].

Some studies have attempted to integrate anti-stigma education into school curricula. Ma *et al*[146], for example, designed a teaching component covering knowledge, experience, and action, as well as education strategies, direct contact, and indirect contact, to reduce the stigma of mental illness among medical students[146]. Case discussions and systematic curriculum instruction[147,148] can be used to achieve intervention goals[149]. Both educational and direct contact strategies, as well as educational and indirect contact strategies, have demonstrated practical intervention effects[150-168]. However, in a study of an adolescent population, educational strategies alone were superior to combined education and contact strategies in reducing mental illness stigma[165]. The researchers suggest that the young age of the intervention subjects may have contributed to this result and that contact strategies should be used with caution when intervening with adolescents. As a result, more evidence is needed to determine whether the combined strategy is superior to the single intervention strategy in various age groups.

Fifth, systematic intervention strategies emphasize the importance of considering the interplay of factors at different levels when intervening with stigma and integrating different types of intervention



**Table 3 Intervention of comprehensive strategies**

Strategy	Type	Content	Target	Objective
[143]Lecture	Education + direct contact	Video + personal experience + questions and answers (50 min)	Undergraduate	Increase knowledge of depression and help-seeking willingness
[144]Three-stage intervention paradigm	Education + direct contact + indirect contact	Dance-drama + lectures + direct contact (2 h)	Undergraduate	Increase understanding and tolerance toward patients with mental illness; Decrease negative review, negative labels and social distance toward patients with mental illness
[145]Filmed theatrical intervention based on a personal narrative	Education + indirect contact	Filmed version of a one-woman stage play performed by a recognized educator and speaker on the lived experience of mental illness and recovery (50-min)	Health-care service providers, university students in a health-care-related course, people with BD and their friends and family members and the general public	Decrease stigma toward people with BD among health-care service provider (significant/sustainable); others (more limited)
[146]Course	Education + direct contact + indirect contact	Lecture + watch videos/movie + in-class discussion + direct contact + roleplay + action project (18 wk)	Occupational therapy students	Decrease stigma toward mental illness and social distance
[147]Course	Education + direct contact + indirect contact	Teaching (42 h) + movie (3/4 h) + contact + presentations (2 h) + clinical correlations (6 h)	Medical students	Increase student confidence; Decrease stigma toward mental illness
[148]Course	Education + direct contact + indirect contact	Face-to-face (45 min) + video-based contact (40 min) + educational lecture (90-min)	Medical students	Decrease stigma toward mental illness
[149]Case discussion	Education + direct contact + indirect contact	Document patient strengths and treatment recommendations; Group presentation discussion (10 wk)	Primary care providers	Increase willingness to help and hope for recovery; Decrease negative stereotypes
[150]Course	Education + direct contact	Personal experience + questions and answers (69-90 min)	Pharmacy students	Decrease stigma toward mental illness
[151]Workshop	Education + direct contact	Workshop (4 d): Knowledge teaching + personal experience + questions and answers	Public Health Services administrators	Decrease stigma toward patients with mental illness
[152]Workshop	Education + direct contact	Workshop (1 h): Knowledge teaching + personal experience	Social work students	Decrease stigma toward mental illness
[153]Course	Education + direct contact	Panel discussion (2 h) + visit rehabilitation center + Small group discussion (20 min)	Medical students	Decrease stigma toward patients with mental illness and psychiatry
[154]Curriculum	Education + direct contact	Courses + clinical practice + stigma views and policy (14 d)	Community mental health staff	Increase related knowledge; Decrease stigma
[155]Education program	Education + direct contact	Enhancing contact model: Psychoeducational training (4 wk) + single and group family contact (8 wk)	Family caregivers of persons with schizophrenia	Decrease stigma toward mental illness
[156]Education program	Education + direct contact	Lectures (2 h) + contact with people with lived experience of mental illness (1 h)	Community health and care staff	Decrease stigma toward mental illness
[157]Course	Education + indirect contact	Teaching + videos (3 h)	Psychology students	Improve attitude toward schizophrenia
[158]Virtual program	Education + indirect contact	Three consecutive interventions (14 h/2 d): 2 sessions (2.5 h) + e-contact (3.5 h) + project-based learning (5.5 h)	University students	Decrease stigma toward mental disorders
[159]Educational video	Education + indirect contact	Social contact film with general mental health education (30 min/2 mo intervals in 1 year)	University students	Decrease mental illness-related stigma
[160]Course	Education + indirect contact	Interactive workshop-style sessions (3.5 h/15 wk)	Primary care physicians	Increase skill and confidence; Decrease stigma toward mental illness
[161]Educational	Education +	Video-based contact intervention	Primary care nurses	Improve attitude toward patients with

Video	indirect contact	(VBCI): Elements of psychoeducation and interviews (5 min)		mental illness in short term
[163]Educational video	Education + indirect contact	YouTube videos (5-10 min)	Muslim adults aged 18 and over	Decrease stigma toward mental illness
[164]Lecture	Education + direct contact	Personal experience + Interaction	High school students	Decrease stigma toward mental illness
[165]Education program	Education + direct contact	Educational program (1 d)	Adolescents	Education decreases stigma toward mental illness; Direct contact fails
[166]Course	Education + direct contact	Education with PBL: Problem-based learning model + direct contact	Nursing students	Improve positive attitude toward mental illness
[167]Online focus group	Education + direct contact + indirect contact	Education + relationship strategies + media contact	Senior undergraduate nursing students	Minimize the negative impact on the people they take care of
[168]Sessions	Education + direct contact	Five sessions (3 d)	Caregivers of mental health service users	Improve family relations and understanding of mental illness

BD: Bipolar disorder.

strategies into a unified framework[162]. According to Heijnders and Van Der Meij[169], the effects obtained by employing a single level of intervention strategy or intervening with a single target group are frequently insufficient and necessitate the integration of multiple factors[169]. As a result, Cook *et al* [170] developed a three-level ecosystem model that divides stigma intervention strategies into individual, interpersonal, and structural levels, arguing that interventions at any level can affect the effectiveness of interventions at other levels[170]. Individual-level interventions concentrate on enhancing coping mechanisms used by stigmatized individuals or altering the attitudes and conduct of stigma abusers. By enhancing engagement and contact between the group of stigma abusers and the group being stigmatized, interpersonal interventions aim to reduce intergroup barriers and foster greater mutual understanding. Through the creation of pertinent laws and regulations to lessen prejudice and discrimination, intervention at the structural level is primarily seen from the perspective of the socio-political environment.

Additionally, Heijnders and Van Der Meij[169] suggested a five-level stigma intervention program (individual level, interpersonal level, social level, organizational and institutional level, and governmental level)[169]. In particular, professional counselling and therapy can be used at the individual level to help the stigmatized population cope with stigma; interventions at the interpersonal level concentrate on improving how the stigmatized population interacts with the elements of their environment in order to gain care and support; and interventions at the social level focus on changing public attitudes through education, advocacy, or contact strategies. At the organizational and institutional levels, specialized training or the development of internal systems can be adopted to reduce stigma in a group, organization, or institution; at the government level, interventions focus on the establishment of relevant laws and policies, as well as the use of coercive measures, to reduce stigmatized speech and behavior.

Sixth, increasing researchers have attempted to incorporate psychotherapeutic approaches and techniques into stigma interventions and have achieved promising results (see Table 4). Hong *et al*[171] used mindfulness training to intervene with self-stigma in depressed patients and got excellent and long-lasting results[171]. It was found by Clinton *et al*[172] that Positive Empathy Intervention (imagining positive contact with the stigmatized person and establishing a positive emotional connection) was superior to traditional contact methods in terms of stigma reduction[172]. Cognitive-behavioral Intervention and Wise Intervention have been shown to increase perceived social support and reduce self-stigma in individuals[173,174]. Furthermore, Visual Information Image Interventions [175] and Work Integration[176] assist individuals in gaining correct knowledge, changing perceptions, and enhancing social inclusion, thereby reducing stigma.

Art Therapy is currently widely applied in the work practice of mental health education in schools. When intervening with the stigma of mental illness in adolescents, art therapy methods such as film, drama, and role-playing can be adopted[177,178]. Picture storytelling or fiction writing can enable the individual's indirect contact with an AIDS patient[179]. The Petkari's study[180] discovered that watching a series of movies about mental illness (over 10 wk) accelerated college students' agency, compassion, and proximity toward patients[180]. Acceptance and Commitment Therapy (ACT), based on relational framework theory, has been dubbed the "Third Wave" of behavioral therapy[181]. In interventions for weight stigma[182], substance use disorder stigma[183], and homosexuality stigma [181], ACT has shown significant positive effects. Furthermore, Narrative Therapy[184], Written

**Table 4 Other intervention strategies**

Strategy	Objective
[171]Positive thinking training	Intervene with self-stigma in depressed patients
[172]Positive empathy intervention	Establish a positive emotional connection with stigmatized person to reduce stigma
[173]Cognitive-behavioral intervention	Reduce self-stigma associated with mental illness
[174]Wise intervention	Increase perceived social support; Reduce personal discrimination
[175]Visual information image intervention	Increase sense of support and engagement; Reduce stigma
[176]Work integration	Reduce perceived stigma of workers with serious mental illness
[177-180]Art therapy	Reduce mental illness stigma of adolescents
ACT	Reduce weight stigma[182], substance use disorder stigma[183], and homosexuality stigma[181]
[184]Narrative Therapy	Reduce the level of stigma for patients with permanent enterostomies
[185]Written expression-based emotional intervention	Reduce negative emotion
[186]Sand tray therapy	Improve the development of self-esteem in sexually abused adolescents
[187]CBGT	Reduce the level of stigma in the population of people living with HIV
[188]Intergenerational choir	Reduce the stigma of Alzheimer's patients and increasing positive attitudes
[189]FSM	Reduce stigma and promote mental health help-seeking behavior

ACT: Acceptance and commitment therapy; CBGT: Compassion-based group therapy; FSM: Fresh start mindset framing.

Expression-based Emotional Intervention[185], Sand Tray Therapy[186], Compassion-Based Group Therapy[187], Intergenerational Choir[188], and Fresh Start Mindset Framing[189] can be used as stigma interventions.

To summarize, in the intervention of psychiatrists' occupational stigma, protest, education, contact, integrated and systematic intervention strategies, and multiple approaches such as positive thinking training, cognitive behavioral therapy, ACT, and art therapy are available to be adopted. In order to achieve optimal results, the appropriate intervention method must be chosen based on the characteristics of the intervention target. It should be noted that while the above intervention strategies have been applied to various types of stigma interventions with positive results, more evidence is needed to support whether the strategies mentioned above are applicable to psychiatrists' occupational stigma interventions and how they can be implemented to achieve significant intervention effects.

## DISCUSSION

In summary, previous studies have yielded results regarding the concept, measurement, and intervention of psychiatrists' occupational stigma. Nevertheless, there is room for improvement in this field. This review aimed to elaborate on the thinking and practices of related issues. Therefore, future research should consider improving on the four aspects outlined below.

First, clarify and refine the concept of psychiatrists' occupational stigma. Although various scholars have defined the concept of occupational stigma, related research has focused more on dirty work. Indeed, there is a paucity of research specifically on psychiatrists' occupational stigma. Based on Ashforth and Kreiner's research[43], this paper elaborates on the sources and dimensions of psychiatrists' occupational stigma, namely physically, socially, and morally. This theoretical framework allows for the development of relevant quantitative research and intervention studies. Although Ashforth and Kreiner's concept[43] of occupational stigma has gained widespread acceptance, it is not formally classified other than according to the work content of an occupation. As such, it does not reflect the cognitive, emotional or behavioral components of occupational stigma. In addition, the understanding of occupational stigma is not consistent across disciplines. Future research should combine theories from other disciplines (including individual cognitive models, social identity theory, self-verification perspectives, and other conceptual models) to further explore and extend the conceptual connotations of psychiatrists' occupational stigma. Furthermore, the theoretical framework of occupational stigma should be combined with statistical analysis to determine the multiple dimensions of psychiatrists' occupational stigma. This study provides such a theoretical basis for future measurement and intervention studies.

Second, develop specific tools to measure psychiatrists' occupational stigma. Lately, as public awareness of the harm of occupational stigma has increased, relevant measurement tools have been refined. However, some existing instruments are not sufficiently reliable nor valid, and tools specifically designed to assess psychiatrists' occupational stigma are lacking. As no consensus exists on the conceptual and operationalization scope of occupational stigma, there is inconsistency in developing relevant dimensional and measurement scales. Furthermore, most tools lack rigorous cross-cultural consistency. Future research should consider the following: (1) Define the dimensional scale and classification of psychiatrists' occupational stigma based on a multidisciplinary synthesis; (2) Develop special assessment tools for different stigma types (public stigma and self-stigma) and cohorts (psychiatrists, psychiatric students, mental illness patients, patients' families, and the public); (3) Expand the sample scope across different races, countries and age ranges to determine the impact of cross-cultural backgrounds and generational effects on the results; and (4) Based on traditional self-reporting questionnaires, adopt more indirect survey methods such as virtual reality technology, videos, or games allowing for measurement methods with higher ecological validity and aligned to life situations that yield a realistic and contextualized understanding.

Third, improve intervention strategies for psychiatrists' occupational stigma. Intervention strategies specifically applicable to psychiatrists' occupational stigma are currently lacking. Initially, when intervening for psychiatrists' occupational stigma, other types of stigma intervention strategies may be considered. However, undoubtedly these could lead to biases in the effectiveness of the intervention. Therefore, future studies should test whether existing intervention strategies are suitable for psychiatrists. Follow-up horizontal comparison and longitudinal studies can be conducted on the effects of the three common intervention strategies (protest, education, and contact), as well as integrated, systematic, or other strategies, seeking to find the most appropriate traditional intervention strategies and setting for psychiatrists. It is necessary to acknowledge that stigma may exacerbate or impede such processes as psychological[190,191] and behavioral responses[192,193] or social relationships, intensifying stress and burnout that could result in mental health disorders. Some studies have demonstrated that educational interventions which provide in-depth information about the negative effects of stigma on mental health professionals can be effective in decreasing stigma, especially for general healthcare professionals with little or no formal mental health training. Alternatively, future research should develop unique, simple, and effective intervention strategies tailored to the characteristics of psychiatrists. Combining intervention studies with experimental studies could identify simple and accessible ways to reduce occupational public stigma directed toward, and self-stigma experienced by, psychiatrists. Importantly also consider that the effects of a particular intervention may not be uniform among psychiatrists from different countries, cultural backgrounds, or years of practice. Thus, when formulating intervention strategies, full consideration should be given to differences in intervention targets.

Fourth, identify cross-cultural consistency or differences in psychiatrists' occupational stigma. Self-evidently, psychiatrists' occupational stigma can vary culturally. Future research should explore the consistency or differences in occupational stigma concepts, measurements, and interventions among psychiatrists in cross-cultural settings. For example, in Chinese culture, traditional ideas conveyed across millennia, such as Confucianism, Taoism, Buddhism, and folklore, have influenced Chinese thinking and behavior towards self-regulation; this combined with strong family values and a face-saving culture, deems mental illness as both a personal sin and a family shame[194]. Other regions may have different stigma levels toward mental illness[195], so cultural traditions may influence the inception of psychiatrists' occupational stigma. Is it possible that perceptions of psychiatrists' occupational stigma differ across cultures? Does this influence the measurement and treatment of psychiatrists' occupational stigma? Such interrogations have yet to be confirmed through in-depth research.

## LIMITATIONS

It should be acknowledged that there were certain deficiencies in the process of screening and synthesizing many studies in this literature review. First, the selection criterion, which only considered English and French literature, was limiting. Therefore, it is possible that relevant studies that satisfied other inclusion criteria were excluded. Thus, overall integrity is somewhat lacking. Second, the literature search was carried out by both professors and students. Irrelevant studies, duplicates, or those arising from incorrect search results were excluded. However, given the excessive literature search results, no secondary duplication test was undertaken. Therefore, it is impossible to determine whether excluded studies should have been included, indicating a lack of rigor. Finally, this review was based on the authors' analyses and synthesis of the literature; although the study seeks to remain objective, it contains some subjectivity.



## CONCLUSION

By surveying the existing literature, this literature review has proposed a theoretical reference of the concept, measurement, and intervention methods for psychiatrists' occupational stigma. Psychiatrists' occupational stigma is a complex concept that should be interpreted in multiple dimensions. Psychiatrists are associated with three types of stigma (physical, social, and moral taint) because of the dangers of their work environment, their exposure to and treatment of high-stigma groups, or their use of controversial or aggressive treatments. Currently, there is no occupational stigma scale applicable specifically to psychiatrists. Relevant quantitative research could achieve this by adapting other occupational stigma scales. **Table 1** summarizes eight possible categories of occupational stigma measurement tools for psychiatrists, including four types for public stigma and another four for self-stigma. Currently, there are few studies on occupational stigma interventions for psychiatrists. Therefore, a theoretical reference for identifying relevant intervention practices for psychiatrists' occupational stigma is required. This study has classified such stigma intervention strategies into six categories: Protest, education (**Table 2**), contact, comprehensive strategies (**Table 3**), systemic strategies, and means of incorporating psychotherapeutic approaches (**Table 4**).

Given that research on psychiatrists' occupational stigma has received insufficient attention and discussion in the academic community, this study provided a theoretical basis and support for future practical research. The theoretical significance of this review lies in that it refines the concept and structure of psychiatrists' occupational stigma, expands the general research field of occupational stigma, and encourages the mutual discussion of multi-disciplinary occupational stigma theories. This study further outlines relevant empirical research for the development of specialized measurement tools and creative implementations of effective interventions to reduce psychiatrists' occupational stigma, thereby promoting the healthy development of psychiatry and physician-patient relationships.

## FOOTNOTES

**Author contributions:** Shi XL wrote the article and conducted the literature analysis; Li LY contributed to the data collection and approved the final version of the article; Fan ZG contributed to the article framework guidance and revision.

**Conflict-of-interest statement:** All the authors report no relevant conflicts of interest for this article.

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** China

**ORCID number:** Xiao-Li Shi 0000-0001-5248-9299; Lu-Yao Li 0000-0002-4364-1751; Zhi-Guang Fan 0000-0001-9959-4432.

**S-Editor:** Li L

**L-Editor:** Filipodia

**P-Editor:** Li L

## REFERENCES

- 1 Kreiner G, Mihelcic CA, Mikolon S. Stigmatized Work and Stigmatized Workers. *Annu Rev Organ Psychol Organ Behav* 2022; **9**: 95-120 [DOI: [10.1146/annurev-orgpsych-012420-091423](https://doi.org/10.1146/annurev-orgpsych-012420-091423)]
- 2 Hughes EC. Men and their work. Quid Pro, LLC., 1958
- 3 Kreiner GE, Ashforth BE, Sluss DM. Identity Dynamics in Occupational Dirty Work: Integrating Social Identity and System Justification Perspectives. *Organ Sci* 2006; **17**: 619-636 [DOI: [10.1287/orsc.1060.0208](https://doi.org/10.1287/orsc.1060.0208)]
- 4 Link BG, Phelan JC. Conceptualizing Stigma. *Annu Rev Sociol* 2001; **27**: 363-385 [DOI: [10.1146/annurev.soc.27.1.363](https://doi.org/10.1146/annurev.soc.27.1.363)]
- 5 Fujishiro K, Xu J, Gong F. What does "occupation" represent as an indicator of socioeconomic status?: exploring occupational prestige and health. *Soc Sci Med* 2010; **71**: 2100-2107 [PMID: [21041009](https://pubmed.ncbi.nlm.nih.gov/21041009/) DOI: [10.1016/j.socscimed.2010.09.026](https://doi.org/10.1016/j.socscimed.2010.09.026)]
- 6 Feingold JH, Drossman DA. Deconstructing stigma as a barrier to treating DGBI: Lessons for clinicians. *Neurogastroenterol Motil* 2021; **33**: e14080 [PMID: [33484225](https://pubmed.ncbi.nlm.nih.gov/33484225/) DOI: [10.1111/nmo.14080](https://doi.org/10.1111/nmo.14080)]
- 7 Sharma S, Lal Gautam P, Sharma S, Kaur A, Bhatia N, Singh G, Kaur P, Kumar A. Questionnaire-based Evaluation of Factors Leading to Patient-physician Distrust and Violence against Healthcare Workers. *Indian J Crit Care Med* 2019; **23**: 302-309 [PMID: [31406431](https://pubmed.ncbi.nlm.nih.gov/31406431/) DOI: [10.5005/jp-journals-10071-23203](https://doi.org/10.5005/jp-journals-10071-23203)]

- 8 **Gaebele W**, Zäske H, Zielasek J, Cleveland HR, Samjeske K, Stuart H, Arboleda-Florez J, Akiyama T, Baumann AE, Gureje O, Jorge MR, Kastrup M, Suzuki Y, Tasman A, Fidalgo TM, Jarema M, Johnson SB, Kola L, Krupchanka D, Larach V, Matthews L, Mellisop G, Ndeti DM, Okasha TA, Padalko E, Spurgeoun JA, Tyszkowska M, Sartorius N. Stigmatization of psychiatrists and general practitioners: results of an international survey. *Eur Arch Psychiatry Clin Neurosci* 2015; **265**: 189-197 [PMID: [25190351](#) DOI: [10.1007/s00406-014-0530-8](#)]
- 9 **Catthoor K**, Hutsebaut J, Schrijvers D, De Hert M, Peuskens J, Sabbe B. Preliminary study of associative stigma among trainee psychiatrists in Flanders, Belgium. *World J Psychiatry* 2014; **4**: 62-68 [PMID: [25250223](#) DOI: [10.5498/wjp.v4.i3.62](#)]
- 10 **Lai JYM**, Chan KW, Lam LW. Defining who you are not: The roles of moral dirtiness and occupational and organizational disidentification in affecting casino employee turnover intention. *J Bus Res* 2013; **66**: 1659-1666 [DOI: [10.1016/j.jbusres.2012.12.012](#)]
- 11 **Baran BE**, Rogelberg SG, Carello Lopina E, Allen JA, Spitzmüller C, Bergman M. Shouldering a silent burden: The toll of dirty tasks. *Hum Relat* 2012; **65**: 597-626 [DOI: [10.1177/0018726712438063](#)]
- 12 **Huang B**, Ma L, Huang L. My Work Is Meaningless: The Consequences of Perceived Occupational Stigma for Employees in High-Prestige Occupations. *Front Psychol* 2022; **13**: 715188 [PMID: [35572310](#) DOI: [10.3389/fpsyg.2022.715188](#)]
- 13 **Daniela RE**, Andrea S, Peter F. Psychiatrists' self-stigma, the DGPPN guideline for psychosocial interventions, and contemporary treatment strategies. *Eur Arch Psychiatry Clin Neurosci* 2015; **265**: 171-2 [PMID: [25708457](#) DOI: [10.1007/s00406-015-0586-0](#)]
- 14 **Yao H**, Wang P, Tang YL, Liu Y, Liu T, Liu H, Chen Y, Jiang F, Zhu J. Burnout and job satisfaction of psychiatrists in China: a nationwide survey. *BMC Psychiatry* 2021; **21**: 593 [PMID: [34819029](#) DOI: [10.1186/s12888-021-03568-6](#)]
- 15 **Shanafelt TD**, West CP, Sinsky C, Trockel M, Tutty M, Satele DV, Carlasare LE, Dyrbye LN. Changes in Burnout and Satisfaction With Work-Life Integration in Physicians and the General US Working Population Between 2011 and 2017. *Mayo Clin Proc* 2019; **94**: 1681-1694 [PMID: [30803733](#) DOI: [10.1016/j.mayocp.2018.10.023](#)]
- 16 **Zhang Y**, Feng X. The relationship between job satisfaction, burnout, and turnover intention among physicians from urban state-owned medical institutions in Hubei, China: a cross-sectional study. *BMC Health Serv Res* 2011; **11**: 235 [PMID: [21943042](#) DOI: [10.1186/1472-6963-11-235](#)]
- 17 **Tateno M**, Jovanović N, Beezhold J, Uehara-Aoyama K, Umene-Nakano W, Nakamae T, Uchida N, Hashimoto N, Kikuchi S, Wake Y, Fujisawa D, Ikari K, Otsuka K, Takahashi K, Okugawa G, Watanabe N, Shirasaka T, Kato TA. Suicidal ideation and burnout among psychiatric trainees in Japan. *Early Interv Psychiatry* 2018; **12**: 935-937 [PMID: [28786526](#) DOI: [10.1111/eip.12466](#)]
- 18 **West CP**, Dyrbye LN, Shanafelt TD. Physician burnout: contributors, consequences and solutions. *J Intern Med* 2018; **283**: 516-529 [PMID: [29505159](#) DOI: [10.1111/joim.12752](#)]
- 19 **Patel RS**, Bachu R, Adikey A, Malik M, Shah M. Factors Related to Physician Burnout and Its Consequences: A Review. *Behav Sci (Basel)* 2018; **8** [PMID: [30366419](#) DOI: [10.3390/bs8110098](#)]
- 20 **Pinel EC**, Paulin N. Stigma Consciousness at Work. *Basic Appl Soc Psychol* 2005; **27**: 345-352 [DOI: [10.1207/s15324834basp2704\\_7](#)]
- 21 Wildes VJ. Stigma in Food Service Work: How it Affects Restaurant Servers' Intention to Stay in the Business or Recommend a Job to Another. *Tour Hosp Res* 2005; **5**: 213-233 DOI: [10.1057/palgrave.thr.6040022](#)
- 22 **Jiang F**, Hu L, Rakofsky J, Liu T, Wu S, Zhao P, Hu G, Wan X, Liu H, Liu Y, Tang YL. Sociodemographic Characteristics and Job Satisfaction of Psychiatrists in China: Results From the First Nationwide Survey. *Psychiatr Serv* 2018; **69**: 1245-1251 [PMID: [30301449](#) DOI: [10.1176/appi.ps.201800197](#)]
- 23 **Liang D**, Mays VM, Hwang WC. Integrated mental health services in China: challenges and planning for the future. *Health Policy Plan* 2018; **33**: 107-122 [PMID: [29040516](#) DOI: [10.1093/heapol/czz137](#)]
- 24 **Lambert TW**, Turner G, Fazel S, Goldacre MJ. Reasons why some UK medical graduates who initially choose psychiatry do not pursue it as a long-term career. *Psychol Med* 2006; **36**: 679-684 [PMID: [16426488](#) DOI: [10.1017/S0033291705007038](#)]
- 25 **Maidment R**, Livingston G, Katona C, McParland M, Noble L. Change in attitudes to psychiatry and intention to pursue psychiatry as a career in newly qualified doctors: a follow-up of two cohorts of medical students. *Med Teach* 2004; **26**: 565-569 [PMID: [15763837](#) DOI: [10.1080/01421590410001711562](#)]
- 26 **Ndeti DM**, Khasakhala L, Ongecha-Owuor F, Kuria M, Mutiso V, Syanda J, Kokonya D. Attitudes toward psychiatry: a survey of medical students at the University of Nairobi, Kenya. *Acad Psychiatry* 2008; **32**: 154-159 [PMID: [18349338](#) DOI: [10.1176/appi.ap.32.2.154](#)]
- 27 **Niaz U**, Hassan S, Hussain H, Saeed S. Attitudes towards Psychiatry in Pre-Clinical and Post-Clinical Clerkships in Different Medical Colleges of Karachi. *Pak J Med Sci* 2003; **19**: 253-263
- 28 **Niedermier JA**, Bornstein R, Brandemihl A. The junior medical student psychiatry clerkship: curriculum, attitudes, and test performance. *Acad Psychiatry* 2006; **30**: 136-143 [PMID: [16609120](#) DOI: [10.1176/appi.ap.30.2.136](#)]
- 29 **Holmes D**, Tumiel-Berhalter LM, Zayas LE, Watkins R. "Bashing" of medical specialties: students' experiences and recommendations. *Fam Med* 2008; **40**: 400-406 [PMID: [18773777](#)]
- 30 **Motlová LB**, Janoušková M, Formánek T, Goetz M, Holub D, Hubeňák J, Kašpárek T, Látlavová K, Papežová H, Světlák M, Šilhán P, Trančík P, Vevera J, Balon R. Medical Students' Career Choice and Attitudes Toward Psychiatry: Case of the Czech Republic. *Acad Psychiatry* 2020; **44**: 751-755 [PMID: [33000447](#) DOI: [10.1007/s40596-020-01311-z](#)]
- 31 **Cutler JL**, Harding KJ, Mozian SA, Wright LL, Pica AG, Masters SR, Graham MJ. Discrediting the notion "working with 'crazies' will make you 'crazy'": addressing stigma and enhancing empathy in medical student education. *Adv Health Sci Educ Theory Pract* 2009; **14**: 487-502 [PMID: [18766453](#) DOI: [10.1007/s10459-008-9132-4](#)]
- 32 **Lindsey W**, John S, Erick M. Factors Influencing Medical School Graduates Toward a Career in Psychiatry: Analysis from the 2011–2013 Association of American Medical Colleges Graduation Questionnaire. *Acad Psychiatry* 2015; **40**: 255-260 [PMID: [25697263](#) DOI: [10.1007/s40596-015-0287-z](#)]
- 33 **Balon R**, Franchini GR, Freeman PS, Hassenfeld IN, Keshavan MS, Yoder E. Medical Students' Attitudes and Views of Psychiatry: 15 Years Later. *Acad Psychiatry* 1999; **23**: 30-36 [DOI: [10.1007/BF03340033](#)]

- 34 **Möller-Leimkühler AM**, Möller HJ, Maier W, Gaebel W, Falkai P. EPA guidance on improving the image of psychiatry. *Eur Arch Psychiatry Clin Neurosci* 2016; **266**: 139-154 [PMID: [26874959](#) DOI: [10.1007/s00406-016-0678-5](#)]
- 35 **Sun KS**, Lam TP, Lam KF, Lo TL. Barriers and facilitators for psychiatrists in managing mental health patients in Hong Kong-Impact of Chinese culture and health system. *Asia Pac Psychiatry* 2018; **10** [PMID: [28371455](#) DOI: [10.1111/appy.12279](#)]
- 36 **Passmore K**, Leung WC. Defensive practice among psychiatrists: a questionnaire survey. *Postgrad Med J* 2002; **78**: 671-673 [PMID: [12496324](#) DOI: [10.1136/pmj.78.925.671](#)]
- 37 **Reuveni I**, Pelov I, Reuveni H, Bonne O, Canetti L. Cross-sectional survey on defensive practices and defensive behaviours among Israeli psychiatrists. *BMJ Open* 2017; **7**: e014153 [PMID: [28320795](#) DOI: [10.1136/bmjopen-2016-014153](#)]
- 38 **Studdert DM**, Mello MM, Sage WM, DesRoches CM, Peugh J, Zapert K, Brennan TA. Defensive medicine among high-risk specialist physicians in a volatile malpractice environment. *JAMA* 2005; **293**: 2609-2617 [PMID: [15928282](#) DOI: [10.1001/jama.293.21.2609](#)]
- 39 **Beale C**. Magical thinking and moral injury: exclusion culture in psychiatry. *BJPsych Bull* 2022; **46**: 16-19 [PMID: [34517935](#) DOI: [10.1192/bjb.2021.86](#)]
- 40 **Helingoe KE**, Cullum S, Ng L. Stigma in Consultation Liaison Psychiatry: A Qualitative Study of Trainee Psychiatrists. *Australas Psychiatry* 2022; **30**: 658-662 [PMID: [35139666](#) DOI: [10.1177/10398562211064221](#)]
- 41 **Ashforth BE**, Kreiner GE. Dirty Work and Dirtier Work: Differences in Countering Physical, Social, and Moral Stigma. *Manag Organ Rev* 2014; **10**: 81-108 [DOI: [10.1111/more.12044](#)]
- 42 **Ashforth BE**, Kreiner GE, Clark MA, Fugate M. Normalizing Dirty Work: Managerial Tactics for Countering Occupational Taint. *Acad Manage J* 2007; **50**: 149-174 [DOI: [10.5465/amj.2007.24162092](#)]
- 43 **Ashforth BE**, Kreiner GE. 'How Can You Do It?': Dirty Work and the Challenge of Constructing a Positive Identity. *Acad Manage Rev* 1999; **24**: 413 [DOI: [10.2307/259134](#)]
- 44 **Benjamin O**, Bernstein D, Motzafi-Haller P. Emotional Politics in Cleaning Work: The Case of Israel. *Hum Relat* 2011; **64**: 337-357 [DOI: [10.1177/0018726710378383](#)]
- 45 **Carden P**. Rising from the Dead: Delimiting Stigma in the Australian Funeral Industry. *Health Sociol Rev* 2001; **10**: 79-87 [DOI: [10.5172/hesr.2001.10.2.79](#)]
- 46 **Johnson CC**, Vega L, Kohalmi AL, Roth JC, Howell BR, Van Hasselt VB. Enhancing Mental Health Treatment for the Firefighter Population: Understanding Fire Culture, Treatment Barriers, Practice Implications, and Research Directions. *Prof Psychol Res Pract* 2020; **51**: 304-311 [DOI: [10.1037/pro0000266](#)]
- 47 **Maibvise C**, Shongwe M, Jele V, Dlamini P, Chiviya W. Perceptions about tuberculosis and perceived tuberculosis-related stigma and associated factors among the mining community in Eswatini. *Afr Health Sci* 2022; **22**: 551-559 [PMID: [36032435](#) DOI: [10.4314/ahs.v22i1.64](#)]
- 48 **Tracy SJ**. The Construction of Correctional Officers: Layers of Emotionality behind Bars. *Qual Inq* 2004; **10**: 509-533 [DOI: [10.1177/1077800403259716](#)]
- 49 **Lohiniva AL**, Kamal W, Benkirane M, Numair T, Abdelrahman M, Saleh H, Zahran A, Talaat M, Kandeel A. HIV Stigma Toward People Living With HIV and Health Providers Associated With Their Care: Qualitative Interviews With Community Members in Egypt. *J Assoc Nurses AIDS Care* 2016; **27**: 188-198 [PMID: [26718817](#) DOI: [10.1016/j.jana.2015.11.007](#)]
- 50 **Lim N**, Paul AM. Stigma on a Spectrum: Differentiated Stigmatization of Migrant Domestic Workers' Romantic Relationships in Singapore. *Gend Place Cult* 2021; **28**: 22-44 [DOI: [10.1080/0966369X.2019.1710474](#)]
- 51 **Li Y**, Song Y, Wang M, Huan T-C (T. C). The Influence of Tour Guides' Service Quality on Tourists' Tour Guide Stigma Judgment: An Asian Perspective. *J Hosp Tour Manag* 2021; **48**: 551-560 [DOI: [10.1016/j.jhtm.2021.08.011](#)]
- 52 **Aniteye P**, O'Brien B, Mayhew SH. Stigmatized by association: challenges for abortion service providers in Ghana. *BMC Health Serv Res* 2016; **16**: 486 [PMID: [27612453](#) DOI: [10.1186/s12913-016-1733-7](#)]
- 53 **Stardust Z**, Treloar C, Cama E, Kim J. 'I Wouldn't Call the Cops if I was being Bashed to Death': Sex Work, Whore Stigma and the Criminal Legal System. *Int J Crime Justice Soc Democr* 2021; **10** [DOI: [10.5204/ijcsd.1894](#)]
- 54 **Wang S**. Intimate Situations, and the Circulation Of Same-Sex Affect: Monetizing Affective Encounters on Blued. *Sexualities* 2020; **23**: 934-950 [DOI: [10.1177/1363460719872724](#)]
- 55 **Vines M**, Linders A. The Dirty Work of Poker: Impression Management and Identity. *Deviant Behav* 2016; **37**: 1064-1076 [DOI: [10.1080/01639625.2016.1169740](#)]
- 56 **Stenger S**, Roulet TJ. Pride Against Prejudice? The Stakes of Concealment and Disclosure of a Stigmatized Identity for Gay and Lesbian Auditors. *Work Employ Soc* 2018; **32**: 257-273 [DOI: [10.1177/0950017016682459](#)]
- 57 **Zhang R**, Wang MS, Toubiana M, Greenwood R. Stigma Beyond Levels: Advancing Research on Stigmatization. *Acad Manag Ann* 2021; **15**: 188-222 [DOI: [10.5465/annals.2019.0031](#)]
- 58 **Katschnig H**. Psychiatry's contribution to the public stereotype of schizophrenia: Historical considerations. *J Eval Clin Pract* 2018; **24**: 1093-1100 [PMID: [30112785](#) DOI: [10.1111/jep.13011](#)]
- 59 **Pescosolido BA**, Manago B, Monahan J. Evolving Public Views On The Likelihood Of Violence From People With Mental Illness: Stigma And Its Consequences. *Health Aff (Millwood)* 2019; **38**: 1735-1743 [PMID: [31589533](#) DOI: [10.1377/hlthaff.2019.00702](#)]
- 60 **Terkelsen TB**, Larsen IB. Fear, danger and aggression in a Norwegian locked psychiatric ward: Dialogue and ethics of care as contributions to combating difficult situations. *Nurs Ethics* 2016; **23**: 308-317 [PMID: [25552587](#) DOI: [10.1177/0969733014564104](#)]
- 61 **Kelly EL**, Fenwick K, Brekke JS, Novaco RW. Well-Being and Safety Among Inpatient Psychiatric Staff: The Impact of Conflict, Assault, and Stress Reactivity. *Adm Policy Ment Health* 2016; **43**: 703-716 [PMID: [26377816](#) DOI: [10.1007/s10488-015-0683-4](#)]
- 62 **Chen Y**, Wang P, Zhao L, He Y, Chen N, Liu H, Liu Y, Liu T, Tang YL, Jiang F, Zhu J. Workplace Violence and Turnover Intention Among Psychiatrists in a National Sample in China: The Mediating Effects of Mental Health. *Front Psychiatry* 2022; **13**: 855584 [PMID: [35782425](#) DOI: [10.3389/fpsy.2022.855584](#)]

- 63 **Alhassan RK, Poku KA.** Experiences of Frontline Nursing Staff on Workplace Safety and Occupational Health Hazards in Two Psychiatric Hospitals in Ghana *BMC Public Health* 2018; **18**: 701 [PMID: [29875015](#) DOI: [10.1186/s12889-018-5620-5](#)]
- 64 **Eisele F, Flammer E, Steinert T.** Incidents of aggression in German psychiatric hospitals: Is there an increase? *PLoS One* 2021; **16**: e0245090 [PMID: [33400702](#) DOI: [10.1371/journal.pone.0245090](#)]
- 65 **Kılınçel O, Erzin G, Kılınçel Ş, Hun Şenol S, Ceylan D, Acar M, Gürcan A, Çapraz N.** Profile of Young Psychiatrists: A cross-sectional survey study from Turkey (eng). *J Clin Psychiatry* 2021; **24**: 278-287 [DOI: [10.5505/kpd.2021.15428](#)]
- 66 **Farid Ali A, Khalid AS, Muhammad Ajmal Z.** Violence Against Medical Staff: Prevalence and Effects of Violence Against Psychiatrists in Kuwait. *KMJ-Kuwait Med J* 2009; **41**: 322-326
- 67 **Lin X, Rosenheck R, Sun B, Xie G, Zhong G, Tan C, Li Z, Yu M, He H.** Associative stigma experienced by mental health professionals in China and the United States. *Soc Psychiatry Psychiatr Epidemiol* 2019; **54**: 745-753 [PMID: [30542960](#) DOI: [10.1007/s00127-018-1643-6](#)]
- 68 **Shibre T, Negash A, Kullgren G, Kebede D, Alem A, Fekadu A, Fekadu D, Madhin G, Jacobsson L.** Perception of stigma among family members of individuals with schizophrenia and major affective disorders in rural Ethiopia. *Soc Psychiatry Psychiatr Epidemiol* 2001; **36**: 299-303 [PMID: [11583460](#) DOI: [10.1007/s001270170048](#)]
- 69 **Ebsworth SJ, Foster JLH.** Public perceptions of mental health professionals: stigma by association? *J Ment Health* 2017; **26**: 431-441 [PMID: [27596719](#) DOI: [10.1080/09638237.2016.1207228](#)]
- 70 **Rüsch N, Angermeyer MC, Corrigan PW.** Mental illness stigma: concepts, consequences, and initiatives to reduce stigma. *Eur Psychiatry* 2005; **20**: 529-539 [PMID: [16171984](#) DOI: [10.1016/j.eurpsy.2005.04.004](#)]
- 71 **Corrigan PW, Miller FE.** Shame, Blame, and Contamination: A Review of the Impact of Mental Illness Stigma on Family Members. *J Ment Health* 2004; **13**: 537-548 [DOI: [10.1080/09638230400017004](#)]
- 72 **Rössler W.** The stigma of mental disorders: A millennia-long history of social exclusion and prejudices. *EMBO Rep* 2016; **17**: 1250-1253 [PMID: [27470237](#) DOI: [10.15252/embr.201643041](#)]
- 73 **Zolezzi M, Alamri M, Shaar S, Rainkie D.** Stigma associated with mental illness and its treatment in the Arab culture: A systematic review. *Int J Soc Psychiatry* 2018; **64**: 597-609 [PMID: [30019976](#) DOI: [10.1177/0020764018789200](#)]
- 74 **Lloyd CEM.** Contending with Spiritual Reductionism: Demons, Shame, and Dividualising Experiences Among Evangelical Christians with Mental Distress. *J Relig Health* 2021; **60**: 2702-2727 [PMID: [33991287](#) DOI: [10.1007/s10943-021-01268-9](#)]
- 75 **Ciftci A, Jones N, Corrigan PW.** Mental Health Stigma in the Muslim Community. *J Muslim Ment Health* 2013; **7** [DOI: [10.3998/jmmh.10381607.0007.102](#)]
- 76 **Hwang K-K, Han K-H.** Face and Morality in Confucian society. In: Bond MH, editor. *Oxford Handbook of Chinese Psychology*. 1st ed. Oxford University Press, 2012: 479-498 [DOI: [10.1093/oxfordhb/9780199541850.013.0029](#)]
- 77 **Yang LH, Kleinman A.** 'Face' and the embodiment of stigma in China: the cases of schizophrenia and AIDS. *Soc Sci Med* 2008; **67**: 398-408 [PMID: [18420325](#) DOI: [10.1016/j.socscimed.2008.03.011](#)]
- 78 **Thornicroft G, Brohan E, Rose D, Sartorius N, Leese M; INDIGO Study Group.** Global pattern of experienced and anticipated discrimination against people with schizophrenia: a cross-sectional survey. *Lancet* 2009; **373**: 408-415 [PMID: [19162314](#) DOI: [10.1016/S0140-6736\(08\)61817-6](#)]
- 79 **Kudva KG, El Hayek S, Gupta AK, Kurokawa S, Bangshan L, Armas-Villavicencio MVC, Oishi K, Mishra S, Tiensuntisook S, Sartorius N.** Stigma in mental illness: Perspective from eight Asian nations. *Asia Pac Psychiatry* 2020; **12**: e12380 [PMID: [31922363](#) DOI: [10.1111/appy.12380](#)]
- 80 **Völm B, Nedopil N, editor.** The Use of Coercive Measures in Forensic Psychiatric Care: Legal, Ethical and Practical Challenges. Switzerland: Springer Cham, 2016 [DOI: [10.1007/978-3-319-26748-7](#)]
- 81 **McLaughlin P, Giacco D, Priebe S.** Use of Coercive Measures during Involuntary Psychiatric Admission and Treatment Outcomes: Data from a Prospective Study across 10 European Countries. *PLoS One* 2016; **11**: e0168720 [PMID: [28033391](#) DOI: [10.1371/journal.pone.0168720](#)]
- 82 **Teichert M, Schäfer I, Lincoln TM.** [Where there's a Will, there's a Way? A nationwide Online-Survey of Psychiatrists about the Use of Alternatives to Coercive Measures]. *Psychiatr Prax* 2016; **43**: 101-106 [PMID: [25526502](#) DOI: [10.1055/s-0034-1387431](#)]
- 83 **Gowda GS, Lepping P, Ray S, Noorthoorn E, Nanjgowda RB, Kumar CN, Math SB.** Clinician attitude and perspective on the use of coercive measures in clinical practice from tertiary care mental health establishment - A cross-sectional study. *Indian J Psychiatry* 2019; **61**: 151-155 [PMID: [30992609](#) DOI: [10.4103/psychiatry.IndianJPsychiatry\\_336\\_18](#)]
- 84 **Voskes Y, Evenblij K, Noorthoorn E, Porz R, Widdershoven G.** [Moral case deliberation about coercion in psychiatry. Dilemmas, value and implementation]. *Psychiatr Prax* 2014; **41**: 364-370 [PMID: [25295977](#) DOI: [10.1055/s-0034-1370292](#)]
- 85 **Hui A, Middleton H, Völm B.** Coercive Measures in Forensic Settings: Findings from the Literature. *Int J Forensic Ment Health* 2013; **12**: 53-67 [DOI: [10.1080/14999013.2012.740649](#)]
- 86 **Ma ML, He LP.** Electroconvulsive therapy plays an irreplaceable role in treatment of major depressive disorder. *World J Clin Cases* 2022; **10**: 5515-5517 [PMID: [35812666](#) DOI: [10.12998/wjcc.v10.i16.5515](#)]
- 87 **Gong J, Cui LB, Zhao YS, Liu ZW, Yang XJ, Xi YB, Liu L, Liu P, Sun JB, Zhao SW, Liu XF, Jia J, Li P, Yin H, Qin W.** The correlation between dynamic functional architecture and response to electroconvulsive therapy combined with antipsychotics in schizophrenia. *Eur J Neurosci* 2022; **55**: 2024-2036 [PMID: [35388553](#) DOI: [10.1111/ejn.15664](#)]
- 88 **Bahji A.** The Rise, Fall, and Resurgence of Electroconvulsive Therapy. *J Psychiatr Pract* 2022; **28**: 440-444 [PMID: [36355582](#) DOI: [10.1097/PRA.0000000000000666](#)]
- 89 **Richmond J, Boynton MH, Ozawa S, Muessig KE, Cykert S, Ribisl KM.** Development and Validation of the Trust in My Doctor, Trust in Doctors in General, and Trust in the Health Care Team Scales. *Soc Sci Med* 2022; **298**: 114827 [PMID: [35255277](#) DOI: [10.1016/j.socscimed.2022.114827](#)]
- 90 **Fan Z, Chen H, Wu H, Zhang X.** Patient Toward Physician Occupational Stigma Scale: Development of the Chinese Version. *Psychol Res Behav Manag* 2022; **15**: 2117-2127 [PMID: [35983020](#) DOI: [10.2147/PRBM.S375032](#)]
- 91 **Dempsey B, Favier M, Mullally A, Higgins MF.** Exploring providers' experience of stigma following the introduction of



- more liberal abortion care in the Republic of Ireland. *Contraception* 2021; **104**: 414-419 [PMID: 33864811 DOI: 10.1016/j.contraception.2021.04.007]
- 92 **Uvais NA**, Shihabudheen P, Hafi NAB. Perceived Stress and Stigma Among Doctors Working in COVID-19-Designated Hospitals in India. *Prim Care Companion CNS Disord* 2020; **22** [PMID: 32731315 DOI: 10.4088/PCC.20br02724]
  - 93 **Low LF**, Purwaningrum F. Negative stereotypes, fear and social distance: a systematic review of depictions of dementia in popular culture in the context of stigma. *BMC Geriatr* 2020; **20**: 477 [PMID: 33203379 DOI: 10.1186/s12877-020-01754-x]
  - 94 **Shiu C**, Chen WT, Hung CC, Huang EP, Lee TS. COVID-19 stigma associates with burnout among healthcare providers: Evidence from Taiwanese physicians and nurses. *J Formos Med Assoc* 2022; **121**: 1384-1391 [PMID: 34654583 DOI: 10.1016/j.jfma.2021.09.022]
  - 95 **Lucas J**, Phelan J. Influence and Social Distance Consequences Across Categories of Race and Mental Illness. *Soc Ment Health* 2018; **9**: 215686931876112 [DOI: 10.1177/2156869318761125]
  - 96 **Crowley F**, Doran J. COVID-19, Occupational Social Distancing and Remote Working Potential: An Occupation, Sector and Regional Perspective. *Reg Sci Policy Pract* 2020; **12** [DOI: 10.1111/rsp3.12347]
  - 97 **Ta TMT**, Böge K, Cao TD, Schomerus G, Nguyen TD, Dettling M, Mungee A, Martensen LK, Diefenbacher A, Angermeyer MC, Hahn E. Public attitudes towards psychiatrists in the metropolitan area of Hanoi, Vietnam. *Asian J Psychiatr* 2018; **32**: 44-49 [PMID: 29207317 DOI: 10.1016/j.ajp.2017.11.031]
  - 98 **Shantz A**, Booth JE. Service Employees and Self-Verification: The Roles of Occupational Stigma Consciousness and Core Self-Evaluations. *Hum Relat* 2014; **67**: 1439-1465 [DOI: 10.1177/0018726713519280]
  - 99 **Zhao Y**, Zhan J. Platform Riders' Occupational Stigma Consciousness and Workplace Deviant Behavior: the Mediating Role of Self-depletion. *Balt J Manag* 2022; **17**: 233-249 [DOI: 10.1108/BJM-06-2021-0215]
  - 100 **Mostafa A**, Mostafa NS, Ismail N. Validity and Reliability of a COVID-19 Stigma Scale Using Exploratory and Confirmatory Factor Analysis in a Sample of Egyptian Physicians: E16-COVID19-S. *Int J Environ Res Public Health* 2021; **18** [PMID: 34069754 DOI: 10.3390/ijerph18105451]
  - 101 **Oktar D**, Aydoğan S, Sungur S, Önsüz M, Metintas S, Kosger F, Altinoz E. Reliability and Validity of the Perception of Stigma due to COVID-19 in Physicians. *Eur J Public Health* 2021; **31**: 275 [DOI: 10.1093/eurpub/ckab164.718]
  - 102 **Riley R**, Buszewicz M, Kokab F, Teoh K, Gopfert A, Taylor AK, Van Hove M, Martin J, Appleby L, Chew-Graham C. Sources of work-related psychological distress experienced by UK-wide foundation and junior doctors: a qualitative study. *BMJ Open* 2021; **11**: e043521 [PMID: 34162634 DOI: 10.1136/bmjopen-2020-043521]
  - 103 **Wijeratne C**, Johnco C, Draper B, Earl J. Doctors' reporting of mental health stigma and barriers to help-seeking. *Occup Med (Lond)* 2021; **71**: 366-374 [PMID: 34534344 DOI: 10.1093/occmed/kqab119]
  - 104 **Zarzycki MZ**, Goetz Z, Flaga-Luczkiwicz M. What do medical students think about medical doctors with mental health issues? *Psychol Health Med* 2020; **25**: 623-629 [PMID: 32019331 DOI: 10.1080/13548506.2020.1724309]
  - 105 **Clough BA**, Ireland MJ, March S. Development of the SOSS-D: a scale to measure stigma of occupational stress and burnout in medical doctors. *J Ment Health* 2019; **28**: 26-33 [PMID: 28868957 DOI: 10.1080/09638237.2017.1370642]
  - 106 **Clough BA**, Hill M, Delaney M, Casey LM. Development of a measure of stigma towards occupational stress for mental health professionals. *Soc Psychiatry Psychiatr Epidemiol* 2020; **55**: 941-951 [PMID: 31897577 DOI: 10.1007/s00127-019-01820-9]
  - 107 **Fan Z**, Cong X, Tao M, Wu S, Gao P. Development of the Chinese Version of the Physician Internalized Occupational Stigma Scale (PIOSS). *Psychol Res Behav Manag* 2022; **15**: 3445-3459 [PMID: 36471790 DOI: 10.2147/PRBM.S386724]
  - 108 **Healey LV**. The psychometric properties of the Forensic Stigma Scale (FSS). *Int J Law Psychiatry* 2022; **83**: 101804 [PMID: 35772282 DOI: 10.1016/j.ijlp.2022.101804]
  - 109 **Verhaeghe M**, Bracke P. Associative stigma among mental health professionals: implications for professional and service user well-being. *J Health Soc Behav* 2012; **53**: 17-32 [PMID: 22382718 DOI: 10.1177/0022146512439453]
  - 110 **Gaebel W**, Zäske H, Cleveland HR, Zielasek J, Stuart H, Arboleda-Florez J, Akiyama T, Gureje O, Jorge MR, Kastrup M, Suzuki Y, Tasman A, Sartorius N. Measuring the stigma of psychiatry and psychiatrists: development of a questionnaire. *Eur Arch Psychiatry Clin Neurosci* 2011; **261** Suppl 2: S119-S123 [PMID: 21947511 DOI: 10.1007/s00406-011-0252-0]
  - 111 **Mittal D**, Sullivan G, Chekuri L, Allee E, Corrigan PW. Empirical Studies of Self-Stigma Reduction Strategies: A Critical Review of the Literature. *Psychiatr Serv* 2012; **63**: 974-981 [PMID: 22855130 DOI: 10.1176/appi.ps.201100459]
  - 112 **Jeffrey JB**, Christine B, Trisha BJ, Melissa Z. Addressing mental health stigma among young adolescents: evaluation of a youth-led approach. *Health Soc Work* 2014; **39**: 73-80 [PMID: 24946423 DOI: 10.1093/hsw/hlu008]
  - 113 **Hartog K**, Hubbard CD, Krouwer AF, Thornicroft G, Kohrt BA, Jordans MJD. Stigma reduction interventions for children and adolescents in low- and middle-income countries: Systematic review of intervention strategies. *Soc Sci Med* 2020; **246**: 112749 [PMID: 31978636 DOI: 10.1016/j.socscimed.2019.112749]
  - 114 **Poole AH**. "Tearing the Shroud of Invisibility": Communities of Protest Information Practices and the Fight for LGBTQ Rights in US Librarianship. *Libr Q* 2020; **90**: 530-562 [DOI: 10.1086/710255]
  - 115 **Corrigan PW**, Penn DL. Lessons from social psychology on discrediting psychiatric stigma. *Am Psychol* 1999; **54**: 765-776 [PMID: 10510666 DOI: 10.1037/0003-066X.54.9.765]
  - 116 **Makita M**, Mas-Bleda A, Morris S, Thelwall M. Mental Health Discourses on Twitter during Mental Health Awareness Week. *Issues Ment Health Nurs* 2021; **42**: 437-450 [PMID: 32926796 DOI: 10.1080/01612840.2020.1814914]
  - 117 **Koteyko N**, Atanasova D. Mental Health Advocacy on Twitter: Positioning in Depression Awareness Week tweets. *Discourse Context Media* 2018; **25**: 52-59 [DOI: 10.1016/j.dcm.2018.04.007]
  - 118 **Laura JA**, Sarah EG, Racheal AR. Key Ingredients—Target Groups, Methods and Messages, and Evaluation—of Local-Level, Public Interventions to Counter Stigma and Discrimination: A Lived Experience Informed Selective Narrative Literature Review. *Community Ment Health J* 2017; **54**: 312-333 [PMID: 29185150 DOI: 10.1007/s10597-017-0189-5]
  - 119 **Esmacili AA**, Saeidi A, Mohammadi Y, Raeesoon M. The effect of group psychological training on the attitude of family caregivers of patients with schizophrenia. *Heliyon* 2022; **8**: e09817 [PMID: 35874056 DOI: 10.1016/j.heliyon.2022.e09817]

- 120 Li J, Fan Y, Zhong HQ, Duan XL, Chen W, Evans-Lacko S, Thornicroft G. Effectiveness of an anti-stigma training on improving attitudes and decreasing discrimination towards people with mental disorders among care assistant workers in Guangzhou, China. *Int J Ment Health Syst* 2019; **13**: 1 [PMID: 30622627 DOI: 10.1186/s13033-018-0259-2]
- 121 Li J, Li J, Huang Y, Thornicroft G. Mental health training program for community mental health staff in Guangzhou, China: effects on knowledge of mental illness and stigma. *Int J Ment Health Syst* 2014; **8**: 49 [PMID: 25520750 DOI: 10.1186/1752-4458-8-49]
- 122 Tomar N, Ghezzi MA, Brinkley-Rubinstein L, Wilson AB, Van Deinse TB, Burgin S, Cuddeback GS. Statewide mental health training for probation officers: improving knowledge and decreasing stigma. *Health Justice* 2017; **5**: 11 [PMID: 29143153 DOI: 10.1186/s40352-017-0057-y]
- 123 Yanos PT, Roe D, West ML, Smith SM, Lysaker PH. Group-based treatment for internalized stigma among persons with severe mental illness: findings from a randomized controlled trial. *Psychol Serv* 2012; **9**: 248-258 [PMID: 22545821 DOI: 10.1037/a0028048]
- 124 Ke S, Lai J, Sun T, Yang MM, Wang JC, Austin J. Healthy young minds: the effects of a 1-hour classroom workshop on mental illness stigma in high school students. *Community Ment Health J* 2015; **51**: 329-337 [PMID: 25017811 DOI: 10.1007/s10597-014-9763-2]
- 125 Ojio Y, Yamaguchi S, Ohta K, Ando S, Koike S. Effects of biomedical messages and expert-recommended messages on reducing mental health-related stigma: a randomised controlled trial. *Epidemiol Psychiatr Sci* 2019; **29**: e74 [PMID: 31753045 DOI: 10.1017/S2045796019000714]
- 126 Friedrich B, Evans-Lacko S, London J, Rhydderch D, Henderson C, Thornicroft G. Anti-stigma training for medical students: the Education Not Discrimination project. *Br J Psychiatry Suppl* 2013; **55**: s89-s94 [PMID: 23553700 DOI: 10.1192/bjp.bp.112.114017]
- 127 McCormack Z, Gilbert JL, Ott C, Plake KS. Mental health first aid training among pharmacy and other university students and its impact on stigma toward mental illness. *Curr Pharm Teach Learn* 2018; **10**: 1342-1347 [PMID: 30527363 DOI: 10.1016/j.cptl.2018.07.001]
- 128 Sreeram A, Cross WM, Townsin L. Anti-stigma initiatives for mental health professionals-A systematic literature review. *J Psychiatr Ment Health Nurs* 2022; **29**: 512-528 [PMID: 35500153 DOI: 10.1111/jpm.12840]
- 129 Martínez-Martínez C, Sánchez-Martínez V, Sales-Orts R, Dinca A, Richart-Martínez M, Ramos-Pichardo JD. Effectiveness of direct contact intervention with people with mental illness to reduce stigma in nursing students. *Int J Ment Health Nurs* 2019; **28**: 735-743 [PMID: 30693628 DOI: 10.1111/inm.12578]
- 130 Corrigan PW, Morris SB, Michaels PJ, Rafacz JD, Rüsch N. Challenging the public stigma of mental illness: a meta-analysis of outcome studies. *Psychiatr Serv* 2012; **63**: 963-973 [PMID: 23032675 DOI: 10.1176/appi.ps.201100529]
- 131 Thornicroft G, Mehta N, Clement S, Evans-Lacko S, Doherty M, Rose D, Koschorke M, Shidhaye R, O'Reilly C, Henderson C. Evidence for effective interventions to reduce mental-health-related stigma and discrimination. *Lancet* 2016; **387**: 1123-1132 [PMID: 26410341 DOI: 10.1016/S0140-6736(15)00298-6]
- 132 Yamaguchi S, Wu SI, Biswas M, Yate M, Aoki Y, Barley EA, Thornicroft G. Effects of short-term interventions to reduce mental health-related stigma in university or college students: a systematic review. *J Nerv Ment Dis* 2013; **201**: 490-503 [PMID: 23719324 DOI: 10.1097/NMD.0b013e31829480df]
- 133 Corrigan PW, Vega E, Larson J, Michaels PJ, McClintock G, Krzyzanowski R, Gause M, Buchholz B. The California schedule of key ingredients for contact-based antistigma programs. *Psychiatr Rehabil J* 2013; **36**: 173-179 [PMID: 23834612 DOI: 10.1037/prj0000006]
- 134 Corrigan PW, Michaels PJ, Vega E, Gause M, Larson J, Krzyzanowski R, Botcheva L. Key ingredients to contact-based stigma change: a cross-validation. *Psychiatr Rehabil J* 2014; **37**: 62-64 [PMID: 24417232 DOI: 10.1037/prj0000038]
- 135 Janoušková M, Tušková E, Weissová A, Trancík P, Pasz J, Evans-Lacko S, Winkler P. Can video interventions be used to effectively destigmatize mental illness among young people? A systematic review. *Eur Psychiatry* 2017; **41**: 1-9 [PMID: 28049074 DOI: 10.1016/j.eurpsy.2016.09.008]
- 136 Adu J, Oudshoorn A, Anderson K, Marshall CA, Stuart H. Social Contact: Next Steps in an Effective Strategy to Mitigate the Stigma of Mental Illness. *Issues Ment Health Nurs* 2022; **43**: 485-488 [PMID: 34652979 DOI: 10.1080/01612840.2021.1986757]
- 137 Nakanishi E, Tamachi M, Hashimoto T. Effectiveness of a Co-Production with Dialogue Program for Reducing Stigma against Mental Illness: A Quasi-Experimental Study with a Pre- and Post-Test Design. *Int J Environ Res Public Health* 2022; **19** [PMID: 36361212 DOI: 10.3390/ijerph192114333]
- 138 Hogg B, Moreno-Alcázar A, Tóth MD, Serbanescu I, Aust B, Leduc C, Paterson C, Tsantilla F, Abdulla K, Cerga-Pashoja A, Cresswell-Smith J, Fanaj N, Meksi A, Ni Dhálaigh D, Reich H, Ross V, Sanches S, Thomson K, Van Audenhove C, Pérez V, Arensman E, Purebl G, Amann BL; MENTUPP consortium members. Supporting employees with mental illness and reducing mental illness-related stigma in the workplace: an expert survey. *Eur Arch Psychiatry Clin Neurosci* 2023; **273**: 739-753 [PMID: 35867155 DOI: 10.1007/s00406-022-01443-3]
- 139 Qin S, Sheehan L, Yau E, Chen Y, Wang Y, Deng H, Lam C, Chen Z, Zhao L, Gan S, Mao H, Mo X, Yang L, Zhang F, Corrigan P, CBPR Team. Adapting and Evaluating a Strategic Disclosure Program to Address Mental Health Stigma Among Chinese. *Int J Ment Health Addict* 2022 [DOI: 10.1007/s11469-022-00911-0]
- 140 Conceição V, Rothes I, Gusmão R. The effects of a video-based randomized controlled trial intervention on depression stigma and help-seeking attitudes in university students. *Psychiatry Res* 2022; **308**: 114356 [PMID: 34972028 DOI: 10.1016/j.psychres.2021.114356]
- 141 Winkler P, Janoušková M, Kožený J, Pasz J, Mladá K, Weissová A, Tušková E, Evans-Lacko S. Short video interventions to reduce mental health stigma: a multi-centre randomised controlled trial in nursing high schools. *Soc Psychiatry Psychiatr Epidemiol* 2017; **52**: 1549-1557 [PMID: 29101447 DOI: 10.1007/s00127-017-1449-y]
- 142 Clement S, van Nieuwenhuizen A, Kassam A, Flach C, Lazarus A, de Castro M, McCrone P, Norman I, Thornicroft G. Filmed v. live social contact interventions to reduce stigma: randomised controlled trial. *Br J Psychiatry* 2012; **201**: 57-64 [PMID: 22157800 DOI: 10.1192/bjp.bp.111.093120]
- 143 Tan GTH, Shahwan S, Abdin E, Lau JH, Goh CMJ, Ong WJ, Samari E, Kwok KW, Chong SA, Subramaniam M.

- Recognition of Depression and Help-Seeking Preference Among University Students in Singapore: An Evaluation of the Impact of Advancing Research to Eliminate Mental Illness Stigma an Education and Contact Intervention. *Front Psychiatry* 2021; **12**: 582730 [PMID: 34054591 DOI: 10.3389/fpsy.2021.582730]
- 144 **Ahuja K**, Dhillon M, Juneja A, Sharma B. Breaking Barriers: An Education and Contact Intervention to Reduce Mental Illness Stigma among Indian College Students. *Psychosoc Interv* 2017; **26** [DOI: 10.1016/j.psi.2016.11.003]
  - 145 **Hawke LD**, Michalak EE, Maxwell V, Parikh SV. Reducing stigma toward people with bipolar disorder: impact of a filmed theatrical intervention based on a personal narrative. *Int J Soc Psychiatry* 2014; **60**: 741-750 [PMID: 24351967 DOI: 10.1177/0020764013513443]
  - 146 **Ma HI**, Hsieh CE. An Anti-Stigma Course for Occupational Therapy Students in Taiwan: Development and Pilot Testing. *Int J Environ Res Public Health* 2020; **17** [PMID: 32756448 DOI: 10.3390/ijerph17155599]
  - 147 **Papish A**, Kassam A, Modgill G, Vaz G, Zanussi L, Patten S. Reducing the stigma of mental illness in undergraduate medical education: a randomized controlled trial. *BMC Med Educ* 2013; **13**: 141 [PMID: 24156397 DOI: 10.1186/1472-6920-13-141]
  - 148 **Fernandez A**, Tan KA, Knaak S, Chew BH, Ghazali SS. Effects of Brief Psychoeducational Program on Stigma in Malaysian Pre-clinical Medical Students: A Randomized Controlled Trial. *Acad Psychiatry* 2016; **40**: 905-911 [PMID: 27527730 DOI: 10.1007/s40596-016-0592-1]
  - 149 **Flanagan EH**, Buck T, Gamble A, Hunter C, Sewell I, Davidson L. "Recovery Speaks": A Photovoice Intervention to Reduce Stigma Among Primary Care Providers. *Psychiatr Serv* 2016; **67**: 566-569 [PMID: 26766754 DOI: 10.1176/appi.ps.201500049]
  - 150 **Patten SB**, Remillard A, Phillips L, Modgill G, Szeto ACh, Kassam A, Gardner DM. Effectiveness of contact-based education for reducing mental illness-related stigma in pharmacy students. *BMC Med Educ* 2012; **12**: 120 [PMID: 23216787 DOI: 10.1186/1472-6920-12-120]
  - 151 **Villani M**, Kovess-Masfety V. Could a short training intervention modify opinions about mental illness? A case study on French health professionals. *BMC Psychiatry* 2017; **17**: 133 [PMID: 28390399 DOI: 10.1186/s12888-017-1296-0]
  - 152 **Rubio-Valera M**, Aznar-Lou I, Vives-Collet M, Fernández A, Gil-Girbau M, Serrano-Blanco A. Reducing the Mental Health-Related Stigma of Social Work Students: A Cluster RCT. *Res Soc Work Pract* 2018; **28**: 164-172 [DOI: 10.1177/1049731516641492]
  - 153 **Amsalem D**, Gothelf D, Dorman A, Goren Y, Tene O, Shelef A, Horowitz I, Dunsky LL, Rogev E, Klein EH, Mekori-Domachevsky E, Fischel T, Levkovitz Y, Martin A, Gross R. Reducing Stigma Toward Psychiatry Among Medical Students: A Multicenter Controlled Trial. *Prim Care Companion CNS Disord* 2020; **22** [PMID: 32135042 DOI: 10.4088/PCC.19m02527]
  - 154 **Li J**, Li J, Thornicroft G, Yang H, Chen W, Huang Y. Training community mental health staff in Guangzhou, China: evaluation of the effect of a new training model. *BMC Psychiatry* 2015; **15**: 263 [PMID: 26503370 DOI: 10.1186/s12888-015-0660-1]
  - 155 **Ran MS**, Wang YZ, Lu PY, Weng X, Zhang TM, Deng SY, Li M, Luo W, Wong IY, Yang LH, Thornicroft G, Lu L. Effectiveness of enhancing contact model on reducing stigma of mental illness among family caregivers of persons with schizophrenia in rural China: A cluster randomized controlled trial. *Lancet Reg Health West Pac* 2022; **22**: 100419 [PMID: 35257121 DOI: 10.1016/j.lanwpc.2022.100419]
  - 156 **Zhang W**, Henderson C, Magnusdottir E, Chen W, Ma N, Ma H, Thornicroft G. Effect of a contact-based education intervention on reducing stigma among community health and care staff in Beijing, China: Pilot randomized controlled study. *Asian J Psychiatr* 2022; **73**: 103096 [PMID: 35430494 DOI: 10.1016/j.ajp.2022.103096]
  - 157 **Magliano L**, Rinaldi A, Costanzo R, De Leo R, Schioppa G, Petrillo M, Read J. Improving psychology students' attitudes toward people with schizophrenia: A quasi-randomized controlled study. *Am J Orthopsychiatry* 2016; **86**: 253-264 [PMID: 26963178 DOI: 10.1037/ort0000161]
  - 158 **Rodríguez-Rivas ME**, Cangas AJ, Fuentes-Olavarria D. Controlled Study of the Impact of a Virtual Program to Reduce Stigma Among University Students Toward People With Mental Disorders. *Front Psychiatry* 2021; **12**: 632252 [PMID: 33633613 DOI: 10.3389/fpsy.2021.632252]
  - 159 **Koike S**, Yamaguchi S, Ojio Y, Ohta K, Shimada T, Watanabe K, Thornicroft G, Ando S. A randomised controlled trial of repeated filmed social contact on reducing mental illness-related stigma in young adults. *Epidemiol Psychiatr Sci* 2018; **27**: 199-208 [PMID: 27989255 DOI: 10.1017/S2045796016001050]
  - 160 **Beaulieu T**, Patten S, Knaak S, Weinerman R, Campbell H, Lauria-Horner B. Impact of Skill-Based Approaches in Reducing Stigma in Primary Care Physicians: Results from a Double-Blind, Parallel-Cluster, Randomized Controlled Trial. *Can J Psychiatry* 2017; **62**: 327-335 [PMID: 28095259 DOI: 10.1177/0706743716686919]
  - 161 **Ng YP**, Rashid A, O'Brien F. Determining the effectiveness of a video-based contact intervention in improving attitudes of Penang primary care nurses towards people with mental illness. *PLoS One* 2017; **12**: e0187861 [PMID: 29131841 DOI: 10.1371/journal.pone.0187861]
  - 162 **Hatzenbuehler ML**, Phelan JC, Link BG. Stigma as a fundamental cause of population health inequalities. *Am J Public Health* 2013; **103**: 813-821 [PMID: 23488505 DOI: 10.2105/AJPH.2012.301069]
  - 163 **Hankir A**, Abi Rached M, Zaman R. Pioneering an Innovative Intervention to Reduce Mental Health Related Stigma in Muslim Communities: A Protocol. *Psychiatr Danub* 2021; **33**: 69-73 [PMID: 34862896]
  - 164 **Chen SP**, Koller M, Krupa T, Stuart H. Contact in the Classroom: Developing a Program Model for Youth Mental Health Contact-Based Anti-stigma Education. *Community Ment Health J* 2016; **52**: 281-293 [PMID: 26429792 DOI: 10.1007/s10597-015-9944-7]
  - 165 **Chisholm K**, Patterson P, Torgerson C, Turner E, Jenkinson D, Birchwood M. Impact of contact on adolescents' mental health literacy and stigma: the SchoolSpace cluster randomised controlled trial. *BMJ Open* 2016; **6**: e009435 [PMID: 26895983 DOI: 10.1136/bmjopen-2015-009435]
  - 166 **Duman ZÇ**, Güntüşen NP, İnan FŞ, Ince SÇ, Sari A. Effects of two different psychiatric nursing courses on nursing students' attitudes towards mental illness, perceptions of psychiatric nursing, and career choices. *J Prof Nurs* 2017; **33**: 452-459 [PMID: 29157575 DOI: 10.1016/j.profnurs.2017.06.005]



- 167 **Valentim O**, Moutinho L, Laranjeira C, Querido A, Tomás C, Longo J, Carvalho D, Gomes J, Morgado T, Correia T. "Looking beyond Mental Health Stigma": An Online Focus Group Study among Senior Undergraduate Nursing Students. *Int J Environ Res Public Health* 2023; **20** [PMID: 36901609 DOI: 10.3390/ijerph20054601]
- 168 **Monnapula-Mazabane P**, Petersen I. Feasibility and acceptability of a mental health stigma intervention for low-income South African caregivers: A qualitative investigation. *S Afr J Psychiatr* 2022; **28**: 1824 [PMID: 35402016 DOI: 10.4102/sajpsychiatry.v28i0.1824]
- 169 **Heijnders M**, Van Der Meij S. The fight against stigma: an overview of stigma-reduction strategies and interventions. *Psychol Health Med* 2006; **11**: 353-363 [PMID: 17130071 DOI: 10.1080/13548500600595327]
- 170 **Cook JE**, Purdie-Vaughns V, Meyer IH, Busch JTA. Intervening within and across levels: a multilevel approach to stigma and public health. *Soc Sci Med* 2014; **103**: 101-109 [PMID: 24513229 DOI: 10.1016/j.soescimed.2013.09.023]
- 171 **Hong S**, Satyshur MD, Burnett-Zeigler I. The association of mindfulness and depression stigma among African American women participants in a mindfulness-based intervention: A pilot study. *Transcult Psychiatry* 2022; 13634615221076709 [PMID: 35505619 DOI: 10.1177/13634615221076709]
- 172 **Clinton AJ**, Pollini RA. Using Positive Empathy Interventions to Reduce Stigma Toward People Who Inject Drugs. *Front Psychol* 2021; **12**: 616729 [PMID: 34305698 DOI: 10.3389/fpsyg.2021.616729]
- 173 **Young DK-W**. Cognitive Behavioral Therapy Group for Reducing Self-Stigma for People With Mental Illness. *Res Soc Work Pract* 2018; **28**: 827-837 [DOI: 10.1177/1049731516681849]
- 174 **Lu T**, Guo Z, Li H, Zhang X, Ren Z, Yang W, Wei L, Huang L. Effects of Wise Intervention on Perceived Discrimination Among College Students Returning Home From Wuhan During the COVID-19 Outbreak. *Front Psychol* 2021; **12**: 689251 [PMID: 34163414 DOI: 10.3389/fpsyg.2021.689251]
- 175 **Siegel J**, Yassi A, Rau A, Buxton JA, Wouters E, Engelbrecht MC, Uebel KE, Nophale LE. Workplace interventions to reduce HIV and TB stigma among health care workers - Where do we go from here? *Glob Public Health* 2015; **10**: 995-1007 [PMID: 25769042 DOI: 10.1080/17441692.2015.1021365]
- 176 **Villotti P**, Zaniboni S, Corbière M, Guay S, Fraccaroli F. Reducing perceived stigma: Work integration of people with severe mental disorders in Italian social enterprise. *Psychiatr Rehabil J* 2018; **41**: 125-134 [PMID: 29698004 DOI: 10.1037/prj0000299]
- 177 **Gaiha SM**, Salisbury TT, Usmani S, Koschorke M, Raman U, Petticrew M. Effectiveness of arts interventions to reduce mental-health-related stigma among youth: a systematic review and meta-analysis. *BMC Psychiatry* 2021; **21**: 364 [PMID: 34294067 DOI: 10.1186/s12888-021-03350-8]
- 178 **Zeppegno P**, Gramaglia C, Feggi A, Lombardi A, Torre E. The effectiveness of a new approach using movies in the training of medical students. *Perspect Med Educ* 2015; **4**: 261-263 [PMID: 26346496 DOI: 10.1007/s40037-015-0208-6]
- 179 **Teti M**, Schulhoff AM, Koegler E, Saffran L, Bauerband LA, Shaffer V. Exploring the Use of Photo-Stories and Fiction Writing to Address HIV Stigma Among Health Professions Students. *Qual Health Res* 2019; **29**: 260-269 [PMID: 30095044 DOI: 10.1177/1049732318790939]
- 180 **Petkari E**. Building Beautiful Minds: Teaching Through Movies to Tackle Stigma in Psychology Students in the UAE. *Acad Psychiatry* 2017; **41**: 724-732 [PMID: 28577114 DOI: 10.1007/s40596-017-0723-3]
- 181 **Yadavaia JE**, Hayes SC. Acceptance and Commitment Therapy for Self-Stigma Around Sexual Orientation: A Multiple Baseline Evaluation. *Cogn Behav Pract* 2012; **19**: 545-559 [DOI: 10.1016/j.cbpra.2011.09.002]
- 182 **Levin ME**, Potts S, Haeger J, Lillis J. Delivering Acceptance and Commitment Therapy for Weight Self-Stigma Through Guided Self-Help: Results From an Open Pilot Trial. *Cogn Behav Pract* 2018; **25**: 87-104 [DOI: 10.1016/j.cbpra.2017.02.002]
- 183 **Gul M**, Aqeel M. Acceptance and Commitment Therapy for Treatment of Stigma and Shame in Substance Use Disorders: A Double-Blind, Parallel-Group, Randomized Controlled Trial. *J Subst Use* 2020; **26**: 4, 413-419 [DOI: 10.1080/14659891.2020.1846803]
- 184 **Sun L**, Liu X, Weng X, Deng H, Li Q, Liu J, Luan X. Narrative therapy to relieve stigma in oral cancer patients: A randomized controlled trial. *Int J Nurs Pract* 2022; **28**: e12926 [PMID: 33817894 DOI: 10.1111/ijn.12926]
- 185 **Ironson G**, Bira L, Hylton E. Positive and negative emotional expression measured from a single written essay about trauma predicts survival 17 years later in people living with HIV. *J Psychosom Res* 2020; **136**: 110166 [PMID: 32559504 DOI: 10.1016/j.jpsychores.2020.110166]
- 186 **Tornero MDLA**, Capella C. Change during Psychotherapy through Sand Play Tray in Children That Have Been Sexually Abused. *Front Psychol* 2017; **8**: 617 [PMID: 28522978 DOI: 10.3389/fpsyg.2017.00617]
- 187 **Skinta MD**, Lezama M, Wells G, Dilley JW. Acceptance and Compassion-Based Group Therapy to Reduce HIV Stigma. *Cogn Behav Pract* 2015; **22**: 481-490 [DOI: 10.1016/j.cbpra.2014.05.006]
- 188 **Harris PB**, Caporella CA. An intergenerational choir formed to lessen Alzheimer's disease stigma in college students and decrease the social isolation of people with Alzheimer's disease and their family members: a pilot study. *Am J Alzheimers Dis Other Demen* 2014; **29**: 270-281 [PMID: 24413542 DOI: 10.1177/1533317513517044]
- 189 **Lee Y**, Yoon HJ, Joo J. Role of Fresh Start Mindset Framing in Reducing Stigma and Promoting Mental Health Help-Seeking Behavior. *J Consum Behav* 2023; **cb.2155** [DOI: 10.1002/cb.2155]
- 190 **Tajti J**, Szok D, Csáti A, Szabó Á, Tanaka M, Vécsei L. Exploring Novel Therapeutic Targets in the Common Pathogenic Factors in Migraine and Neuropathic Pain. *Int J Mol Sci* 2023; **24** [PMID: 36835524 DOI: 10.3390/ijms24044114]
- 191 **Tanaka M**, Szabó Á, Vécsei L. Integrating Armchair, Bench, and Bedside Research for Behavioral Neurology and Neuropsychiatry: Editorial. *Biomedicine* 2022; **10** [PMID: 36551755 DOI: 10.3390/biomedicine10122999]
- 192 **Battaglia S**, Cardelluccio P, Di Fazio C, Nazzi C, Fracasso A, Borgomaneri S. Stopping in (e)motion: Reactive action inhibition when facing valence-independent emotional stimuli. *Front Behav Neurosci* 2022; **16**: 998714 [PMID: 36248028 DOI: 10.3389/fnbeh.2022.998714]
- 193 **Tanaka M**, Szabó Á, Spekter E, Polyák H, Tóth F, Vécsei L. Mitochondrial Impairment: A Common Motif in Neuropsychiatric Presentation? The Link to the Tryptophan-Kynurenine Metabolic System. *Cells* 2022; **11** [PMID: 36010683 DOI: 10.3390/cells11162607]



- 194 **Lam C**, Tsang H, Corrigan P, Lee Y-T, Angell B, Shi K, H. J, Larson J. Chinese Lay Theory and Mental Illness Stigma: Implications for Research and Practices. *J Rehabil* 2010; **76**: 35-40
- 195 **Ran MS**, Hall BJ, Su TT, Prawira B, Breth-Petersen M, Li XH, Zhang TM. Stigma of mental illness and cultural factors in Pacific Rim region: a systematic review. *BMC Psychiatry* 2021; **21**: 8 [PMID: [33413195](#) DOI: [10.1186/s12888-020-02991-5](#)]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

**Telephone:** +1-925-3991568

**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

**Help Desk:** <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

