

# World Journal of *Psychiatry*

*World J Psychiatry* 2024 January 19; 14(1): 1-193



**EDITORIAL**

- 1 Unlocking the power of physical activity in easing psychological distress  
*Liu XQ, Wang X*

**MINIREVIEWS**

- 8 Effects of psychological intervention on negative emotions and psychological resilience in breast cancer patients after radical mastectomy  
*Wang J, Kang DX, Zhang AJ, Li BR*

**ORIGINAL ARTICLE****Clinical and Translational Research**

- 15 Association between inflammatory bowel disease and all-cause dementia: A two-sample Mendelian randomization study  
*Liao OL, Xie SY, Ye J, Du Q, Lou GC*

**Retrospective Cohort Study**

- 26 Effects of ulinastatin combined with dexmedetomidine on cognitive dysfunction and emergence agitation in elderly patients who underwent total hip arthroplasty  
*Huo QF, Zhu LJ, Guo JW, Jiang YA, Zhao J*

**Retrospective Study**

- 36 Survey and clinical considerations of gender identity in lower primary school children  
*Zhang YL, Zhang HM, Xu JX, Zhou QY, Wang H, Pan XC*
- 44 Improvement of the nutritional support management system for patients in intensive care units  
*Zhang YY, Wang CY, Guo DX, Gao HN, Jin XS, Wu YL, Chen LH, Feng ZX*
- 53 Assessing myocardial indices and inflammatory factors to determine anxiety and depression severity in patients with chronic heart failure  
*Zhang L, Wang Q, Cui HS, Luo YY*
- 63 Postpartum quality of life and mental health in women with heart disease: Integrated clinical communication and treatment  
*Liu JL, Wang Q, Qu DY*
- 76 Clinicopathological features, psychological status, and prognosis of 33 patients with occult breast cancer  
*Wang HM, Yu AY, Li LL, Ma LY, Cao MH, Yang YL, Qin XB, Tang JJ, Han ZX*

**Observational Study**

- 88 Des-Arg(9) bradykinin as a causal metabolite for autism spectrum disorder  
*Huang ZY, Lyu ZP, Li HG, You HZ, Yang XN, Cha CH*
- 102 Performance of the walking trail making test in older adults with white matter hyperintensities  
*Zhao HY, Zhang ZQ, Huang YH, Li H, Wei FY*
- 111 Embracing different languages and local differences: Co-constructive patient simulation strengthens host countries' clinical training in psychiatry  
*Çamlı ŞE, Yavuz BE, Gök MF, Yazgan I, Yazgan Y, Brand-Gothelf A, Gothelf D, Amsalem D, Martin A*
- 119 Postpartum depression and partner support during the period of lactation: Correlation research and its influencing factors  
*Ruan JM, Wu LJ*
- 128 Abnormalities of electroencephalography microstates in patients with depression and their association with cognitive function  
*Peng RJ, Fan Y, Li J, Zhu F, Tian Q, Zhang XB*
- 141 Analysis of influencing factors of anxiety and depression in patients with periodontitis  
*Kong Y*
- 148 Relationship between physical activity and specific working memory indicators of depressive symptoms in university students  
*Zhao Q, Wang X, Li SF, Wang P, Wang X, Xin X, Yin SW, Yin ZS, Mao LJ*

**Basic Study**

- 159 Nutritional epigenetics education improves diet and attitude of parents of children with autism or attention deficit/hyperactivity disorder  
*Dufault RJ, Adler KM, Carpenter DO, Gilbert SG, Crider RA*

**META-ANALYSIS**

- 179 Global epidemiology of mental disorder in atrial fibrillation between 1998-2021: A systematic review and meta-analysis  
*Zhang S, Zhang N, Liu L, Zheng W, Ma ZL, Qiao SY, Zhao YL, Wei YH, Wu G, Yu QT, Deng B, Shen L*

**ABOUT COVER**

Editorial Board Member of *World Journal of Psychiatry*, Subho Chakrabarti, MD, FRCPsych, FAMS, Professor, Department of Psychiatry, Postgraduate Institute of Medical Education and Research, Chandigarh 16002, India. [chakrabarti.subho@pgimer.edu.in](mailto:chakrabarti.subho@pgimer.edu.in)

**AIMS AND SCOPE**

The primary aim of *World Journal of Psychiatry* (*WJP*, *World J Psychiatry*) is to provide scholars and readers from various fields of psychiatry with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

*WJP* mainly publishes articles reporting research results and findings obtained in the field of psychiatry and covering a wide range of topics including adolescent psychiatry, biological psychiatry, child psychiatry, community psychiatry, ethnopsychology, psychoanalysis, psychosomatic medicine, etc.

**INDEXING/ABSTRACTING**

The *WJP* is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports/Science Edition, PubMed, PubMed Central, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for *WJP* as 3.1; IF without journal self cites: 2.9; 5-year IF: 4.2; Journal Citation Indicator: 0.52; Ranking: 91 among 155 journals in psychiatry; and Quartile category: Q3.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: *Yu-Xi Chen*; Production Department Director: *Xu Guo*; Editorial Office Director: *Jia-Ping Yan*.

**NAME OF JOURNAL**

*World Journal of Psychiatry*

**ISSN**

ISSN 2220-3206 (online)

**LAUNCH DATE**

December 31, 2011

**FREQUENCY**

Monthly

**EDITORS-IN-CHIEF**

Ting-Shao Zhu

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/2220-3206/editorialboard.htm>

**PUBLICATION DATE**

January 19, 2024

**COPYRIGHT**

© 2024 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>



## Retrospective Cohort Study

# Effects of ulinastatin combined with dexmedetomidine on cognitive dysfunction and emergence agitation in elderly patients who underwent total hip arthroplasty

Qi-Fan Huo, Li-Juan Zhu, Jian-Wei Guo, Yan-An Jiang, Jing Zhao

**Specialty type:** Psychiatry

**Provenance and peer review:**

Unsolicited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): 0  
Grade B (Very good): 0  
Grade C (Good): C, C  
Grade D (Fair): 0  
Grade E (Poor): 0

**P-Reviewer:** Trzeciak P, Poland;  
Yirci R, Turkey

**Received:** August 30, 2023

**Peer-review started:** August 30, 2023

**First decision:** September 13, 2023

**Revised:** October 11, 2023

**Accepted:** December 5, 2023

**Article in press:** December 5, 2023

**Published online:** January 19, 2024



Qi-Fan Huo, Li-Juan Zhu, Jian-Wei Guo, Yan-An Jiang, Jing Zhao, Department of Anesthesiology, Shaanxi Provincial People's Hospital, Xi'an 710068, Shaanxi Province, China

**Corresponding author:** Jing Zhao, MM, Doctor, Department of Anesthesiology, Shaanxi Provincial People's Hospital, No. 256 Youyi West Road, Beilin District, Xi'an 710068, Shaanxi Province, China. [zhaojingsxxa@163.com](mailto:zhaojingsxxa@163.com)

## Abstract

### BACKGROUND

With the continuous growth of the modern elderly population, the risk of fracture increases. Hip fracture is a common type of fracture in older people. Total hip arthroplasty (THA) has significant advantages in relieving chronic pain and promoting the recovery of hip joint function.

### AIM

To investigate the effect of ulinastatin combined with dexmedetomidine (Dex) on the incidences of postoperative cognitive dysfunction (POCD) and emergence agitation in elderly patients who underwent THA.

### METHODS

A total of 397 patients who underwent THA from February 2019 to August 2022. We conducted a three-year retrospective cohort study in Shaanxi Provincial People's Hospital. Comprehensive demographic data were obtained from the electronic medical record system. We collected preoperative, intraoperative, and postoperative data. One hundred twenty-nine patients who were administered Dex during the operation were included in the Dex group. One hundred fifty patients who were intravenously injected with ulinastatin 15 min before anesthesia induction were included in the ulinastatin group. One hundred eighteen patients who were administered ulinastatin combined with Dex during the operation were included in the Dex + ulinastatin group. The patients' perioperative conditions, hemodynamic indexes, postoperative Mini-Mental State Examination (MMSE) scores, Ramsay score, incidence of POCD, and serum inflammatory cytokines were evaluated.

### RESULTS

There was a significant difference in the 24 h visual analogue scale score among



the three groups, and the score in the Dex + ulinastatin group was the lowest ( $P < 0.05$ ). Compared with the Dex and ulinastatin group, the MMSE scores of the Dex + ulinastatin group were significantly increased at 1 and 7 d after the operation (all  $P < 0.05$ ). Compared with those in the Dex and ulinastatin groups, incidence of POCD, levels of serum inflammatory cytokines in the Dex + ulinastatin group were significantly decreased at 1 and 7 d after the operation (all  $P < 0.05$ ). The observer's assessment of the alertness/sedation score and Ramsay score of the Dex + ulinastatin group were significantly different from those of the Dex and ulinastatin groups on the first day after the operation (all  $P < 0.05$ ).

## CONCLUSION

Ulinastatin combined with Dex can prevent the occurrence of POCD and emergence agitation in elderly patients undergoing THA.

**Key Words:** Ulinastatin; Dexmedetomidine; Postoperative cognitive dysfunction; Inflammatory cytokines; Total hip arthroplasty

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** In this study, we found a significant difference in the score of 24 h visual analogue scale among the three groups, and the score in the dexmedetomidine (Dex)+ ulinastatin group was the lowest. Compared with the Dex and ulinastatin group, the Mini-Mental State Examination scores of the Dex + ulinastatin group were significantly increased at 1 and 7 d after the operation. Compared with the Dex and ulinastatin group, the incidence of postoperative cognitive dysfunction (POCD), the levels of serum inflammatory cytokines of the Dex + ulinastatin group were significantly decreased at 1 and 7 d after the operation. The observer's assessment of the alertness/sedation score and Ramsay score of the Dex + ulinastatin group were significantly different from those of the Dex and ulinastatin groups on the first day after the operation. We observed that ulinastatin combined with dexmedetomidine can prevent the occurrence of POCD and the emergence of agitation in elderly patients undergoing total hip arthroplasty.

**Citation:** Huo QF, Zhu LJ, Guo JW, Jiang YA, Zhao J. Effects of ulinastatin combined with dexmedetomidine on cognitive dysfunction and emergence agitation in elderly patients who underwent total hip arthroplasty. *World J Psychiatry* 2024; 14(1): 26-35

**URL:** <https://www.wjgnet.com/2220-3206/full/v14/i1/26.htm>

**DOI:** <https://dx.doi.org/10.5498/wjp.v14.i1.26>

## INTRODUCTION

Older adults are prone to fall because of their slow response and unstable gait. With the continuous growth of the elderly population, the risk of fracture has increased. Hip fracture is a common type of fracture in older people[1]. Traditional conservative treatment is prone to delay healing and cause muscle atrophy, joint stiffness, and other problems, seriously affecting the quality of life of elderly patients[2]. With the improvement of modern surgical technology, an increasing number of elderly patients with fractures choose surgical treatment[3]. Total hip arthroplasty (THA) has significant advantages in relieving chronic pain and promoting the recovery of hip joint function[4]. However, most elderly patients have chronic diseases, and their tolerance to surgery is poor, so they are prone to various complications in the perioperative period, such as postoperative delirium, cognitive dysfunction, and pressure sores[5]. The occurrence of postoperative complications is closely related to the physical stress response, so corresponding intervention measures should be taken perioperatively to reduce postoperative stress reactions and the occurrence of complications in elderly patients.

Dexmedetomidine (Dex) is a specific  $\alpha_2$  adrenoceptor agonist that is effective for surgical analgesia and sedation[6]. Dex can effectively relieve postoperative pain and reduce the stress reaction in elderly patients undergoing medullary joint replacement[7]. Emerging evidence suggests that intraoperative application of Dex can reduce the risk of postoperative delirium and cognitive dysfunction[8]. Remarkably, it has been shown recently that Dex can protect the nervous system by reducing surgical stress and inflammatory responses[9]. Ulinastatin is a trypsin inhibitor that can effectively inhibit the decomposition of many enzymes and the release of inflammatory factors[10]. As reported previously, ulinastatin reduced neuronal apoptosis by inhibiting inflammatory factors such as interleukin (IL)-6 and C-reactive protein (CRP), thus relieving postoperative cognitive impairment and reducing the incidence of postoperative cognitive dysfunction (POCD) [11]. However, it is unclear whether the combination of ulinastatin and Dex can further reduce the incidence of POCD in elderly patients undergoing THA. Therefore, the aim of the current study is to evaluate the effect of ulinastatin combined with Dex on the incidences of POCD and emergence agitation. We also investigated the observer's assessment of alertness/sedation (OAA/S) scale scores of the elderly patients undergoing THA.

## MATERIALS AND METHODS

### Study design and patients

We conducted a three-year (February 2019 to August 2022) retrospective cohort study in Shaanxi Provincial People's Hospital. Comprehensive demographic data were obtained from the electronic medical record system. The Ethics Committee of the Shaanxi Provincial People's Hospital permitted this study, which was carried out according to the Declaration of Helsinki. We retrospectively reviewed the medical records of patients who underwent unilateral THA.

The inclusion criteria in this retrospective study were as follows: (1) Patients undergoing THA under general anesthesia in our hospital; (2) Patients aged 65 to 80 years old; (3) Patients with an American Society of Anesthesiologists (ASA) class I-III; and (4) Patients with a body mass index (BMI) 18-25 kg/m<sup>2</sup>. We excluded: (1) Patients with delirium or those diagnosed with dementia, depression, schizophrenia or epilepsy, Parkinson's disease, hypertension, coronary heart disease, diabetes, severe liver and kidney dysfunction, or hearing impairment in the past; and (2) Patients with severe hepatorenal dysfunction, severe malnutrition, blood coagulation disorders, and immune system diseases.

### Data collection

Preoperative information of qualified patients, including sex, age, BMI, ASA classification, mental disease, and cognitive damage, was obtained from the medical records. Intraoperative statistics were collected from anesthesia records, including anesthesia and surgery method, operation time, intraoperative data, blood loss volume, urine volume, and Dex and ulinastatin dosage. Record the heart rate (HR) and mean arterial pressure (MAP) of three groups of patients before anesthesia, 10 min of anesthesia, and 60 min of anesthesia. Postoperative data, including visual analogue scale (VAS) scores[12] at 24 h after the operation, Ramsay scores, Mini-Mental State Examination (MMSE) scores, OAA/S scale scores, serum CRP and IL-6, were collected. The Ramsay score is 1-6 points, with 1 point indicating restlessness; 2 points: Clear and able to follow commands; 3 points: Sleepy, able to follow instructions; 4 points: Shallow sleep state, able to awaken; 5 points: Call response is slow; 6 points: Deep sleep state, no response; the higher the score, the better the sedative effect. The MMSE score[13] included 10 points for orientation, 9 points for language and spatial structure, 5 points for attention and calculation, 3 points for recognition memory, and 3 points for memory, for a total of 30 points. Those with a score 2 points lower than those before anesthesia were considered to have cognitive impairment. The VAS score was evaluated by a visual analog scale, with a total score of 10. The higher the score, the more severe the pain. The OAA/S sedation score[14] is divided into five levels: Level 1: The patient is unconscious and does not respond to gentle tapping on the body but responds to noxious stimuli. Level 2: The patient's level of consciousness is vague but consciousness is not wholly lost, and there is no response when the name is called loudly; Level 3: The level of consciousness is vague but consciousness is not wholly lost, and there is a response to tapping the body; Level 4: The patient's level of consciousness is vague, and there is a response when calling the name; Level 5: The patient is conscious and can respond to the questions asked by the nursing staff.

### Outcome measures and statistical analysis

By examining the patients' electronic medical records, 576 elderly patients who underwent THA in our hospital from February 2019 to August 2022 were selected. Among them, 129 patients who were administered Dex 0.3 µg/kg/h during the operation were included in the Dex group. One hundred fifty patients who were intravenously injected with 5000 U/kg ulinastatin 15 min before anesthesia were included in the ulinastatin group. During the operation, one hundred eighteen patients who were administered 5000 U/kg ulinastatin (15 min before anesthesia) combined with Dex 0.3 µg/kg/h were included in the Dex + ulinastatin group. Data were evaluated by SPSS 22.0 (IBM Corporation). The categorical data are indicated as *n* (%), and continuous variables are indicated as the mean ± SD. Continuous variables were compared with one-way ANOVA or the Mann-Whitney *U* test. Categorical variables were compared with Fisher's exact test. *P* < 0.05 was considered statistically significant.

## RESULTS

### Patient inclusion process

The original investigation of our hospital's electronic medical system showed that 576 patients underwent THA from February 2019 to August 2022. Fifty-eight patients who were either under age 65 years or over 80 years were disqualified. After excluding 63 patients complicated with severe chronic physical diseases and 58 patients with incomplete data, 397 patients were eligible for this study. A flowchart describing the selection of patients included in this retrospective study is shown in Figure 1.

### Patient characteristics

The Dex cohort (129 cases) consisted of patients who received Dex during the operation. The ulinastatin cohort (150 cases) consisted of patients who received ulinastatin 15 min before anesthesia. In the Dex + ulinastatin group, 118 patients received Dex (during the operation) combined with ulinastatin (15 min before anesthesia). The descriptive statistics between the three cohorts were not statistically significant (all *P* > 0.05, Table 1).

### Comparison of perioperative conditions

Table 2 shows no significant difference in operation duration, intraoperative blood loss volume, operation time,

**Table 1 Descriptive statistics of the demographics of patients.**

	Dex group (n = 129)	Ulinastatin group (n = 150)	Dex + ulinastatin group (n = 118)	<i>F</i> / $\chi^2$	<i>P</i> value
Sex, <i>n</i>				1.240	0.092
Male	62	74	53		
Female	67	76	65		
Age (yr), mean $\pm$ SD	71.72 $\pm$ 2.45	71.23 $\pm$ 4.91	72.25 $\pm$ 2.65	1.217	0.301
Weight (kg), mean $\pm$ SD	59.12 $\pm$ 6.90	61.21 $\pm$ 7.13	59.23 $\pm$ 6.18	0.680	0.509
BMI, mean $\pm$ SD	24.80 $\pm$ 4.69	24.72 $\pm$ 5.04	24.90 $\pm$ 4.82	1.207	0.304
Intraoperative hypotension, <i>n</i>	8	7	9	2.447	0.099
Marital status, <i>n</i>				0.669	0.095
Single	10	7	6		
Married	119	143	112		
ASA classification (level III)	5	6	4	0.519	0.082

BMI: Body mass index; ASA classification: American Society of Anesthesiologists classification; Dex: Dexmedetomidine.

**Table 2 Comparison of perioperative conditions among the three groups**

	Dex group (n = 129)	Ulinastatin group (n = 150)	Dex + ulinastatin group (n = 118)	<i>F</i> value	<i>P</i> value
Intraoperative blood loss (mL)	33.12 $\pm$ 3.75	34.27 $\pm$ 3.81	32.73 $\pm$ 3.85	2.297	0.107
Operation duration (min)	206.45 $\pm$ 36.32	201.65 $\pm$ 40.03	199.85 $\pm$ 40.38	0.690	0.082
Preoperative MMSE score	28.70 $\pm$ 0.82	28.84 $\pm$ 0.92	28.52 $\pm$ 0.84	0.379	0.686
VAS score 24 h after the operation	3.25 $\pm$ 0.37	3.30 $\pm$ 0.52	3.13 $\pm$ 0.25	3.878	0.024
Time of anesthesia (min)	217.18 $\pm$ 42.79	220.51 $\pm$ 38.94	223.35 $\pm$ 41.67	1.550	0.980
Infusion volume (mL)	1558.72 $\pm$ 170.56	1489.54 $\pm$ 187.46	1580.67 $\pm$ 191.88	0.673	0.058
Urine volume (mL)	174.87 $\pm$ 42.57	165.38 $\pm$ 56.16	168.48 $\pm$ 58.16	1.097	0.779

Dex: Dexmedetomidine; MMSE: Mini-Mental State Examination.

preoperative MMSE scores, infusion volume, total blood loss volume, or urine volume among the three groups (all  $P > 0.05$ ). There were significant differences in VAS scores at 24 h after the operation among the three groups, and the score in the Dex + ulinastatin group was the lowest ( $P < 0.05$ ).

### Comparison of MMSE scores

Compared with those of the Dex and ulinastatin groups, the MMSE scores of the Dex + ulinastatin group were significantly increased at 1 and 7 d after the operation (all  $P < 0.05$ , Table 3 and Figure 2A).

### Comparison of the incidence of POCD

Compared with that in the Dex and ulinastatin groups, the incidence of POCD in the Dex + ulinastatin group significantly decreased at 1 and 7 d after the operation (all  $P < 0.05$ , Table 4 and Figure 2B).

### Comparison of serum inflammatory cytokines

Compared with those in the Dex and ulinastatin groups, the levels of serum inflammatory cytokines (CRP and IL-6) in the Dex + ulinastatin group were significantly decreased at 1 and 7 d after the operation (all  $P < 0.05$ , Figure 3).

### Comparison of OAA/S sedation score and Ramsay score on the first day after the operation

As shown in Table 5, OAA/S sedation score and Ramsay score of the Dex + ulinastatin group were significantly different from those of the Dex and ulinastatin groups on the first day after the operation (all  $P < 0.05$ ).

### Comparison of hemodynamic indexes among the three groups

Compared with those of the Dex and ulinastatin groups, HR and MAP of the Dex + ulinastatin group significantly



**Table 3 Comparison of Mini-Mental State Examination scores among the three groups**

	Dex group (n = 129)	Ulinastatin group (n = 150)	Dex + ulinastatin group (n = 118)
Preoperative MMSE score	28.70 ± 0.82	28.84 ± 0.92	28.52 ± 0.84
One day after the operation	21.87 ± 6.35 <sup>a</sup>	20.72 ± 5.45 <sup>b</sup>	23.70 ± 2.46
Seven days after the operation	26.29 ± 2.86 <sup>a</sup>	25.16 ± 3.90 <sup>b</sup>	27.60 ± 1.76

<sup>a</sup>*P* < 0.05, dexmedetomidine + ulinastatin *vs* dexmedetomidine.  
<sup>b</sup>*P* < 0.01, dexmedetomidine + ulinastatin *vs* ulinastatin.  
MMSE: Mini-Mental State Examination; Dex: Dexmedetomidine.

**Table 4 Comparison of the incidence of postoperative cognitive dysfunction among the three groups**


	Dex group (n = 129)	Ulinastatin group (n = 150)	Dex + ulinastatin group (n = 118)
One day after the operation	19 (14.73%) <sup>a</sup>	28 (18.67%) <sup>b</sup>	6 (5.08%)
Seven days after the operation	12 (9.30%) <sup>a</sup>	18 (12.00%) <sup>b</sup>	2 (1.69%)

<sup>a</sup>*P* < 0.05, dexmedetomidine + ulinastatin *vs* dexmedetomidine.  
<sup>b</sup>*P* < 0.01, dexmedetomidine + ulinastatin *vs* ulinastatin.  
Dex: Dexmedetomidine.

**Table 5 Comparison of observer's assessment of the alertness/sedation scores and Ramsay score on the first day after the operation among the three groups**


	Dex group (n = 129)	Ulinastatin group (n = 150)	Dex + ulinastatin group (n = 118)	F value	P value
OAA/S sedation score	3.60 ± 0.61	3.51 ± 0.60	2.21 ± 0.40	9.501	< 0.001
Ramsay score	1.98 ± 0.23	2.02 ± 0.33	3.21 ± 1.02	15.698	< 0.001

Dex: Dexmedetomidine; OAA/S: Observer's assessment of the alertness/sedation.




DOI: 10.5498/wjp.v14.i1.26 Copyright ©The Author(s) 2024.

**Figure 1** Flow diagram describing the selection of patients involved in this retrospective study. Dex: Dexmedetomidine.



**Figure 2 Comparison of Mini-Mental State Examination scores and the incidence of postoperative cognitive dysfunction among the three groups.** A: Mini-Mental State Examination scores; B: The incidence of postoperative cognitive dysfunction. <sup>a</sup> $P < 0.05$ , dexmedetomidine + ulinastatin vs dexmedetomidine. <sup>b</sup> $P < 0.01$ , dexmedetomidine + ulinastatin vs ulinastatin. MMSE: Mini-Mental State Examination; POCD: Postoperative cognitive dysfunction; Dex: Dexmedetomidine.



**Figure 3 Comparison of serum inflammatory cytokines among the three groups.** A: Serum C-reactive protein; B: Serum interleukin-6. <sup>a</sup> $P < 0.05$ , dexmedetomidine + ulinastatin vs dexmedetomidine. <sup>b</sup> $P < 0.01$ , dexmedetomidine + ulinastatin vs ulinastatin. Dex: Dexmedetomidine; CRP: C-reactive protein; IL-6: Interleukin-6.

reduced on the 10 minutes of anesthesia and 60 min of anesthesia (all  $P < 0.05$ , Table 6).

## DISCUSSION

THA is a standard approach for the surgical treatment of hip paralysis that mainly involves replacement of the damaged hip joint with an artificial hip[15]. The efficacy and safety of THA have improved in recent years. However, patients still have a particular risk of postoperative complications, mainly due to various stress reactions caused by bleeding intraoperatively, performing the operation, and making the surgical incision[16]. Postoperative pain is a common complication after hip arthroplasty, and pain can cause symptoms such as increased HR and high blood pressure[17]. Effective analgesic measures should be taken to prevent various complications caused by postoperative stress reactions[18].

Dex is an  $\alpha_2$  receptor agonist with a high affinity for  $\alpha_2$  receptors and is often used in sedation therapy in the intensive care unit and surgery. The drug's half-life is 6 min after intravenous injection, excreted in urine after extensive metabolism, and the clearance half-life is two hours[19]. Because the drug is injected through the kidney, patients with impaired renal function may have more adverse reactions[20]. Therefore, patients with hepatic and renal dysfunction were excluded from this study. Ulinastatin is isolated in and excreted through human urine, and widely used because of its broad spectrum of enzyme inhibition[21]. Previous studies have shown that ulinastatin can inhibit the organic inflammatory response and improve learning and memory dysfunction by reducing the apoptosis of nerve cells, which is beneficial in reducing the incidence of postoperative cognitive impairment in elderly patients[22]. Combining two drugs with the same effect has a synergistic effect. However, the effect of the combination of Dex and ulinastatin on elderly patients undergoing THA remains unknown. Therefore, this retrospective cohort study investigated the effect of Dex combined with ulinastatin on the incidences of POCD and emergence agitation in elderly patients who underwent THA.

As reported previously, ulinastatin has an excellent anti-inflammatory response, reduces the level of inflammatory mediators, relieves postoperative pain, and reduces the postoperative stress response. The clearance rate of the drug in elderly patients is slower than that in young patients, and because the kidney does not metabolize the drug, it can be used even if the patient has impaired renal function[23]. In this study, we observed that the combination of ulinastatin and Dex could effectively reduce the stress response and improve the quality of life of patients after the operation. The VAS scores of the Dex + ulinastatin group were lower than those of the ulinastatin or Dex group at 24 h after the operation, indicating that the analgesic scheme of ulinastatin combined with Dex could effectively relieve postoperative pain

**Table 6 Comparison of hemodynamic indexes among the three groups**

Item		Dex group (n = 129)	Ulinastatin group (n = 150)	Dex + ulinastatin group (n = 118)	F value	P value
HR (times/min)	T0	78.02 ± 3.28	78.66 ± 3.49	78.51 ± 3.64	1.140	0.323
	T1	84.32 ± 5.47	85.74 ± 5.48	81.07 ± 5.55	7.084	0.001
	T2	85.16 ± 5.39	86.63 ± 5.73	80.39 ± 5.85	18.882	< 0.001
MAP (mmHg)	T0	91.72 ± 8.74	92.04 ± 8.38	91.69 ± 8.77	0.033	0.965
	T1	96.31 ± 7.05	97.72 ± 7.37	92.11 ± 6.38	7.417	0.001
	T2	95.87 ± 7.85	96.48 ± 7.69	91.05 ± 7.94	8.289	< 0.001

Dex: Dexmedetomidine; HR: Heart rate; MAP: Mean arterial pressure; T0: Before anesthesia; T1: 10 min of anesthesia; T2: 60 min of anesthesia.

symptoms in elderly patients who underwent total joint replacement. The Ramsay score and OAA/S sedation score mainly reflects the sedation of patients after the operation[24]. In this study, OAA/S sedation score and Ramsay score of the Dex + ulinastatin group were significantly different from those of the Dex and ulinastatin groups on the first day after the operation, meanwhile, compared with those of the Dex and ulinastatin groups, HR and MAP of the Dex + ulinastatin group significantly reduced on the 10 min of anesthesia and 60 min of anesthesia, indicating that preemptive analgesia with ulinastatin combined with Dex could effectively reduce the postoperative stress response and improve the sedation effect.

Previous studies have shown that systemic inflammatory response syndrome caused by surgical trauma, postoperative pain, body stress, and the release of different types of cytokines play an essential role in the development of perioperative neurocognitive impairment[25]. The body's immune system can be activated by surgical trauma, leading to a robust inflammatory response[26]. Therefore, the levels of inflammatory factors in surgical patients' central nervous system and peripheral tissues increase. At the same time, surgery can lead to enhanced production of proinflammatory cytokines, such as plasma IL-6 and IL-1 $\beta$ , and the degree of increase is related to the decline in cognitive function[27]. Changes in cerebral oxygenation during THA may cause different degrees of nerve injury and induce POCD[28]. Therefore, how to reduce the incidence of POCD in elderly patients undergoing THA and improve their quality of life in their later years is particularly important. Our results found that compared with those of the Dex and ulinastatin groups, the MMSE scores of the Dex + ulinastatin group were significantly increased at 1 and 7 d after the operation, suggesting that preinjection of Dex and ulinastatin might play a significant role in preventing POCD and that the combined infusion is more helpful in preventing the occurrence of early POCD. A prophylactic combination of Dex and ulinastatin during knee arthroplasty in older adults is beneficial in reducing the incidence of POCD. The possible mechanism is that Dex inhibits excessive stress responses, such as inhibiting the concentration of plasma cortisol and reducing the damage of catecholamines to brain nerves. At the same time, ulinastatin protects the brain by inhibiting the release of proinflammatory cytokines (IL-6 and CRP).

This study also has some limitations. At present, multiscale combined evaluation is recommended for the diagnosis of POCD. This study used the MMSE scale to evaluate cognitive function, and some postoperative patients with cognitive impairment may not have been identified. Therefore, whether there are other mechanisms by which Dex combined with ulinastatin reduces the incidence of POCD in elderly patients undergoing THA should be further studied.

## CONCLUSION

In summary, ulinastatin combined with Dex can reduce the incidence of POCD in elderly patients undergoing THA, and the mechanism may be related to the reduction in plasma levels of CRP and IL-6. In addition, we also observed that applying ulinastatin combined with Dex can reduce postoperative pain and improve the postoperative stress response in elderly patients undergoing THA.

## ARTICLE HIGHLIGHTS

### Research background

Most elderly patients are prone to various complications in the perioperative period, such as postoperative delirium and cognitive dysfunction. Cognitive dysfunction seriously affects the postoperative rehabilitation of patients, prolongs hospital stay, increases the incidence of postoperative complications and mortality, in addition to a serious decline in personal quality of life, but also increases the burden of the family and society. Many researchers believe that advanced age and "major surgery" are important risk factors for cognitive dysfunction in patients undergoing non-cardiac surgery. Therefore, cognitive dysfunction is an important issue in the medical field at present, and the study of the occurrence factors, pathogenesis, effective prevention, and treatment of cognitive dysfunction is an important topic in the field of

anesthesiology.

### Research motivation

It is unclear whether the combination of ulinastatin and dexmedetomidine (Dex) can further reduce the incidence of postoperative cognitive dysfunction (POCD) in elderly patients undergoing total hip arthroplasty (THA). Therefore, the current study aims to evaluate the effect of ulinastatin combined with Dex on POCD and emergence agitation. Therefore, solving these problems could effectively improve the postoperative stress response and improve the sedation effect for elderly patients undergoing THA. This is the first study to explore the effect of ulinastatin combined with Dex on POCD and emergence agitation in elderly patients undergoing THA.

### Research objectives

In view of the high incidence of POCD in elderly orthopedic patients, this study will compare the effects of different anesthetic methods on the early cognitive function of elderly patients after THA, and explore the related factors, to provide some guidance for the anesthetic mode and anesthetic management of clinical orthopedic surgery in the future. The purpose of this study is to provide clinical basis for the selection of anesthesia in elderly patients undergoing hip arthroplasty.

### Research methods

In this study, we collected the postoperative data, including visual analogue scale (VAS) scores at 24 h after operation, postoperative stress response indicators [Mini-Mental State Examination (MMSE) scores, hemodynamic indexes, observer's assessment of alertness/sedation (OAA/S) scale and Ramsay score], inflammatory cytokines [C-reactive protein (CRP) and interleukin (IL)-6]. A total of 129 patients administrated with Dex 0.3 µg/kg/h during the operation were included in the Dex group. One hundred fifty patients who were intravenously injected 5000 U/kg of ulinastatin 15 min before anesthesia were included in the ulinastatin group. One hundred eighteen patients who were administrated with 5000 U/kg of ulinastatin (15 min before anesthesia) combined with Dex 0.3 µg/kg/h during the operation were included in the Dex + ulinastatin group.

### Research results

The findings of this study demonstrated that ulinastatin combined with Dex could reduce the incidence of POCD in elderly patients undergoing THA, and the mechanism might be associated with the lessening of plasma levels of CRP and IL-6. Moreover, the combination of ulinastatin and Dex can reduce postoperative pain and postoperative stress response in elderly patients undergoing THA. Our results validated that the combination therapy can effectively improve the cognitive function of nervous system after operation and improved the quality of life for patients.

### Research conclusions

This is the first study explored the efficacy of ulinastatin combined with Dex on POCD and emergence agitation in elderly patients undergoing THA. We collected the postoperative data, including VAS scores at 24 h after operation, postoperative stress response indicators (MMSE scores, hemodynamic indexes, OAA/S scale, CRP, and IL-6). The results of this study not only demonstrate that combination therapy can reduce postoperative cognitive impairment, but also find that the mechanism may be related to the reduction of plasma levels of CRP and IL-6.

### Research perspectives

In future research, we will focus on the specific mechanisms related to the combined therapy through *in vitro* and *in vivo* experiments.

---

## FOOTNOTES

**Author contributions:** Huo QF and Guo JW initiated the project and designed the experiment; Zhu LJ and Jiang YN conducted clinical data collection; Huo QF and Zhu LJ performed postoperative follow-up and recorded data; Guo JW, Jiang YN, and Zhao J conducted a number of collation and statistical analysis; Zhao J wrote the original manuscript and revised the paper; and all authors reviewed and approved the paper, and approved the final manuscript.

**Institutional review board statement:** This study was approved by the Ethics Committee of Shaanxi Provincial People's Hospital.

**Informed consent statement:** The informed consent was approved to waive.

**Conflict-of-interest statement:** All the authors report no relevant conflicts of interest for this article.

**Data sharing statement:** All data generated or analyzed during this study are included in this published article.

**STROBE statement:** The authors have read the STROBE Statement-checklist of items, and the manuscript was prepared and revised according to the STROBE Statement-checklist of items.

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers.

It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** China

**ORCID number:** Jing Zhao [0009-0001-2373-685X](https://orcid.org/0009-0001-2373-685X).

**S-Editor:** Wang JJ

**L-Editor:** A

**P-Editor:** Cai YX

## REFERENCES

- Meinberg E, Ward D, Herring M, Miclau T. Hospital-based Hip fracture programs: Clinical need and effectiveness. *Injury* 2020; **51** Suppl 2: S2-S4 [PMID: [32386840](https://pubmed.ncbi.nlm.nih.gov/32386840/) DOI: [10.1016/j.injury.2020.03.046](https://doi.org/10.1016/j.injury.2020.03.046)]
- Pech-Ciau BA, Lima-Martínez EA, Espinosa-Cruz GA, Pacho-Aguilar CR, Huchim-Lara O, Alejos-Gómez RA. [Hip fracture in the elderly: epidemiology and costs of care]. *Acta Ortop Mex* 2021; **35**: 341-347 [PMID: [35139593](https://pubmed.ncbi.nlm.nih.gov/35139593/)]
- Alexiou KI, Roushias A, Varitimidis SE, Malizos KN. Quality of life and psychological consequences in elderly patients after a hip fracture: a review. *Clin Interv Aging* 2018; **13**: 143-150 [PMID: [29416322](https://pubmed.ncbi.nlm.nih.gov/29416322/) DOI: [10.2147/CIA.S150067](https://doi.org/10.2147/CIA.S150067)]
- Scott CEH, Clement ND, Davis ET, Haddad FS. Modern total hip arthroplasty: peak of perfection or room for improvement? *Bone Joint J* 2022; **104-B**: 189-192 [PMID: [35094584](https://pubmed.ncbi.nlm.nih.gov/35094584/) DOI: [10.1302/0301-620X.104B2.BJJ-2022-0007](https://doi.org/10.1302/0301-620X.104B2.BJJ-2022-0007)]
- Stibolt RD Jr, Patel HA, Huntley SR, Lehtonen EJ, Shah AB, Naranje SM. Total hip arthroplasty for posttraumatic osteoarthritis following acetabular fracture: A systematic review of characteristics, outcomes, and complications. *Chin J Traumatol* 2018; **21**: 176-181 [PMID: [29773451](https://pubmed.ncbi.nlm.nih.gov/29773451/) DOI: [10.1016/j.cjtee.2018.02.004](https://doi.org/10.1016/j.cjtee.2018.02.004)]
- Wen G, Xin N. Dexmedetomidine promotes the activity of breast cancer cells through miR-199a/HIF-1α axis. *Transl Cancer Res* 2021; **10**: 4817-4828 [PMID: [35116334](https://pubmed.ncbi.nlm.nih.gov/35116334/) DOI: [10.21037/ter-21-1937](https://doi.org/10.21037/ter-21-1937)]
- Li X, Li X, Wang R, Hua T, Li G. Application of dexmedetomidine in gynecological laparoscopic surgery and its effect on S-100 β protein and cognitive function in patients. *Minerva Med* 2022; **113**: 599-601 [PMID: [34586767](https://pubmed.ncbi.nlm.nih.gov/34586767/) DOI: [10.23736/S0026-4806.21.07723-5](https://doi.org/10.23736/S0026-4806.21.07723-5)]
- Deiner S, Luo X, Lin HM, Sessler DI, Saager L, Sieber FE, Lee HB, Sano M; and the Dexlirium Writing Group, Jankowski C, Bergese SD, Candiotti K, Flaherty JH, Arora H, Shander A, Rock P. Intraoperative Infusion of Dexmedetomidine for Prevention of Postoperative Delirium and Cognitive Dysfunction in Elderly Patients Undergoing Major Elective Noncardiac Surgery: A Randomized Clinical Trial. *JAMA Surg* 2017; **152**: e171505 [PMID: [28593326](https://pubmed.ncbi.nlm.nih.gov/28593326/) DOI: [10.1001/jamasurg.2017.1505](https://doi.org/10.1001/jamasurg.2017.1505)]
- Muñoz-Leyva F, Jack JM, Bhatia A, Chin KJ, Gandhi R, Perlas A, Jin R, Chan V. No Benefits of Adding Dexmedetomidine, Ketamine, Dexamethasone, and Nerve Blocks to an Established Multimodal Analgesic Regimen after Total Knee Arthroplasty. *Anesthesiology* 2022; **137**: 459-470 [PMID: [35867857](https://pubmed.ncbi.nlm.nih.gov/35867857/) DOI: [10.1097/ALN.0000000000004326](https://doi.org/10.1097/ALN.0000000000004326)]
- Lv B, Jiang XM, Wang DW, Chen J, Han DF, Liu XL. Protective Effects and Mechanisms of Action of Ulinastatin against Cerebral Ischemia-Reperfusion Injury. *Curr Pharm Des* 2020; **26**: 3332-3340 [PMID: [32124689](https://pubmed.ncbi.nlm.nih.gov/32124689/) DOI: [10.2174/1381612826666200303114955](https://doi.org/10.2174/1381612826666200303114955)]
- Zhang YH, Guo XH, Zhang QM, Yan GT, Wang TL. Serum CRP and urinary trypsin inhibitor implicate postoperative cognitive dysfunction especially in elderly patients. *Int J Neurosci* 2015; **125**: 501-506 [PMID: [25105909](https://pubmed.ncbi.nlm.nih.gov/25105909/) DOI: [10.3109/00207454.2014.949341](https://doi.org/10.3109/00207454.2014.949341)]
- Han Y, Wang M, Shen J, Zhang Z, Zhao M, Huang J, Chen Y, Chen Z, Hu Y, Wang Y. Differential efficacy of methylcobalamin and alpha-lipoic acid treatment on symptoms of diabetic peripheral neuropathy. *Minerva Endocrinol* 2018; **43**: 11-18 [PMID: [27901334](https://pubmed.ncbi.nlm.nih.gov/27901334/) DOI: [10.23736/S0391-1977.16.02505-0](https://doi.org/10.23736/S0391-1977.16.02505-0)]
- Lauretani F, Ticinesi A, Gionti L, Prati B, Nouvenne A, Tana C, Meschi T, Maggio M. Short-Physical Performance Battery (SPPB) score is associated with falls in older outpatients. *Aging Clin Exp Res* 2019; **31**: 1435-1442 [PMID: [30515724](https://pubmed.ncbi.nlm.nih.gov/30515724/) DOI: [10.1007/s40520-018-1082-y](https://doi.org/10.1007/s40520-018-1082-y)]
- Lim TW, Choi YH, Kim JY, Choi JB, Lee SK, Youn EJ, Lee JS. Efficacy of the bispectral index and Observer's Assessment of Alertness/Sedation Scale in monitoring sedation during spinal anesthesia: A randomized clinical trial. *J Int Med Res* 2020; **48**: 300060519893165 [PMID: [31875756](https://pubmed.ncbi.nlm.nih.gov/31875756/) DOI: [10.1177/0300060519893165](https://doi.org/10.1177/0300060519893165)]
- Fleivas DA, Tsantes AG, Mavrogenis AF. Direct Anterior Approach Total Hip Arthroplasty Revisited. *JBJS Rev* 2020; **8**: e0144 [PMID: [32304500](https://pubmed.ncbi.nlm.nih.gov/32304500/) DOI: [10.2106/JBJS.RVW.19.00144](https://doi.org/10.2106/JBJS.RVW.19.00144)]
- Moerenhout K, Derome P, Laflamme GY, Leduc S, Gaspard HS, Benoit B. Direct anterior versus posterior approach for total hip arthroplasty: a multicentre, prospective, randomized clinical trial. *Can J Surg* 2020; **63**: E412-E417 [PMID: [33009898](https://pubmed.ncbi.nlm.nih.gov/33009898/) DOI: [10.1503/cjs.012019](https://doi.org/10.1503/cjs.012019)]
- Goh GS, Parvizi J. Nerve Injuries Following Total Hip Arthroplasty: The Influence of Surgical Approach. *Orthop Clin North Am* 2022; **53**: 129-137 [PMID: [35365257](https://pubmed.ncbi.nlm.nih.gov/35365257/) DOI: [10.1016/j.ocl.2021.12.002](https://doi.org/10.1016/j.ocl.2021.12.002)]
- Urish KL, Giori NJ, Lemons JE, Mihalko WM, Hallab N. Trunnion Corrosion in Total Hip Arthroplasty-Basic Concepts. *Orthop Clin North Am* 2019; **50**: 281-288 [PMID: [31084829](https://pubmed.ncbi.nlm.nih.gov/31084829/) DOI: [10.1016/j.ocl.2019.02.001](https://doi.org/10.1016/j.ocl.2019.02.001)]
- Persson NDÅ, Uusalo P, Nedergaard M, Lohela TJ, Lilius TO. Could dexmedetomidine be repurposed as a glymphatic enhancer? *Trends Pharmacol Sci* 2022; **43**: 1030-1040 [PMID: [36280451](https://pubmed.ncbi.nlm.nih.gov/36280451/) DOI: [10.1016/j.tips.2022.09.007](https://doi.org/10.1016/j.tips.2022.09.007)]
- Eizaga Rebollar R, García Palacios MV, Fernández Riobó MC, Torres Morera LM. Dexmedetomidine and perioperative analgesia in children. *Rev Esp Anestesiología Reanim (Engl Ed)* 2022; **69**: 487-492 [PMID: [36100555](https://pubmed.ncbi.nlm.nih.gov/36100555/) DOI: [10.1016/j.redare.2022.08.003](https://doi.org/10.1016/j.redare.2022.08.003)]
- Wei X, Zhu X, Jiang L, Long M, Du Y. Recent research progress on the role of ulinastatin in chronic kidney disease. *Nephrology (Carlton)* 2021; **26**: 708-714 [PMID: [34050574](https://pubmed.ncbi.nlm.nih.gov/34050574/) DOI: [10.1111/nep.13906](https://doi.org/10.1111/nep.13906)]
- Chen L, Jin S, Yang M, Gui C, Yuan Y, Dong G, Zeng W, Zeng J, Hu G, Qiao L, Wang J, Xi Y, Sun J, Wang N, Wang M, Xing L, Yang Y, Teng Y, Hou J, Bi Q, Cai H, Zhang G, Hong Y, Zhang Z. Integrated Single Cell and Bulk RNA-Seq Analysis Revealed Immunomodulatory Effects of Ulinastatin in Sepsis: A Multicenter Cohort Study. *Front Immunol* 2022; **13**: 882774 [PMID: [35634310](https://pubmed.ncbi.nlm.nih.gov/35634310/) DOI: [10.3389/fimm.2022.882774](https://doi.org/10.3389/fimm.2022.882774)]



- 10.3389/fimmu.2022.882774]
- 23 **Hong Y**, Meng S, Wang S, Liu T, Liu J. Ulinastatin Alleviates Repetitive Ketamine Exposure-Evoked Cognitive Impairment in Adolescent Mice. *Neural Plast* 2022; **2022**: 6168284 [PMID: 36545238 DOI: 10.1155/2022/6168284]
- 24 **Sieber F**, Neufeld K, Oh ES, Gottschalk A, Wang NY. Effect of baseline cognitive impairment on association between predicted propofol effect site concentration and Bispectral index or sedation score. *BMC Anesthesiol* 2020; **20**: 129 [PMID: 32466776 DOI: 10.1186/s12871-020-01043-5]
- 25 **Liao YL**, Zhou XY, Ji MH, Qiu LC, Chen XH, Gong CS, Lin Y, Guo YH, Yang JJ. S100A9 Upregulation Contributes to Learning and Memory Impairments by Promoting Microglia M1 Polarization in Sepsis Survivor Mice. *Inflammation* 2021; **44**: 307-320 [PMID: 32918665 DOI: 10.1007/s10753-020-01334-6]
- 26 **Bortolotti P**, Faure E, Kipnis E. Inflammasomes in Tissue Damages and Immune Disorders After Trauma. *Front Immunol* 2018; **9**: 1900 [PMID: 30166988 DOI: 10.3389/fimmu.2018.01900]
- 27 **Wang J**, Zhou Y, Li K, Li X, Guo M, Peng M. A Noradrenergic Lesion Attenuates Surgery-Induced Cognitive Impairment in Rats by Suppressing Neuroinflammation. *Front Mol Neurosci* 2021; **14**: 752838 [PMID: 34916906 DOI: 10.3389/fnmol.2021.752838]
- 28 **Lin R**, Zhang F, Xue Q, Yu B. Accuracy of regional cerebral oxygen saturation in predicting postoperative cognitive dysfunction after total hip arthroplasty: regional cerebral oxygen saturation predicts POCD. *J Arthroplasty* 2013; **28**: 494-497 [PMID: 23151365 DOI: 10.1016/j.arth.2012.06.041]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

**Telephone:** +1-925-3991568

**E-mail:** [office@baishideng.com](mailto:office@baishideng.com)

**Help Desk:** <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

