

World Journal of *Psychiatry*

World J Psychiatry 2024 April 19; 14(4): 489-599



EDITORIAL

- 489 Mindfulness training in medical education as a means to improve resilience, empathy, and mental health in the medical profession

Vidal EIO, Ribeiro LFA, Carvalho-Filho MA, Fukushima FB

REVIEW

- 494 Adolescent suicide risk factors and the integration of social-emotional skills in school-based prevention programs

Liu XQ, Wang X

ORIGINAL ARTICLE**Case Control Study**

- 507 Psychiatric outcomes in outpatients affected by long COVID: A link between mental health and persistence of olfactory complaint

Metelkina-Fernandez V, Dumas LE, Vandersteen C, Chirio D, Gros A, Fernandez A, Askenazy F, Manera V

Retrospective Cohort Study

- 513 Clarifying the relationship and analyzing the influential factors of bronchial asthma in children with attention-deficit hyperactivity disorder

Wang GX, Xu XY, Wu XQ

Retrospective Study

- 523 Relationship between plasma risperidone concentrations and clinical features in chronic schizophrenic patients in China

Xu JW, Guan XB, Wang XY, Feng Y, Zhang Q, Zhu JJ, Chen JH

Observational Study

- 533 Analysis of acupoint massage combined with touch on relieving anxiety and pain in patients with oral implant surgery

Qu JH, Shou CC, He X, Wang Q, Fang YX

- 541 Resilience provides mediating effect of resilience between fear of progression and sleep quality in patients with hematological malignancies

Tian Y, Wang YL

- 553 Nurse anesthetists' perceptions and experiences of managing emergence delirium: A qualitative study

Xin Y, Lin FC, Huang C, He B, Yan YL, Wang S, Zhang GM, Li R

Basic Study

- 563** Tanshinone IIA improves Alzheimer's disease *via* RNA nuclear-enriched abundant transcript 1/microRNA-291a-3p/member RAS oncogene family Rab22a axis
Yang LX, Luo M, Li SY

SYSTEMATIC REVIEWS

- 582** Outcomes of long-acting injectable antipsychotics use in pregnancy: A literature review
Pejčić AV, Stefanović SM, Milosavljević MN, Janjić VS, Folić MM, Folić ND, Milosavljević JZ

ABOUT COVER

Editorial Board Member of *World Journal of Psychiatry*, Sujita Kumar Kar, MD, Additional Professor, Department of Psychiatry, King George's Medical University, Lucknow-226003, UP, India. drsujita@gmail.com

AIMS AND SCOPE

The primary aim of *World Journal of Psychiatry (WJP, World J Psychiatry)* is to provide scholars and readers from various fields of psychiatry with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJP mainly publishes articles reporting research results and findings obtained in the field of psychiatry and covering a wide range of topics including adolescent psychiatry, biological psychiatry, child psychiatry, community psychiatry, ethnopsychology, psychoanalysis, psychosomatic medicine, etc.

INDEXING/ABSTRACTING

The *WJP* is now abstracted and indexed in Science Citation Index Expanded (SCIE, also known as SciSearch®), Current Contents/Clinical Medicine, Journal Citation Reports/Science Edition, PubMed, PubMed Central, Reference Citation Analysis, China Science and Technology Journal Database, and Superstar Journals Database. The 2023 Edition of Journal Citation Reports® cites the 2022 impact factor (IF) for *WJP* as 3.1; IF without journal self cites: 2.9; 5-year IF: 4.2; Journal Citation Indicator: 0.52; Ranking: 91 among 155 journals in psychiatry; and Quartile category: Q3.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Si Zhao; Production Department Director: Xu Guo; Cover Editor: Jia-Ping Yan.

NAME OF JOURNAL

World Journal of Psychiatry

ISSN

ISSN 2220-3206 (online)

LAUNCH DATE

December 31, 2011

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Ting-Shao Zhu

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2220-3206/editorialboard.htm>

PUBLICATION DATE

April 19, 2024

COPYRIGHT

© 2024 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Observational Study

Analysis of acupoint massage combined with touch on relieving anxiety and pain in patients with oral implant surgery

Jin-Hong Qu, Cheng-Cheng Shou, Xin He, Qin Wang, Yue-Xia Fang

Specialty type: Psychiatry

Provenance and peer review:

Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0

Grade B (Very good): 0

Grade C (Good): C

Grade D (Fair): 0

Grade E (Poor): 0

P-Reviewer: Kappelmann N, Germany

Received: December 19, 2023

Peer-review started: December 19, 2023

First decision: January 9, 2024

Revised: January 13, 2024

Accepted: March 7, 2024

Article in press: March 7, 2024

Published online: April 19, 2024



Jin-Hong Qu, Department of Stomatology, Hangzhou Lin'an Traditional Chinese Medicine Hospital, Hangzhou 311300, Zhejiang Province, China

Cheng-Cheng Shou, Department of Software R&D, Hangzhou Broadlink Technology Co, Ltd., Hangzhou 311300, Zhejiang Province, China

Xin He, Department of Stomatology, Hangzhou Red Cross Hospital, Hangzhou 310003, Zhejiang Province, China

Qin Wang, Department of Cardiovascular, Hangzhou Lin'an Traditional Chinese Medicine Hospital, Hangzhou 311300, Zhejiang Province, China

Yue-Xia Fang, Department of Traditional Chinese Medicine Development, Hangzhou Lin'an Traditional Chinese Medicine Hospital, Hangzhou 311300, Zhejiang Province, China

Corresponding author: Yue-Xia Fang, RN, Associate Chief Nurse, Department of Traditional Chinese Medicine Development, Hangzhou Lin'an Traditional Chinese Medicine Hospital, No. 288 Xishui North Road, Jinbei Street, Lin'an District, Hangzhou 311300, Zhejiang Province, China. 710125030@qq.com

Abstract

BACKGROUND

Oral implant surgery is an effective procedure for artificial implants in missing tooth areas under local anesthesia. Because patients under local anesthesia are conscious during this procedure, compared with general anesthesia-related operations, they are more likely to experience negative emotions, such as anxiety and tension. These emotional reactions result in shivering and chills in the limbs, leading to poor doctor-patient cooperation and even avoidance of treatment. In traditional Chinese medicine, it is believed that acupoint massage regulates blood and Qi, dredge menstruation, and relieve pain, which is beneficial for patients' emotional adjustment; however, there are few related clinical studies.

AIM

To observe the changes in anxiety and pain in patients with oral implant after acupoint massage combined with touch therapy.

METHODS

One hundred patients undergoing oral implantation in our hospital between May

2020 and May 2023 were randomly divided into control and study groups, according to a random number table, with 50 patients in each group. The control group received routine intervention, and the study group received acupoint massage combined with touch on the basis of the control group. Anxiety [assessed using the Modified Dental Anxiety Scale (MDAS)], pain severity, blood pressure, heart rate, and satisfaction were compared between the two groups.

RESULTS

Before intervention, the difference in MDAS score between the two groups was not significant ($P > 0.05$), while after the intervention, the MDAS scores decreased in both groups compared with those before the intervention ($P < 0.05$); the MDAS score of the study group was lower than that of the control group, with a statistically significant difference ($P < 0.05$). The degree of pain in the intervention group was significantly lower than that in the control group ($P < 0.05$). Before the intervention, there were no significant differences in systolic and diastolic blood pressures or heart rate between the two groups ($P > 0.05$). The systolic and diastolic blood pressures and heart rate in the intervention group, during and after the intervention, were significantly lower than those in the control group ($P < 0.05$). The total degree of satisfaction in the study group was significantly higher than that in the control group ($P < 0.05$).

CONCLUSION

Acupoint massage combined with touch better relieves anxiety and pain in patients undergoing dental implant surgery, improving the perioperative comfort of these patients and ensuring safety and a smooth operation.

Key Words: Oral implant; Acupoint massage; Touch; Anxiety; Degree of pain

©The Author(s) 2024. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Traditional Chinese medicine believes that acupoint massage can regulate blood and Qi, dredge menstruation, and relieve pain, all of which are beneficial for patients' emotional adjustment. However, patients undergoing oral implant surgery are in a state of local anesthesia and consciousness, and can feel tension and pain, which may lead to the interruption of treatment. Therefore, acupoint massage combined with touch may improve treatment efficacy.

Citation: Qu JH, Shou CC, He X, Wang Q, Fang YX. Analysis of acupoint massage combined with touch on relieving anxiety and pain in patients with oral implant surgery. *World J Psychiatry* 2024; 14(4): 533-540

URL: <https://www.wjgnet.com/2220-3206/full/v14/i4/533.htm>

DOI: <https://dx.doi.org/10.5498/wjp.v14.i4.533>

INTRODUCTION

Oral implants are a new technology developed jointly by bionics and tissue engineering, which integrates materials science, oral surgery, medical cosmetology, prosthodontics, and other multidisciplinary technologies, and have gradually become the primary treatment for oral diseases, such as missing anterior teeth and severe alveolar bone absorption[1-4]. Compared with traditional restorative approaches, oral implant technology has received favorable reviews from clinical doctors and patients in terms of aesthetics and oral function restoration[5-7]. Oral implants are effective in edentulous areas under local anesthesia. However, it should be noted that patients under local anesthesia are conscious. Compared to general anesthesia-related surgery, patients are more likely to experience negative emotions, such as anxiety and progression. These emotional reactions often lead to limb tremors and colds. Additionally, they are relatively sensitive to the operating environment and equipment, and many patients experience aggravated pain. To a certain extent, these phenomena affect doctor-patient cooperation and even prevent treatment, which is not conducive to smooth operation[8-12]. To relieve adverse psychological emotions and pain levels in patients undergoing oral implant therapy, in traditional Chinese medicine, it is believed that acupoint massage is beneficial to the emotional management of patients by regulating blood and Qi, dredging channels, and relieving pain[13,14]. Studies have shown that[15] acupoint massage has certain antidepressant and anxiety effects in patients with diabetic retinopathy, and the operation is simple and easy to use; Another study[16] confirmed that touch therapy combined with acupoint massage has a significant effect in reducing the pain of patients during percutaneous laser disc decompression and improving the medical experience of patients. Therefore, it is presumed that acupoint massage combined with touch can relieve anxiety and pain of patients undergoing oral implant surgery. Therefore, in this study, acupoint massage combined with touch therapy was performed on patients with oral implants to observe changes in the degree of anxiety and pain, with the aim of helping patients benefit from it.

MATERIALS AND METHODS

General data

Study participants were selected from 100 patients with oral implants admitted to our hospital between May 2020 and May 2023. The participants were divided into two groups, according to the random number table: 50 cases in the control group, including 15 males and 35 females (age was between 22–50 years, with an average of 35.25 ± 5.08 years). There were 50 cases in the study group, including 18 males and 32 females (age was between 20–58 years, with an average of 36.15 ± 5.15 years). There was no significant difference in age or sex between the two groups ($P > 0.05$).

Inclusion and exclusion criteria

Inclusion criteria: (1) All participants who received oral implant-related treatment for the first time; and (2) all the participants signed the consent form voluntarily.

Exclusion criteria: (1) Patients with abnormal coagulation function or those who took anticoagulant drugs recently; (2) patients with periodontal tissue lesions; (3) systemic diseases, such as liver and kidney dysfunction and osteoporosis; (4) medication history, including anti-inflammatory drugs, glucocorticoids, and immunomodulatory agents within one month before enrollment; (5) complications of infectious diseases and ineffective treatment; and (6) combined cachexia.

Method

Control group: The patients received conventional intervention: They were informed of the operating procedures, possible problems in the treatment, and corresponding solutions, and specific explanations were given for the questions raised by the patients. The patients communicated with each other during the operation interval, and anxiety and progressive emotions were relieved by verbal reassurance.

Study group: Acupoint massage combined with touch was performed in the study group on the basis of the control group, preparation work: Environment: The indoor temperature was controlled at 24–26 °C, and the humidity was controlled at about 70% (heating or air conditioning is required in winter and summer); Light songs or soft music can be played in the room to relax the patients and make them in a relaxed and quiet state. Before operation, the operator should wash his hands with warm water to ensure that the palm temperature is appropriate. Patients were placed in a comfortably in the supine position, as far as possible. One arm of the patient was held alternately and gently with both hands from top to bottom, and the upper arm to the wrist was massaged. During this process, the middle finger and thumb of one hand were placed at the Neiguan and Waiguan points, and the Hegu and Shenmen points were gently pressed by rotating the two fingers for approximately 3 min. We can also press the feet, massage the Sanyinjiao point on the ankle with emphasis, and adopt the manipulation of twisting, rotating, tamping and moving, 20 times per method. The opposite hand was massaged using the same technique for approximately 20 min. The patient's reaction, expression and pulse changes was observed during the massage, and a pressing force was adjusted if the patient experienced it, or hold patient's hand in time or tap the patient's left front upper limb shoulder, or press the patient's forehead outward from the center of the eyebrow, and ask patient to breathe deeply at the same time. During this process, the massage and touch areas should be far from the operation area so as not to affect the procedure of the operator.

Outcome indicators

Anxiety: The Modified Dental Anxiety Scale (MDAS)[17] was used to evaluate and compare the anxiety of the patients in the two groups before and after the intervention. The MDAS contains four items; each item is assigned according to a 1–5 grading system, with a score range is 4–20 points. The higher the score, the more serious the dental anxiety. When the total score is ≥ 13 points, the patient is assessed to also have dental anxiety.

Pain severity: Pain was assessed using the World Health Organization pain classification[18], and pain during dental implant surgery was compared between the two groups. The assessment was mainly based on the subjective feelings of the patients: No pain was rated as grade 0; mild pain: Slight pain sensation, but bearable; moderate pain: The pain sensation was slightly aggravated, but all of them were within the tolerance range of the patient; and severe pain: the pain was severe and beyond the patient's tolerance, and the patient started shouting.

Blood pressure and heart rate: Blood pressure and heart rate were measured and compared between the two groups using an upright sphygmomanometer before, during, and after the intervention. Radial arterial pulse was also recorded.

Satisfaction: A self-made scale was used to assess satisfaction of the patients (Cronbach's α coefficient was 0.89, and re-test validity was 0.88). The assessment content of the scale mainly included the development of clinical work, work suggestions, and work attitudes. The full score on both sides was 100 points, with a score of ≥ 90 points was very satisfactory, 70–89 points was generally satisfactory, and < 70 points was unsatisfactory.

Statistical analysis

Statistical analysis was performed using SPSS version 24.0. All measurement data were subjected to Shapiro-Wilk normality test, and the measurement data was represented by mean \pm SD in line with a normal distribution. An independent sample *t* test was used for intergroup comparisons, and analysis of variance was used for multi-time point comparisons. Count data were expressed as percentage using χ^2 test, and rank data were tested using rank-sum test, with the test level of $\alpha = 0.05$.

RESULTS

Comparison of anxiety (MDAS score)

Before the intervention, the difference in the MDAS anxiety scores between the two groups was not significant ($P > 0.05$). After the intervention, the MDAS scores of the two groups were lower than those before the intervention ($P < 0.05$). Moreover, the MDAS score of the intervention group was significantly lower than that of the control group ($P < 0.05$) (Table 1).

Comparison of the degree of pain

The degree of pain in the intervention group was significantly lower than that in the control group ($P < 0.05$) (Table 2).

Comparison of blood pressure and heart rate

Before the intervention, there were no significant differences in systolic blood pressure, diastolic blood pressure, or heart rate between the two groups ($P > 0.05$). Compared with the control group, the systolic blood pressure, diastolic blood pressure, and heart rate of patients in the study group, during and after the intervention, were significantly lower ($P < 0.05$) (Table 3).

Comparison of degree of satisfaction

The total degree of satisfaction in the intervention group was significantly higher than that in the control group ($P < 0.05$) (Table 4).

DISCUSSION

Anxiety, tension, and other emotions of patients undergoing oral implantations are triggered by many factors. For example, many patients undergoing oral implantations face unfamiliar medical specialties and treatment experiences due to lack of adequate cognition of dental treatment and understanding of related medical interventions. This triggers a series of negative emotions, such as tension and anxiety, leading to increased secretion of endogenous epinephrine and sympathetic nerve excitability. The final manifestations in these patients include increased blood pressure, heart rate, sweating, chest congestion, and palpitations. The patients were skeptical about the treatment or even evaded it, which affected the smooth development of oral treatment and oral health. Deteriorating oral health could adversely cause anxiety in patients, thereby forming a vicious circle[19-21]. Therefore, it is particularly important to implement effective measures to alleviate anxiety in patients undergoing oral implant surgery.

Touch is a non-drug intervention method. Regular, scientific, and orderly mild touch is performed on a patient's body with both hands. Combined with energy technology, touch is an auxiliary method for adjusting and balancing a patient's energy field, thereby affecting patient's physical, psychological, and emotional aspects through intentional guidance[22-24]. In traditional Chinese medicine, it is believed that the body and mind are integrated, internal organs correspond to five aspirations and five emotions of the body, and the running channels of Qi and blood are fundamentally meridians and collaterals. Acupoint massage, as an external therapy in traditional Chinese medicine, can achieve the purpose of harmonize Qi, blood, and the body and mind by stimulating specific acupoints to follow the meridians. Modern pharmacology indicates that acupoint massage can increase nervous system excitability, accelerate metabolism, and improve the immune system[25,26]. The results of this study showed that the degree of pain was lower in the experimental group than that in the control group. The systolic blood pressure, diastolic blood pressure, and heart rate of patients in the study group were lower during and after the intervention, and the overall satisfaction was high, suggesting that acupoint massage combined with touch was beneficial in relieving the degree of pain, improving anxiety, and improving satisfaction. The reasons for this findings may be as follows: (1) Touch can increase the production of natural morphine-like substances, such as dynorphin and enkephalin in the brain, which are transported to the whole body through the body conduction system, to help the body relieve pain and promote emotional excitement[27,28]; (2) touch can reduce the expression of norepinephrine and cortisol, which in turn, improves the stress levels of the body and relieves tension and anxiety[29,30]; and (3) Shenmen and Neiguan points are acupoints in the heart and pericardium meridians, which have the effects of tranquilizing and relieving anxiety. The large intestine meridian runs from the upper arm to the head, and the Hegu point is the source point of the large intestine meridian, and is of great significance in stimulating the healthy Qi of the body for resist stasis[31]. Touch and acupoint massages exhibit strong operability, good randomness, and repeatability. Additionally, acupoint massage uses hand-to-point instead of a needle, which can integrate the respective advantages of the two methods to achieve the effects of moving Qi and activating blood, dredging meridians and collaterals, relieving local pain in patients, and tranquilizing the mind[32,33]. The frontal lobe is the emotion generating center of the human brain, the left frontal lobe is mainly responsible for regulating and generating positive emotions, while the right frontal lobe is responsible for regulating and generating negative emotions. Changes in frontal lobe function are important factors for anxiety. Fan *et al*[34]'s study found that transcutaneous electrical acupoint stimulation of Neiguan can effectively relieve anxiety, with the most obvious changes in the left and right frontal lobes, suggesting that transcutaneous electrical acupoint stimulation of Neiguan can alleviate anxiety through promoting the brain functional activities of the frontal lobe, but it is different from the acupoint massage stimulation in this study. Whether the mechanism of improving anxiety and depression is consistent still needs further study. He *et al*[35] compared the anxiety and pain levels of 36 cases of oral implant patients receiving conventional nursing care, and 36 cases of oral implant patients receiving acupoint massage in combination with touch. Their results showed that the anxiety and pain

Table 1 Comparison of anxiety (Modified Dental Anxiety Scale score) between the two groups (mean \pm SD, point)

Group	Pre-intervention	Post-intervention	<i>t</i> value	<i>P</i> value
Control group (<i>n</i> = 50)	17.12 \pm 1.05	11.85 \pm 0.58	31.066	< 0.001
Study group (<i>n</i> = 50)	17.15 \pm 1.07	8.25 \pm 0.45	54.216	< 0.001
<i>t</i> value	0.142	34.676		
<i>P</i> value	0.888	< 0.001		

MDAS: Modified Dental Anxiety Scale.

Table 2 Comparison of pain severity between the two groups, *n* (%)

Group	Painless	Mild pain	Moderate pain	Severe pain
Control group (<i>n</i> = 50)	3 (6.00)	25 (50.00)	20 (40.00)	2 (4.00)
Study group (<i>n</i> = 50)	42 (84.00)	8 (16.00)	0	0
<i>Z</i> value	7.890			
<i>P</i> value	< 0.001			

Table 3 Comparison of blood pressure and heart rate between the two groups (mean \pm SD)

Group	Pre-intervention	Under intervention	Post-intervention	<i>F</i> value	<i>P</i> value
Systolic blood pressure (mmHg)					
Control group (<i>n</i> = 50)	130.12 \pm 10.20	138.25 \pm 15.52	128.25 \pm 10.75	9.208	< 0.001
Study group (<i>n</i> = 50)	130.25 \pm 10.22	120.12 \pm 8.25	115.25 \pm 6.85	40.027	< 0.001
<i>t</i> value	0.064	7.294	7.211		
<i>P</i> value	0.949	< 0.001	< 0.001		
Diastolic blood pressure (mmHg)					
Control group (<i>n</i> = 50)	85.25 \pm 5.15	90.25 \pm 8.15	88.25 \pm 8.25	5.900	< 0.001
Study group (<i>n</i> = 50)	85.23 \pm 5.12	75.12 \pm 5.50	72.12 \pm 4.50	92.253	< 0.001
<i>t</i> value	0.020	10.881	12.137		
<i>P</i> value	0.985	< 0.001	< 0.001		
Heart rate (beats/min)					
Control group (<i>n</i> = 50)	115.25 \pm 8.10	120.25 \pm 5.25	95.25 \pm 10.12	134.211	< 0.001
Study group (<i>n</i> = 50)	115.20 \pm 8.05	92.25 \pm 5.85	82.25 \pm 5.80	322.693	< 0.001
<i>t</i> value	0.031	25.189	7.881		
<i>P</i> value	0.975	< 0.001	< 0.001		

Table 4 Comparison of satisfaction between the two groups, *n* (%)

Group	Very satisfied	General satisfaction	Dissatisfied	Total satisfaction
Control group (<i>n</i> = 50)	12 (24.00)	32 (64.00)	6 (12.00)	44 (88.00)
Study group (<i>n</i> = 50)	48 (96.00)	2 (4.00)	0	50 (100.00)
χ^2 value				4.433
<i>P</i> value				0.035

levels of patients with oral implants receiving acupoint massage in combination with touch were significantly lower than those of patients receiving conventional nursing care. Additionally, their findings suggest that a combination of acupoint massage and touch could alleviate pain and anxiety of patients, which were consistent with the conclusions of this study. Liang *et al*[36] compared the efficacy of 50 cases of adult dental anxiety patients receiving conventional nursing care, and 50 cases of adult dental anxiety patients receiving psychological intervention combined with traditional Chinese medicine acupoint massage. Their results showed that the total effective rate of the treatment group (64.00%) was higher than that of the control group (22.00%), suggesting that psychological intervention combined with acupoint massage can alleviate the anxiety and depression of dental anxiety patients, which is also consistent with the conclusion of this study.

CONCLUSION

Acupoint massage combined with touch is conducive for alleviating anxiety and pain severity of patients undergoing oral implant surgery, thereby improving the perioperative comfort of these patients and ensuring a safe and smooth operation. However, it should still be noted that due to the small sample size and single center research, the credibility of the research conclusion needs to be verified by increasing the sample size and carrying out multi center research in the future.

ARTICLE HIGHLIGHTS

Research background

Oral implant surgery generally involves local anesthesia, and patients are in a conscious state. Under normal circumstances, patients are prone to negative emotions, such as nervousness, and may resist treatment and other uncooperative situations, which are not conducive for a smooth operation. In traditional Chinese medicine, it is believed that acupoint massage can regulate blood and Qi, dredge menstruation, and relieve pain, and its application in oral implant surgery may have certain effects in relieving pain.

Research motivation

At present, there are few clinical studies on the application of acupoint massage combined with touch in patients undergoing oral implant surgery; therefore, this study applied acupoint massage combined with touch in patients undergoing oral implant surgery to observe its influence on anxiety and pain.

Research objectives

This study aimed to observe the changes in anxiety and pain in patients with oral implants using acupoint massage combined with touch.

Research methods

One hundred patients undergoing oral implantation were randomly divided into control and study groups, according to a random number table, with 50 patients in each group. Anxiety [assessed using the Modified Dental Anxiety Scale (MDAS)], pain severity, blood pressure, heart rate, and satisfaction were compared between the two groups.

Research results

The MDAS score and degree of pain were lower in the study group than those in the control group. The systolic blood pressure, diastolic blood pressure, and heart rate of patients in the study group were lower during and after the intervention, and the overall satisfaction was high.

Research conclusions

Acupoint massage combined with touch is better in relieving anxiety of patients undergoing dental implant surgery, thereby relieving their pain and improving perioperative comfort.

Research perspectives

It should be noted that due to the small sample size and single center research, the credibility of the research conclusion needs to be verified by increasing the sample size and carrying out multi center research in the future.

FOOTNOTES

Co-first authors: Jin-Hong Qu and Qin Wang.

Author contributions: Qu JH and Wang Q contributed equally to this work as co-first authors; Qu JH, Shou CC, He X, Wang Q, and Fang YX designed the research study; Qu JH, Shou CC, He X, Wang Q, and Fang YX performed the research; Qu JH and Wang Q contributed new reagents and analytic tools; Qu JH and Wang Q analyzed the data and wrote the manuscript; and all authors have read and approve

the final manuscript.

Institutional review board statement: The study was reviewed and approved by the Hangzhou Lin'an Traditional Chinese Medicine Hospital Institutional Review Board (Approval No. lazyyIIK20180508001).

Informed consent statement: All study participants, or their legal guardian, provided informed written consent.

Conflict-of-interest statement: No conflicts of interest.

Data sharing statement: No additional data are available.

STROBE statement: The authors have read the STROBE Statement – checklist of items, and the manuscript was prepared and revised according to the STROBE Statement – checklist of items.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/Licenses/by-nc/4.0/>

Country/Territory of origin: China

ORCID number: Jin-Hong Qu 0009-0003-1931-4923; Cheng-Cheng Shou 0009-0006-9299-4488; Xin He 0009-0006-5306-3082; Qin Wang 0009-0007-5008-4176; Yue-Xia Fang 0009-0009-3734-4943.

S-Editor: Wang JL

L-Editor: A

P-Editor: Zhang YL

REFERENCES

- 1 Baas W, O'Connor B, Welliver C, Stahl PJ, Stember DS, Wilson SK, Köhler TS. Worldwide trends in penile implantation surgery: data from over 63,000 implants. *Transl Androl Urol* 2020; **9**: 31-37 [PMID: 32055463 DOI: 10.21037/tau.2019.09.26]
- 2 DeRose JJ Jr, Mancini DM, Chang HL, Argenziano M, Dagenais F, Ailawadi G, Perrault LP, Parides MK, Taddei-Peters WC, Mack MJ, Glower DD, Yerokun BA, Atluri P, Mullen JC, Puskas JD, O'Sullivan K, Sledz NM, Tremblay H, Moquete E, Ferket BS, Moskowitz AJ, Iribarne A, Gelijns AC, O'Gara PT, Blackstone EH, Gillinov AM; CTSN Investigators. Pacemaker Implantation After Mitral Valve Surgery With Atrial Fibrillation Ablation. *J Am Coll Cardiol* 2019; **73**: 2427-2435 [PMID: 31097163 DOI: 10.1016/j.jacc.2019.02.062]
- 3 Imamura T, Nnanabu J, Rodgers D, Raikhehlkar J, Kalantar S, Smith B, Nguyen A, Chung B, Narang N, Ota T, Song T, Burkhoff D, Jeevanandam V, Kim G, Sayer G, Uriel N. Hemodynamic Effects of Concomitant Mitral Valve Surgery and Left Ventricular Assist Device Implantation. *ASAIO J* 2020; **66**: 355-361 [PMID: 30985304 DOI: 10.1097/MAT.0000000000000999]
- 4 Briasoulis A, Yokoyama Y, Kuno T, Ueyama H, Shetty S, Alvarez P, Malik AH. In-Hospital Outcomes of Left Ventricular Assist Device Implantation and Concomitant Valvular Surgery. *Am J Cardiol* 2020; **132**: 87-92 [PMID: 32753267 DOI: 10.1016/j.amjcard.2020.06.067]
- 5 Jayaraj A, DeMartino RR, Bower TC, Oderich GS, Gloviczki P, Kalra M, Duncan AA, Fleming MD. Outcomes Following Inferior Mesenteric Artery Reimplantation During Elective Aortic Aneurysm Surgery. *Ann Vasc Surg* 2020; **66**: 65-69 [PMID: 31953141 DOI: 10.1016/j.avsg.2019.12.035]
- 6 Moskowitz G, Hong KN, Giustino G, Gillinov AM, Ailawadi G, DeRose JJ Jr, Iribarne A, Moskowitz AJ, Gelijns AC, Egorova NN. Incidence and Risk Factors for Permanent Pacemaker Implantation Following Mitral or Aortic Valve Surgery. *J Am Coll Cardiol* 2019; **74**: 2607-2620 [PMID: 31753204 DOI: 10.1016/j.jacc.2019.08.1064]
- 7 Ryu JI, Kim HY, Kwon YD. Is implant surgery a risk factor for osteonecrosis of the jaw in older adult patients with osteoporosis? A national cohort propensity score-matched study. *Clin Oral Implants Res* 2021; **32**: 437-447 [PMID: 33452837 DOI: 10.1111/clr.13713]
- 8 Lin CS, Wu SY, Yi CA. Association between Anxiety and Pain in Dental Treatment: A Systematic Review and Meta-analysis. *J Dent Res* 2017; **96**: 153-162 [PMID: 28106507 DOI: 10.1177/0022034516678168]
- 9 D'Amico G, Giannini F, Massucci M, Tebaldi M, Cafaro A, Ielasi A, Sgura F, De Marco F, Stefanini GG, Ciardetti M, Versaci F, Latini RA, Saccà S, Ghiringhelli S, Picchi A, Cerrito M, Gaspardone A, Tarantini G. Usefulness of Coronary Sinus Reducer Implantation for the Treatment of Chronic Refractory Angina Pectoris. *Am J Cardiol* 2021; **139**: 22-27 [PMID: 32998007 DOI: 10.1016/j.amjcard.2020.09.045]
- 10 Fonteyne E, Van Doorne L, Becue L, Matthys C, Bronckhorst E, De Bruyn H. Speech evaluation during maxillary mini-dental implant overdenture treatment: A prospective study. *J Oral Rehabil* 2019; **46**: 1151-1160 [PMID: 31271663 DOI: 10.1111/joor.12852]
- 11 Hilton TJ, Funkhouser E, Ferracane JL, Schultz-Robins M, Gordan VV, Bramblett BJ, Snead RM Jr, Manning W, Remakel JR; National Dental PBRN Collaborative Group. Recommended treatment of cracked teeth: Results from the National Dental Practice-Based Research Network. *J Prosthet Dent* 2020; **123**: 71-78 [PMID: 31202547 DOI: 10.1016/j.prosdent.2018.12.005]
- 12 Sturm JJ, Vicario-Quinones F, Shavit SS, Lalwani AK. Is Unilateral Cochlear Implantation Cost-Effective for the Treatment of Bilateral Sensorineural Hearing Loss? *Laryngoscope* 2021; **131**: 460-461 [PMID: 32339258 DOI: 10.1002/lary.28703]
- 13 Grupper A, Schwartz D, Baruch R, Schwartz IF, Nakache R, Goykhman Y, Katz P, Lebedinsky A, Nachmany I, Lubezky N, Aouizerate J, Shashar M, Katchman H. Kidney transplantation in patients with inflammatory bowel diseases (IBD): analysis of transplantation outcome and IBD activity. *Transpl Int* 2019; **32**: 730-738 [PMID: 30793376 DOI: 10.1111/tri.13415]
- 14 Su X, Yan B, Wang L, Lv J, Cheng H, Chen Y. Oral Anticoagulant Agents in Patients With Atrial Fibrillation and CKD: A Systematic Review and Pairwise Network Meta-analysis. *Am J Kidney Dis* 2021; **78**: 678-689.e1 [PMID: 33872690 DOI: 10.1053/j.ajkd.2021.02.328]

- 15 **Wu R**, Han Y, Li JF, Zhou XL, Feng AL. [Application of acupoint massage combined with decompression ball therapy on anxiety and depression in patients with diabetic retinopathy]. *Hushi Jinxu Zazhi* 2023; **38**: 1505-1509 [DOI: [10.16821/j.cnki.hsjx.2023.16.014](https://doi.org/10.16821/j.cnki.hsjx.2023.16.014)]
- 16 **Sun HJ**, Zhang KY. [Effect of acupoint massage combined with massage on pain relief in patients undergoing percutaneous laser disc decompression]. *Shanxi Yiyao Zazhi* 2012; **41**: 917-948
- 17 **Kamiya K**, Shimizu K, Igarashi A, Kitazawa Y, Kojima T, Nakamura T, Ichikawa K; Japan ICL Study Group. Piggyback implantable collamer lens implantation for the correction of residual refractive errors after cataract surgery: a multicenter study. *Acta Ophthalmol* 2019; **97**: e946-e947 [PMID: [30298628](https://pubmed.ncbi.nlm.nih.gov/30298628/) DOI: [10.1111/aos.13937](https://doi.org/10.1111/aos.13937)]
- 18 **Chen J**, Wang JL. The International Association for The Study of Pain (IASP) Revision and Systematic Classification of Chronic Pain for WHO ICD-11. *Zhongguo Tengtong Yixue Zazhi* 2019; **25**: 323-330 [DOI: [10.3969/j.issn.1006-9852.2019.05.002](https://doi.org/10.3969/j.issn.1006-9852.2019.05.002)]
- 19 **Jin JQ**, Cui HM, Han Y, Su S, Liu HW. Multifactor analysis of patients with oral sensory complaints in a case-control study. *Chin Med J (Engl)* 2020; **133**: 2822-2828 [PMID: [33273331](https://pubmed.ncbi.nlm.nih.gov/33273331/) DOI: [10.1097/CM9.0000000000001190](https://doi.org/10.1097/CM9.0000000000001190)]
- 20 **Dimitriou TD**, Verykoui E, Papatrantaflou J, Konsta A, Kazis D, Tsolaki M. Non-Pharmacological interventions for the anxiety in patients with dementia. A cross-over randomised controlled trial. *Behav Brain Res* 2020; **390**: 112617 [PMID: [32428636](https://pubmed.ncbi.nlm.nih.gov/32428636/) DOI: [10.1016/j.bbr.2020.112617](https://doi.org/10.1016/j.bbr.2020.112617)]
- 21 **Lee PL**, Tam KW, Yeh ML, Wu WW. Acupoint stimulation, massage therapy and expressive writing for breast cancer: A systematic review and meta-analysis of randomized controlled trials. *Complement Ther Med* 2016; **27**: 87-101 [PMID: [27515882](https://pubmed.ncbi.nlm.nih.gov/27515882/) DOI: [10.1016/j.ctim.2016.06.003](https://doi.org/10.1016/j.ctim.2016.06.003)]
- 22 **Zheng Q**, Wang R, Shi Y, Sun Q. Effects of acupoint massage combined with relaxation therapy on patients with postoperative fatigue syndrome after lumbar surgery. *Medicine (Baltimore)* 2021; **100**: e25849 [PMID: [34106628](https://pubmed.ncbi.nlm.nih.gov/34106628/) DOI: [10.1097/MD.00000000000025849](https://doi.org/10.1097/MD.00000000000025849)]
- 23 **Qu X**, Liu H, Yang Y, Liu L, Shen X, Liu S. The effects of laser stimulation at acupoint ST36 on anxiety-like behaviors and anterior cingulate cortex c-Fos expression in a rat post-traumatic stress disorder model. *Lasers Med Sci* 2021; **36**: 279-287 [PMID: [32333335](https://pubmed.ncbi.nlm.nih.gov/32333335/) DOI: [10.1007/s10103-020-03026-x](https://doi.org/10.1007/s10103-020-03026-x)]
- 24 **Rapaport MH**, Schettler PJ, Larson ER, Dunlop BW, Rakofsky JJ, Kinkead B. Six vs twelve weeks of Swedish massage therapy for generalized anxiety disorder: Preliminary findings. *Complement Ther Med* 2021; **56**: 102593 [PMID: [33197668](https://pubmed.ncbi.nlm.nih.gov/33197668/) DOI: [10.1016/j.ctim.2020.102593](https://doi.org/10.1016/j.ctim.2020.102593)]
- 25 **Havner RD**, Lapid MI, Dockter TJ, McCue SA, Stelpflug AJ, Bigelow ML, Robsahm MM, Elwood T, Strand JJ, Bauer BA, Cutshall SM, Sloan JA, Walton MP, Whitford KJ. Impact of Massage Therapy on the Quality of Life of Hospice Patients and Their Caregivers: A Pilot Study. *J Palliat Care* 2022; **37**: 41-47 [PMID: [33213233](https://pubmed.ncbi.nlm.nih.gov/33213233/) DOI: [10.1177/0825859720975991](https://doi.org/10.1177/0825859720975991)]
- 26 **Kim YJ**, Kim HR, Jung YH, Park YH, Seo SW. Effects of Electrical Automatic Massage on Cognition and Sleep Quality in Alzheimer's Disease Spectrum Patients: A Randomized Controlled Trial. *Yonsei Med J* 2021; **62**: 717-725 [PMID: [34296549](https://pubmed.ncbi.nlm.nih.gov/34296549/) DOI: [10.3349/ymj.2021.62.8.717](https://doi.org/10.3349/ymj.2021.62.8.717)]
- 27 **Hsu CH**, Chi CC, Chen PS, Wang SH, Tung TH, Wu SC. The effects of aromatherapy massage on improvement of anxiety among patients receiving palliative care: A systematic review of randomized controlled trials. *Medicine (Baltimore)* 2019; **98**: e14720 [PMID: [30817616](https://pubmed.ncbi.nlm.nih.gov/30817616/) DOI: [10.1097/MD.00000000000014720](https://doi.org/10.1097/MD.00000000000014720)]
- 28 **Sözen KK**, Karabulut N. Efficacy of Hand and Foot Massage in Anxiety and Pain Management Following Laparoscopic Cholecystectomy: A Controlled Randomized Study. *Surg Laparosc Endosc Percutan Tech* 2020; **30**: 111-116 [PMID: [31855924](https://pubmed.ncbi.nlm.nih.gov/31855924/) DOI: [10.1097/SLE.0000000000000738](https://doi.org/10.1097/SLE.0000000000000738)]
- 29 **Zhu K**, Shen Y, Zhu Y, Li L, You Y. Effect of acupoint therapies on prostatitis: A systematic review protocol. *Medicine (Baltimore)* 2020; **99**: e18967 [PMID: [32028405](https://pubmed.ncbi.nlm.nih.gov/32028405/) DOI: [10.1097/MD.00000000000018967](https://doi.org/10.1097/MD.00000000000018967)]
- 30 **Wu X**, Jiang W, Shi C, Qian H, Fan X, Zhou P. The clinical efficacy of acupoint sticking combined with massage to treatment functional dyspepsia: A protocol of systematic review and meta-analysis. *Medicine (Baltimore)* 2020; **99**: e23869 [PMID: [33371173](https://pubmed.ncbi.nlm.nih.gov/33371173/) DOI: [10.1097/MD.00000000000023869](https://doi.org/10.1097/MD.00000000000023869)]
- 31 **Maratos FA**, Duarte J, Barnes C, McEwan K, Sheffield D, Gilbert P. The physiological and emotional effects of touch: Assessing a hand-massage intervention with high self-critics. *Psychiatry Res* 2017; **250**: 221-227 [PMID: [28167436](https://pubmed.ncbi.nlm.nih.gov/28167436/) DOI: [10.1016/j.psychres.2017.01.066](https://doi.org/10.1016/j.psychres.2017.01.066)]
- 32 **Jamali Z**, Vatandoost M, Erfanparast L, Aminabadi NA, Shirazi S. The relationship between children's media habits and their anxiety and behaviour during dental treatment. *Acta Odontol Scand* 2018; **76**: 161-168 [PMID: [29105531](https://pubmed.ncbi.nlm.nih.gov/29105531/) DOI: [10.1080/00016357.2017.1396493](https://doi.org/10.1080/00016357.2017.1396493)]
- 33 **Pasyar N**, Rambod M, Kahkhaee FR. The Effect of Foot Massage on Pain Intensity and Anxiety in Patients Having Undergone a Tibial Shaft Fracture Surgery: A Randomized Clinical Trial. *J Orthop Trauma* 2018; **32**: e482-e486 [PMID: [30444801](https://pubmed.ncbi.nlm.nih.gov/30444801/) DOI: [10.1097/BOT.0000000000001320](https://doi.org/10.1097/BOT.0000000000001320)]
- 34 **Fan FJ**, Bai Y, Ji HF. [Explore the Influence of Transcutaneous Electrical Acupoint Stimulation Neiguan Point on Anxiety Based on Electroencephalogram Nonlinear Dynamic Analysis Method]. *Jiliang Xuebao* 2022; **43**: 133-139 [DOI: [10.3969/j.issn.1000-1158.2022.01.21](https://doi.org/10.3969/j.issn.1000-1158.2022.01.21)]
- 35 **He J**, Huang X, Lei Y, Yuan Q. [Effect of touch combined with acupoint massage on relieving anxiety and pain in patients during oral implant surgery]. *Zhongguo Shiyong Huli Zazhi* 2019; **35**: 1311-1314 [DOI: [10.3760/cma.j.issn.1672-7088.2019.17.008](https://doi.org/10.3760/cma.j.issn.1672-7088.2019.17.008)]
- 36 **Liang JB**, Liang QQ, Lv JB. [Effect of psychological intervention combined with traditional Chinese medicine acupoint massage in the treatment of dental anxiety]. *Zhongyiyao Daobao* 2013; **12**: 151-153



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: office@baishideng.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

