

World Journal of *Transplantation*

World J Transplant 2020 January 18; 10(1): 1-28



REVIEW

- 1 Therapeutics administered during *ex vivo* liver machine perfusion: An overview
Buchwald JE, Xu J, Bozorgzadeh A, Martins PN
- 15 Machine perfusion in abdominal organ transplantation: Current use in the Netherlands
Rijkse E, IJzermans JN, Minnee RC

ABOUT COVER

Editorial Board Member of *World Journal of Transplantation*, Maria Irene Bellini, FEBS, MD, PhD, Surgeon, Renal Transplant, Belfast Health and Social Care Trust, Belfast BT97AB, United Kingdom

AIMS AND SCOPE

The primary aim of *World Journal of Transplantation* (WJT, *World J Transplant*) is to provide scholars and readers from various fields of transplantation with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJT mainly publishes articles reporting research results obtained in the field of transplantation and covering a wide range of topics including bone transplantation, brain tissue transplantation, corneal transplantation, descemet stripping endothelial keratoplasty, fetal tissue transplantation, heart transplantation, kidney transplantation, liver transplantation, lung transplantation, pancreas transplantation, skin transplantation, etc..

INDEXING/ABSTRACTING

The WJT is now abstracted and indexed in PubMed, PubMed Central, China National Knowledge Infrastructure (CNKI), and Superstar Journals Database.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: Yan-Xia Xing, Production Department Director: Yun-Xiaojuan Wu, Editorial Office Director: Jia-Ping Yan.

NAME OF JOURNAL

World Journal of Transplantation

ISSN

ISSN 2220-3230 (online)

LAUNCH DATE

December 24, 2011

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Maurizio Salvadori, Sami Akbulut, Vassilios Papalois

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2220-3230/editorialboard.htm>

PUBLICATION DATE

January 18, 2020

COPYRIGHT

© 2021 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Therapeutics administered during *ex vivo* liver machine perfusion: An overview

Julianna E Buchwald, Jing Xu, Adel Bozorgzadeh, Paulo N Martins

ORCID number: Julianna E Buchwald 0000-0001-5893-471X; Jing Xu 0000-0001-9391-6146; Adel Bozorgzadeh 0000-0002-0790-9562; Paulo N Martins 0000-0001-9333-0233.

Author contributions: Buchwald JE, Xu J and Martins PN drafted the manuscript; Bozorgzadeh A reviewed the manuscript; all authors contributed to editing and approved the final manuscript version.

Conflict-of-interest statement: The authors have no conflict of interests to disclose.

Open-Access: This is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Specialty type: Transplantation

Julianna E Buchwald, Jing Xu, Adel Bozorgzadeh, Paulo N Martins, Division of Transplantation, Department of Surgery, University of Massachusetts Medical School, Worcester, MA 01655, United States

Corresponding author: Paulo N Martins, MD, PhD, Assistant Professor, Attending Doctor, Surgeon, Department of Surgery, Transplant Division, University of Massachusetts Medical School, 55 North Lake Avenue, Worcester, MA 01655, United States. paulo.martins@umassmemorial.org

Abstract

Although the use of extended criteria donors has increased the pool of available livers for transplant, it has also introduced the need to develop improved methods of protection against ischemia-reperfusion injury (IRI), as these "marginal" organs are particularly vulnerable to IRI during the process of procurement, preservation, surgery, and post-transplantation. In this review, we explore the current basic science research investigating therapeutics administered during *ex vivo* liver machine perfusion aimed at mitigating the effects of IRI in the liver transplantation process. These various categories of therapeutics are utilized during the perfusion process and include invoking the RNA interference pathway, utilizing defatting cocktails, and administering classes of agents such as vasodilators, anti-inflammatory drugs, human liver stem cell-derived extracellular vesicles, and δ -opioid agonists in order to reduce the damage of IRI. *Ex vivo* machine perfusion is an attractive alternative to static cold storage due to its ability to continuously perfuse the organ, effectively deliver substrates and oxygen required for cellular metabolism, therapeutically administer pharmacological or cytoprotective agents, and continuously monitor organ viability during perfusion. The use of administered therapeutics during machine liver perfusion has demonstrated promising results in basic science studies. While novel therapeutic approaches to combat IRI are being developed through basic science research, their use in clinical medicine and treatment in patients for liver transplantation has yet to be explored.

Key Words: Therapeutics; Liver transplantation; *Ex vivo* machine perfusion; Ischemia reperfusion injury; Organ preservation; Extended criteria donors

©The Author(s) 2020. Published by Baishideng Publishing Group Inc. All rights reserved.

Received: September 8, 2019

Peer-review started: September 8, 2019

First decision: October 14, 2019

Revised: October 26, 2019

Accepted: December 6, 2019

Article in press: December 6, 2019

Published online: January 18, 2020

P-Reviewer: Morimatsu H,
Soriano-Ursúa MA, Zhang ZX

S-Editor: Ma RY

L-Editor: A

P-Editor: Xing YX



Core tip: The use of extended criteria donors has increased the donor pool of available livers for transplant but has also introduced other hurdles in protecting these vulnerable organs against ischemia-reperfusion injury (IRI). Current basic science research is aimed at mitigating the effects of IRI during the transplantation process by administering therapeutics during *ex vivo* liver machine perfusion. Of interest include therapeutics aimed at invoking the RNA interference pathway, utilizing defatting cocktails, and administering classes of agents such as vasodilators and anti-inflammatory drugs to reduce the damage of IRI following liver procurement and transplantation for ultimate preservation of the organ.

Citation: Buchwald JE, Xu J, Bozorgzadeh A, Martins PN. Therapeutics administered during *ex vivo* liver machine perfusion: An overview. *World J Transplant* 2020; 10(1): 1-14

URL: <https://www.wjgnet.com/2220-3230/full/v10/i1/1.htm>

DOI: <https://dx.doi.org/10.5500/wjt.v10.i1.1>

INTRODUCTION

The overall increasing success of liver transplantation over the last several years has unfortunately introduced one of the most significant hurdles to date - longer waiting lists and increased mortality while on the waiting list. In an effort to combat the organ shortage, transplant centers have extended the criteria for donors often considered for transplantation. Common categories of extended criteria donors (ECDs) now being included in the context of the donor liver pool include donation after cardiac death (DCD), hepatic steatosis, donors of advanced age, organs that have experienced prolonged normothermic and cold storage, and donors with an increased infectious risk. The inclusion of ECD in the donor pool has increased access to previously deemed un-transplantable organs by 77% while reducing the mortality of those on the waitlist by over 50%^[1].

While inclusion of ECDs has positively impacted the pool of livers available for transplant, the new criteria has also highlighted the need for improved methods to ameliorate ischemia-reperfusion injury (IRI) in these less than optimal organs due to a weakened defense against ischemia-reperfusion injury during the transplantation process^[2]. Ischemia-reperfusion injury occurs when blood supply to an organ is inhibited and then later restored, with this process resulting in oxidative damage, cell death, and generation of reactive oxygen species (ROS)^[3]. The hepatic molecular pathways involved in IRI are complex with liver sinusoidal endothelial cells and hepatocytes as the initial targets for cell death as a result of ATP depletion. Neutrophils and macrophages then accumulate in the liver leading to ROS generation while hepatic stellate cells then become activated to aid in recovery, ultimately leading to fibrosis of the allograft^[4-6].

Targeting specific candidates implicated in hepatic IRI therefore becomes challenging due to the complex molecular pathways that become activated. Some of the activated pathways and molecules include the complement cascade, the innate immune response and toll-like receptors (TLRs), CD4 T lymphocytes, inflammatory cytokines propagating the post-inflammatory response, nuclear factor κ B (NF- κ B) leading to production of TNF- α , adhesion molecules, apoptotic pathway activation, and ROS production and release^[7,8]. As it will be discussed, basic science research focused on hepatic IRI has attempted to target many key mediators implicated in the IRI cascade. Most studies rely on using a combination of therapies that block multiple, perhaps redundant, reperfusion injury pathways in order to achieve a significant reduction in injury and overall improvement in graft function^[9].

There currently exists no established clinical therapies to avoid IRI, and the main method implemented to reduce IRI relies on limiting the cold preservation period and re-warming of the organ^[2]. Other methods of graft protection prior to transplantation include immunosuppressive agents and modulation of the immune response. Immunosuppressive therapies, particularly in the setting of kidney transplantation, has proven to be advantageous when the donor is treated prior to graft procurement or when the graft is directly treated during perfusion or cold storage^[10].

While static cold storage (SCS) remains the gold standard for liver preservation, *ex vivo* machine perfusion (MP) preservation of the liver is gaining attention from both

basic scientists and transplant surgeons alike. *Ex vivo* MP is an attractive alternative to SCS due to its ability to continuously perfuse the organ microcirculation, effectively deliver substrates and oxygen required for cellular metabolism, therapeutically administer pharmacological or cytoprotective agents, and continuously monitor organ viability during perfusion^[2].

CURRENT EX VIVO LIVER MP CLINICAL TRIALS

Several recent liver transplantation clinical studies demonstrate the logistical feasibility and safety of MP in the hospital setting. Ravikumar *et al*^[11] published the first Phase 1 normothermic perfusion trial of 20 patients who underwent NMP liver transplantations demonstrating decreased AST levels compared with controls during the first 7 d and a 95% one-year patient survival rate in the NMP group. Although there was no statistical difference in the primary outcome, this study reported the first use of NMP as logistically feasible and safe for use in the clinic^[11]. Selzner *et al*^[12] report the use of normothermic *ex vivo* liver perfusion in 10 human liver grafts using an albumin-based Steen solution with comparable outcomes to traditional SCS post-liver transplantation. In addition, Czigany *et al*^[13] report an ongoing open-label, phase 2 randomized controlled trial using HMP in liver transplantation from ECDs although primary and secondary endpoints and extent of IRI have not yet been published.

Another recent advancement in human liver MP involved the first randomized controlled trial of 220 liver transplantations performed by Nasralla *et al*^[14] at normothermic preservation conditions and demonstrated a 50% lower level of graft damage compared to the traditional cold static method of preservation in addition to 50% fewer discarded organs in the normothermic machine perfused group. This trial describes the novel expansion of normothermic human liver MP from experimental bench studies to introduction into clinical practice and demonstrates its benefit over the traditional cold method of preservation. While a larger study is needed to determine the impact of NMP on liver graft and patient survival, preliminary results indicate an exciting future for normothermic liver preservation.

Although the purpose of this review is not to highlight every current clinical trial involving *ex vivo* liver MP to date, it is worth noting that the number of both prospective and retrospective studies investigating the role of HMP *vs* SCS in human liver transplantation is increasing but still remains limited. These studies have been conducted in a variety of countries including the United States, Switzerland, the Netherlands, and the United Kingdom^[15]. The clinical use of *ex vivo* liver MP may be viewed as a limitation due to it being in its infancy as a standard therapy in liver transplantation. However, the basic science advances that will be highlighted in this review continue to expand the applications of *ex vivo* liver MP closer to its acceptance as a more reliable, efficacious method of organ preservation for liver transplantation.

BASIC SCIENCE EX VIVO LIVER MP THERAPEUTICS

In this paper, we review basic science advances made in the area of therapeutics administered specifically during *ex vivo* liver machine preservation transplantation models (Table 1) and their importance in extending the donor criteria for liver transplantation in a clinical setting. While therapeutics for this review were only considered in the context of the liver, administered therapeutics during MP of additional organs such as the lungs, heart, and kidneys are currently being explored in an effort to reduce IRI during transplantation and increase the available organs suitable for transplantation. Therefore, the impact of therapeutics administered during MP to mitigate IRI during and post-organ transplantation holds tremendous potential to increase the donor pools of many transplantable organs while simultaneously reducing the waiting time for those hoping to gain a second chance at life.

RNA interference and its therapeutic role in the liver during *ex vivo* MP

One of the most recent basic science advances in liver MP therapeutics includes utilization of the RNA interference (RNAi) pathway to silence specific genes implicated in IRI. RNAi selectively silences genes by the RNA-induced silencing complex (RISC) upon hybridization with the target mRNA, subsequently leading to degradation of the mRNA by Argonaute, an RNase H enzyme. If there are mismatches between the RNA complex and the target mRNA, silencing can occur at the post-

Table 1 Major categories of ex vivo machine perfusion therapeutics in liver transplantation

Category	Agents
RNAi pathway	siRNA
	Anti-sense oligonucleotide (Miravirsen)
Defatting cocktail	Variable
Vasodilators	Prostacyclin
	BQ123 and Verapamil
	Prostaglandin E1
Others	Anti-inflammatory agents
	Human liver stem cells extracellular vesicles
	δ-opioid agonist (Enkephalin)
	NLRP3 inflammasome inhibitor MCC950

RNAi: Ribonucleic acid interference; siRNA: Small interfering ribonucleic acid; NLRP3: Nucleotide-binding domain leucine-rich repeat containing family pyrin domain containing 3.

transcriptional level leading to translational repression or exonucleolytic degradation^[16].

The overall RNAi mechanism contains several unique regulatory RNA molecules including microRNA (miRNA), small interfering RNA (siRNA), and short-hairpin RNA (shRNA). There have been only a handful of reports of the RNAi pathway being implemented in the context of animal liver transplantation. For example, Li *et al*^[17] employed a rat liver transplantation model to study the effects of Fas siRNA on IRI. Hydrodynamic injection of 200 nmol/kg Fas siRNA transfection of the penile vein was performed 48 h before liver procurement. Measurements from the recipient rats demonstrated reduced ALT levels, decreased apoptotic index levels, and reduced Fas mRNA and protein levels 24 h after blood reperfusion.

Contreras *et al*^[18] delivered caspase-8 or caspase-3 siRNA in an *in vivo* C57BL/6 mouse model *via* the portal vein by high-volume injection 1 hour before induction of ischemia for 90 min. Results demonstrated a reduction in caspase-8 and caspase-3 gene expression of greater than 60% following siRNA injection. In addition, siRNA-treated mice showed improved survival for greater than 30 d when treated with caspase-8 siRNA (30%) and caspase-3 siRNA (50%) compared to controls where all of the mice died within five days after being subjected to total liver ischemia.

In addition, Wu *et al*^[19] targeted interleukin-1 receptor-associated kinase-4 using shRNA (IRAK-4-shRNA) in a rat liver transplantation model to prevent IRI. IRAK-4 is implicated in the downstream signaling pathways of lipopolysaccharide (LPS) activation of IRI in addition to its role in Toll-like receptor (TLR) and IL-1R mediated innate immune responses and was therefore selected as an ideal candidate for shRNA targeting to prevent IRI^[19-23]. In this study, rat liver grafts were perfused *via* the portal vein with a plasmid expressing IRAK-4-shRNA for 4 min during the cold ischemia time and then stored in University of Wisconsin (UW) perfusion solution for a period of 6 h prior to transplantation. Post-liver transplantation results indicated improved liver function, preserved tissue architecture, decreased IRAK-4 mRNA and protein levels, decreased NF-κB, TNFα, IL-6, and IL-1B levels over a period of 180 minutes post-reperfusion^[19].

While these previous studies demonstrated the utilization of the RNAi pathway to selectively target genes implicated in IRI by introducing siRNA or shRNA using hydrodynamic injection, our group most recently reported the first use of *ex vivo* liver MP as a method of siRNA delivery prior to rat liver transplantation^[24]. In this study, siRNA targeting the Fas receptor was added directly to the perfusion solution and MP of the liver was maintained at either hypothermic (4 °C) or normothermic (37 °C) conditions using a closed loop perfusion circuit^[24]. The Fas siRNA construct was conjugated to invivofectamine lipid nanoparticles and the siRNA-lipid complexes were perfused *via* the portal vein for 4 h. Confocal imaging studies revealed Fas siRNA distribution throughout the liver sinusoids and central veins in both the hypothermic and normothermic conditions^[24].

This study demonstrated for the first time siRNA uptake and distribution in the

liver following *ex vivo* MP at hypothermic and normothermic conditions. While future studies will examine the effects of siRNA uptake and delivery using MP in a rat transplant model, this report highlights the exciting use of RNAi in the context of *ex vivo* liver preservation.

In addition, our group utilized a similar *ex vivo* normothermic machine preservation system as aforementioned to silence the p53 tumor suppressor gene in a rat liver damage model. Rats were injected with p53 siRNA conjugated to invivofectamine prior to initiation of liver damage^[25]. To induce liver damage, the liver hilum was clamped for 15 minutes. Confocal microscopy studies revealed p53 siRNA uptake into the machine perfused liver with reduced levels of the inflammatory cytokines IL-1, IL-6, and TNF α compared with controls^[25].

The utilization of the RNAi pathway in the context of machine preservation of the liver is a new concept with promising therapeutic value. Although administering therapeutic siRNA to human donors before procurement is clinically feasible, the cost of therapy and potential for side effects increase^[26]. The therapeutic dosage of siRNA delivery to the donor liver could be efficiently delivered with little to no side effects to other organs as the siRNA therapeutic concentration would be based upon the weight of the donor liver rather than the weight of the whole donor at the time of procurement^[26]. In addition, preliminary studies demonstrate the uptake of siRNA in both hypothermic and normothermic temperatures in a rat model suggesting that perfusion temperatures have little to no effect on uptake while future studies will be aimed at addressing the effect of temperature on siRNA efficacy in silencing its target in a transplant model. Therefore, inclusion of siRNA in the MP solution for liver transplant models demonstrates promising and exciting results while also holding tremendous therapeutic value for future use in the operating room.

It is also worth noting that anti-sense oligonucleotide (ASO) therapies have also recently been used against hepatitis C virus (HCV) infection. Mir-122 is a known miRNA of the liver and is a target of miravirsin, which is currently in Phase 2 clinical trials for treatment of HCV^[27]. Miravirsin has been shown to nearly completely eradicate HCV presence in cell culture. In addition, miravirsin was shown to sequester miR-122 (a necessary agent for HCV infection) in pig liver without inducing harmful effects^[28]. Miravirsin is a recent exciting advancement in liver-targeted therapies as it could significantly increase the donor pool by allowing the transplantation of HCV⁺ livers.

Although the concept of gene silencing using RNAi in liver transplantation MP models is in its earliest stages of development and optimization, it holds exciting potential as a future therapeutic to combat the damage caused by IRI as a result of liver transplantation.

Defatting cocktails to reduce steatosis during MP of the liver

The increasing prevalence of obesity and metabolic syndrome defined as insulin resistance, hyperlipidemia, hypertension, and hyperglycemia, have contributed to potential donors developing hepatic steatosis. Hepatic steatosis is defined as having intrahepatic triacylglycerol (TAG) of at least 5% of the total liver weight or 5% of hepatocytes containing lipid vacuoles without a patient history of secondary contributing factors including viral infection, excess alcohol intake, or drug treatments^[29]. Reports indicate that 33% of the United States adult population has nonalcoholic hepatic steatosis^[30].

Potential donors with hepatic steatosis are now included in the ECD donor pool, however this criterion has implications in post-transplant outcomes. An analysis of the Scientific Registry of Transplant Recipients reported that liver allografts with greater than 30% macrovesicular steatosis were independently predictive of reduced 1-year graft survival^[31].

Furthermore, steatotic livers are particularly susceptible to IRI, increasing the risk of postoperative morbidity and mortality after liver surgeries and liver transplantations^[32]. Recent basic science evidence also provides support that steatosis exacerbates the effects of IRI. Chu *et al*^[33] demonstrated in a rat model of liver steatosis that the steatotic-IRI livers had elevated ALT levels, evidence of histological injury, and impaired mitochondrial complex-1 function after partial hepatic normothermic ischemia compared to the lean liver controls. Liss *et al*^[34] provided evidence in a murine model of hepatic IRI that steatosis increases plasma ALT, inflammatory cytokine levels such as TNF- α and IL-6, and necroptosis markers RIPK1, RIPK3, and MLKL compared with low-fat diet controls.

Gehrau *et al*^[35] explored the effect of IRI on immune response pathways in human graft biopsies classified based upon the degree of graft steatosis. The results showed that compared with non-steatotic control grafts, the steatotic grafts had significant

post-transplant innate immune response activation of IL-6, IL-8, and IL-10, macrophage production of nitric oxide (NO) and ROS, and neutrophil and leukocyte recruitment around the sites of hepatocyte lipid accumulation^[35]. Ramachandran *et al.*^[36] also demonstrated that NF κ B P65 is associated with the inflammatory pathway implicated in IRI and necrosis in rat steatotic liver transplantation. Multidrug donor preconditioning of steatotic rat liver grafts has also been reported to abolish the IRI inflammatory mechanism while preventing an increase in parenchymal cell death following cold storage and reperfusion^[37].

A systematic review examining the animal model experimental studies investigating hepatic steatosis and IRI found that livers with > 30% macrovesicular steatosis were associated with a lower graft and recipient survival rate as a result of the effects of IRI^[38].

More recently, liver MP has been investigated as a method of steatotic liver preservation and has shown promising results. Bessems *et al.*^[39] compared the traditional method of cold storage *vs* hypothermic MP for rat donor steatotic liver preservation. After 24 h of either hypothermic cold storage or MP, results demonstrated reduced levels of AST and LDH and increased bile production, ammonia clearance, urea production, and ATP levels after MP *vs* cold storage.

Vairetti *et al.*^[40] examined the effects of rat liver preservation using MP at 20 °C in steatotic livers compared to the SCS method of preservation. Results demonstrated that the adenosine triphosphate/adenosine diphosphate ratio and bile production were higher and oxidative stress and biliary enzymes were lower in the machine preservation-treated steatotic livers compared with the SCS method of preservation^[40]. Additionally, there was a 2-fold increase in TNF α levels and caspase-3 activity in the SCS steatotic livers compared with the machine perfused livers^[40]. These findings suggest that MP at 20 °C improves rat steatotic liver preservation compared with the SCS method.

Subnormothermic machine preservation has also been investigated in the context of macrosteatotic rat livers and has shown to reduce parenchymal ALT, mitochondrial glutamate dehydrogenase release while protecting against steatotic-induced sinusoidal microvascular alterations and preserving mitochondrial structure^[41]. Thus, implementation of defatting protocols seeks to efficiently decrease the proportion of macrosteatotic hepatocytes while ensuring viability and functionality in the remaining hepatocytes^[42].

In an effort to mitigate the detrimental effects of hepatic steatosis on transplantation outcomes, several animal studies have investigated the role of defatting protocols to reduce the intrahepatic TAG content prior to transplantation. These protocols may span a period of days to weeks and rely on a change in diet to alter the fat content in livers prior to transplantation.

The concept of defatting steatotic livers holds significant therapeutic and clinical potential as defatting in humans has shown to decrease steatosis. For example, one study investigated the implementation of a protein-rich (1000 kcal/d) diet, exercise (600 kcal/d), and the lipid-lower drug bezafibrate (400 mg/d) for 2-8 wk in 11 candidates for living-donor liver transplantation^[43]. Results demonstrated significantly improved body weight, BMI, and steatosis allowing for the transplantation of 7 of the treated liver grafts to recipients^[43]. Post-transplant tests demonstrated liver functioning with no significant differences in measured functional parameters^[43]. Additionally, a study of 120 consecutive living donors with non-alcoholic fatty liver disease of $\geq 30\%$ or an estimated donor-recipient weight ratio of < 0.8 demonstrated that following diet and exercise modifications leading to $\geq 10\%$ total cholesterol reduction and $\geq 5\%$ weight reduction, an improvement in steatosis of $\geq 20\%$ was seen in the 120 donors^[44].

The aforementioned studies demonstrate the feasibility of reducing fat content using established defatting protocols that are reliant on a diet change prior to liver transplantation, however they do not reflect a viable approach for liver grafts that are procured and intended for transplant, typically requiring a time frame of less than 12 h^[45,46]. While the use of defatting protocols to reduce steatosis in donor livers for transplantation has provided initial promising results for expanding the liver donor pool, further experimental animal studies investigating the use of *ex vivo* perfusion of donor livers to reduce steatosis remains limited.

Jamieson *et al.*^[47] demonstrated in a porcine model that agents involved in peroxisome proliferation for lipid export, visfatin to reduce triglyceride (TG) levels, and forskolin to stimulate oxidation of lipids and ketogenesis decreased hepatocyte TG levels by 31% in 48 h of normothermic MP. Periportal hepatocytes were "defatted" compared to hepatocytes near the perivenous region with an overall increase in bile production^[47]. Additionally, Nagrath *et al.*^[48] utilized a perfusate medium supplemented with defatting agents (forskolin, GW7647, hypericin, scoparone,

visfatin, and GW501516) on steatotic livers from obese Zucker rats. Following *ex vivo* normothermic perfusion for 180 min with the defatting perfusate medium, the TG content decreased by 65% and produced elevated bile levels compared to the control perfusion^[48].

Liu *et al*^[49] reported the use of subnormothermic (20 °C) MP supplemented with a defatting cocktail for 6 h in obese Zucker rats. Results demonstrated a significant increase in very low density lipoprotein (VLDL) and TG content in the perfusate in groups with and without the defatting cocktail. Additionally, the oxygen uptake rate, VLDL and TG secretion, and venous resistance were also similar in both groups^[49]. This study demonstrates the process of lipid export during subnormothermic MP.

In another study, the addition of carvedilol, a beta- and alpha-adrenergic blocking agent, used commonly in the setting of ischemic heart disorders and hypertension, to UW solution prevented rat hepatic injury associated with IRI in both steatotic and non-steatotic livers after 2 h of normothermic perfusion^[50-52].

As basic science methods aimed at reducing hepatic steatosis prior to transplantation are constantly being explored and optimized, clinical studies investigating the role of defatting cocktails in MP is relatively limited. Since the field of liver transplantation has only recently established MP as the superior preservation method compared to SCS, we expect more clinical studies focused on the addition of therapeutic agents to the perfusate, such as defatting cocktails, in the near future.

Boteon *et al*^[53] has recently utilized pharmacological agents in the setting of *ex situ* normothermic MP to enhance lipid metabolism in steatotic human donor livers discarded for transplantation. Using a previously published cocktail of drugs by Nagrath *et al*^[48] and supplemented with L-carnitine, Boteon *et al*^[53] added this defatting cocktail to the perfusate at normothermic conditions (37 °C). Results showed that within 6 h of NMP supplemented with the pharmacologic defatting cocktail, the steatotic livers had enhanced lipid metabolism with decreased TG content, decreased vascular resistance of the portal vein with increased flow, decreased lactate levels, and decreased tissue expression of markers implicated in IRI such as CD14 and CD11b and decreased cytokine profiles of TNF- α and IL1 β ^[53]. Most notably, following pharmacologic NMP treatment, all 5 treated livers were considered transplantable based upon viability criteria established by the authors^[48,53].

This particular study highlights the potential power of therapeutics administered during MP to salvage previously deemed untransplantable livers while addressing the liver donor pool shortage. To our knowledge, there exists no clinical trial to date using defatting protocols with NMP in the setting of human liver transplantation, as the pharmacological defatting agents used at the bench have not yet been approved for use in humans in a clinical setting^[54]. While Boteon *et al*^[55] report cytotoxicity results of the defatting cocktail in the setting of primary human hepatocytes on non-parenchymal liver cells, an exciting future for approved pharmacologic defatting therapeutics for human liver perfusion and transplantation surely exists.

Vasodilator administration during liver machine preservation

While IRI induces damage at the cellular level as highlighted, microvasculature is also disrupted. IRI insult affects endothelial cells by disrupting the normal barrier function, vascular tone, and expression of adhesion molecules^[56]. Nitric oxide is reduced during reperfusion and can therefore no longer control vasodilation during this period^[56]. Additionally, capillaries become occluded due to the activation of inflammatory and coagulation cascades^[56]. Therefore, several labs have examined therapeutic agents in MP to preserve the microvascular integrity of the liver.

In a study designed to address the susceptibility of uncontrolled non-heart-beating donors to warm IRI during liver transplantation, the authors devised a new rat model technique for liver grafts using short oxygenated warm *ex vivo* perfusion (SOWP) and prostaglandin E1 (PGE1)^[57,58]. PGE1 has been shown to have vasodilative and hepatoprotective effects such as reducing hepatocytic degeneration, central and portal ICAM-1 expression, central and sinusoidal VCAM-1 expression, central and portal P-selectin expression, and portal and sinusoidal E-selectin expression in the context of reperfusion^[57]. Results from the SOWP and PGE1 study showed increased total bile production during reperfusion to the same level as the heart-beating donor grafts^[58]. Additionally, PGE1 supplementation to the SOWP buffer decreased AST, ALT, and TNF α levels following 1 h of reperfusion^[58]. Necrosis and apoptosis were examined *via* histology and TUNEL staining and demonstrated significantly reduced levels following PGE1 treatment with SOWP^[58]. This study revealed that SOWP and PGE1 treatment before cold preservation improves the functioning of liver grafts following warm IRI^[58].

In a similar SOWP study, Maida *et al*^[59] performed rat liver transplantations

following a 6 h cold preservation period in order to determine the *in vivo* effects of SOWP supplemented with PGE1 in DCD rats. Results indicated that in the PGE1-treated SOWP groups, serum liver enzymes, cellular damage, and intercellular adhesion molecule 1 levels were significantly decreased compared to the control group^[59].

Nassar *et al*^[60] investigated the role of a prostacyclin analog, epoprostenol sodium, in a pig DCD model during NMP for 10 h after 60 min of warm ischemia time and demonstrated lower levels of AST, ALT, LDH, increased bile production, and preserved hepatic architecture compared with the control groups. Echeverri *et al*^[61] studied the safety and efficacy of BQ123 (endothelin1 antagonist), epoprostenol (prostacyclin analogue), and verapamil (calcium channel antagonist) in a pig transplantation model using normothermic *ex vivo* liver MP. Livers treated with BQ123 and verapamil demonstrated increased hepatic artery flow and reduced hepatocyte injury compared with controls^[61].

Addition of vasodilators to the perfusate solution during MP in animal transplantation models demonstrates significant therapeutic potential. While IRI incites damage to the microvasculature, vasodilators such as PGE1 impart protective effects on the vasculature such as preserving adhesion and selectin expression for maintenance of vascular integrity.

Bae *et al*^[62] have also explored the use of α -tocopherol with Vasosol in a DCD rodent model as an additive in the perfusate during HMP to reduce inflammatory and apoptotic markers implicated in reperfusion injury. Addition of α -tocopherol to the HMP solution reduced ALT levels during reperfusion and also reduced levels of inflammatory cytokines IL-6, TNF- α , and MCP-1, and caspase 3/7 as a result of reducing cytochrome C mRNA levels^[62].

Clinical trials investigating the role of vasodilator additives to machine perfusate also highlight the importance of therapeutics in the human liver transplantation setting. The first human clinical trial using HMP for liver transplantation was completed by Guarrera *et al*^[63] and compared HMP-preserved livers in 20 adults with a SCS matched group^[45,63,64]. A Vasosol solution was used as the HMP perfusate which included added antioxidants, metabolic substrates, and vasodilators including nitroglycerin and prostaglandin E1^[45,63]. Results indicated significantly reduced peak levels of AST, ALT, total bilirubin, and serum creatinine in the HMP preserved group. Molecular analysis from this clinical study revealed that HMP attenuated expression of inflammatory cytokines, oxidation markers, adhesion molecules and chemokines, and apoptosis and CD68 positive macrophages compared with the SCS group^[64]. The overall early graft dysfunction rates were 5% in the HMP groups compared with 25% in the control group, and the mean hospital stay was also shorter in the HMP group compared with the SCS group suggesting that HMP with vasodilator additives in liver transplantation is a safe and future method of liver perfusion and preservation^[45].

OTHER THERAPEUTICS IMPLEMENTED IN LIVER MP

Additional classes of therapeutics used in the setting of liver MP include anti-inflammatory agents, human liver stem cell-derived extracellular vesicles (HLSC-EV), and more recently, the δ -opioid agonist, enkephalin. While these therapeutics will not be discussed extensively in this review, they are summarized in Table 2 and are important to mention given their potential therapeutic role in the liver transplantation field.

Goldaracena *et al*^[9] reported improved liver function and lower inflammation in a pig transplantation model using anti-inflammatory agents (alprostadil, n-acetylcysteine, carbon monoxide, sevoflurane) added to perfusate in NMP. Rigo *et al*^[65] demonstrated, for the first time, reuptake of HLSC-EV in *ex vivo* rat liver perfusion. Treated livers demonstrated reduced necrosis and apoptosis, lower hypoxia-induced markers, and superior liver function (lower AST and LDH).

Recently, Beal *et al*^[66] demonstrated the efficacy of enkephalin, a δ -opioid agonist, to reduce oxidative stress in a rat *ex vivo* perfusion model. Treated livers had lower AST and malondialdehyde levels, in addition to higher ATP and glutathione levels in the perfusate. Yu *et al*^[67] also most recently reported that the selective NLRP3 inflammasome inhibitor mcc950 added to the perfusate of an HMP system and intravenously injected in a pig liver transplantation model demonstrated improved outcomes in DCD organs compared with controls.

Table 2 Other major therapeutic additives in models of ex vivo liver machine perfusion

Ref.	Ex vivo perfusion type	Therapeutic	Problem	Animal	Model	Sample size	Ex vivo perfusion time (h)	Outcomes
Goldaracena <i>et al</i> ^[9] , 2016	SNMP	Anti-inflammatory agents (Alprostadil, n-acetylcysteine, carbon monoxide, sevoflurane)	IRI	Pig	Transplantation	5	4	During EVLP: lower AST, TNF- α , IL-6 Lower HA levels, β -galactosidase and higher IL-10 (nonsignificant) After transplantation: Lower bilirubin, lower IL-6, lower cleaved caspase 3 staining, intact sinusoidal endothelial cell lining Lower AST, TNF- α , HA, ALP and higher IL-10 (nonsignificant)
Rigo <i>et al</i> ^[65] , 2018	NMP	HLSC-EV	IRI	Rat	EVLP	9	4	HLSC-EV uptake in treated livers Reduced necrosis and apoptosis on histology, lower Suzuki tissue injury score, lower apoptosis, lower AST and LDH, lower HIF-1 α & TGF- β 1 (hypoxia induced markers) NMP had low hematocrit to induce hypoxia
Beal <i>et al</i> ^[66] , 2019	NMP	Enkephalin (δ -opioid agonist)	IRI	Rat	EVLP	6	4	10 μ mol/L determined to be optimal concentration in an <i>in vitro</i> model: Lower ALT and MDA; better preservation of structural architecture; decreased caspase-3 expression; decreased TUNEL staining; decreased phosphorylation of p38 and JNK; increased expression of p-Akt, PI3K, p-Bad and Bcl-2
Yu <i>et al</i> ^[67] , 2019	HMP	NLRP3 Inflammasome Inhibitor mcc950	IRI	Pig	Transplantation	6	2	All reduced in HMP-postop group with added mcc950 in perfusate and IV administration of mcc950 After transplantation: TNF- α , IL-1 β , β -galactosidase, post-reperfusion serum ALT and AST, MDA, apoptosis staining, caspase-1 levels

SNMP: Subnormothermic machine perfusion; NMP: Normothermic machine perfusion; HMP: Hypothermic machine perfusion; IRI: Ischemia-reperfusion injury; EVLP: *Ex vivo* liver perfusion; AST: Aspartate aminotransferase; HA: Hyaluronic acid; ALP: Alkaline phosphatase; HLSC-EV: Human liver stem cells-derived extracellular vesicles; LDH: Lactate dehydrogenase; ALT: Alanine aminotransferase; MDA: Malondialdehyde; IV: Intravenous.

CONCLUSION

The use of administered therapeutics during machine liver perfusion has demonstrated promising results in basic science studies and initial clinical reports indicate their safety and efficacy in human liver transplantation. While novel therapeutic approaches to combat IRI are being developed through basic science research, their use in clinical medicine and treatment in patients for liver transplantation has yet to be fully explored. The number of human clinical trials investigating the use of liver MP is increasing, however the information obtained from these trials remains limited until additional robust studies are performed. A proposed summary of the advantages and limitations of the aforementioned MP therapeutics is offered in Table 3. While the therapies mentioned are relatively recent advances in the

Table 3 Summary of proposed advantages and limitations of liver machine perfusion therapeutics

Therapeutic	Advantages	Limitations/future considerations
RNAi pathway	Selectively targets and silences/degrades specific genes	Most beneficial/effective siRNA target against liver IRI in transplantation remains to be identified
	Mechanism of siRNA silencing pathway is generally understood	Potential for administration of multiple siRNA constructs each with a different target
	Organ-specific uptake	Requires design of siRNA against target mRNA and validation of target silencing
	Permits imaging studies visualizing tissue uptake and distribution	
Defatting cocktails	Ability to restore liver function by defatting	Steatotic livers are already predisposed to IRI ^[32]
	Aimed at enhancing natural lipid metabolism <i>via</i> lipid export, reduction of triglyceride levels, and stimulation of lipid oxidation and ketogenesis	Mechanisms of glucose and lipid control in liver remain poorly defined ^[69]
		Undefined consensus for quantifying degree of steatosis ^[70]
		Need for perfusate exchange protocol as secreted triglycerides recirculate causing further increase in lipid deposition ^[47]
Vasodilators	Focused on improving intrinsic function of liver to improve blood flow <i>via</i> smooth muscle relaxation and vasodilation	Kinetics of defatting may surpass average timeframe of liver transplantation ^[48]
	Act to increase arterial flow and decrease post-sinusoidal resistance ^[71]	Effects of vasodilators in marginal grafts remains unclear ^[61]
Anti-inflammatory agents	Some agents also act as vasodilators ^[68,72,73]	Combination of agents does not allow for specific identification of most beneficial agent(s)
	Act to scavenge free radicals to prevent IRI ^[74]	Mechanisms of anti-inflammatory agents remains unexplored in context of <i>ex vivo</i> liver perfusion ^[9]
	Ability to protect other cell types such as endothelial cells ^[75,76]	Combination of agents does not allow for determination of which specific agents were beneficial ^[9]
HLSC-EV	Regenerative and hepatoprotective properties ^[77,78]	Unknown mechanism of hypoxic protection ^[65]
	Diverse differentiating capabilities ^[77]	Timing of EV uptake during NMP currently unknown ^[65]
	Contain mRNA and miRNA subsets that modulate activity of target cells ^[79]	
	May serve as option for liver diseases without need for stem cells transplantation ^[65]	
δ-opioid agonist (Enkephalin)	Protects against oxidative stress ^[66]	Unknown therapeutic role in setting of post-perfusion liver transplant model with measured outcomes of graft function ^[66]
	Prevention of mitochondrial dysfunction <i>via</i> opioid receptor signaling ^[66]	
	Protection against IRI by slowing cellular metabolism ^[80,81]	Unknown role in cold ischemia conditions for liver transplant models ^[66]
NLRP3 inflammasome inhibitor (mcc950)	Blocks NLRP3-inflammasome activation preventing inflammatory liver damage ^[82,83]	mcc950 half-life is 3.27 h, while NLRP3 inflammasome activation lasts for several days after reperfusion ^[84,85]
	Reduces apoptosis post liver transplantation ^[67]	Additional mcc950 inhibition studies involving <i>in vitro</i> and <i>in vivo</i> models needed ^[67]

HLSC-EV: Human liver stem cells extracellular vesicles; NLRP3: Nucleotide-binding domain leucine-rich repeat containing family pyrin domain containing 3.

field of *ex vivo* liver MP, potential future directions are also included in Table 3 to indicate areas of further exploration. It is also worth noting that in terms of future considerations, the aforementioned studies in this review commonly employed multiple therapeutic agents in the perfusion solution, and several studies highlighted the need to delineate which agents out of the administered cocktail indeed conferred the most protective effect on the liver during *ex vivo* perfusion. MP therapeutics in liver transplantation therefore has tremendous potential to increase the liver donor pool and decrease the waiting time for lifesaving organs.

REFERENCES

- Renz JF**, Kin C, Kinkhabwala M, Jan D, Varadarajan R, Goldstein M, Brown R Jr, Emond JC. Utilization of extended donor criteria liver allografts maximizes donor use and patient access to liver transplantation. *Ann Surg* 2005; **242**: 556-563; discussion 563-565 [PMID: [16192816](#) DOI: [10.1097/01.sla.0000183973.49899.b1](#)]
- Zhai Y**, Petrowsky H, Hong JC, Busuttill RW, Kupiec-Weglinski JW. Ischaemia-reperfusion injury in liver transplantation--from bench to bedside. *Nat Rev Gastroenterol Hepatol* 2013; **10**: 79-89 [PMID: [23229329](#) DOI: [10.1038/nrgastro.2012.225](#)]
- Chouchani ET**, Pell VR, Gaude E, Aksentijević D, Sundier SY, Robb EL, Logan A, Nadtochiy SM, Ord ENJ, Smith AC, Eyassu F, Shirley R, Hu CH, Dare AJ, James AM, Rogatti S, Hartley RC, Eaton S, Costa ASH, Brookes PS, Davidson SM, Duchon MR, Saeb-Parsy K, Shattock MJ, Robinson AJ, Work LM, Frezza C, Krieg T, Murphy MP. Ischaemic accumulation of succinate controls reperfusion injury through mitochondrial ROS. *Nature* 2014; **515**: 431-435 [PMID: [25383517](#) DOI: [10.1038/nature13909](#)]
- Dar WA**, Sullivan E, Bynon JS, Eltzschig H, Ju C. Ischaemia reperfusion injury in liver transplantation: Cellular and molecular mechanisms. *Liver Int* 2019; **39**: 788-801 [PMID: [30843314](#) DOI: [10.1111/liv.14091](#)]
- Peralta C**, Jiménez-Castro MB, Gracia-Sancho J. Hepatic ischemia and reperfusion injury: effects on the liver sinusoidal milieu. *J Hepatol* 2013; **59**: 1094-1106 [PMID: [23811302](#) DOI: [10.1016/j.jhep.2013.06.017](#)]
- Stewart RK**, Dangi A, Huang C, Murase N, Kimura S, Stolz DB, Wilson GC, Lentsch AB, Gandhi CR. A novel mouse model of depletion of stellate cells clarifies their role in ischemia/reperfusion- and endotoxin-induced acute liver injury. *J Hepatol* 2014; **60**: 298-305 [PMID: [24060854](#) DOI: [10.1016/j.jhep.2013.09.013](#)]
- Luedde T**, Assmus U, Wüstefeld T, Meyer zu Vilsendorf A, Roskams T, Schmidt-Supprian M, Rajewsky K, Brenner DA, Manns MP, Pasparakis M, Trautwein C. Deletion of IKK2 in hepatocytes does not sensitize these cells to TNF-induced apoptosis but protects from ischemia/reperfusion injury. *J Clin Invest* 2005; **115**: 849-859 [PMID: [15776110](#) DOI: [10.1172/JCI23493](#)]
- Konishi T**, Lentsch AB. Hepatic Ischemia/Reperfusion: Mechanisms of Tissue Injury, Repair, and Regeneration. *Gene Expr* 2017; **17**: 277-287 [PMID: [28893351](#) DOI: [10.3727/105221617X15042750874156](#)]
- Goldaracena N**, Echeverri J, Spetzler VN, Kathis JM, Barbas AS, Louis KS, Adeyi OA, Grant DR, Selzner N, Selzner M. Anti-inflammatory signaling during ex vivo liver perfusion improves the preservation of pig liver grafts before transplantation. *Liver Transpl* 2016; **22**: 1573-1583 [PMID: [27556578](#) DOI: [10.1002/lt.24603](#)]
- Martins PN**, Chandraker A, Tullius SG. Modifying graft immunogenicity and immune response prior to transplantation: potential clinical applications of donor and graft treatment. *Transpl Int* 2006; **19**: 351-359 [PMID: [16623870](#) DOI: [10.1111/j.1432-2277.2006.00301.x](#)]
- Ravikumar R**, Jassem W, Mergental H, Heaton N, Mirza D, Perera MT, Quaglia A, Holroyd D, Vogel T, Coussios CC, Friend PJ. Liver Transplantation After Ex Vivo Normothermic Machine Preservation: A Phase I (First-in-Man) Clinical Trial. *Am J Transplant* 2016; **16**: 1779-1787 [PMID: [26752191](#) DOI: [10.1111/ajt.13708](#)]
- Selzner M**, Goldaracena N, Echeverri J, Kathis JM, Linares I, Selzner N, Serrick C, Marquez M, Sapisochin G, Renner EL, Bhat M, McGilvray ID, Lilly L, Greig PD, Tsien C, Cattral MS, Ghanekar A, Grant DR. Normothermic ex vivo liver perfusion using steen solution as perfusate for human liver transplantation: First North American results. *Liver Transpl* 2016; **22**: 1501-1508 [PMID: [27339754](#) DOI: [10.1002/lt.24499](#)]
- Czigany Z**, Schöning W, Ulmer TF, Bednarsch J, Amygdalos I, Cramer T, Rogiers X, Popescu I, Botea F, Froněk J, Kroy D, Koch A, Tacke F, Trautwein C, Tolba RH, Hein M, Koek GH, Dejong CHC, Neumann UP, Lurje G. Hypothermic oxygenated machine perfusion (HOPE) for orthotopic liver transplantation of human liver allografts from extended criteria donors (ECD) in donation after brain death (DBD): a prospective multicentre randomised controlled trial (HOPE ECD-DBD). *BMJ Open* 2017; **7**: e017558 [PMID: [29018070](#) DOI: [10.1136/bmjopen-2017-017558](#)]
- Nasralla D**, Coussios CC, Mergental H, Akhtar MZ, Butler AJ, Ceresa CDL, Chiochia V, Dutton SJ, García-Valdecasas JC, Heaton N, Imber C, Jassem W, Jochmans I, Karani J, Knight SR, Kocabayoglu P, Malagò M, Mirza D, Morris PJ, Pallan A, Paul A, Pavel M, Perera MTPR, Pirenne J, Ravikumar R, Russell L, Upponi S, Watson CJE, Weissenbacher A, Ploeg RJ, Friend PJ; Consortium for Organ Preservation in Europe. A randomized trial of normothermic preservation in liver transplantation. *Nature* 2018; **557**: 50-56 [PMID: [29670285](#) DOI: [10.1038/s41586-018-0047-9](#)]
- Zhang Y**, Zhang Y, Zhang M, Ma Z, Wu S. Hypothermic machine perfusion reduces the incidences of early allograft dysfunction and biliary complications and improves 1-year graft survival after human liver transplantation: A meta-analysis. *Medicine (Baltimore)* 2019; **98**: e16033 [PMID: [31169745](#) DOI: [10.1097/MD.00000000000016033](#)]
- Carthew RW**, Sontheimer EJ. Origins and Mechanisms of miRNAs and siRNAs. *Cell* 2009; **136**: 642-655 [PMID: [19239886](#) DOI: [10.1016/j.cell.2009.01.035](#)]
- Li X**, Zhang JF, Lu MQ, Yang Y, Xu C, Li H, Wang GS, Cai CJ, Chen GH. Alleviation of ischemia-reperfusion injury in rat liver transplantation by induction of small interference RNA targeting Fas. *Langenbecks Arch Surg* 2007; **392**: 345-351 [PMID: [17235585](#) DOI: [10.1007/s00423-006-0142-5](#)]

- 18 **Contreras JL**, Vilatoba M, Eckstein C, Bilbao G, Anthony Thompson J, Eckhoff DE. Caspase-8 and caspase-3 small interfering RNA decreases ischemia/reperfusion injury to the liver in mice. *Surgery* 2004; **136**: 390-400 [PMID: [15300206](#) DOI: [10.1016/j.surg.2004.05.015](#)]
- 19 **Wu Y**, Liu Y, Li M, Liu Z, Gong J. IRAK-4-shRNA Prevents Ischemia/Reperfusion Injury Via Different Perfusion Periods Through the Portal Vein After Liver Transplantation in Rat. *Transplant Proc* 2016; **48**: 2803-2808 [PMID: [27788821](#) DOI: [10.1016/j.transproceed.2016.06.058](#)]
- 20 **Kim TW**, Staschke K, Bulek K, Yao J, Peters K, Oh KH, Vandenburg Y, Xiao H, Qian W, Hamilton T, Min B, Sen G, Gilmour R, Li X. A critical role for IRAK4 kinase activity in Toll-like receptor-mediated innate immunity. *J Exp Med* 2007; **204**: 1025-1036 [PMID: [17470642](#) DOI: [10.1084/jem.20061825](#)]
- 21 **Bahia MS**, Kaur M, Silakari P, Silakari O. Interleukin-1 receptor associated kinase inhibitors: potential therapeutic agents for inflammatory- and immune-related disorders. *Cell Signal* 2015; **27**: 1039-1055 [PMID: [25728511](#) DOI: [10.1016/j.cellsig.2015.02.025](#)]
- 22 **Chen Y**, Liu Z, Liang S, Luan X, Long F, Chen J, Peng Y, Yan L, Gong J. Role of Kupffer cells in the induction of tolerance of orthotopic liver transplantation in rats. *Liver Transpl* 2008; **14**: 823-836 [PMID: [18508376](#) DOI: [10.1002/lt.21450](#)]
- 23 **Liu ZJ**, Yan LN, Li SW, You HB, Gong JP. Glycine blunts transplantative liver ischemia-reperfusion injury by downregulating interleukin 1 receptor associated kinase-4. *Acta Pharmacol Sin* 2006; **27**: 1479-1486 [PMID: [17049125](#) DOI: [10.1111/j.1745-7254.2006.00413.x](#)]
- 24 **Gillooly AR**, Perry J, Martins PN. First Report of siRNA Uptake (for RNA Interference) During Ex Vivo Hypothermic and Normothermic Liver Machine Perfusion. *Transplantation* 2019; **103**: e56-e57 [PMID: [30418428](#) DOI: [10.1097/TP.0000000000002515](#)]
- 25 **Moore C**, Thijssen M, Wang X, Mandrekar P, Xiaofei E, Abdi R, Porte R, Bozorgzadeh A, Leuvenink H, Kowalik T, Martins P. Gene Silencing with p53 si-RNA Downregulates Inflammatory Markers in the Liver: Potential Utilization during Normothermic Machine Preservation. *Am J Transplant* 2017; **17** Suppl 3
- 26 **Thijssen MF**, Brüggenwirth IMA, Gillooly A, Khvorova A, Kowalik TF, Martins PN. Gene Silencing With siRNA (RNA Interference): A New Therapeutic Option During Ex Vivo Machine Liver Perfusion Preservation. *Liver Transplantation* 2019; **25**: 140-151 [DOI: [10.1002/lt.25383](#)]
- 27 **Gebert LF**, Rebhan MA, Crivelli SE, Denzler R, Stoffel M, Hall J. Miravirsin (SPC3649) can inhibit the biogenesis of miR-122. *Nucleic Acids Res* 2014; **42**: 609-621 [PMID: [24068553](#) DOI: [10.1093/nar/gkt852](#)]
- 28 **Goldaracena N**, Spetzler VN, Echeverri J, Kath JM, Cherepanov V, Persson R, Hodges MR, Janssen HL, Selzner N, Grant DR, Feld JJ, Selzner M. Inducing Hepatitis C Virus Resistance After Pig Liver Transplantation-A Proof of Concept of Liver Graft Modification Using Warm Ex Vivo Perfusion. *Am J Transplant* 2017; **17**: 970-978 [PMID: [27805315](#) DOI: [10.1111/ajt.14100](#)]
- 29 **Nassir F**, Rector RS, Hammoud GM, Ibdah JA. Pathogenesis and Prevention of Hepatic Steatosis. *Gastroenterol Hepatol (NY)* 2015; **11**: 167-175 [PMID: [27099587](#)]
- 30 **Mehta SR**, Thomas EL, Bell JD, Johnston DG, Taylor-Robinson SD. Non-invasive means of measuring hepatic fat content. *World J Gastroenterol* 2008; **14**: 3476-3483 [PMID: [18567074](#) DOI: [10.3748/wjg.14.3476](#)]
- 31 **Spitzer AL**, Lao OB, Dick AA, Bakthavatsalam R, Halldorson JB, Yeh MM, Upton MP, Reyes JD, Perkins JD. The biopsied donor liver: incorporating macrosteatosis into high-risk donor assessment. *Liver Transpl* 2010; **16**: 874-884 [PMID: [20583086](#) DOI: [10.1002/lt.22085](#)]
- 32 **Tashiro H**, Kuroda S, Mikuriya Y, Ohdan H. Ischemia-reperfusion injury in patients with fatty liver and the clinical impact of steatotic liver on hepatic surgery. *Surg Today* 2014; **44**: 1611-1625 [PMID: [24078000](#) DOI: [10.1007/s00595-013-0736-9](#)]
- 33 **Chu MJ**, Premkumar R, Hickey AJ, Jiang Y, Delahunt B, Phillips AR, Bartlett AS. Steatotic livers are susceptible to normothermic ischemia-reperfusion injury from mitochondrial Complex-I dysfunction. *World J Gastroenterol* 2016; **22**: 4673-4684 [PMID: [27217699](#) DOI: [10.3748/wjg.v22.i19.4673](#)]
- 34 **Liss KHH**, McCommis KS, Chambers KT, Pietka TA, Schweitzer GG, Park SL, Nalbantoglu I, Weinheimer CJ, Hall AM, Finck BN. The impact of diet-induced hepatic steatosis in a murine model of hepatic ischemia/reperfusion injury. *Liver Transpl* 2018; **24**: 908-921 [PMID: [29729104](#) DOI: [10.1002/lt.25189](#)]
- 35 **Gehrau RC**, Mas VR, Dumur CI, Suh JL, Sharma AK, Cathro HP, Maluf DG. Donor Hepatic Steatosis Induce Exacerbated Ischemia-Reperfusion Injury Through Activation of Innate Immune Response Molecular Pathways. *Transplantation* 2015; **99**: 2523-2533 [PMID: [26285018](#) DOI: [10.1097/TP.0000000000000857](#)]
- 36 **Ramachandran S**, Liaw JM, Jia J, Glasgow SC, Liu W, Csontos K, Upadhyaya GA, Mohanakumar T, Chapman WC. Ischemia-reperfusion injury in rat steatotic liver is dependent on NFkB P65 activation. *Transpl Immunol* 2012; **26**: 201-206 [PMID: [22286145](#) DOI: [10.1016/j.trim.2012.01.001](#)]
- 37 **von Heesen M**, Seibert K, Hülser M, Scheuer C, Wagner M, Menger MD, Schilling MK, Moussavian MR. Multidrug donor preconditioning protects steatotic liver grafts against ischemia-reperfusion injury. *Am J Surg* 2012; **203**: 168-176 [PMID: [21782153](#) DOI: [10.1016/j.amjsurg.2011.01.026](#)]
- 38 **Chu MJ**, Hickey AJ, Phillips AR, Bartlett AS. The impact of hepatic steatosis on hepatic ischemia-reperfusion injury in experimental studies: a systematic review. *Biomed Res Int* 2013; **2013**: 192029 [PMID: [24062999](#) DOI: [10.1155/2013/192029](#)]
- 39 **Bessemis M**, Doorschodt BM, Kolkert JL, Vetelainen RL, van Vliet AK, Vreeling H, van Marle J, van

- Gulik TM. Preservation of steatotic livers: a comparison between cold storage and machine perfusion preservation. *Liver Transpl* 2007; **13**: 497-504 [PMID: [17394146](#) DOI: [10.1002/lt.21039](#)]
- 40 **Vairetti M**, Ferrigno A, Carlucci F, Tabucchi A, Rizzo V, Boncompagni E, Neri D, Gringeri E, Freitas I, Cillo U. Subnormothermic machine perfusion protects steatotic livers against preservation injury: a potential for donor pool increase? *Liver Transpl* 2009; **15**: 20-29 [PMID: [19109848](#) DOI: [10.1002/lt.21581](#)]
- 41 **Okamura Y**, Hata K, Tanaka H, Hirao H, Kubota T, Inamoto O, Kageyama S, Tamaki I, Yermek N, Yoshikawa J, Uemoto S. Impact of Subnormothermic Machine Perfusion Preservation in Severely Steatotic Rat Livers: A Detailed Assessment in an Isolated Setting. *Am J Transplant* 2017; **17**: 1204-1215 [PMID: [27860296](#) DOI: [10.1111/ajt.14110](#)]
- 42 **Nativ NI**, Maguire TJ, Yarmush G, Brasaemle DL, Henry SD, Guarrera JV, Berthiaume F, Yarmush ML. Liver defatting: an alternative approach to enable steatotic liver transplantation. *Am J Transplant* 2012; **12**: 3176-3183 [PMID: [23057797](#) DOI: [10.1111/j.1600-6143.2012.04288.x](#)]
- 43 **Nakamuta M**, Morizono S, Soejima Y, Yoshizumi T, Aishima S, Takasugi S, Yoshimitsu K, Enjoji M, Kotoh K, Taketomi A, Uchiyama H, Shimada M, Nawata H, Maehara Y. Short-term intensive treatment for donors with hepatic steatosis in living-donor liver transplantation. *Transplantation* 2005; **80**: 608-612 [PMID: [16177634](#) DOI: [10.1097/01.tp.0000166009.77444.f3](#)]
- 44 **Jin YJ**, Kim KM, Hwang S, Lee SG, Ha TY, Song GW, Jung DH, Kim KH, Yu E, Shim JH, Lim YS, Lee HC, Chung YH, Lee Y, Suh DJ. Exercise and diet modification in non-obese non-alcoholic fatty liver disease: analysis of biopsies of living liver donors. *J Gastroenterol Hepatol* 2012; **27**: 1341-1347 [PMID: [22554085](#) DOI: [10.1111/j.1440-1746.2012.07165.x](#)]
- 45 **Guarrera JV**, Henry SD, Samstein B, Odeh-Ramadan R, Kinkhabwala M, Goldstein MJ, Ratner LE, Renz JF, Lee HT, Brown RS Jr, Emond JC. Hypothermic machine preservation in human liver transplantation: the first clinical series. *Am J Transplant* 2010; **10**: 372-381 [PMID: [19958323](#) DOI: [10.1111/j.1600-6143.2009.02932.x](#)]
- 46 **Guarrera JV**, Henry SD, Chen SW, Brown T, Nachber E, Arrington B, Boykin J, Samstein B, Brown RS Jr, Emond JC, Lee HT. Hypothermic machine preservation attenuates ischemia/reperfusion markers after liver transplantation: preliminary results. *J Surg Res* 2011; **167**: e365-e373 [PMID: [20451921](#) DOI: [10.1016/j.jss.2010.01.038](#)]
- 47 **Jamieson RW**, Zilveti M, Roy D, Hughes D, Morovat A, Coussios CC, Friend PJ. Hepatic steatosis and normothermic perfusion-preliminary experiments in a porcine model. *Transplantation* 2011; **92**: 289-295 [PMID: [21681143](#) DOI: [10.1097/TP.0b013e318223d817](#)]
- 48 **Nagrath D**, Xu H, Tanimura Y, Zuo R, Berthiaume F, Avila M, Yarmush R, Yarmush ML. Metabolic preconditioning of donor organs: defatting fatty livers by normothermic perfusion ex vivo. *Metab Eng* 2009; **11**: 274-283 [PMID: [19508897](#) DOI: [10.1016/j.ymben.2009.05.005](#)]
- 49 **Liu Q**, Berendsen T, Izamis ML, Uygun B, Yarmush ML, Uygun K. Perfusion defatting at subnormothermic temperatures in steatotic rat livers. *Transplant Proc* 2013; **45**: 3209-3213 [PMID: [24182786](#) DOI: [10.1016/j.transproceed.2013.05.005](#)]
- 50 **Abshagen U**. A new molecule with vasodilating and beta-adrenoceptor blocking properties. *J Cardiovasc Pharmacol* 1987; **10** Suppl 11: S23-S32 [PMID: [2454364](#)]
- 51 **Prichard BN**, Tomlinson B. Progress in antihypertensive therapy with a multiple-action drug. *Drugs* 1988; **36** Suppl 6: 20-25 [PMID: [2908301](#) DOI: [10.2165/00003495-198800366-00005](#)]
- 52 **Ben Mosbah I**, Roselló-Catafau J, Alfany-Fernandez I, Rimola A, Parellada PP, Mitjavila MT, Lojek A, Ben Abdennebi H, Boillot O, Rodés J, Peralta C. Addition of carvedilol to University Wisconsin solution improves rat steatotic and nonsteatotic liver preservation. *Liver Transpl* 2010; **16**: 163-171 [PMID: [20104484](#) DOI: [10.1002/lt.21968](#)]
- 53 **Boteon YL**, Attard J, Boteon APCS, Wallace L, Reynolds G, Hubscher S, Mirza DF, Mergental H, Bhogal RH, Afford SC. Manipulation of Lipid Metabolism During Normothermic Machine Perfusion: Effect of Defatting Therapies on Donor Liver Functional Recovery. *Liver Transpl* 2019; **25**: 1007-1022 [PMID: [30821045](#) DOI: [10.1002/lt.25439](#)]
- 54 **Raigani S**, Markmann JF, Yeh H. Rehabilitation of Discarded Steatotic Livers Using Ex Situ Normothermic Machine Perfusion: A Future Source of Livers for Transplantation. *Liver Transpl* 2019; **25**: 991-992 [PMID: [31077626](#) DOI: [10.1002/lt.25490](#)]
- 55 **Boteon YL**, Wallace L, Boteon APCS, Mirza DF, Mergental H, Bhogal RH, Afford S. An effective protocol for pharmacological defatting of primary human hepatocytes which is non-toxic to cholangiocytes or intrahepatic endothelial cells. *PLoS One* 2018; **13**: e0201419 [PMID: [30044872](#) DOI: [10.1371/journal.pone.0201419](#)]
- 56 **Seal JB**, Gewertz BL. Vascular dysfunction in ischemia-reperfusion injury. *Ann Vasc Surg* 2005; **19**: 572-584 [PMID: [15981128](#) DOI: [10.1007/s10016-005-4616-7](#)]
- 57 **Hafez T**, Moussa M, Nesim I, Baligh N, Davidson B, Abdul-Hadi A. The effect of intraportal prostaglandin E1 on adhesion molecule expression, inflammatory modulator function, and histology in canine hepatic ischemia/reperfusion injury. *J Surg Res* 2007; **138**: 88-99 [PMID: [17174338](#) DOI: [10.1016/j.jss.2006.05.009](#)]
- 58 **Hara Y**, Akamatsu Y, Maida K, Kashiwadate T, Kobayashi Y, Ohuchi N, Satomi S. A new liver graft preparation method for uncontrolled non-heart-beating donors, combining short oxygenated warm perfusion and prostaglandin E1. *J Surg Res* 2013; **184**: 1134-1142 [PMID: [23688794](#) DOI: [10.1016/j.jss.2013.04.030](#)]
- 59 **Maida K**, Akamatsu Y, Hara Y, Tokodai K, Miyagi S, Kashiwadate T, Miyazawa K, Kawagishi N, Ohuchi N. Short Oxygenated Warm Perfusion With Prostaglandin E1 Administration Before Cold

- Preservation as a Novel Resuscitation Method for Liver Grafts From Donors After Cardiac Death in a Rat In Vivo Model. *Transplantation* 2016; **100**: 1052-1058 [PMID: 26950723 DOI: 10.1097/TP.0000000000001127]
- 60 **Nassar A**, Liu Q, Farias K, D'Amico G, Buccini L, Urcuyo D, Kelly D, Hashimoto K, Eghtesad B, Uso TD, Miller C, Quintini C. Role of vasodilation during normothermic machine perfusion of DCD porcine livers. *Int J Artif Organs* 2014; **37**: 165-172 [PMID: 24619899 DOI: 10.5301/ijao.5000297]
- 61 **Echeverri J**, Goldaracena N, Kathis JM, Linares I, Roizales R, Kollmann D, Hamar M, Urbanellis P, Ganesh S, Adeyi OA, Tazari M, Selzner M, Selzner N. Comparison of BQ123, Epoprostenol, and Verapamil as Vasodilators During Normothermic Ex Vivo Liver Machine Perfusion. *Transplantation* 2018; **102**: 601-608 [PMID: 29189484 DOI: 10.1097/TP.0000000000002021]
- 62 **Bae C**, Pichardo EM, Huang H, Henry SD, Guarrera JV. The benefits of hypothermic machine perfusion are enhanced with Vasosol and α -tocopherol in rodent donation after cardiac death livers. *Transplant Proc* 2014; **46**: 1560-1566 [PMID: 24880463 DOI: 10.1016/j.transproceed.2013.12.050]
- 63 **Guarrera JV**, Karim NA. Liver preservation: is there anything new yet? *Curr Opin Organ Transplant* 2008; **13**: 148-154 [PMID: 18685295 DOI: 10.1097/MOT.0b013e3282f63930]
- 64 **Henry SD**, Nachber E, Tulipan J, Stone J, Bae C, Reznik L, Kato T, Samstein B, Emond JC, Guarrera JV. Hypothermic machine preservation reduces molecular markers of ischemia/reperfusion injury in human liver transplantation. *Am J Transplant* 2012; **12**: 2477-2486 [PMID: 22594953 DOI: 10.1111/j.1600-6143.2012.04086.x]
- 65 **Rigo F**, De Stefano N, Navarro-Tableros V, David E, Rizza G, Catalano G, Gilbo N, Maione F, Gonella F, Roggio D, Martini S, Patrono D, Salizzoni M, Camussi G, Romagnoli R. Extracellular Vesicles from Human Liver Stem Cells Reduce Injury in an Ex Vivo Normothermic Hypoxic Rat Liver Perfusion Model. *Transplantation* 2018; **102**: e205-e210 [PMID: 29424767 DOI: 10.1097/TP.0000000000002123]
- 66 **Beal EW**, Kim JL, Reader BF, Akateh C, Maynard K, Washburn WK, Zweier JL, Whitson BA, Black SM. [D-Ala², D-Leu⁵] Enkephalin Improves Liver Preservation During Normothermic Ex Vivo Perfusion. *J Surg Res* 2019; **241**: 323-335 [PMID: 31071481 DOI: 10.1016/j.jss.2019.04.010]
- 67 **Yu Y**, Cheng Y, Pan Q, Zhang YJ, Jia DG, Liu YF. Effect of the Selective NLRP3 Inflammasome Inhibitor MCC950 on Transplantation Outcome in a Pig Liver Transplantation Model With Organs From Donors After Circulatory Death Preserved by Hypothermic Machine Perfusion. *Transplantation* 2019; **103**: 353-362 [PMID: 30247318 DOI: 10.1097/TP.0000000000002461]
- 68 **Lee LY**, Kaizu T, Toyokawa H, Zhang M, Ross M, Stolz DB, Huang C, Gandhi C, Geller DA, Murase N. Carbon monoxide induces hypothermia tolerance in Kupffer cells and attenuates liver ischemia/reperfusion injury in rats. *Liver Transpl* 2011; **17**: 1457-1466 [PMID: 21850691 DOI: 10.1002/lt.22415]
- 69 **Postic C**, Girard J. Contribution of de novo fatty acid synthesis to hepatic steatosis and insulin resistance: lessons from genetically engineered mice. *J Clin Invest* 2008; **118**: 829-838 [PMID: 18317565 DOI: 10.1172/JCI34275]
- 70 **Imber CJ**, St Peter SD, Handa A, Friend PJ. Hepatic steatosis and its relationship to transplantation. *Liver Transpl* 2002; **8**: 415-423 [PMID: 12004340 DOI: 10.1053/jlts.2002.32275]
- 71 **Nakai T**, Tanimura H, Hirokawa F, Tamaki T. Altered hepatic hemodynamics and improved liver function following intrahepatic vascular infusion of prostaglandin E1. *J Gastroenterol* 1998; **33**: 362-367 [PMID: 9658315 DOI: 10.1007/s005350050097]
- 72 **Vollmar B**, Menger MD. The hepatic microcirculation: mechanistic contributions and therapeutic targets in liver injury and repair. *Physiol Rev* 2009; **89**: 1269-1339 [PMID: 19789382 DOI: 10.1152/physrev.00027.2008]
- 73 **Tomiyama K**, Ikeda A, Ueki S, Nakao A, Stolz DB, Koike Y, Afrazi A, Gandhi C, Tokita D, Geller DA, Murase N. Inhibition of Kupffer cell-mediated early proinflammatory response with carbon monoxide in transplant-induced hepatic ischemia/reperfusion injury in rats. *Hepatology* 2008; **48**: 1608-1620 [PMID: 18972563 DOI: 10.1002/hep.22482]
- 74 **Winbladh A**, Björnsson B, Trulsson L, Bojmar L, Sundqvist T, Gullstrand P, Sandström P. N-acetyl cysteine improves glycogenesis after segmental liver ischemia and reperfusion injury in pigs. *Scand J Gastroenterol* 2012; **47**: 225-236 [PMID: 22242616 DOI: 10.3109/00365521.2011.643480]
- 75 **Annecke T**, Rehm M, Bruegger D, Kubitz JC, Kemming GI, Stoeckelhuber M, Becker BF, Conzen PF. Ischemia-reperfusion-induced unmeasured anion generation and glycocalyx shedding: sevoflurane versus propofol anesthesia. *J Invest Surg* 2012; **25**: 162-168 [PMID: 22583012 DOI: 10.3109/08941939.2011.618524]
- 76 **Knaak JM**, Spetzler VN, Goldaracena N, Boehnert MU, Bazerbach F, Louis KS, Adeyi OA, Minkovich L, Yip PM, Keshavjee S, Levy GA, Grant DR, Selzner N, Selzner M. Subnormothermic ex vivo liver perfusion reduces endothelial cell and bile duct injury after donation after cardiac death pig liver transplantation. *Liver Transpl* 2014; **20**: 1296-1305 [PMID: 25179693 DOI: 10.1002/lt.23986]
- 77 **Herrera MB**, Bruno S, Buttiglieri S, Tetta C, Gatti S, Deregibus MC, Bussolati B, Camussi G. Isolation and characterization of a stem cell population from adult human liver. *Stem Cells* 2006; **24**: 2840-2850 [PMID: 16945998 DOI: 10.1634/stemcells.2006-0114]
- 78 **Herrera MB**, Fonsato V, Bruno S, Grange C, Gilbo N, Romagnoli R, Tetta C, Camussi G. Human liver stem cells improve liver injury in a model of fulminant liver failure. *Hepatology* 2013; **57**: 311-319 [PMID: 22829291 DOI: 10.1002/hep.25986]
- 79 **Quesenberry PJ**, Aliotta J, Deregibus MC, Camussi G. Role of extracellular RNA-carrying vesicles

- in cell differentiation and reprogramming. *Stem Cell Res Ther* 2015; **6**: 153 [PMID: [26334526](#) DOI: [10.1186/s13287-015-0150-x](#)]
- 80 **Inuo H**, Eguchi S, Yanaga K, Hamada T, Yamanouchi K, Okudaira S, Kanematsu T. Protective effects of a hibernation-inducer on hepatocyte injury induced by hypothermic preservation. *J Hepatobiliary Pancreat Surg* 2007; **14**: 509-513 [PMID: [17909722](#) DOI: [10.1007/s00534-007-1214-9](#)]
- 81 **Yamanouchi K**, Yanaga K, Okudaira S, Eguchi S, Furui J, Kanematsu T. [D-Ala2, D-Leu5] enkephalin (DADLE) protects liver against ischemia-reperfusion injury in the rat. *J Surg Res* 2003; **114**: 72-77 [PMID: [13678701](#) DOI: [10.1016/s0022-4804\(03\)00196-3](#)]
- 82 **Coll RC**, Robertson AA, Chae JJ, Higgins SC, Muñoz-Planillo R, Inerra MC, Vetter I, Dungan LS, Monks BG, Stutz A, Croker DE, Butler MS, Haneklaus M, Sutton CE, Núñez G, Latz E, Kastner DL, Mills KH, Masters SL, Schroder K, Cooper MA, O'Neill LA. A small-molecule inhibitor of the NLRP3 inflammasome for the treatment of inflammatory diseases. *Nat Med* 2015; **21**: 248-255 [PMID: [25686105](#) DOI: [10.1038/nm.3806](#)]
- 83 **Mridha AR**, Wree A, Robertson AAB, Yeh MM, Johnson CD, Van Rooyen DM, Haczezyi F, Teoh NC, Savard C, Ioannou GN, Masters SL, Schroder K, Cooper MA, Feldstein AE, Farrell GC. NLRP3 inflammasome blockade reduces liver inflammation and fibrosis in experimental NASH in mice. *J Hepatol* 2017; **66**: 1037-1046 [PMID: [28167322](#) DOI: [10.1016/j.jhep.2017.01.022](#)]
- 84 **Liu H**, Lo CM, Yeung OWH, Li CX, Liu XB, Qi X, Ng KTP, Liu J, Ma YY, Lam YF, Lian Q, Chan SC, Man K. NLRP3 inflammasome induced liver graft injury through activation of telomere-independent RAPI/KC axis. *J Pathol* 2017; **242**: 284-296 [PMID: [28378341](#) DOI: [10.1002/path.4901](#)]
- 85 **Kim HY**, Kim SJ, Lee SM. Activation of NLRP3 and AIM2 inflammasomes in Kupffer cells in hepatic ischemia/reperfusion. *FEBS J* 2015; **282**: 259-270 [PMID: [25327779](#) DOI: [10.1111/febs.13123](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

Telephone: +1-925-3991568

E-mail: bpgoffice@wjgnet.com

Help Desk: <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

