

World Journal of *Transplantation*

World J Transplant 2022 February 18; 12(2): 21-26



LETTER TO THE EDITOR

- 21 Immunosuppressive regimens and outcomes of inflammatory bowel disease patients requiring kidney transplantation
Singh U, Singh B, Bellini MI
- 24 Assessment of advanced age candidates for liver transplantation warrants more caution
Parente A, Ronca V

ABOUT COVER

Associate Editor of *World Journal of Transplantation*, Sanem Guler Cimen, MD, MSc, Associate Professor, Director, Surgeon, Department of General Surgery, Health Sciences University, Ankara 6500, Turkey. s.cimen@dal.ca

AIMS AND SCOPE

The primary aim of *World Journal of Transplantation (WJT, World J Transplant)* is to provide scholars and readers from various fields of transplantation with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJT mainly publishes articles reporting research results obtained in the field of transplantation and covering a wide range of topics including bone transplantation, brain tissue transplantation, corneal transplantation, descemet stripping endothelial keratoplasty, fetal tissue transplantation, heart transplantation, kidney transplantation, liver transplantation, lung transplantation, pancreas transplantation, skin transplantation, etc..

INDEXING/ABSTRACTING

The *WJT* is now abstracted and indexed in PubMed, PubMed Central, Scopus, China National Knowledge Infrastructure (CNKI), and Superstar Journals Database.

RESPONSIBLE EDITORS FOR THIS ISSUE

Production Editor: *Lin-YuTong Wang*; Production Department Director: *Xu Guo*; Editorial Office Director: *Jia-Ping Yan*.

NAME OF JOURNAL

World Journal of Transplantation

ISSN

ISSN 2220-3230 (online)

LAUNCH DATE

December 24, 2011

FREQUENCY

Monthly

EDITORS-IN-CHIEF

Maurizio Salvadori, Sami Akbulut, Vassilios Papalois, Atul C Mehta

EDITORIAL BOARD MEMBERS

<https://www.wjgnet.com/2220-3230/editorialboard.htm>

PUBLICATION DATE

February 18, 2022

COPYRIGHT

© 2022 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION ETHICS

<https://www.wjgnet.com/bpg/GerInfo/288>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE


<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>



Immunosuppressive regimens and outcomes of inflammatory bowel disease patients requiring kidney transplantation

Urvashi Singh, Baljit Singh, Maria Irene Bellini

ORCID number: Urvashi Singh 0000-0003-2176-5750; Baljit Singh 0000-0001-9713-0247; Maria Irene Bellini 0000-0003-0730-4923.

Author contributions: Singh B and Bellini MI designed the study, provided guidance regarding inflammatory bowel disease and kidney transplantation respectively and proofread the manuscript before submission; Singh U performed literature review and wrote manuscript; all authors have read and approve the final manuscript.

Conflict-of-interest statement: There is no conflict of interest to be declared by any of the authors.

Country/Territory of origin: United Kingdom

Specialty type: Transplantation

Provenance and peer review: Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): B
Grade C (Good): C, C
Grade D (Fair): D
Grade E (Poor): 0

Urvashi Singh, Department of Surgery, Wythenshawe Hospital, Manchester University Hospitals NHS Foundation Trust, Manchester M23 9LT, United Kingdom

Baljit Singh, Department of Colorectal Surgery, University Hospitals Leicester, Leicester LE1 5WW, United Kingdom

Maria Irene Bellini, Department of Surgical Sciences, Sapienza University, Roma 00161, Italy

Corresponding author: Urvashi Singh, MBBS, MSc, Department of Surgery, Wythenshawe Hospital, Manchester University Hospitals NHS Foundation Trust, Southmoor Road, Manchester M23 9LT, United Kingdom. usingh01@qub.ac.uk

Abstract

Patients with inflammatory bowel disease (IBD) can develop extra-renal complications and as a result, suffer from end stage renal failure requiring kidney transplantation (KT). A brief review of available literature revealed that IBD patients undergoing KT have shorter overall survival rates compared to their controls. Literature reporting steroid regimens and survival outcomes specific to IBD and post kidney transplant are scarce and these studies have small sample sizes thus making it difficult to draw accurate conclusions. Further research is required in the form of a randomized controlled study to clarify the effect and mechanism of steroid immunosuppression on the prognosis of renal transplant recipients and explore new treatment schemes.

Key Words: Inflammatory bowel disease; Kidney transplantation; Steroids; Immunosuppression; Kidney failure; Ulcerative colitis; Crohn's disease

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Patients with inflammatory bowel disease (IBD) can develop extra-renal complications and as a result, suffer from end stage renal failure requiring kidney transplantation (KT). A brief review of available literature revealed that IBD patients undergoing KT have shorter overall survival rates compared to their controls. We highlight through our paper, previously reported survival outcomes and immunosuppressive regimens used in this cohort of patients through a brief literature review.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Received: October 5, 2021

Peer-review started: October 5, 2021

First decision: December 16, 2021

Revised: December 28, 2021

Accepted: January 20, 2022

Article in press: January 20, 2022

Published online: February 18, 2022

P-Reviewer: Gisbert JP, Velikova TV, Zhou Y

S-Editor: Wang JJ

L-Editor: A

P-Editor: Wang JJ



Citation: Singh U, Singh B, Bellini MI. Immunosuppressive regimens and outcomes of inflammatory bowel disease patients requiring kidney transplantation. *World J Transplant* 2022; 12(2): 21-23

URL: <https://www.wjgnet.com/2220-3230/full/v12/i2/21.htm>

DOI: <https://dx.doi.org/10.5500/wjt.v12.i2.21>

TO THE EDITOR

The recent systematic review published by Aref *et al*[1] titled 'Does steroid-free immunosuppression improve the outcome in kidney transplant recipients compared to conventional protocols?' provided thought provoking insight into the impact of steroid-free immunosuppression on the outcome of kidney transplant recipients. Further to the authors conclusions in their paper, we aim to highlight the effect of steroids and steroid free regimens on the outcomes of inflammatory bowel disease (IBD) patients who were kidney transplant receipts.

IBD is comprised of Crohn's disease (CD) and ulcerative colitis (UC). Patients with IBD can develop extra-intestinal manifestations. These include peripheral arthritis, oral aphthous ulcers, erythema nodosum, episcleritis, pyoderma gangrenosum, primary sclerosing cholangitis and uveitis[2]. Patients with IBD can develop renal manifestations of the disease, including nephrolithiasis, glomerulonephritis, tubulointerstitial nephritis, and secondary amyloidosis[3]. A study reported that the incidence of end stage renal disease (ESRD) in patients with CD was 5 times higher than cross matched controls[3]. Kidney injury can also result from dehydration, long term malnutrition and side effects of IBD medical therapy. These can all contribute to chronic kidney disease and eventually ESRD warranting kidney dialysis and transplantation[3].

Studies reporting IBD patients requiring kidney transplantation (KT) are scarce. However, existing literature discussing IBD, and post KT outcomes reports similar survival rates for IBD patients post transplantation. In a recent detailed study, in which 12 IBD patients (7 CD patients and 5 UC patients) underwent KT due to immunoglobulin A (IgA) nephropathy and polycystic kidney disease, the estimated survival of IBD patients was reported to be 80.8% *vs* 96.8% in patients without IBD ($P = 0.001$)[4]. Treatment with infliximab or a dalimumab resulted in stable disease or improvement in kidney transplant patients affected by mild to moderate IBD. Eleven out of 12 patients were on maintenance immunosuppression with low dose corticosteroids (5 mg prednisolone daily), calcineurin inhibitors (tacrolimus), and anti-metabolite (mycophenolic acid in nine and mycophenolate mofetil in two); the twelfth patient was kept on low-dose corticosteroids and tacrolimus only. IBD course remained stable in the whole transplant group, but resulted in an increased risk of mortality and hospitalization, due to a higher infection rate[4].

Data on immunosuppression and steroid regimens specific to IBD and the post-KT period remains poorly reported. In a study, six patients (5 CD patients and 1 UC patient) out of 1537 patients with IBD, underwent KT for kidney failure secondary to amyloidosis, IgA nephropathy, oxalate nephropathy, haemolytic uraemic syndrome, and chronic kidney failure of unknown origin[5]. Five of the six patients received steroid therapy after transplantation, yet specific immunosuppressive regimens are not reported pre and post transplantation. The study discusses the outcomes of IBD patients post liver transplantation together with KT, hence it is not possible to comment on the post KT alone. However, the study does report an 84% survival rate during a total follow up of 103.0 mo and median follow up of 33 mo after solid organ transplantation. One male patient also developed papillary renal cell carcinoma in the transplanted kidney in this study. No graft rejection was reported[5]. In a different prospective cohort study that followed 26 patients with IBD and systemic ascorbic acid (AA) amyloidosis between 1989 and 2010, an 83% survival rate 15 years post transplantation was reported. In this study all patients had renal dysfunction as result of AA amyloidosis[6]. However, only six patients required renal transplantation due to ESRD. Four patients had deceased donor transplants and two patients had live-related transplants. There were five functioning grafts at census 0.8, 3.2, 4.2, 20.1 and 24.6 years after transplantation. One graft failure was reported at 14.5 years after renal transplantation due to recurrence of amyloidosis and sustained chronic inflammatory activity. The study notes that patients were provided with steroid regimens however does not provide specific details about whether these regimens were supplemented with other immunosuppressants[6]. Specific reporting regarding immunosuppressive regimens is warranted for IBD patients before and after KT.

In conclusion, IBD is an immunomediated disease that is associated with kidney disease and can cause ESRD in patients. From the available literature, it is suggested that patients with IBD that undergo KT have shorter overall survival rates compared to their controls. Reported data is scarce and inconclusive due to the small patient cohort sizes. Further research is required in the form of a randomized controlled study to clarify the effect and mechanism of steroid immunosuppression on the prognosis of renal transplant recipients and explore new treatment schemes.

REFERENCES

- 1 **Aref A**, Sharma A, Halawa A. Does steroid-free immunosuppression improve the outcome in kidney transplant recipients compared to conventional protocols? *World J Transplant* 2021; **11**: 99-113 [PMID: [33954088](#) DOI: [10.5500/wjt.v11.i4.99](#)]
- 2 **Annese V**. A Review of Extraintestinal Manifestations and Complications of Inflammatory Bowel Disease. *Saudi J Med Med Sci* 2019; **7**: 66-73 [PMID: [31080385](#) DOI: [10.4103/sjmms.sjmms_81_18](#)]
- 3 **Park S**, Chun J, Han KD, Soh H, Choi K, Kim JH, Lee J, Lee C, Im JP, Kim JS. Increased end-stage renal disease risk in patients with inflammatory bowel disease: A nationwide population-based study. *World J Gastroenterol* 2018; **24**: 4798-4808 [PMID: [30479466](#) DOI: [10.3748/wjg.v24.i42.4798](#)]
- 4 **Grupper A**, Schwartz D, Baruch R, Schwartz IF, Nakache R, Goykhman Y, Katz P, Lebedinsky A, Nachmany I, Lubezky N, Aouizerate J, Shashar M, Katchman H. Kidney transplantation in patients with inflammatory bowel diseases (IBD): analysis of transplantation outcome and IBD activity. *Transpl Int* 2019; **32**: 730-738 [PMID: [30793376](#) DOI: [10.1111/tri.13415](#)]
- 5 **Schnitzler F**, Friedrich M, Stallhofer J, Schönermarck U, Fischereder M, Habicht A, Karbalai N, Wolf C, Angelberger M, Olszak T, Beigel F, Tillack C, Göke B, Zachoval R, Denk G, Guba M, Rust C, Grüner N, Brand S. Solid Organ Transplantation in Patients with Inflammatory Bowel Diseases (IBD): Analysis of Transplantation Outcome and IBD Activity in a Large Single Center Cohort. *PLoS One* 2015; **10**: e0135807 [PMID: [26288187](#) DOI: [10.1371/journal.pone.0135807](#)]
- 6 **Sattianayagam PT**, Gillmore JD, Pinney JH, Gibbs SD, Wechalekar AD, Gilbertson JA, Rowczenio D, Hawkins PN, Lachmann HJ. Inflammatory bowel disease and systemic AA amyloidosis. *Dig Dis Sci* 2013; **58**: 1689-1697 [PMID: [23371008](#) DOI: [10.1007/s10620-012-2549-x](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

