

69951_Auto_Edited-zyl.docx

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 69951

Manuscript Type: CASE REPORT

1
Migration of a Hem-o-Lok Clip to the renal pelvis after laparoscopic partial nephrectomy(LPN) : a case report

A case of foreign body stones surgery in kidney

Ji Sun, Liwei Zhao, Xuliang Wang, Jiaguo Huang, Yi Fan

Abstract

BACKGROUND

Hem-o-Lok clip (HOLC) has been widely used in laparoscopic surgery due to their ease of application and secure clamping, though the rare complications associated with this technique should not be ignored. This rare complication of laparoscopic partial nephrectomy (LPN) consists of the clip migrating into the renal pelvis and acting as a nidus for stone formation.

CASE SUMMARY

The case described here involved a 63-year-old female who was found with stones in the right kidney and upper ureter during a recent reexamination following laparoscopic partial nephrectomy. We performed percutaneous nephrolithotomy (PCNL) for her, but during the operation, it was found that the center of the stone within the kidney was a HOLC, which was removed with forceps. For this reason, we speculate that the HOLC, which was employed to halt tumor wound bleeding, spontaneously drifted into the renal pelvis and formed kidney stones, with the clip being initially misdiagnosed as a kidney stone.

CONCLUSION

By reviewing related case reports, we concluded that in order to prevent complications related to HOLC, loose clips should be actively searched for and retrieved from the wound during urinary tract surgery, while the deployment of clips in close proximity of anastomotic stoma of collecting systems should be avoided.

Key Words: Partial nephrectomy; Foreign body stones; Hem-o-Lok (surgical instruments); Case report

Sun J, Zhao L, Wang X, Huang J, Fan Y. Migration of a Hem-o-Lok Clip to the renal pelvis after laparoscopic partial nephrectomy(LPN) : a case report . *World J Clin Cases* 2021; In press

Core Tip: The present case involves a patient who was found to have spontaneously drifted Hem-o-Lok into the renal pelvis and formed stones after laparoscopic partial nephrectomy. We concluded that in order to prevent complications related to Hem-o-Lok, loose clips should be actively sought and retrieved from the wound during urinary system surgery, with the deployment of clips in close proximity to the anastomotic stoma of collecting systems should be avoided.

INTRODUCTION

The nonabsorbable Hem-o-lok polymer ligating clip (HOLC) was introduced in 1999 and brought a novel and efficient ligation method for surgery, gradually replacing titanium clips and endovascular stapling devices, and is favored by surgeons^[1]. With its unique locking mechanism, this kind of polymer clip potentially reduces the risk of displacement from tissue. In addition, its narrow profile, good tactile feedback with closure, ability to remove the clip, and no artifact production on imaging techniques also render it increasingly widely employed, particularly within various laparoscopic surgical procedures. Within the urinary system, HOLC is commonly used for ligating the ureter, renal artery and renal vein, and halting kidney or prostate excision bed bleeding, and thus plays an important role in the popularization of laparoscopy in urology. Through a urologist's perspective, HOLC has unique advantages in avoiding thermal injury during dissection of the neurovascular bundle in LPN, laparoscopic radical prostatectomy (LRP) and even robot-assisted laparoscopic radical prostatectomy (RALP).

However, a central issue remains regarding HOLC stability. Although such novel devices for tissue and blood vessel occlusions are widely accepted, few data are available to support their efficacy and safety. Bleeding incidents caused by HOLC-

induced damage during operation or emergency shedding post-operation have been reported^[2]. However, it is rare to report that foreign bodies are formed due to the long-term displacement of clips post-operatively, which leads to related complications. This report presents a case of spontaneous migration of a surgical clip into the collecting system after LPN, which led to the occurrence of renal stones and misdiagnosis prior to surgery.

10

CASE PRESENTATION

Chief complaints

A 63-year-old female was admitted to our service due to urinary ultrasound revealing right renal and ureteral calculi.

History of present illness

In the past four years, the patient did not have any urinary tract symptoms. Until 48 h prior to admission, she had been examined with urinary ultrasound for follow-up of the renal cell carcinoma and found that the right kidney stone and upper ureteral stone was found in the B-ultrasound. She had no fever, no lower back pain, no urinary tract irritation, no gross hematuria or any other uncomfortable symptoms.

History of past illness

Four years ago, LPN for a minute right renal mass was performed in our hospital. During surgery, a mass of 2.5 cm x 2.3 cm in the lower pole of the right kidney was identified. The renal tumor was completely removed along the edge of the renal tumor (0.5 cm margin), and the capsule was intact. Post-resection, a fissure was found at the ureteropelvic junction. Therefore, a 3-0 Vicryl was applied to repair the fissure, and a 2-0 Vicryl was applied to the edge of the renal parenchyma. Furthermore, surgical bed hemostasis had been achieved by automatic non-absorbing surgical clips. The post-operative course was un-eventful, and pathology confirmed this to be renal cell

carcinoma of the right kidney (T1a / N0 / M0). Consequently, the patient was discharged from hospital and instructed to have regular re-examinations.

12

Personal and family history

There is no family history of urinary calculi and tumors.

Physical examination

Physical examination showed positive percussion pain of the right kidney and no other positive signs.

Laboratory examinations

No obvious abnormalities were found within hematological indicators and routine urinalyses for this particular patient.

Imaging examinations

B-ultrasound demonstrated that the right kidney collecting system was separated and a plurality of liquid-segregated dark areas could be visualized, with a strong echo of 1.1 × 0.8 cm within, and the upper part of the right kidney ureter was dilated, with a strong echo of 2.1 × 1.1cm within. Physical examination revealed right costovertebral angle tenderness, and the rest had no positive signs. The abdominal computed tomography (CT) scan demonstrated that the lower pole of the right kidney was absent post-surgery, with a high-density fringe, and two opacities could be seen in the right renal pelvis (A:1295HU) and proximal ureter (B:1335HU), respectively (Figure 1). A preliminary diagnosis of right hydronephrosis with nephroureteral calculi was concluded.

FINAL DIAGNOSIS

The final revised diagnosis was a right renal foreign body with calculus formation and right hydronephrosis with ureteral calculus.

TREATMENT

Consequently, we decided to perform percutaneous nephrolithotomy for this patient. During surgery, there were stones in the lower and middle-right renal calices and the upper part of the right ureter, which were both approximately 1.5cm x 1.5cm in size. Following an initial renal pelvic stone fragmentation with the ultrasound device progression stopped, it was found that the center of the stones was a white rectangular foreign body, surprisingly (Figure 2A). This was consequently removed with a forceps (Figure 2B). After careful *in vitro* examination, it was confirmed as a medium-sized surgical clip (Figure 2C).

OUTCOME AND FOLLOW-UP

Post-operatively, the patient had urinary tract infection, which improved following antibiotic treatment, and she was eventually discharged from hospital. Presently, the patient is followed up regularly, with no obvious abnormality found in the collecting system of both kidneys.

DISCUSSION

Intrarenal foreign bodies are extremely rare-expected circumstances, and gastrointestinal tract is one of the most common routes for reaching the kidney. Such foreign bodies are typically torn objects in food, such as fish bones, needles, pins, hairgrips and toothpicks^[3]. Such clinical manifestations also vary, including purulent cutaneous fistulae, renal and perirenal pseudotumors, and the formation of kidney stones^[4-7]. These foreign bodies can also remain asymptomatic for years, as reported in this case. A proper follow-up with radiological evaluation is helpful to ensure a proper diagnosis prior to symptom manifestations. When evaluating patients who have had surgical experience, a high degree of suspicion index is essential. In this case, the diagnosis was not suspected preoperatively, since the foreign body had no radiopaque markers and was accompanied with stones, it could not be described by typical X-ray manifestations or computed tomography alone.

Foreign bodies due to surgical instruments being left behind after kidney surgery are rarely reported in the literature, and spontaneous metastasis to the intrarenal site is even rarer. From January 1999 to July 2020, a total of 262 adverse events involving Hem-o-Lok clips were reported to the Maude database. Due to consequent medical and legal issues, we believe that such adverse events are obviously underreported. Among them, within general surgical literature, it has been reported that the clip shifted to the common bile duct after laparoscopic cholecystectomy, and induced gallstone formation or cholangitis^[8,9]. Since Banks first reported the migration of this class of surgical clip into the bladder after LRP in 2008^[10], such related adverse events for HOLC have gradually increased within urinary system. In addition, the majority of reported complications with HOLC are urethral erosion, bladder neck contracture and subsequent stone formation^[10-14], though were finally visualized and removed by endoscopy.

The collecting system migration of suture material post-LPN has also been reported. Park *et al* reported a case of ureteral migration of a HOLC after two years of laparoscopic right partial nephrectomy, and it was removed by using a ureteroscopic stone basket device^[15]. Dasgupta reported a HOLC that migrated into the renal pelvis, with renal calculus formation following laparoscopic dismembered pyeloplasty within a few weeks^[16], and a similar case was also reported after 6 years of retroperitoneoscopic left pelvolithotomy^[17]. In both reported cases, the HOLC was removed by percutaneous nephroscope. With the increasing deployment of HOLC, similar complications have also appeared in robotic surgery. Blumenthal and his colleague^[11] reported the first case of HOLC migration into the vesicourethral anastomosis and urethra after RALRP (robot assisted laparoscopic radical prostatectomy), while Yadav *et al*^[18] presented a HOLC that migrated into the subrenal calyx of the left kidney, presenting as a renal stone, which was employed to close the mesenteric window that formed during bilateral robotic pyeloplasty.

Unfortunately, clips consisting of differing materials cannot avoid similar situations. Miller *et al*^[19] reported that Lapra-Ty absorbable suture clips (Ethicon

EndoSurgery™) migrated from a laparoscopic partial nephrectomy bed into the collecting system at six weeks after LPN, causing renal colic. Massoud^[20] also reported that a metal surgical clip was displaced into the ureter after open partial nephrectomy (OPN), and all of which were passed spontaneously. Furthermore, within urologic literature, it has been reported that surgical clips, and even absorbable gastrointestinal staples, can act as a nidus for stone formation when in contact with urine^[21]. Msezane *et al*^[22] reviewed differing sealants and laparoscopic instruments that can be used to stop bleeding of the renal parenchyma within LPN and concluded that there is no gold-standard single agent or combination of products that can be applied to all cases. A combined approach, utilizing manual suturing and hemostatic technology, may be the best strategy to achieve hemostasis.

Although there existed non-common cases related to HOLC migrating into the renal pelvis with renal calculus formation, yet it is unclear how a HOLC become dislodged and migrated into the collecting system. According to the surgical experience of this case four years ago, we analyzed that the persistent tensile force of the suture line may facilitate HOLC migration into the renal pelvis from the fissure at the ureteropelvic junction before the sutures absorbed and embedded, and it can act as a nidus for stone formation when in contact with urine for extended time periods. By reviewing related case reports of laparoscopic surgery within renal pelvis or collecting system, we believe that the use of HOLC instead of traditional knots -when suturing incisions within renal pelvis or collecting system - leads to a higher risk of HOLC migration. However, more research must be performed to uncover how and why HOLC migrates into the collection system.

Due to the limitation of warm ischemia time in partial nephrectomy, HOLC can be used for vascular pedicle control or rapid suture stabilization, and there is no better substitute presently. Therefore, risk reduction for this rare complication is still a presently challenging. Surgical or endoscopic removal of the foreign body is mandatory, in view of the high complication rate and significant clinical symptoms. Several previous reports include the removal of foreign bodies by percutaneous

nephrostomy or endoscopic maneuvers. But prevention of HOLC migration is clearly better than complication treatments, so the surgeon must be aware of the possibility of clip migration. Meanwhile, all loose clips should be actively sought and retrieved from the wound during urinary system surgery, and the use of clips in close proximity to the anastomotic stoma of collecting systems should also be avoided.

CONCLUSION

In summary, we concluded that in order to prevent complications related to HOLC, loose clips should be actively sought and retrieved from the wound during urinary system surgery, and the use of clips in close proximity to the anastomotic stoma of collecting systems should be avoided.

13%

SIMILARITY INDEX

PRIMARY SOURCES

- 1

Kyung Seo Park, Young Jin Sim, Han Jung. "Migration of a Hem-o-Lok Clip to the Ureter Following Laparoscopic Partial Nephrectomy Presenting With Lower Urinary Tract Symptoms", International Neurourology Journal, 2013

Crossref

49 words — 2%

- 2

Meredith Miller, James Kyle Anderson, Margaret S. Pearle, Jeffrey A. Cadeddu. "Resorbable clip migration in the collecting system after laparoscopic partial nephrectomy", Urology, 2006

Crossref

42 words — 2%

- 3

www.scielo.br

Internet

31 words — 1%

- 4

Yu, Chih-Chin, Cheng-Kuang Yang, and Yen-Chuan Ou. "Three Types of Intravesical Hem-o-Lok Clip Migration After Laparoscopic Radical Prostatectomy", Journal of Laparoendoscopic & Advanced Surgical Techniques, 2015.

Crossref

20 words — 1%

- 5

Kai Huang, Zhi-Qiang Jiang. "Hem-o-lok", Chinese Medical Journal, 2016

Crossref

18 words — 1%

- 6

utsouthwestern.influent.utsystem.edu

Internet

18 words — 1%

-
- 7 www.ncbi.nlm.nih.gov 18 words — 1%
Internet
-
- 8 elischolar.library.yale.edu 16 words — 1%
Internet
-
- 9 Ranan Dasgupta. "<i>Case Report</i>: Percutaneous Nephrolithotomy for a Stone on a Hem-o-Lok[®] Clip", Journal of Endourology, 03/2008 15 words — 1%
Crossref
-
- 10 Suo-Yuan Li, Ye Miao, Liang Cheng, Ye-Feng Wang, Zhi-Qiang Li, Yu-Bo Liu, Tian-Ming Zou, Jun Shen. "Surgical treatment of delayed cervical infection and incomplete quadriplegia with fish-bone ingestion: A case report", World Journal of Clinical Cases, 2021 15 words — 1%
Crossref
-
- 11 Saleh Binsaleh. "Evaluation of Hemostatic Aids in Laparoscopic Nephron-Sparing Surgery", The Scientific World JOURNAL, 2011 14 words — 1%
Crossref
-
- 12 Meng Zhang, Ling Jin, Yan-Long Duan, Jing Yang et al. "Diagnosis and treatment of pediatric anaplastic lymphoma kinase-positive large B-cell lymphoma: A case report", World Journal of Clinical Cases, 2021 13 words — 1%
Crossref
-
- 13 Yu Seob Shin. "Floating Hem-o-Lok Clips in the Bladder without Stone Formation after Robot-Assisted Laparoscopic Radical Prostatectomy", Korean Journal of Urology, 2012 12 words — 1%
Crossref

EXCLUDE QUOTES ON

EXCLUDE MATCHES < 12 WORDS

EXCLUDE BIBLIOGRAPHY ON