

IgA nephropathy treatment with traditional Chinese medicine: A case report

By Ying-Yu Zhang

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 72162

Manuscript Type: CASE REPORT

IgA nephropathy treatment with traditional Chinese medicine: A case report

Zhang YY *et al.* TCM treatment of IgAN

Ying-Yu Zhang, Yan-Lin Chen, Lan Yi, Kun Gao

Abstract

BACKGROUND

IgA nephropathy (IgAN) is a common primary glomerular disease that leads to end-stage renal disease with poor therapy efficacy. Traditional Chinese Medicine (TCM) is effective in the treatment of IgAN and has the potential to become an alternative treatment for IgAN. Professor Yan-Qin Zou is a nephropathy expert, a National Chinese Medicine Master, and an heir to the Menghe School of Medicine.

CASE SUMMARY

A 28-year-old man had positive urinary protein and elevated serum creatinine (Scr) results and was diagnosed with IgAN 2-3 years, respectively, before outpatient visit. He attended the outpatient department at our hospital in 2017. Professor Zou used the following methods to treat the patient: Invigorating the spleen and tonifying the kidney, removing dampness and clearing turbidity, quickening the blood and transforming stasis, and freeing vessels and regulating collaterals. She adjusted the prescription in accordance with the patient's symptoms. After 6 mo of treatment, the symptoms had resolved and serological indexes were also decreased [Scr from 288.5 to 188.6 $\mu\text{mol/L}$, blood urea nitrogen (BUN) from 10.9 to 9.5 mmol/L, serum uric acid (UA) from 612 to 503 $\mu\text{mol/L}$]. During follow-up, BUN, Scr, and UA levels remained stable.

CONCLUSION

Professor Zou's therapeutic strategy to treat IgAN using TCM was efficacious and a good reference for application.

Key Words: Traditional Chinese medicine; IgA nephropathy; Primary glomerular disease; Case report

Zhang YY, Chen YL, Yi L, Gao K. IgA nephropathy treatment with traditional Chinese medicine: A case report. *World J Clin Cases* 2022; In press

Core Tip: IgA nephropathy (IgAN) is a common primary glomerular disease leading to end-stage renal disease with poor prognosis. Traditional Chinese Medicine (TCM) is an alternative treatment for IgAN. Professor Yan-Qin Zou, one of the National Chinese Medicine Masters, has accumulated rich and unique experience in TCM diagnosis and treatment of nephropathy. This case report analyses an IgAN patient receiving TCM treatment with satisfying clinical efficacy. Professor Zou's therapeutic strategy to treat IgAN using TCM was efficacious and a good reference for clinical application.

INTRODUCTION

IgA nephropathy (IgAN) is one of the most common forms of primary nephritis and is characterized by mesangial proliferation and diffused IgA deposition in the mesangium^[1]. In China, IgAN accounts for 45.26% of primary glomerulonephritis cases^[2]. To date, corticosteroids, immunosuppressive agents, and symptomatic treatment have been the main therapeutic strategies, but the long-term treatment efficacy and safety need to be improved^[3]. How to preserve the renal function remains a major challenge for nephrologists^[4]. A new target, sodium-dependent glucose transporters 2 (SGLT2), has reportedly shown efficacy in IgAN, but the high incidence of urinary tract infections has limited the use of SGLT2 inhibitors^[5]. Resultingly, increasing attention has been paid to Traditional Chinese Medicine (TCM).

TCM has a long history in the treatment of kidney diseases in Eastern Asia^[6]. Over recent decades, the efficacy and safety of TCM in treating kidney diseases have improved considerably^[7,8], as documented studies have confirmed^[9,10]. TCM thus has the potential to become an alternative treatment for IgAN^[11,12]. However, high quality studies remain scarce. Professor Yan-Qin Zou, a National Chinese Medicine Master and the fourth-generation heir of the Fei Boxiong family, is a well-known nephrology expert in TCM. Professor Zou has been engaged in clinical work for nearly 60 years and has

accumulated rich and unique experience in treating nephropathy using TCM methods^[13].

Therapies that successfully delay the progression of IgAN remain limited. Once the condition has progressed to renal failure, there are no effective methods to stop the decline of kidney function^[14]. This case report describes a patient suffering from renal failure due to IgAN. He received TCM treatment with improvement in renal function, thus demonstrating Professor Zou's therapeutic strategy for treating IgAN. We hope this report could provide a valuable reference for the treatment of IgAN with TCM.

CASE PRESENTATION

Chief complaints

The patient's chief complaint was positive urinary protein and elevated serum creatinine (Scr) levels.

History of present illness

A 28-year-old man had positive urinary protein (++) and elevated Scr (149 $\mu\text{mol/L}$) test results in 2014. He underwent renal biopsy in February 2015 and was diagnosed with "IgAN, mesangial proliferative glomerulonephritis with sclerosis". Since then, the patient has been taking losartan (100 mg daily) and ketosteril (α -ketoacid tablets; 2.52 g/3 times daily). He first visited the outpatient department at our hospital on June 1, 2017. At this visit, malaise, waist soreness, acid reflux, and bubbles in the urine were the main clinical manifestations. The patient also complained of frequent urination at night. His tongue was light red with a yellow coating, his pulse was slippery (meaning that his pulse was beating freely and smoothly like the movement of abacus beads), and his pharynx was red.

The patient visited the outpatient department again on August 2, 2017. At this visit, waist soreness and fatigue were resolved. The patient's stomach discomfort was also alleviated. However, he still had a sore throat and spermatorrhea.

At his 3th visit on September 28, 2017, serological indexes were stable. Clinical manifestations such as waist soreness, fatigue, pharynx discomfort, spermatorrhea, and epigastric discomfort were unchanged.

At the 4th visit on October 26, 2017, serological indexes had decreased significantly. The patient's epigastric discomfort had improved, but waist soreness remained.

At the patient's 5th visit on November 23, 2017, his clinical symptoms including waist soreness, fatigue, spermatorrhea, and epigastric discomfort were all resolved, and the serological indexes had also decreased.

5 *History of past illness*

The patient has no history of a relevant illness.

Personal and family history

The patient has no family history of a similar illness.

10 *Physical examination*

On admission, the patient's blood pressure was 120/80 mmHg, there were no abnormalities detected during the physical examination including the cardiopulmonary system. There was no edema in either lower limb.

Laboratory examinations

The serological and urinary indexes are shown in Table 1.

Imaging examinations

Renal ultrasound showed no obvious abnormality.

Pathological analysis

Under light microscopy, a renal biopsy sample contained 25 glomeruli, among which there were 15 glomeruli with global sclerosis and no crescent. The remaining glomeruli

were enlarged with approximately 90-110 cells/glomerulus. There were some mesangial areas with severe matrix hyperplasia with 2-3 cells/mesangium. The basement membrane was normal and the structure of capillary loop remained open. Complete sclerosis was seen in some glomeruli. Masson Stain: Negative.

There were moderate lesions in the tubulointerstitium with indistinct tubular structure, no tubular atrophy, some swollen and degenerated tubular epithelial cells and some vacuolar degeneration. Focal fibrosis (40%) and more inflammatory cell infiltration could be seen in the interstitial area.

Immunofluorescence showed classic prominent IgA deposits.

FINAL DIAGNOSIS

Accordingly, a diagnosis of IgAN (mesangial proliferative glomerulonephritis with sclerosis), Hass III (Hass, 1997), M0E0S0T1 (Oxford, 2009) was considered.

TREATMENT

The symptoms that were present at the patient's first visit are described in TCM as kidney and spleen deficiency, dampness, turbidity, blood stasis, and collateral disharmony. After differentiation, Professor Zou initiated TCM therapy, namely by invigorating the spleen and tonifying the kidney, removing dampness and clearing turbidity, ¹ quickening the blood and transforming stasis, and freeing vessels and regulating collaterals.

The patient's TCM prescription included the following: (1) Taizishen (*Pseudostellaria heterophylla*) 15 g; (2) Shenghuangqi (¹³ *Radix astragali*) 30 g; (3) Chaobaizhu (³ fried *Rhizoma atractylodis macrocephalae*) 10 g; (4) Shengyiyiren (*Semen coicis*) 30 g; (5) Fuling (*Poria*) 30 g; (6) Fushen (*Poria cocos*) 30 g; (7) Jiangcan (*Bombyx batryticatus*) 15 g; (8) Niubangzi (*Fructus arctii*) 10 g; (9) Huangshukuihua (*Abelmoschus manihot*) 30 g; (10) Shiwei (*Pyrrosia shearerii*) 20 g; (11) Chuanduan (*Dipsacus asper*) 15 g; (12) Sangjisheng (³ *Ramulus taxilli*) 15 g; (13) Yinchen (*Herba artemisiae scopariae*) 30 g; (14) Tufuling (*Smilax glabra*) 30 g; (15) Shengpuhuang (*Pollen typhae*) 30 g; (16) Danshen (¹⁵ *Radix salviae*)

miltiorrhizae) 20 g; (17) Chishao (*Radix paeoniae rubra*) 15 g; (18) Honghua (*Flos carthami*) 10 g; (19) Wulingzhi (*Trogopterus xanthipes*) 30 g; (20) Shudahuang (*cooked Radix et rhizoma rhei*) 12 g; (21) Cheqianzi (*Semen plantaginis*) 30 g; (22) Donglingcao (*Rabdosia rubescens*) 10 g; (23) Yumixu (*Zea mays*) 30 g; (24) Bixie (*Dioscoreae hypoglaucæ*) 20 g; and (25) Walengzi (*Arca subcrenata*) 40 g. A dose of Chinese medicinal herbs is decocted twice, and the juice is mixed together. The herbs were taken twice a day.

At the patient's 2th visit, his clinical symptoms were resolved, and the serological indexes were improved. Professor Zou strengthened the therapy of clearing damp-heat and turbidity along with relieving pharynx discomfort and detoxifying in accordance with the abovementioned treatment principles. The patient's prescription included the prescription from visit 1, but the amount of tufuling (*Smilax glabra*) was increased to 50 g, shudahuang (*cooked Radix et rhizoma rhei*) was increased to 20 g, and the following were added: (1) Zisuye (*Folium perilla leaf*) 30 g; (2) Tusizi (*Semen cuscute*) 15 g; (3) Xuanshen (*Radix scrophulariae*) 10 g; (4) Jinyinhua (*Flos lonicerae*) 10 g; and (5) Baihuasheshecao (*Oldenlandia diffusa*) 30 g.

At the patient's 3th visit, Professor Zou strengthened the therapy of tonifying the kidney while maintaining the power of discharging turbidity and detoxifying. In addition to the initial prescription, duzhong (*Cortex eucommiae*) 15 g; huainiuxi (*Radix achyranthis bidentatae*) 10 g; nvzhenzi (*Fructus ligustri lucidi*) 20 g and gouqizi (*Fructus lycii*) 20 g were added to tonify both kidney Yin and kidney Yang. Jixuecao (*Centella asiatica*) 30 g; baihuasheshecao (*Oldenlandia diffusa*) 30 g and shenggancao (*Radix glycyrrhizae*) 6 g were added to purge turbidity and aid detoxification, which are effective for reducing creatinine. Jiangcan (*Bombyx batryticatus*); niubangzi (*Fructus arctii*); donglingcao (*Rabdosia rubescens*); bixie (*Dioscoreae hypoglaucæ*) and walengzi (*Arca subcrenata*) were removed when the abovementioned drugs were added at the third visit. Because shudahuang (*cooked Radix et rhizoma rhei*) can also help to quicken the blood, transform stasis, and free the vessels, chishao (*Radix paeoniae rubra*) and honghua (*Flos carthami*) were removed because there was enough power to promote blood circulation in the previous treatment.

At the 4th visit, Professor Zou strengthened the therapy to tonify the kidney and clear the pharynx, adding duanlonggu (*Os draconis*) 40 g; duanmuli (*Concha ostreae*) 40 g; gouji (*Cibotium barometz*) 20 g and tusizi (*Semen cuscutae*) 15 g to the last prescription.

At the 5th visit, the patient's clinical symptoms were resolved, and the serological indexes were also decreased. Over the following 5 mo, Professor Zou prescribed the original treatment principles (*i.e.*, Invigorating the spleen and tonifying the kidney, removing dampness and clearing turbidity, quickening the blood and transforming stasis, and freeing vessels and regulating collaterals).

OUTCOME AND FOLLOW-UP

The patient's symptoms and discomfort were resolved. As shown in Figures 1, during the 1-year follow-up, renal function was improved as evidenced by the stable level of BUN, Scr, serum cystatin C and UA. As shown in Figure 2, urinary red blood cell remained negative, and urinary protein maintained negative or trace findings.

DISCUSSION

A typical clinical manifestation of IgAN is hematuria with or without proteinuria, hypertension, and impaired renal function^[1]. Some patients with IgAN gradually progress to chronic renal failure 10-20 years after their diagnosis^[15]. The efficacy of conventional therapy is limited because of some side effects. Treatment using TCM is worth considering because it has shown good safety and therapeutic effects.

TCM pathogenesis and IgAN therapy

IgAN is in the edema and hematuria category in TCM^[16]. In ancient times, doctors had a comprehensive theory on pathogenesis, diagnosis, and therapy for kidney diseases, as recorded in the Yellow Emperor's Canon of Medicine Plain Conversation: "All dampness, swelling and turgidity is ascribed to the spleen." Professor Zou proposes that IgAN pathogenesis is a mixture of deficiency and excess. Patients with IgAN are susceptible to outer pathogen infection because of lung Qi deficiency, in accordance

with TCM theory. Qi deficiency in the kidney, spleen, and lung is the internal basis that contributes to damp-heat production as a pathological factor throughout the course of IgAN. Additionally, the long-term existence of the dampness pathogen suppresses Qi and leads to Qi stagnation. Qi stagnation makes it difficult for blood to circulate smoothly. In summary, the main cause of IgAN is derived from kidney, spleen and lung deficiency, which is accompanied by dampness, damp-heat, blood stasis and wind-pathogen. Improving deficiency and eliminating excess pathogenic factors constituted Professor Zou's main clinical therapeutic strategy.

Case study

In this case, Professor Zou initially distinguished between deficiency syndrome and excess syndrome, giving consideration to both improving the deficiency and expelling pathogenic factors. The patient was treated on the basis of the disease stage and its course. In the early stage of therapy, pathogenic factors were prominent. Professor Zou strengthened the therapy of purging turbidity and detoxifying to gradually eliminate the damp-heat and blood stasis from the body. Malaise and waist soreness in the morning indicated kidney and spleen deficiency, while acid reflux, increased Scr and UA levels and bubbles in the urine demonstrated turbidity and dampness. She used treatment measures to tonify the kidney and spleen, purge turbidity and clear dampness, promote blood circulation and remove blood stasis. After the first treatment, the waist and stomach symptoms were relieved and serology indexes were improved. However, the patient's throat discomfort and an increase in spermatorrhea demonstrated persisting kidney deficiency. At the following visits, the deficiency syndrome was the main target. Professor Zou strengthened the therapy of invigorating the kidney, spleen and lung to improve Qi. Throughout the disease treatment, Professor Zou focused on tonifying the kidney and spleen and clearing away damp-heat. After almost 6 months of treatment, the patient's symptoms were resolved, Scr and UA levels were significantly lower. There was an increase in the UA level on January 18, 2018, which may have resulted from the previous cold or the patient's overeating. These were

not recorded and overall, the UA level trended downwards. Over the subsequent follow-up, renal function remained stable.

Common compatibility and dosage

For herb selection, Professor Zou prefers mild and gentle herbs to invigorate the kidney and spleen because warm and hot drugs can easily hurt Yin, and greasy drugs can easily cause dampness stagnation. She uses drugs such as: (1) Taizishen (*Pseudostellaria heterophylla*); (2) Shenghuangqi (*Radix astragali*); (3) Chaobaizhu (fried *Rhizoma atractylodis macrocephalae*); (4) Shengyiyiren (*Semen coicis*), (5) Fuling (*Poria*); and (6) Fushen (*Poria cocos*). Furthermore, the mechanism of kidney diseases in TCM theory is the disorder of Yin and Yang in the lower abdomen (xia-jiao), and it is difficult for the herbs to reach the disease center because xia-jiao is located in the lower part of the body^[17]. However, the course of kidney disease is often long and persistent. Professor Zou used a combination of herbs and larger drug doses to balance Yin and Yang in the lower abdomen. The following herbs are often used for purging turbidity and detoxifying: (1) Huangshukuihua (*Abelmoschus manihot*); (2) Shiwei (*Pyrrosia shearereri*); (3) Yinchen (*Herba artemisiae scopariae*); (4) Tufuling (*Smilax glabra*); (5) Shudahuang (cooked *Radix et rhizoma rhei*); (6) Cheqianzi (*Semen plantaginis*); (6) Donglingcao (*Rabdosia rubescens*); (7) Yumixu (*Zea mays*); (8) Bixie (*Dioscoreae hypoglaucae*); (9) Baihuasheshecao (*Oldenlandia diffusa*); (10) Xuanshen (*Radix scrophulariae*); and (11) Jinyinhua (*Flos lonicerae*). For proteinuria, drugs such as: (1)Jiangcan (*Bombyx batryticatus*); (2) Niubangzi (*Fructus arctii*); and (3) Huangshukuihua (*Abelmoschus manihot*) are commonly used^[18-20]. Modern pharmacological research suggests that the major effects of herbal medicine are related to anti-inflammatory, antioxidative, antifibrotic and immunomodulatory pathways^[6,12,21].

Characteristics and comparison with similar treatment

Most TCM practitioners consider that Qi and Yin deficiencies are the leading causes of IgAN and propose that the treatment of IgAN should be based on reinforcing Qi and

replenishing Yin^[22]. In contrast to the general TCM treatment, Professor Zou suggests that kidney deficiency is the most fundamental internal factor in the occurrence of IgAN and the treatment of IgAN should be based on tonifying the kidney and harmonizing other viscera. In addition, the spleen also plays an important role in the pathogenesis of IgAN. In TCM theory, the spleen governs the transportation and transformation of nutrients and water and it controls the function of spleen-qi to elevate food nutrients^[23]. Tonifying the spleen is conducive to controlling proteinuria and improving metabolism. Recently, some powerful drugs, such as leigongteng (*Tripterygium wilfordii*), have been proven to be protective against kidney diseases^[23]. However, although these powerful drugs may have obvious short-term effects, they may easily hurt healthy Qi. Professor Zou advocates mild drugs to gradually dissipate the pathogenic factors without side effects. Unique academic thought and exquisite herbal selection reflect the power of Professor Zou as a National Chinese Medicine Master.

Limitations

Herein, we report only one case, which has limited reference value to reflect Professor Zou's treatment experience. However, we followed the patient for nearly 1 year and did not observe any subsequent disease relapse. Finally, 24-hour urine protein quantification is also an important laboratory indicator for IgAN, but we did not have access to these results because the patient did not undergo this test in the outpatient setting.

CONCLUSION

IgAN is a prevalent type of glomerular disease in most Western and Asian countries. TCM therapy uses natural medicines that can reduce proteinuria and Scr and improve symptoms such as back pain in a long-term regimen. Professor Zou is a TCM specialist in treatment of IgAN. Her main therapeutic strategies for treating this disease focus on improving deficiency and eliminating excess pathogenic factors. This case provides a specific method and an effective reference for clinical application of TCM to treat IgAN.

Our findings indicate that TCM can play an important role in the treatment of IgAN and is worth consideration as a worldwide treatment option.

ACKNOWLEDGEMENTS

We thank Wei Li for editing the English text of a draft of this manuscript and Pengcheng Sun for helping to revise the manuscript.

IgA nephropathy treatment with traditional Chinese medicine: A case report

ORIGINALITY REPORT

9%

SIMILARITY INDEX

PRIMARY SOURCES

1 Guo Jing, Chen Hongdong, Zhao Xiyan, Zhao Linhua, Tong Xiaolin. "Diabetic kidney disease treated with a modified Shenzhuo formula derived from Traditional Chinese Medicine: a case report", Journal of Traditional Chinese Medicine, 2017 60 words — 2%

Crossref

2 "Essentials of Chinese Medicine", Springer Science and Business Media LLC, 2010 29 words — 1%

Crossref

3 Gu, Sherman, Angela WH Yang, Charlie CL Xue, Chun G Li, Carmen Pang, Weiya Zhang, Hywel C Williams, and Sherman Gu. "Chinese herbal medicine for atopic eczema", Cochrane Database of Systematic Reviews Reviews, 2013. 20 words — 1%

Crossref

4 www.wjgnet.com 19 words — 1%

Internet

5 Cuo-Mao-Ji Zhang, Xiao Wang. "Suspected cerebrovascular air embolism during endoscopic esophageal varices ligation under sedation with fatal outcome: A case report", World Journal of Clinical Cases, 2022 16 words — 1%

Crossref

-
- 6 Yuan-Han Qin, Tian-Biao Zhou, Li-Na Su, Feng-Ying Lei, Wei-Fang Huang, Yan-Jun Zhao. "Association between ACE polymorphism and risk of IgA nephropathy: A meta-analysis", Journal of the Renin-Angiotensin-Aldosterone System, 2011
Crossref 13 words — < 1%
-
- 7 www.dovepress.com
Internet 13 words — < 1%
-
- 8 www.researchsquare.com
Internet 13 words — < 1%
-
- 9 Lihong Zhao, Tianfang Wang, Jian Dong, Angus Chen, Guanying Li. "Liver-stomach disharmony pattern: Theoretical basis, identification and treatment", Journal of Traditional Chinese Medical Sciences, 2018
Crossref 11 words — < 1%
-
- 10 thejns.org
Internet 11 words — < 1%
-
- 11 www.cochrane.org
Internet 11 words — < 1%
-
- 12 Wang Yuanhui, Zhang Lei, Pan Guodong, Liu Jing, Zhang Hongrui, Song Xu, Wang Qiang. "Traditional Chinese Medicine comprehensive therapy for the improvement of motor function in spinal cord injury patients", Journal of Traditional Chinese Medicine, 2016
Crossref 10 words — < 1%
-
- 13 Yubo Guo, Yu Li, Liming Xue, Richele P. Severino, Sihua Gao, Jianzhao Niu, Lu-Ping Qin, Dongwei Zhang, Dieter Brömme. "Salvia miltiorrhiza: An ancient Chinese 9 words — < 1%

14 cmj.org 9 words — < 1%
Internet

15 Li, Hui-qin, Jing-jing Wei, Wan Xia, Ji-huang Li, Ai-ju Liu, Su-bing Yin, Chen Wang, Liang Song, Yan Wang, Guo-qing Zheng, and Ji-ping Fan. "Promoting blood circulation for removing blood stasis therapy for acute intracerebral hemorrhage: a systematic review and meta-analysis", Acta Pharmacologica Sinica, 2015. 8 words — < 1%
Crossref

16 Qiong Zhang. "Making the New World Their Own", Brill, 2015 8 words — < 1%
Crossref

17 www.ncbi.nlm.nih.gov 8 words — < 1%
Internet

18 Pang, Clifford. "Integrative hyperthermia treatments for different types of cancer", Hyperthermia in Oncology, 2015. 7 words — < 1%
Crossref

19 Zhihua Zheng, Yi Wen, Yanhua Yan, Zhiheng Xu, Kechao Nie, Xu Chen, Fengbin Liu, Jinglin Pan, Peiwu Li. "The effect of fortifying the spleen, clearing heat, activating blood method on chronic atrophic gastritis: a real-world study", Cold Spring Harbor Laboratory, 2020 7 words — < 1%
Crossref

