

75087_Auto_Edited.docx

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 75087

Manuscript Type: ORIGINAL ARTICLE

Observational Study

Effect of Osteoarthritic Knee Flexion Deformity Correction by Total Knee Arthroplasty on The Sagittal Spino - Pelvic Alignment in Indian Population - A Prospective Study

Changes in Sagittal Spino-Pelvic Parameters after Total Knee Arthroplasty

Lubaib Karaniveed Puthiyapura, Mantu Jain, Sujit Kumar Tripathy, HariDas MP

Abstract

BACKGROUND

Sagittal alignment of the spine, pelvis, and lower extremities is essential for maintaining a stable and efficient posture and ambulation. Imbalance in any element can result in compensatory changes in the other elements. Knee flexion is a compensatory mechanism for spinopelvic sagittal alignment (SSA) and is markedly affected in severe knee osteoarthritis (OA). The correction of knee flexion deformity (KFD) by total knee arthroplasty (TKA) can lead to complementary changes in the sagittal spinopelvic parameters (SSPs).

AIM

The present study determined the SSP changes in patients with knee OA with or without KFD undergoing TKA.

METHODS

The study was conducted in 32 patients who underwent TKA. A neutral standing whole-spine lateral radiograph was performed before surgery and 3 mo after surgery in these patients. Subjects were divided into two groups (Group 1 obtained $>10^\circ$ corrections in KFD; group B obtained $<10^\circ$ correction). The pelvic tilt (PT), pelvic incidence (PI), sacral slope (SS), lumbar lordosis (LL), and sagittal vertical axis (SVA) were measured.

RESULTS

The median of change in PT, PI, SS, LL, and SVA was 0.20, 1.00, 2.20, -0.40 , and 6.8 mm, respectively. The difference in the change in SSPs between the two groups was statistically non-significant.

CONCLUSION

SSPs such as PI, PT, SS, LL, And SVA do not change significantly following TKA in end-stage knee OA despite a significant correction ($>10^\circ$) in KFD.

Key Words: Knee Osteoarthritis; Total knee replacement; Spino-sagittal parameters

Karaniveed Puthiyapura L, Jain M, Tripathy SK, MP H. Effect of Osteoarthritic Knee Flexion Deformity Correction by Total Knee Arthroplasty on The Sagittal Spino - Pelvic Alignment in Indian Population - A Prospective Study. *World J Clin Cases* 2022; In press

Core Tip: The sagittal alignment of the spine, pelvis, and lower extremities is essential for maintaining a stable and efficient posture and ambulation. Any imbalance in one element can result in compensatory changes in the other. Low back pain arising from hip pathology was termed “Hip Spine Syndrome.” As the sagittal spinopelvic parameters (SSPs) were described, the researchers became inquisitive about documenting these changes in hip pathology and post-surgical correction. After two decades, the “Knee Spine syndrome” was described in a similar logical sequence. The SSPs have become an area of interest with several user-friendly tools to measure. The current papers evaluate these parameters in patients with osteoarthritis knee and undergoing total knee replacement.

1

INTRODUCTION

Osteoarthritis (OA) is a disease characterized by degeneration of cartilage and underlying bone within a joint. Patients with knee OA exhibit fixed flexion deformity (FFD) and varus or valgus coronal deformity in the late stages. Total knee arthroplasty (TKA) is a safe and effective procedure for these patients. It reduces pain, corrects deformities, restores function, and improves quality of life of patients [1].

Sagittal spinal alignment (SSA) of the body is vital for maintaining a stable and efficient posture and ambulation [2-3]. Any imbalance in the SSA can lead to compensatory changes in the pelvis, hip, and knee to ensure a static horizontal gaze

with minimum energy expenditure [4]. Offierski *et al* first described the "hip spine syndrome," wherein hip diseases were associated with lower back pain (LBP) [2]. Additionally, Schwab *et al* observed that altered SSA was mainly responsible for the LBP [4]. Parvizi *et al* noticed a change in the sagittal spinopelvic parameters (SSPs) in patients with hip OA undergoing total hip arthroplasty (THA) [3]. Recently, Murata *et al* described the "Knee spine syndrome" for lumbar spine symptoms caused by degenerative changes in the knee [5]. Lee CS *et al* observed that a change in SSPs compensates knee flexion [6]. Hence, correcting knee flexion deformity (KFD) by TKA could lead to complementary changes in the SSPs. While most patients have improved spinal problems following TKA, few have worsening of spinal problems or recurrence of KFD. Therefore these multiple joints interact and affect each other. However, studies in this regard are scanty [7-8].

Indian people are predisposed to early knee OA due to their habit of squatting and sitting cross-legged [9]. The present study attempted to evaluate and validate changes in SSPs following the correction of KFD by TKA in the Indian population.

MATERIALS AND METHODS

The present prospective observational study was conducted in 32 patients with knee OA admitted for TKA to the Department of Orthopedics at our institute from March 2020 to August 2021 after institutional ethics clearance. Patients with previous spine, hip, or knee surgery; previous spinal fractures, infections, or coronal deformity > 10°; spondylolisthesis; and sagittal foot deformities were excluded from the study. Based on a previous study by Lee SM *et al* [7], with a mean difference of 4 and standard deviation of 8, assuming $\alpha = 0.05$ and power $(1 - \beta) = 80\%$, using the formula $n = 2 (Z [1 - \alpha/2] + Z [1 - \beta])^2 \times SD^2 / d^2$, (SD- standard deviation, d - Effect size), the sample size was calculated to be 32. Informed consent was obtained from all patients. Clinical and radiological examinations of the spine and knee were performed in all patients. A standing neutral whole spine static lateral radiograph (scannogram) was performed preoperatively and after 3 mo of TKA. Radiological SSP was measured on a 36" cassette

orthoscannogram of the whole spine until mid-thigh. The lateral view was obtained using the software Surgimap (Surgimap, New York, NY, USA) [10]. Several studies have validated the accuracy of Surgimap [11-12]. The following SSPs were measured:

Pelvic incidence (PI): Angle between the line joining the center of the bicoxofemoral axis with the mid-point of the S1 endplate and another perpendicular to the S1 at its mid-point ($n = 45 \pm 15$).

Pelvic tilt (PT): Angle between the vertical reference line (VRL) and line joining the hip axis to the center of the superior S1 endplate ($n = 12.6 \pm 4.6$).

Sacral slope (SS): Angle between the superior end of S1 and horizontal ($n = 37.7 \pm 9.7$).

Lumbar lordosis (LL): Angle between the superior endplate of L1 and S1 ($n = 54.6 \pm 10$).

Sagittal vertical axis (SVA): Distance of a line connecting the centroid of the C7 vertebra and posterior superior sacral endplate from the sacral reference ($N < 5$ cm).

(**Figure 1**). Patients were divided into two groups, namely group A comprising patients with KFD correction $> 10^\circ$, and group B comprising patients with KFD correction $< 10^\circ$.

Knee Flexion Deformity (KFD) angle: The knee flexion angle (KFA) is the angle between the long axis of the femur and tibia, with lateral femur epicondyle as the center, and is measured using a goniometer. The average of the two flexion deformity angles was considered for analysis in bilateral cases.

Patients were divided into two groups, Group A with correction of KFD $> 10^\circ$ and Group B with correction of KFD $< 10^\circ$.

Descriptive statistics were used to analyze the demographic profile. Changes in parameters following TKA were analyzed using the Wilcoxon signed-rank test for non-parametric data. The Mann-Whitney *U* test was used to determine the difference between the two groups. Correlation between KFD angle and SSPs was determined using the Spearman's rank test.

RESULTS

The demographic details and baseline clinical and radiological parameters were assessed. No patient was lost to follow-up. Of the 32 patients, 9 were men, and 23 were

women. Bilateral TKA was performed in 22 patients, whereas unilateral TKA was performed in 10 patients. The mean age of the patients was 61.5 ± 7.64 years. The preoperative and postoperative SSPs are presented in **Table 1**. The preoperative KFD was 10.0 (IQR 5–18.75, Range 0–30) and post-operative KFD was 0.0 (IQR 0–5, Range 0–15) with a correction of 5.0 (IQR 1.25–15, Range 0–20), p-value <0.001 . Hence based on a change in KFB, 15 patients qualified in group A and 17 patients in group B. There was no significant difference between the two groups in terms of SSPs (**Table 2**). Similarly, a sub-group analysis of unilateral and bilateral TKA also yielded no significant difference (**Table 3**). No significant difference in SSP's between males and females (**Table 4**).

SVA > 50 mm was labeled as a global sagittal imbalance. Of the 32 patients, 16 exhibited preoperative sagittal imbalance. Of these 16 patients, the sagittal balance was restored only in 7 patients following TKA. No significant correlation was observed between change in KFD and change in SSPs ($p > 0.05$). A case example is illustrated in **Figure 2**.

DISCUSSION

SSA is crucial for maintaining a stable and efficient upright posture and ambulation^[11]. The body requires more energy consumption to maintain equilibrium when the SSA is abnormal. Pathology in the spine, pelvis, or lower extremity can disrupt the postural equilibrium. The close interrelation between these segments ensures compensatory changes in other segments. Offierski and McNab (1983) coined the term "hip spine syndrome" for LBP secondary to a hip pathology^[1]. Schwab *et al* exhibited that alteration in the SSPs was mainly responsible for LBP in this syndrome^[4]. Thus, several studies have evaluated SSPs in patients with hip pathology. However, the results have been equivocal. Parvizi *et al* documented some changes in the SSPs, whereas Ben-Galim *et al*, Eyvazov *et al*, and Eguchi *et al* did not observe any changes in SSPs despite improvement in the Visual Analogue Score (VAS) and Oswestry disability index score for LBP following THA^[3, 13–15].

Murata *et al* described the "knee-spine syndrome" in 2003 and observed a significantly lower LL in patients with KFD more than 50° [5]. Additionally, the limitation of knee extension was more significant in patients with LL < 30°. Thus, degenerative knee joint changes could lead to LBP. Lee CS *et al* used a motion-controlled knee brace bilaterally in a group of young men to simulate knee flexion contractures [6]. The SSPs were examined at 0°, 15°, and 30° of knee flexion. Femoropelvic angle and LL reduced significantly at 15° and 30° of knee flexion compared with full extension. Harato *et al* performed a gait analysis in simulated knee flexion [16]. At 30° flexion, a backward inclination of the pelvis was observed during standing, whereas a forward inclination of the trunk and pelvis was observed during walking. Thus, flexion contracture of the knee could cause changes in SSA, thereby proving the effect of SSPs on KFD. Therefore, the present study evaluated these parameters.

³ The mean age of the patients in the present study was 61.5 years (range: 41–76), which is lesser than that exhibited in studies by Kim *et al* (70.5 years) and Kitagawa *et al* (75 years) [8,17]. This may be due to the habit of squatting, cross-legged sitting, and kneeling among Indian people, which predispose them to OA at an early age [9]. Of the 32 patients in the present study, 22 underwent bilateral TKA, whereas 10 underwent unilateral TKA. The present study was conducted in 23 women and 9 men, whereas the study by Lee *et al* included only women [7].

The median preoperative KFD in the present study was 10° (IQR: 5–18.75, range: 0–30). Complete knee extension was achieved following TKA (IQR: 0°–5°, range: 0°–15°, **p value < 0.001**). Lee *et al*, Kim *et al*, and Kitagawa *et al* also reported a significant reduction in KFD after TKA [7,8,17].

Kim *et al* (2020) defined lumbar flexibility (LF) as the difference in LL in extension and flexion, and pelvic flexibility (PF) as PI. They reported an increase in LL and improvement in SVA following TKA in patients exhibiting better LF. Patients with better PF exhibited an increase in SS [17]. Kitagawa *et al* measured the difference between the PI and LL. A difference of >10° was observed in a majority of their patients (>50%). Additionally, a significant improvement was observed in SSPs following TKA with

increased SS and reduced SVA and PT [8]. However, the present study did not observe a significant change in SSPs (PT, PI, SS, LL, SVA) following TKA ($p > 0.05$). In the present study, 16 patients exhibited global sagittal imbalance ($SVA > 50$ mm). Following TKA, 7 of these patients attained sagittal balance ($SVA < 50$ mm), whereas the sagittal imbalance reduced but persisted in the remaining 9 patients. The present study exhibited a short follow-up period of 3 mo. Longer mobilization could result in more profound changes in the SSPs. The other possibility is that the spinal changes were irreversible, which has been observed in a few studies on SSPs following THA [13,14].

The present study divided the patients into two groups, namely group A comprising patients with KFD correction $> 10^\circ$ and group B comprising patients with KFD correction $< 10^\circ$. Because 5° is too small to measure clinically, 10° was considered the critical angle. No significant difference was observed between these two groups regarding changes in SSPs (p value > 0.05). Lee *et al* reported a significant increase in SS in a few patients with $>10^\circ$ deformity correction. However, a corresponding change in PT was not observed. The authors admitted that it was due to an error while taking the radiograph [7]. PI remained almost the same in all studies because it remains constant for a particular individual unless there is a sacroiliac dissociation.

PI positively correlated with PT, SS, and LL among the preoperative parameters. SVA exhibited a positive correlation with SS. No significant correlation was observed between flexion deformity and preoperative SSPs, whereas a significant positive correlation was observed between postoperative flexion deformity and PI and SS. No significant correlation was observed between correction in flexion deformity and SSP changes. Kitagawa *et al* reported a significant positive correlation of the knee flexion angle with SVA and a negative correlation with SS and LL [8]. The present study conducted a subgroup analysis between patients undergoing unilateral and bilateral TKA. However, the difference was statistically non-significant ($p > 0.05$).

The present study considered TKA and SSPs in the Indian population. The study is novel because literature regarding this aspect among the Indian population is scarce. All patients with concomitant spinal disease or previous surgery were excluded from

the present study to reduce bias. Subgroup analysis was conducted to evaluate the difference between unilateral and bilateral TKA. However, the present study has certain limitations. The lateral radiograph measurements in a neutral standing position of the whole spine was a static capture, not representative of functional performance. Perhaps the degree of pre-operative KFD could have also played a role in the pre-operative SSA. Also, that other factors may more readily determine the SSA, like physical activity exposure related to standing-walking, posture habits in general, and also a long-term body schema memory, establishing a certain deformity permanently (due to a long term knee-flexion contracture), being impossible to reverse even in a three-month post-surgery timeframe. Since the TKA was performed unilaterally in several patients, perhaps a frontal spino-pelvic alignment could have also examined. A small sample size and a short follow-up period prevent the generalization of the study findings. Additionally, LBP was not substantiated using a score despite an improvement in LBP in the follow-up. Further comprehensive multicenter studies with a larger sample size, longer follow-up time, and clinical LBP scores will strengthen the findings of this study.

CONCLUSION

SSPs such as PI, PT, SS, LL, and SVA do not change significantly following TKA in end-stage knee OA despite a significant correction ($>10^\circ$) in KFD. Thus, routine determination of the SSPs may not be required in patients undergoing TKA.

ARTICLE HIGHLIGHTS

Research background

Sagittal spinal alignment (SSA) of the body is vital for maintaining a stable and efficient posture and ambulation. Any imbalance in the SSA can lead to compensatory changes in the pelvis, hip, and knee to ensure a static horizontal gaze with minimum energy expenditure. Offierski *et al* first described the "hip spine syndrome," wherein hip diseases were associated with lower back pain (LBP). Recently, Murata *et al*

described the "Knee spine syndrome" for lumbar spine symptoms caused by degenerative changes in the knee. Additionally, Schwab *et al* observed that altered SSA was mainly responsible for the LBP. Lee CS *et al* observed that a change in SSPs compensates knee flexion [6]. Hence, correcting knee flexion deformity (KFD) by TKA could lead to complementary changes in the SSPs. While most patients have improved spinal problems following TKA, few have worsening of spinal problems or recurrence of KFD. Therefore these multiple joints interact and affect each other. However, studies in this regard are scanty [7-8].

Indian people are predisposed to early knee OA due to their habit of squatting and sitting cross-legged [9]. The present study attempted to evaluate and validate changes in SSPs following the correction of KFD by TKA in the Indian population.

Research motivation

The present study attempted to evaluate and validate changes in SSPs following the correction of KFD by TKA

Research objectives

The present study determined the sagittal spinopelvic parameters changes in patients with knee osteoarthritis with or without knee flexion deformity undergoing total knee arthroplasty.

Research methods

The study was conducted in 32 patients who underwent TKA. A neutral standing whole-spine lateral radiograph was performed before surgery and 3 mo after surgery in these patients. Subjects were divided into two groups (Group 1 obtained $>10^\circ$ corrections in KFD; group B obtained $<10^\circ$ correction). The pelvic tilt (PT), pelvic incidence (PI), sacral slope (SS), lumbar lordosis (LL), and sagittal vertical axis (SVA) were measured.

Research results

The median of change in PT, PI, SS, LL, and SVA was 0.20, 1.00, 2.20, -0.40, and 6.8 mm, respectively. The difference in the change in SSPs between the two groups was statistically non-significant.

Research conclusions

SSPs such as PI, PT, SS, LL, And SVA do not change significantly following TKA in end-stage knee OA despite a significant correction ($>10^\circ$) in KFD.

Research perspectives

The direction of further studies should include a larger sample size, longer follow-up time, and clinical LBP scores to strengthen the findings of this study.

2%

SIMILARITY INDEX

PRIMARY SOURCES

- 1

www.tandfonline.com
Internet

17 words — 1%
- 2

wprim.whooc.org.cn
Internet

16 words — 1%
- 3

K. Ouaz, A. Robier, E. Lescanne, C. Bobillier, S. Morinière, D. Bakhos. "Cancer of the external auditory canal", European Annals of Otorhinolaryngology, Head and Neck Diseases, 2013
Crossref

14 words — < 1%
- 4

Yicheng Li, Xiaogang Zhang, Xiaobin Guo, Wuhuzi Wulamu, Nuerailijiang Yushan, Baochao Ji, Li Cao. "Effective treatment of single-stage revision using intraarticular antibiotic infusion for polymicrobial prosthetic joint infection", The Journal of Arthroplasty, 2021
Crossref

13 words — < 1%

EXCLUDE QUOTES ON
EXCLUDE BIBLIOGRAPHY ON

EXCLUDE SOURCES < 12 WORDS
EXCLUDE MATCHES < 12 WORDS