

75726_Auto_Edited.docx

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 75726

Manuscript Type: CASE REPORT

Primary clear cell sarcoma of soft tissue in the posterior cervical spine invaded the medulla oblongata : A case report

Chen-Chen Liu, Wen-Peng Huang, Jian-Bo Gao

Abstract

BACKGROUND

Clear cell sarcoma (CCS) is a rare and highly malignant soft tissue tumor, usually occurring in the deep soft tissues of the distal tendons and aponeurosis of the extremities, especially the feet and knees. CCS originating in the head and neck is extremely rare. The clinical manifestations of CCS in the head and neck are not typical, and the imaging manifestations have certain characteristics, but the diagnosis still depends on pathological examination and genetic testing.

CASE SUMMARY

The 33-year-old male patient had paroxysmal headache for more than 4 years, accompanied by nausea and vomiting, which could be relieved after rest. Computed tomography angiography (CTA) showed a left paraspinal soft tissue mass. The enhancement showed obvious uneven enhancement with adjacent bone lytic destruction. Magnetic resonance imaging (MRI) examination showed isosignal on T₁-weighted images (T₁WI), slightly high signal on T₂-weighted images (T₂WI), high signal on T₂ fat suppression sequence, significantly high signal on diffusion weighted imaging (DWI), and obviously uneven enhancement. The lesion invaded the anterior medulla oblongata through the left atlantoaxial foramen and compressed the cervical

spinal cord on T₂WI. Primary CCS of soft tissue was diagnosed by pathology and genetic examination.

CONCLUSION

CCS should be considered in the differential diagnosis of soft tissue tumors of the head and neck, and its diagnosis depends on pathological examination and genetic testing.

Key Words: Clear cell sarcoma; Soft tissue; The medulla oblongata; Tomography; X-ray computed; Magnetic resonance imaging; Case report

Liu CC, Huang WP, Gao JB. Primary clear cell sarcoma of soft tissue in the posterior cervical spine invaded the medulla oblongata : A case report. *World J Clin Cases* 2022; In press

Core Tip: Clear cell sarcoma (CCS) is a rare and highly malignant soft tissue tumor, usually occurring in the deep soft tissues of the distal tendons and aponeurosis of the extremities. CCS originating in the head and neck is extremely rare and usually presents as a painless growth mass. In this case, the lesion was located in the head and neck and presented as episodic headache. The lesion was perforated and involved the medulla oblongata. Differential diagnosis of malignant soft tissue tumors of the head and neck should consider CCS in combination with pathological and genetic tests.

INTRODUCTION

Clear cell sarcoma (CCS) is a rare soft tissue tumor that is highly malignant, most of which originate from deep soft tissue at the distal end of the tendon and aponeurosis of the extremities, rarely occurring in the head and neck^[1], and most of its clinical manifestations are painless masses. CT is helpful to find bone changes, MRI can help determine the lesion scope, imaging findings have certain characteristics, can play an auxiliary role, diagnosis still requires pathological and genetic testing. This paper describes the radiographic findings of head and neck CCS in a young male patient.

CASE PRESENTATION

Chief complaints

A 33-year-old male hospitalized with episodic headache for more than 4 years.

History of present illness

4 years ago, the patient had a sudden headache without obvious inducement, presenting dull pain radiating to the left shoulder, lasting for about 10 h each time, accompanied by nausea and vomiting, which could be relieved after rest, without any symptoms such as disturbance of consciousness, unstable walking and numbness of limbs.

History of past illness

The patient was in good health prior to the present illness.

Personal and family history

The patient had no previous or family history of similar illnesses.

Physical examination

The patient's neurological examination was normal and other physical examination showed no abnormalities.

Laboratory examinations

Tumor marker examination showed that cancer antigen 125 was 336.20 U/ mL (normal range ≤ 35 U/ mL), and carbohydrate antigen 19-9 was 13 414.00 U/ mL (normal range ≤ 37 U/mL).

Imaging examinations

Head and neck CTA showed a massive soft tissue mass at the left paraspinal level of 1-2, with a clear boundary (Figure 1A). The mass penetrated inward through the foramen magnum into the spinal canal of the cranio-cervical junction to the front of the cervical spinal cord (Figure 1B). The artery supplying blood to the tumor was tortuous.(Figure 1C). On enhanced scan, the lesion size was 4.0cm × 4.7cm × 3.4cm (anterior and posterior diameter × left and right diameter × upper and lower diameter), showing uneven and obvious enhancement. There was a curved blood supplying artery in the lesion, locally surrounding the left vertebral artery and adjacent bone invasion. MRI examination showed that compared with adjacent muscles, the lesion showed isosignal on T₁WI (Figure 2A), slightly higher signal on T₂WI (Figure 2B), high signal on Tirm fat inhibitory sequence, and significantly higher signal on DWI (Figure 2C). The enhancement showed uneven and obvious enhancement, and the lesion invaded the anterior medulla oblongata through the left atlantoaxial foramina (Figure 2D) with a clear boundary. The protrusion of the mass locally compressed cervical spinal cord on T₂WI(Figure 2E).

Finally, postoperative histopathology showed ovoid tumor cells separated by fibrous tissue into nests or bundles, with atypical mitosis (Figure 3A). Immunohistochemistry:S-100 protein (S-100,+), anti-CD56 antibodies(CD56 ,stove +), synaptophysin (Syn ,+),epithelial membrane antigen(EMA ,±), Ki 67 (40%+), SMARCB1 protein (INI-1 ,+),SRY-related high-mobility group box 10(SOX 10 ,+), human melanin black-45(HMB45 ,+),melanocyte marker (MelanA,+)(Figure 3B and C). Fluorescence in

situ hybridization of molecular pathology: 100 tumor cells were counted, and 54% of the positive cells were in line with EWSR1 gene break(Figure 3D).

FINAL DIAGNOSIS

Combined with pathology and genetic examination, the patient was diagnosed as primary clear cell sarcoma of soft tissue of head and neck.

TREATMENT

After admission, the patient underwent lesion resection of the foramen magnum by his attending physician,Guang-Yi Jiang. To the left of the midline, cut the scalp, skin and muscle, cervical vertebra retractor open, exposed the occipital scales, visible foramen magnum area tumor tissue, tumor gray red, toughening, erosion of bone, good blood supply, due to large tumor size, the tumor tissue was partially excised. The bone flap was reduced and connected and fixed with a titanium plate. An external drainage tube was placed subcutaneously, and the muscle group was sutured. The scalp was sutured layer by layer and then pressed and bandaged. Postoperative pathological examination was carried out. The patient was safely returned to the ward after surgery.

OUTCOME AND FOLLOW-UP

The patient showed no neurological abnormalities after surgery. After 2 mo, the patient received radiotherapy followed by antirotinib targeted therapy. After 6 mo,the patient was lost to follow-up by telephone.

DISCUSSION

CCS is a rare and highly invasive soft tissue tumor, accounting for about 1% of soft tissue sarcomas [2]. The average age of onset of this disease is 35 years old [3], and the most common site is deep soft tissue at the distal tendons and aponeurosis of the extremities, especially the feet and knees. CCS originating in the head and neck is extremely rare. More than half of patients with CCS present with a slow-growing,

painless mass that is difficult to diagnose early. In this case, the patient had chronic headache, which was related to the compression of cervical spinal cord by the larger tumor size. At the same time, cancer antigen 125 and carbohydrate antigen 19-9 was elevated in the patients. Studies have shown that cancer antigen 125 may be a useful marker for the diagnosis of epithelioid sarcoma, and elevated carbohydrate antigen 19-9 is often indicative of tumors of epithelial origin [4]. However, pathologically, the origin of CCS tissue is unclear, and there are no reports about cancer antigen 125, carbohydrate antigen 19-9 and CCS. Microscopically, there are uniformly arranged nests of dense transparent cells and scattered multinucleated giant cells separated by fibrous tissue septum, which is usually contiguous with tendon or aponeurosis. Immunohistochemistry is usually positive for S-100, HMB45, MelanA and Vimentin. The morphological and immunohistochemical characteristics overlap with those of malignant melanoma, but there are differences in genetic changes between the two. CCS mainly involves periodic balanced translocation of EWSR1 and ATF1/CREB1 genes [5]. Risk factors for CCS include ionizing radiation, exposure to chemical substances (vinyl chloride, arsenic), and chronic tissue irritation (lymphedema, foreign body implantation). However, the patient in this case was previously healthy and had no related risk factors or family genetic history. The treatment of CCS is mainly surgical treatment, but it is difficult to achieve extensive resection due to its invasive characteristics, and it is prone to local recurrence and distant metastasis after surgery, with poor prognosis. Currently, the effect of radiotherapy and chemotherapy on the lesions has not been clarified [6]. Literature has reported that the 5-year and 10-year survival rates of CCS are 60%-70% and 40%-50%, respectively [7].

The imaging findings of CCS in the head and neck are rarely reported. CT examination of this case showed uniform soft tissue density without calcification and necrosis. There is also large compression of spinal cord, stronger invasive growth within the spinal canal, adjacent bone dissolve bony destruction, no hardening, no periosteal reaction. Tortuous vessels can be seen in the lesion of arterial stage, locally surrounding the left vertebral artery. The signal was uniform on MR plain scan, equal-intensity on

T₁WI, slightly higher intensity on T₂WI, and significantly higher intensity on DWI. The lesion was perforated and invaded into the anterior medulla oblongata through the left atlantoaxial foramina. The enhanced scan showed uneven and obvious enhancement, which was consistent with the characteristics of dense tumor cells and abundant blood vessels in pathological tissue. The areas with low enhancement may be related to the slow clearance of contrast agent caused by fibrous tissue diaphragm.

CCS in the head and neck should be differentiated from neurofibroma, malignant schwannoma and hemangioma. Neurofibromas are usually solid, with single or multiple cystic changes in the center. There is a hyperactive ring on T₂WI and nerve travel from the tumor. On CT, malignant schwannoma is usually an inhomogeneous mass with unclear boundary, accompanied by bone destruction and facial nerve canal enlargement. On MRI, it shows isosignal on T₁WI and hypersignal with linear septal hyposignal on T₂WI, with cystic lesions and moderate enhancement. Hemangiomas occur in the head and neck of infants and are more common in women. Calcified phlebolith can be seen in the tumor. The density of plain CT scan is similar to that of muscle. On MRI, the signal intensity on T₂WI was uneven and showed significant enhancement.

CONCLUSION

As shown in this case, the clinical manifestations of CCS in the head and neck are not typical, and the imaging manifestations have certain characteristics, which are helpful for the diagnosis of the disease. The diagnosis still requires pathological examination and genetic testing.

7%

SIMILARITY INDEX

PRIMARY SOURCES

- 1

www.wjgnet.com
Internet

32 words — 2%
- 2

Wen-Peng Huang, Li-Na Zhu, Rui Li, Li-Ming Li, Jian-Bo Gao. "Malignant giant cell tumor in the left upper arm soft tissue of an adolescent: A case report", World Journal of Clinical Cases, 2021
Crossref

30 words — 2%
- 3

Yujiao Ye, Ruiqi Tang, Beibei Liu, Yi Li, Yunlu Mo, Yong Zhang. "Prenatal diagnosis and multidisciplinary management: a case report of congenital granular cell epulis and literature review", Journal of International Medical Research, 2021
Crossref

14 words — 1%
- 4

Takamura, Saori, and Yuichi Teraki. "Case of clear cell sarcoma in the left buttock in which serum neuron-specific enolase was a useful marker for monitoring disease progression", The Journal of Dermatology, 2015.
Crossref

13 words — 1%
- 5

pastajesce.it
Internet

12 words — 1%
- 6

www.ncbi.nlm.nih.gov
Internet

11 words — 1%

7 Shaohua Hu, Zhenli Zhao, Zhisheng Wan, Weizhen Bu, Songqiang Chen, Yiqun Lu. "Chemotherapy Combined With Surgery in a Case With Metanephric Adenoma", Frontiers in Pediatrics, 2022 10 words — 1%

Crossref

8 editorialrestauro.com.mx 10 words — 1%

Internet

EXCLUDE QUOTES ON
EXCLUDE BIBLIOGRAPHY ON

EXCLUDE SOURCES < 1%
EXCLUDE MATCHES < 10 WORDS