

76353_Auto_EditedC.docx

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 76353

Manuscript Type: SYSTEMATIC REVIEWS

Assessment tools for the differential diagnosis of neglect: Focusing on egocentric neglect and allocentric neglect

Lee SH *et al.* Assessment tools for neglect

Sang-Hyeok Lee, Byeong-Chan Lim, Chan-Young Jeong, Jun-Hyeok Kim, Woo-Hyuk Jang

Abstract

BACKGROUND

There are very few studies on the differential diagnosis between egocentric neglect (EN) and allocentric neglect (AN).

AIM

To investigate the overall trend of the previously developed assessment tools by conducting a descriptive review of the studies on assessment tools that can perform a differential diagnosis of EN and AN.

METHODS

The data were collected by using databases such as Google Scholar, PubMed, and ScienceDirect. The most commonly used search terms were “neglect”, “stroke”, “egocentric neglect”, and “allocentric neglect”.

RESULTS

Among them, a total of 7 studies that met the inclusion criteria were selected and analyzed. We were able to confirm the research process, test method, and differential diagnosis criteria of the 7 presented assessment tools from 4 studies on paper-based tests and 3 studies on computerized tests. The majority of the tests were carried out *via* the cancellation method using stimuli such as everyday objects or numbers. EN distinguished the left from right based on the test paper, while AN distinguished the left from right based on stimuli. In order to perform differential diagnosis, the difference in the number of left and right responses or non-responses was used based on the EN and AN criteria

CONCLUSION

It was confirmed that all 7 assessment tools can effectively perform differential diagnosis of EN and AN. This study may provide important data that can be used in clinical practice for differential diagnosis and future intervention planning for neglect patients.

Key Words: Egocentric neglect; Allocentric neglect; Brain damage; Assessment tools; Cancellation

Lee SH, Lim BC, Jeong CY, Kim JH, Jang WH. Assessment tools for the differential diagnosis of neglect: Focusing on egocentric neglect and allocentric neglect. *World J Clin Cases* 2022; In press

Core Tip: The ¹ purpose of this study is to investigate the overall trend of the previously developed assessment tools by conducting a descriptive review of the studies on assessment tools that can perform a differential diagnosis of egocentric neglect (EN) and allocentric neglect (AN). It was confirmed that all 7 assessment tools can effectively perform differential diagnosis of EN and AN. This study may provide important data

that can be used in clinical practice for differential diagnosis and future intervention planning for neglect patients.

INTRODUCTION

Neglect is a neurological deficit due to brain damage resulting in difficulty identifying information input from the opposite direction^[1,2]. It is the most frequent serious sequelae following right hemisphere damage^[3]. The main symptom of brain damage is difficulty in recognizing objects or people in the opposite space despite having adequate sensorimotor ability^[4]. These symptoms make it difficult for a person to use their eyes, arms, and legs to search within a neglected space^[5]. They also require assistance for independent daily life due to risk of secondary accidents including falls^[6]. Neglect is classified into sensory neglect and motor neglect based on deficit type, and **personal neglect**, **peri-personal neglect**, and **extra-personal neglect** based on the distance of occurrence^[7]. Due to various symptoms, neglect causes delays in rehabilitation treatment and recovery^[8].

In 2001, Ota *et al*^[9] conducted a study for the development of an assessment tool that can differentiate between two new types of neglect symptoms. The first type of neglect is egocentric neglect (EN), which focuses on an individual and neglects information in the opposite side of the brain damage. The second type of neglect is allocentric neglect (AN), which neglects information in the opposite side of the brain damage, regardless of the object's location^[9,10]. EN is also known as viewer-centered neglect, whereas AN is known as stimulus-centered neglect^[11]. The research of Ota *et al*^[9] led to the development of the apples test and the broken hearts test for better differentiation of the two types of neglect^[12,13].

A study that measured language, memory, number, praxis, extinction, and controlled attention confirmed the difference between EN and AN symptoms. EN patients showed a lower performance in the memory domain, while AN patients showed a lower performance in all other domains^[12]. AN also has a more adverse effect on daily life performance than EN^[14], and AN patients recovered at a slower and more difficult rate than that of the EN patients^[15]. A new treatment method is deemed necessary as the existing neglect treatment had no effect on AN^[16].

Differentiation is important for accurate diagnosis and confirmation of various symptoms in patients with neglect. This is essential for establishing an effective intervention^[7,17]. Studies have been conducted to systematically review treatments, effects, and assessment tools for neglect^[5,18-22], but none have examined the assessment tools that can effectively differentiate between EN and AN.

The purpose of this study is to review the assessment tools that can differentiate between EN and AN, and investigate the overall trend of the Ota test and newly developed assessment tools by analyzing various studies.

MATERIALS AND METHODS

Studies based on the articles on assessment tools that can differentiate between EN and AN were used for this study, such as Google Scholar, PubMed, and ScienceDirect. The search keywords used were “neglect”, “stroke”, “egocentric neglect”, “allocentric neglect”, “viewer-centered neglect”, “stimulus-centered neglect”, “test”, “evaluation”, and “assessment”. The article search yielded 290 articles, among which 7 were selected, excluding duplicated studies that met the exclusion criteria (Figure 1 and Table 1).

RESULTS

Ota *et al*^[9] conducted a study in order to develop the Ota test for the differential diagnosis of EN and AN in 2 stroke patients with neglect following right hemisphere damage. In the development process, the two sub-assessments performed were circle discriminative cancellation (CDC) task and triangle discriminative cancellation (TDC) task. All tests were performed on an A3 paper. For the CDC task, 60 stimuli consisting of complete circle forms (20 stimuli, complete targets) and incomplete circle forms (40 stimuli: 20 left incomplete, 20 right incomplete, incomplete targets) were randomly scattered and placed. TDC task is similar to CDC task, except that the stimuli are in the form of triangles (Figure 2A). This test, which had no time limit, required the subjects to draw a circle to represent complete stimuli and a line to represent incomplete stimuli. Each test began with the paper presented in an upright position, followed by a re-test in

which the paper was presented upside down. One test consisted of four trials (two CDC tasks and two TDC tasks). Results were obtained from all subjects who performed the second test similarly on a different day. The diagnostic methods suggested in this study are as follows. First, EN diagnosis was presented as an omission error by removing a circle in the complete stimuli on the left side of the test paper. Next, AN diagnosis was presented as a case in which the incomplete stimuli on the left were judged as complete stimuli and marked with a circle (false positive response), regardless of the location on the test paper. The study established differentiation by showing that subject 1 was EN and subject 2 was AN^[9]. The limitations of this study are as follows. First, generalization may be difficult due to the small number of study participants. Second, the date interval between the tests and the reason for the four trials in this study was not clearly stated. Third, the severity of neglect cannot be examined because of the missing cut-off score. Therefore, it can only be used to distinguish EN from AN. Lastly, since there were 2 patients with left neglect, the study only showed the left-centered diagnostic methods for EN and AN. However, the diagnostic method for right neglect was not described.

Bickerton *et al*^[12] compared the apples test, which can perform a differential diagnosis of EN and AN, with the star cancellation test, a standardized neglect assessment tool, in a validation study involving the experimental group (25 stroke patients) and the control group (86 normal subjects). All tests were conducted on an A4 paper, and 150 stimuli consisting of complete apple-shaped stimuli (50 stimuli, complete targets) and incomplete apple-shaped stimuli (100 stimuli: 50 left incomplete, 50 right incomplete, incomplete targets) were randomly scattered (Figure 2B). The test paper was divided into 10 invisible cells (5 columns and 2 rows). Each cell received 15 apple-shaped stimuli (3 large apples and 12 small apples), including complete apple-shaped stimuli and left or right incomplete apple-shaped stimuli. The subjects were specifically instructed to mark only the complete apple shape, regardless of size. The test, which had a time limit of 5 min, was performed with a simple preliminary test (up to 2 times) to familiarize the subjects with the test method. The diagnostic methods and cut-off scores proposed in the study are as follows. First, EN is diagnosed by comparing

the correct answers in the left and right columns of the test paper. If the value obtained by subtracting the number of complete apple-shaped stimuli in the left cells from the selected number of complete apple-shaped stimuli in the right cells exceeds 2, it is presented as left side EN, and if it is less than -2, it is presented as right side EN. However, the middle 2 out of 10 cells were not used for scoring. Second, the difference in false positive responses based on the stimuli is used to diagnose AN. If the value obtained by subtracting the number of selected right incomplete apple-shaped stimuli (right incomplete targets) from the number of selected left incomplete apple-shaped stimuli (left incomplete targets) exceeds 1, it is presented as left side AN, and if it is less than -1, it is presented as right side AN. According to the study result, 5 subjects of the experimental group had EN, 2 had AN, and 5 had both EN and AN, thereby suggesting the possibility of differentiation. The apples test was found to be as sensitive and highly validated as the star cancellation test^[12]. However, this study had limitations. First, the study had a relatively small number of subjects. Second, the preliminary test did not mention the practice paper. Third, a time limit was set for the test, but it was not used in the differential diagnosis process. Lastly, exact figures were not presented during the apples test validity verification process.

Demeyere *et al*^[13] conducted a study involving the experimental group (208 stroke patients) and the control group (148 normal subjects) in order to develop the Oxford cognitive screen (OCS) to effectively measure cognitive function. The broken hearts test is a sub-test of OCS that can distinguish between EN and AN. During the development process, the broken hearts test sensitivity was compared to the star cancellation test, a standardized neglect evaluation tool. All tests were conducted on A4 paper, and 150 stimuli consisting of complete heart-shaped stimuli (50 stimuli, complete targets) and incomplete heart-shaped stimuli (100 stimuli: 50 left incomplete, 50 right incomplete, incomplete targets) were randomly scattered and placed. In the test methods, the subjects were instructed to strike through the complete heart-shaped stimuli, regardless of heart size. The test, which had a time limit of 3 min, was performed after test method familiarization *via* a simple preliminary test. The diagnostic methods and cut-off scores

proposed in the study are as follows. First, EN is compared by comparing the correct answers in the test paper's left and right columns. If the value obtained by subtracting the number of complete heart-shaped stimuli in the left cells from the selected number of complete heart-shaped stimuli in the right cells exceeds 3, it is presented as left side EN, and if it is less than -3, it is presented as right side EN. However, the middle 2 out of 10 cells were not used for scoring. Next, the difference of false positive responses based on the stimuli is used to diagnose AN. If the value obtained by subtracting the number of selected right incomplete heart-shaped stimuli (right incomplete targets) from the number of selected left incomplete heart-shaped stimuli (left incomplete targets) exceeds 1, it is presented as left side AN, and if it is less than -1, it is presented as right side AN. Based on the study result, 25% of the experimental group had EN, 11.9% had AN, and 13.6% had both EN and AN. The broken hearts test validation result was also very high at 94.12%^[13]. Although a test time limit was set, it was not used in the differential diagnosis process.

Mizuno *et al*^[24] conducted a study involving the experimental group (3 stroke patients, but only 2 had symptoms of neglect) and the control group (16 normal subjects) to develop the computerised cancellation test (CCT) for the differential diagnosis of EN and AN. During the development process, the conventional behavioral inattention test (BIT-C), a standardized neglect evaluation tool, was also implemented to verify CCT sensitivity. CCT can perform digital tests of circle discriminative cancellation task (CDC task), simple cancellation test, visuomotor search test, and visual search test through a 32-inch touch screen called TouchUbiCom; however, only the CDC task was able to perform a differential diagnosis of EN and AN. The computerized CDC task test presented in CCT is similar to the paper-based CDSC task test developed by Ota *et al*^[9]. The difference is that a person has to touch the complete targets on the screen with a finger, and the result is automatically calculated. According to the study result, subject 1 presented with EN in the experimental group, subject 2 with AN, and the remaining 1 subject without neglect did not present with either EN or AN. Furthermore, CCT was found to be as sensitive as BIT-C^[24]. The limitations of this study

are as follows. First, generalization may be difficult due to the small number of study participants. Second, neglect severity cannot be examined because there is no cut-off score; thus, it can only be used to distinguish EN from AN. Third, requiring a special touch screen for the test may increase the cost. Lastly, exact figures were not presented in the CCT sensitivity verification process.

Montedoro *et al*^[25] conducted a study involving the experimental group (35 stroke patients) and the control group (56 normal subjects) to develop the MonAmour robot test (MRT) for the differential diagnosis of EN and AN. During the development process, it was compared with the bells test, a standardized neglect evaluation tool, to verify MRT sensitivity. MRT used the REAplan® robot equipped with a test screen and a joystick (control handle). The screen is divided into 30 invisible cells (6 rows and 5 columns), and 120 stimuli were randomly placed, with 4 stimuli in each cell. The test employs human-shaped stimuli with raised hands (left, right, and both hands, targets) and 4 instrument-shaped stimuli (distractors) (Figure 2C). In 29 out of 30 cells, 4 instrument-shaped stimuli are presented, and 1 human-shaped stimuli (left hand: 30 times, right hand: 30 times, both hands: 30 times and catch trial: 10 times) and 3 instrument-shaped stimuli are rearranged. The test, which included 100 trials at 7-s intervals, required the subject to push the joystick forward when a person raising both hands appeared, and to pull the joystick back when a person with one hand (left hand, right hand) appeared. The test was performed after familiarization with the test methods through a simple preliminary test (up to 10 times). The diagnostic methods suggested in this study are as follows. First, EN is diagnosed by comparing the mean reaction time on the right area or the number of non-responses (omission error) when human-shaped stimuli (left, right, and both hands) are presented in the left and right columns based on the test screen. If the value obtained by subtracting the number of responses that missed the human-shaped stimuli in the right column from the number of responses that missed the human-shaped stimuli in the left column is 1 or greater, it was presented as left side EN, and if it was -1 or less, it was presented as right side EN. Meanwhile, AN is diagnosed when human-shaped stimuli (including stimuli with

either the left or right hand raised) respond in the opposite direction to the instruction (false positive response). If the value obtained by subtracting the number of opposite reactions to the human-shaped stimuli with a right hand raised (left incomplete targets) from the number of opposite reactions to the human-shaped stimuli with a left hand raised (right incomplete targets) is 1 or greater, it is presented as left side AN, and if it is -1 or less, it is presented as right side AN. Based on the study result, 19 subjects of the experimental group had EN, 2 had AN, and 8 had both EN and AN, thereby suggesting the possibility of differentiation. The verified MRT sensitivity was found to be 84% of the overall standard, confirming a high correlation with the bells test^[25]. The study's limitations include economic burden and space for installation due to the specialized high-priced robot required.

Chen *et al*^[23] conducted a study involving the experimental group (23 stroke or neglect patients) and the control group (186 normal subjects) in order to develop the 3 s spreadsheet test v2 for the EN and AN differential diagnosis. The test was conducted on a letter size paper (8.5 × 11 in) with 140 cells (10 cells per row and 14 cells per column), in which the digit strings served as stimuli. The digit strings had a minimum of 4 digits and a maximum of 9 digits, with digits 0 to 9 being listed repeatedly. The test, which had no time limit, required the subject to find the correct answer 3 (targets) in all digit strings in the cell and strike through (correct answers: 120, other distractors: 720). If the digit strings are an odd number, 3 was not placed in the middle number (Figure 2D). The diagnostic methods and cut-off scores suggested in the study are as follows. First, EN diagnosis was analyzed by the difference in omission errors based on the test paper. If the value obtained by subtracting the number of omissions of the correct stimuli in the left region from the number of omissions of the correct stimuli in the right region exceeded 3, it was presented as left side EN, and if it was less than -3, it was presented as right side EN. For AN diagnosis, the digit strings were divided in half based on the digit strings presented for each cell in the test. If the value obtained by subtracting the number of omissions of the correct stimuli in the right area from the number of omissions of the correct stimuli in the left area exceeded 5, it was presented as left side

AN, and if it was less than -3, it was presented as right side AN. Based on the study result, 3 out of 23 subjects in the experimental group had EN, 2 had AN, and 18 had both EN and AN, thereby suggesting the possibility of differentiation^[23]. The study's limitation is that the subjects may experience high test fatigue due to the 840 stimuli, which is higher compared to the other tests.

Ferraro *et al*^[26] investigated the possibility of replacing the pen and paper test with digital tests, such as Albert's test, line bisection test, and apples test built in the ReMoVES platform for the experimental group (12 neglected patients) and the control group (6 normal patients)^[26]. The ReMoVES platform is a computerized test program developed by the University of Genova, and only the apples test was able to differentiate between EN and AN among the three built-in tests. The computerized apples test presented in the ReMoVES platform is similar to the paper-based apples test studied by Mancuso *et al*^[27]. However, it requires touching the complete apple-shaped stimuli presented on the screen with a finger, and the result is automatically calculated. The study result showed that the paper-based test and computerized test produced similar results in the subject's test performance process, and they can be used interchangeably^[26]. The limitations of this study are as follows. First, personal information (*e.g.*, gender, age, and disease) was not presented for the 12 experimental groups. Finally, generalization is difficult due to the small number of study participants (Figure 2).

DISCUSSION

Comprehensive analysis of 7 assessment tools for differential diagnosis

AN has more adverse effects on cognitive function, activities of daily living, and rehabilitation speed than EN. It was also confirmed that the existing neglect treatment had no significant effect on the AN patient. Therefore, seven assessment tools that can effectively differentially diagnose EN and AN were analyzed.

First, the study results showed that cancellation test type tests were developed in the studies as the most common feature, and the test stimuli presented during the research

process were commonly encountered shapes in everyday life, such as circles, triangles, apples, hearts, numbers^[9,12,13,23-26]. Furthermore, it has been confirmed that the stimuli presented in most studies were complete forms and left or right incomplete^[9,12,13,24-26]. According to the diagnostic methods presented in this study, EN is mainly the difference in the number of correct answers in the left and right areas on the test paper or screen, and AN was presented as the difference in the number of incorrect answers on incomplete forms in the left and right areas based on stimuli (false positive response^[9,12,13,24,26]). This is thought to be presented for accurate differential diagnosis, taking into account the concept of EN neglecting information centered on the individual (self) and AN neglecting information centered on the object (stimuli). However, the MonAmour robot test studied by Montedoro *et al*^[25] showed both the EN and AN diagnostic methods presented as a difference in the number of incorrect answers (non-response, opposite response). In comparison with the other tests, it is considered to be the diagnostic method designed according to the test characteristics, in which the stimuli are newly rearranged for each trial, and must respond to both correct and incorrect answers. According to the study conducted by Chen *et al*^[23] on '3s spreadsheet test v2', both the EN and AN diagnostic methods were presented as the difference in the number of omission of correct answers (omission errors). In comparison with the other tests, it is considered to be the diagnostic method designed according to the test characteristics, in which the stimuli used are presented only as the correct stimuli (targets) and other stimuli (distractors). Furthermore, four articles reviewed the pencil and paper tests^[9,12,13,23], and three articles reviewed the digital^[24-26]. All of the seven presented assessment tools can effectively perform a differential diagnosis of EN and AN, and Ferraro *et al*^[26] confirmed that the paper-based test and the digital test are interchangeable with each other.

The limitations of the studies are as follows. First, they are difficult to generalize due to the small number of study subjects^[9,24,26]. Second, although the diagnostic criteria for EN and AN were presented, a cut-off score to evaluate the severity of neglect was not presented^[9,24]. Third, accurate figures were not presented in the assessment tool

verification process^[12,24]. Fourth, the diagnostic criteria for EN and AN were only focused on the left side, so the diagnosis criteria for right neglect were not presented^[9,24]. Fifth, although the time limit of the test was set, it was not used to determine the degree of neglect^[12,13]. Lastly, there are issues in regard to space due to cost and installation location for the digital tests^[24,25]. In future studies, the following are recommended to complement the limitations of the previous studies: (1) Conduct research with a sufficient number of subjects; (2) Provide a cut-off score required in order to confirm the severity of neglect; (3) Suggest the severity according to the type of neglect by using the time limit of the test; and (4) Consider economic and spatial problems caused by the equipment required for the digital tests.

CONCLUSION

In this study, we reviewed the literature studying assessment tools for the differential diagnosis of EN and AN. Accordingly, 7 tests (pencil and paper: 4 times, digital test: 3 times) were tested and effective, and differential diagnosis can be conducted when the difference in response to various stimuli is used.

In conclusion, these results might offer an easier differential diagnosis of AN, and appropriate intervention at an early stage of injury. In the case of a patient with both EN and AN, it might be possible to seamlessly modify the detailed direction of intervention by determining the improvement of neglect *via* continuous assessment. Finally, the data discussed in this work may provide guidance for developing more convenient and various differential diagnosis methods and new intervention methods for AN as diverse as EN.

ARTICLE HIGHLIGHTS

Research background

There are various types of neglect, and the symptoms are also diverse. However, review studies on the differential diagnosis of the relatively recently discovered allocentric neglect (AN) and the already known egocentric neglect (EN) are lacking.

Research motivation

Compared to EN, AN has a more adverse effect on daily life, and the recovery rate is slower. In addition, the conventional treatment of EN is not effective in the treatment of AN. Therefore, the distinction between AN and EN is very important.

Research objectives

By reviewing the studies on differential diagnosis, we will try to find out the overall trend of the newly developed evaluation tool.

Research methods

A literature search on differential diagnosis was conducted through a search according to appropriate keywords.

Research results

Seven relevant papers were collected (paper-and-pencil 4, digital 3).

Research conclusions

All tests were effective in differential diagnosis of EN and AN.

Research perspectives

A more effective intervention will be possible through an accurate differential diagnosis. It is hoped that more treatments for AN will be developed in the future.

ORIGINALITY REPORT

0%

SIMILARITY INDEX

PRIMARY SOURCES

- 1

"Thursday July 6, 2006 11:30?13:30 Hall 5A Discussion Group Session What future for radiosurgery in epilepsy surgery?", Epilepsia, 12/2006
Crossref

9 words — < 1%
- 2

www.slideshare.net
Internet

8 words — < 1%

EXCLUDE QUOTES OFF
EXCLUDE BIBLIOGRAPHY OFF

EXCLUDE SOURCES OFF
EXCLUDE MATCHES OFF