

74064_Auto_Edited-check.docx

Clinical and Translational Research**Effects of targeted-edited oncogenic insulin-like growth factor-1 receptor with specific-sgRNA on biological behaviors of HepG2 cells**

Yao M *et al.* IGF-1R in HCC

Min Yao, Yin Cai, Zhi-Jun Wu, Ping Zhou, Wen-Li Sai, De-Feng Wang, Li Wang, Deng-Fu Yao

Abstract**BACKGROUND**

Insulin-like growth factor-1 receptor (IGF-1R) was over- expressed in hepatocellular carcinoma (HCC). However, the relationship between *IGF-1R* activation and HCC progression remains identified.

AIM

To investigate the effects of editing *IGF-1R* on biological features of HCC cells.

METHODS

Immunohistochemistry analyzed the expressions of IGF-1R and P-glyco protein (P-gp) in HCC tissues and their distal non-cancerous tissues (Non-Ca). *IGF-1R* was edited with Crispr/Cas9 system, screened specific sgRNAs, and then transfected into HepG2 cells. CCK-8, scratch wound test detected cell proliferation, migration, invasion, and transwell assays, respectively. Alterations of IGF-1R and P-gp were confirmed by

Western blotting. Alterations of anti-cancer drug IC₅₀ values were analyzed at the cell level.

RESULTS

The positive rates of IGF-1R (93.6%, $\chi^2 = 63.947$) or P-gp (88.2%, $\chi^2 = 58.448$) were significantly higher ($P < 0.001$) in the HCC group than those (36.6% in IGF-1R or 26.9% in P-gp) in the Non-Ca group. There were positively correlated between high IGF-1R and P-gp expression, and they were associated with hepatitis B virus infection and vascular invasion of HCC. Abnormal expressions of circulating IGF-1R and P-gp were confirmed and associated with HCC progression. Biological feature alterations of HCC cells transfected with specific sgRNA showed IGF-1R expression down-regulation, cell proliferation inhibition, cell invasion or migration potential decreasing, and enhancing susceptibility of HepG2 cells to anti-cancer drugs.

CONCLUSION

Edited oncogenic *IGF-1R* was useful to inhibit biological behaviors of HepG2 cells.

Key Words: Hepatocellular carcinoma; Insulin-like growth factor-1 receptor; Synergistic effects; Multi-drug resistance; Growth inhibition; Biological behaviors

Yao M, Cai Y, Wu ZJ, Zhou P, Sai WL, Wang DF, Wang L, Yao DF. Effects of targeted-edited oncogenic insulin-like growth factor-1 receptor with specific-sgRNA on biological behaviors of HepG2 cells. *World J Clin Cases* 2022; In press

Core Tip: Abnormal expression of insulin-like growth factor-1 receptor (IGF-1R) was associated with hepatocellular carcinoma (HCC). IGF-1R level was significantly higher in HCC more than that in their noncancerous tissues. Circulating IGF-1R was continue to increase from benign liver disease to HCC. Down-regulating expression of IGF-1R with specific sgRNA was markedly affected on biological behaviors of HCC cells,

including inhibiting cell proliferation, decreasing cell migration or invasion potential, increasing cell apoptosis and enhancing cell susceptibility to anti-tumor drugs, indicated that oncogenic IGF-1R should be a promising targeted-molecule for HCC therapy.

INTRODUCTION

⁶ Hepatocellular carcinoma (HCC) still is one of the most common malignant tumors worldwide, and its increasing incidence might be associated with nonalcoholic fatty liver disease because of effective control of hepatitis B virus (HBV) or hepatitis C virus infection^[1-3]. The limited effective therapies available for advanced HCC patients are considered with aberrant gene transcriptions that govern crucial signal pathways, high migration or invasion potential, or multi-drug resistance (MDR) formation^[4]. Previous studies have indicated that the related-signal molecules of the insulin-like growth factor (IGF) axis play a crucial role in malignant transformation of hepatocytes or HCC progression. Recently, the genomic sequencing studies of IGFs, type 1 IGF receptor (IGF-1R), and IGF binding proteins (IGFBPs) have confirmed that IGFs regulate multiple physiological processes, including mammalian development, metabolism, and growth^[5,6]. IGF-1R attributes to high-energy intake, cell proliferation, or apoptosis, and promotes mitogenic pathway activation *via* tyrosine kinase, conformational changes, tyrosine autophosphorylation, and insulin-receptor substrate proteins^[7,8], suggesting that high IGF-1R level might affect ineffective treatment of HCC^[9,10].

High IGF-1R expression and MDR with increasing P-glyco protein (P-gp) are major obstacles to the successful treatment of HCC with chemotherapy^[11,12]. High IGF-1R mediated mitogenic, differentiating, and anti-apoptotic features were be found in HCC cells but not in mature hepatocytes^[13]. The P-gp, as a member of the superfamily of ATP-binding cassette transporters, is encoded in *MDR 1*. MDR emergence is still a complex problem for effective treatment, and a high P-gp level is closely related to MDR in HCC. Therefore, it is essential to establish a new technology of MDR reversal, selectively inhibit P-gp or interfere with the activation of related genes to improve the

curative effect or prognosis of HCC treatment^[14,15]. Although high IGF-1R was associated with malignant transformation of hepatocytes^[16], *IGF-1R* activation could be inhibited by small hairpin RNA (shRNA)^[9]. However, the editing of *IGF-1R* on biological behaviors or MDR of HCC remain be explored. This study aimed to investigate the clinicopathological features of IGF-1R or P-gp expression in HCC, further analyze *IGF-1R* on effects of biological behaviors of HepG2 cells, and synergistic role with anti-cancer drugs on reversal MDR of HCC.

MATERIALS AND METHODS

HCC tissues

According to the self-controlled method, a total of 93 pairs of HCC- and their distal non-cancerous (3 cm to cancer, Non-Ca) tissues were collected after the HCC patient operation from Feb 2015 to Aug 2016. The prior written informed consent was obtained from all patients according to the Helsinki Declaration of World Medical Association. This study was approved by the Ethics Committee permission (TDFY2013008) at the Affiliated Hospital Nantong University, China. Based on medical records, the cases were 78 males and 15 females with 35-80 years old (average 55.8 years \pm 16.1 years). There were 74 cases with single tumor and 19 cases with multiples; 22 cases with tumor size > 5.0 cm and 71 cases \leq 5.0 cm; 36 cases with and 57 without vascular invasion. Differentiation degree were well ($n = 21$), middle ($n = 49$), and poor ($n = 23$) on the Edmondson grading system. Clinical staging was 58 cases at I-II and 35 at III-IV on the tumor-node-metastasis (TNM) classification of the International Union against Cancer. There were 64.5% (60 of 93) cases with positive HBV surface antigen and 71.0% (66 of 93) with liver cirrhosis. Serum α -fetoprotein (AFP) levels were 34 cases with ≥ 400 $\mu\text{g/L}$ and 59 with < 400 $\mu\text{g/L}$. All cases had regular follow-up from operation to death until Aug 2021. Criteria of HCC diagnosis were set by the Chinese Collaborative Liver Cancer Research Group^[17].

Tissue microarray and immunohistochemistry

Liver sections from HCC or Non-Ca tissues were made for constructing tissue microarray (TMA). They were boiled for antigen retrieval in pH 6.0 citrate buffer and incubated for 2 h with 1st mouse anti-IGF-1R (Abcam, United Kingdom) or anti-P-gp antibodies (Santa Cruz, United States), washing with phosphate-buffered saline (PBS), and then incubated with horseradish peroxidase (HRP) conjugated goat anti-mouse IgG (Dako, Carpinteria, United States) and diaminobenzidine solution, counterstained with Hematoxylin & Xylene. Negative control was used PBS instead of 1st antibodies. Two independent pathologists examined TMA staining. Positive cells (%) were scored into: 0 (negative, 0%), 1 (weak, 1%-33%), 2 (moderate, 34%-66%), and 3 (strong, 67%-100%). Liver IGF-1R or P-gp levels were divided into low (0-1) and high (2-3) scores according to immunohistochemistry (IHC) staining^[18].

Western blotting

Purified proteins from 50 mg of liver tissue or five × 10⁴ cells were quantitatively detected using BCA assay (Beyotime, China) for the specific concentration/per mg wet liver or cell protein analysis. Phenyl-methanesulfonyl fluoride (Beyotime, China) was added in case of protein degradation. Proteins (50 µg/case) from transfected cells were separated on 10% sodium dodecyl sulfate-polyacrylamide gel electrophoresis and transferred to polyvinylidene fluoride membranes (Millipore, United States) and blocked in 5% skimmed milk at room temperature, incubated with antibodies of mouse anti-human IGF-1R or anti-β-actin (Abcam, United States), and anti-human P-gp (Santa Cruz, United States) at 4 °C overnight and goat anti-mouse HRP-conjugated IgG (Dako, United States) at room temperature for 2 h. Protein bands were captured by enhanced chemiluminescence kit (Millipore, United States) and analyzed by Quantity-one software (Bio-Rad, United States).

IGF-1R-sgRNA plasmid and In vitro transfection

Human LO2 cells, HCC Bel-7404, Bel-7402, HepG2, and HeH-7 cell lines were obtained from the Chinese Academy of Sciences (Shanghai, China). HepG2 cell lines were grown

in DMEM (Hyclone, United States) containing 10% FBS (Gibco, United States) at 37 °C in a humidified atmosphere of 5% CO₂. Three candidate sgRNAs targeting human IGF-1R (NM000875)^[22] with Crispr/Cas9 system along with vectors (Lenti-CAS9-puro and Lenti-sgRNA-EGFP) were designed and constructed by GeneChem (Shanghai, China). Their sequences were listed as follows: sgRNA1: 5'-TCAGTACGCCGTTTACGTCA-3' (PCA00469); sgRNA2: 5'-TGTTTCCGAAATTTACCGCA-3' (PCA00470); and sgRNA3: 5'-GGCTCTCTCCCCGTTGTTCC-3' (PCA00471). HepG2 cells were divided into blank control (Con), negative sgRNA-CON244 (sgRNA-Neg), and IGF-1R-sgRNA (sgRNA) groups. Cell transfection, briefly, five × 10⁴ HepG2 cells at 50% confluency in 6-well plates were transfected with Lenti-CAS9-puro vector for 48 h and then screened by puromycin. Following harvest, Lenti-sgRNA-EGFP with different sgRNAs against IGF-1R was transfected into Cas9-transfected cells. Then the transfection efficacy was analyzed under a fluorescence microscope (Olympus, Japan).

SURVEYOR assay

DNA of samples with sgRNAs was harvested for PCR and nuclease digestion for analysis using SURVEYOR Mutation Detection Kits (IDT, United States)^[19]. DNA fragments were electrophoresed on 2% agarose gel.

Cell proliferation and immunofluorescence staining

Cell proliferation were analyzed by CCK-8 assay using the Cell Proliferation Kit I (Dojindo, Japan). According to the manufacturer's instruction (Beyotime, China), the dark blue formazan crystal was formed. Removed the medium carefully, added dimethyl sulfoxide, and detected the A values with Multi-Detection Microplate Reader (Bio-Tek, United States) at 570 nm. Cells (2 × 10⁵) or transfected cells after 48 h were placed on the bottom with glass coverslips for 24 h, fixed in 4% paraformaldehyde in PBS (pH 7.5). Cover slips were first immersed for one h in a blocking solution containing 5% bovine serum albumin in PBS and incubated overnight at 4 °C with mouse antibodies against IGF-1R (CST, United States). DNA was counterstained with

4',6-diamidino-2-phenylindole and observed under inverted fluorescence micro-scope. Levels of IGF-1R expression in culture supernatant were quantitatively detected with an ELISA kit at 450 nm.

Cell apoptosis and flow cytometry

Effective apoptosis of transfected cells was evaluated by the PE-labeled Annexin-V/7-AAD assay according to the manufacturer's protocol. In brief, HepG2 cells (2×10^5) were collected, rinsed twice in cold PBS, and added with binding buffer; then, apoptosis rate was analyzed. After 24 h, they were collecting (including dead cells in culture medium) and processing cells according to the instructions of Annexin V-FITC/PI Double stain apoptosis detection kit (#4101-2, Bestbio, Shanghai, China). The fluorescence intensity on the flow cytometer was collected to immediately check apoptotic rate (Cytotflex Beckman, China). Three independent experiments were performed.

Transwell assay

Transwell chambers (8.0 μ m pore size, Corning, United States) were adopted for testing cell migration and invasion assays. Cells were seeded in the upper chamber after transfection for 24 h. Lower chambers were added 600 μ L medium. After 24 h, cells were fixed with 4% paraformaldehyde and stained with 0.1% crystal violet. The upper chamber cells were wiped with cotton. During the invasion assay, the upper chambers needed to be pretreated with a 0.3% Matrigel matrix (#356234, Corning, United States) and incubated at 37 °C 4 h. Three independent experiments were performed.

Cell migration or invasion assay

Quantitative and qualitative analysis of HepG2 cells migration were assessed by *in vitro* Transwell assay with modified Boyden Chambers and Transwell-coated Matrigel membrane filter (BD Biosciences, Bedford, MA, United States). Cells (5×10^3) from Con, Neg, and MiR groups ($n = 3$ /per group) were plated onto the upper compartment

without FBS or 10% FBS in the lower chamber as a chemoattractant. Fluorescent images of nuclear Hoechst staining (10 µg/mL) were captured at 24 h of incubation in a 5% CO₂ humidified at 37 °C. Percentages of migrated cells in each group were counted from 10 random microscope fields for each sample in 3 independent experiments. The modified Boyden Chambers without the Transwell-precoated Matrigel membrane filter in the above method was performed for cell migration analysis.

Drug sensitivity tests

HCC cells were seeded to 96-well plates (5000 cells/well) and placed in the incubator for 24 h under cell culture conditions. The culture medium is then replaced by serum-free medium containing varying dosages of Sorafenib (Bayer Corporation, Germany) from 0.001 to 1000 µmol/L, and oxaliplatin (Jiangsu AoSaikang pharmaceutical Co. China) from 0.001 to 1000 nmol/L, where appropriate. Dose-response curve and IC₅₀ calculation were analyzed by evaluating the cell viability using the Cell Counting Kit, CCK-8 (Tojindo), 48 h after drug treatment with the nonlinear regression method by GraphPad Prism 5.0 software (San Diego, CA, United States).

Statistical analysis

Continuous variables are expressed as means ± SD. Data were analyzed by using SPSS software (version 20.0). Favorable rates and clinicopathological features of IGF-IR or P-gp expression were evaluated by χ^2 test, Wilcoxon signed-rank test in paired tissue samples; Between-group differences were assessed by *t*-test or one-way analysis of variance. A *P* value less than 0.05 value was considered statistically significant.

RESULTS

IGF-IR or P-gp with HCC patients survival

Comparative analysis of the representational IHC staining of IGF-IR or P-gp from the 93 pairs of HCC or their distal non-cancerous tissues (Non-Ca) is shown in Figure 1. According to the hepatic IGF-IR or P-gp staining, there were the stronger in the HCC

group (**Figure 1A and B**), the lighter or non-expression in the Non-Ca group (**Figure 1 A1 and B1**). Both brown positive staining particles were mainly located in the cytosol with the clearly heterogeneous distribution. Specific concentrations of IGF-1R/per mg wet tissues ($t = 23.451$, $P < 0.001$, **Figure 1 A2**) or P-gp/per mg wet tissues ($t = 19.832$, $P < 0.001$, **Figure 1 B2**) were quantitatively detected, with significantly higher expression in the HCC group more than those in the Non-Ca group. Abnormal expressions of IGF-1R (**Figure 1 A3, 130 kD**) or P-gp (**Figure 1 B3, 142 kD**) were confirmed by Western blotting, with high ratios ($P < 0.001$) from IGF-1R to β -actin (**Figure 1 A3**) or P-gp to β -actin (**Figure 1 B3**) expression in the HCC group more than those in the Non-Ca group. According to the cumulative survival curves, there were shorter survival times in HCC patients with high IGF-1R (**Figure 1C**) or P-gp (**Figure 1D**) expression.

Clinicopathological features of IGF-1R or P-gp expression

The summary of hepatic IGF-1R or P-gp expression and its scores between HCC and D-can tissues are shown in Table 1. The positive rates of IGF-1R expression were 93.6% in the HCC group, with significantly higher ($\chi^2 = 63.947$, $P < 0.001$) than that in the D-can group (36.6%). Furthermore, the higher IGF-1R with 2-3 scores in the HCC group (82.8%) was significantly higher ($Z = 9.682$, $P < 0.001$) than that in the D-can group (9.7%). Similar to IGF-1R expression, the positive rates of P-gp expression in the HCC group (88.2%) was significantly higher ($\chi^2 = 58.448$, $P < 0.001$) more than that in the D-can group (26.9%), and the higher P-gp with 2-3 scores in the HCC group was 74.2% was significantly higher ($Z = 8.941$, $P < 0.001$) more than that in the D-can group (0%). The clinicopathological features of IGF-1R or P-gp expression in HCC are shown in Table 2. The higher expression of IGF-1R in the HCC group was closely related to vascular invasion, HBV infection, and middle or poor differentiation degree, but not to age or gender of patients, AFP level, TNM stage, liver cirrhosis, and tumor size. Also, the abnormal P-gp levels in the HCC group were closely related to vascular invasion, HBV infection, poor differentiation degree, and TNM stage, but not to age or gender of patients, AFP level, liver cirrhosis, and tumor size or number.

Circulating IGF-1R or P-gp levels in liver diseases

Comparative analysis of IGF-1R and P-gp levels in sera of patients with liver diseases are shown in Table 3. Significant differences were found in circulating IGF-1R ($F = 154.501$, $P < 0.001$) or P-gp ($F = 66.182$, $P < 0.001$) levels among different liver disease groups, with high IGF-1R or P-gp levels in the HCC group more than those in the liver cirrhosis or chronic hepatitis or healthy control group. The incidences of IGF-1R over 600 pg/mL or P-gp over nine ng/mL were 86% or 75% in HCC patients, and less than 20% in patients with other diseases, with an increasing tendency from chronic hepatitis, liver cirrhosis to HCC. IGF-1R or P-gp expressions were 93.0% or 87.2% in sera of HCC patients, 25.5% or 26% in sera of cases with liver cirrhosis, and 5% or 8% in sera of cases with chronic hepatitis, respectively. A closely positive correlation ($r = 0.682$, $P < 0.001$) was found between IGF-1R or P-gp expression and liver disease progression.

Editing IGF-1R with cell proliferation inhibition

Human HepG2 cells were divided into control, sgRNA-neg, and sgRNA2 groups. Using the CRISPR/Cas9 system, human IGF-1R in HepG2 cells genetically modified and the editing IGF-1R on effects of HepG2 cell proliferation and P-gp expression are shown in **Figure 2**. The IGF-1R in HepG2 cells was the strongest expression among LO2 cells (**A & A1**), increasing P-gp levels after HepG2 cells plus drug treatment (HepG2/ADM cells, **B & B1**). Comparative analysis of the constructed-sgRNA1-3 and sgRNA-neg plasmids into HepG2 cell lines, the transfected efficiencies were 91% for sgRNA1 (**C1 & D1**), 90% for sgRNA2 (**C2 & D2**), 88% for sgRNA3 (**C3 & D3**), and 86% for sgRNA-neg (**C4 & D4**) plasmids, respectively. The IGF-1R expression in the transfected HepG2 cells were confirmed by Western blotting (**E, Upper**), and the relative ratios of IGF-1R to β -actin were 1.32 ± 0.13 in the control group, 1.14 ± 1.23 in the sgRNA-neg group, 1.01 ± 0.94 in the sgRNA1 group, 0.43 ± 0.79 in the sgRNA2 group, and 0.99 ± 0.82 in the sgRNA3 group (**E, Down**), respectively. Specific sgRNA were chosen to study further, the curves of the transfected HepG2 cell proliferation were significantly inhibited ($P <$

0.01) in a time dependence manner, and the inhibiting rate were 0.31 ± 0.08 on the 1st day, 0.42 ± 0.14 on the 2nd day, and 0.62 ± 0.21 in the 3rd day (**F & G**). Also, the decreasing P-gp level in the HepG2/ADM cells is transfected with sgRNA2 (**H**).

Effects of edited IGF-1R on the biological features of HepG2 cells

Editing IGF-1R on effects of HepG2 cell apoptosis, migration, and invasion are shown in **Figure 3**. HepG2 cells transfected with specific sgRNA2 besides inhibiting proliferation, targeting IGF-1R also directly affected the biological function of HepG2 cells by the analysis of flow cytometry. Apoptotic rates of HepG2 cells in the sgRNA2 group (56.25%) were significantly higher ($P < 0.01$, **Figure 3A Left**) than those in the sgRNA-neg group (5.98%) or control group (5.66%). Edited IGF-1R led to cell cycle arrest in G1 phase of HepG2 cells ($P < 0.01$, **Figure 3A Right**). The numbers of cell migration were 292.3 ± 28.6 in the sgRNA2 group and significantly lower ($F = 391.322$, $P < 0.001$) than those in the sgRNA-neg (564.5 ± 15.8 , $q = 35.241$, $P < 0.001$) or the control (580.5 ± 15.2 , $q = 35.214$, $P < 0.001$) group (**Figure 3B**). The numbers of cell invasion in the sgRNA2 group (59.3 ± 19.1) was significantly lower ($F = 69.510$, $P < 0.001$) than those in the sgRNA-neg (165.5 ± 24.8 , $q = 13.684$, $P < 0.001$) or control (176.1 ± 12.2 , $q = 15.102$, $P < 0.001$) group (**Figure 3C**). There was no significantly differences of cell migration or invasion between the control group and the sgIGF-1R-neg group. These data indicated that edited IGF-1R might be obviously decreasing invasion or migration potential of HepG2 cells.

Synergistic effect of sgRNA with anti-cancer drugs

The synergistic effect of specific sgRNA with anti-cancer drugs on inhibition of HepG2 cell growth is shown in Table 4. HepG2 cells with or without transfected sgRNA were exposed to sorafenib from 0.0 nmol/L to 80.0 nmol/L or oxaliplatin from 0.0 μ mol/L to 40 μ mol/L for 24 h, and cell survival rates were assessed using the CCK-8 assay. IGF-1R expression was down-regulated by sgRNA in a dose-dependent manner, especially at 20 nmol/L with high inhibiting effects on HCC cells. IC₅₀ values were 16.9 nmol/L in

the control group, 16.5 nM/L in the sgRNA-neg group, and 4.4 nmol/L in the sgRNA group. Similar to sorafenib, the specific sgRNA plus oxaliplatin had higher inhibitory effects on the cells. IC₅₀ values were 14.6 nmol/L in the control group, 14.2 nmol/L in the sgRNA-neg group, and 8.2 nmol/L in the sgRNA group. These data indicated that HCC cells transfected with specific sgRNA could be more sensitive to anti-cancer drugs.

DISCUSSION

Accumulating data of basic and clinical studies have indicated that many kinds of signaling are associated with HCC hypoxic microenvironment and oncogenesis^[20]. IGF-1R activation was related to malignant transformation of hepatocytes^[11], specific shRNA for inhibiting *IGF-1R*^[10], IGF-1R tyrosine kinase inhibitors and vitamin K1 enhancing the antitumor effects of regorafenib^[8], and miRNA-187 inhibits HCC growth and metastasis *via* targeting IGF-1R^[9] have been explored. This study confirmed high IGF-1R expression in cancerous tissues that were related to poorly differentiated degree, shorter survival, TNM stage, and lymph node metastasis of HCC; also, serum IGF-1R level in the HCC group was significantly higher than those in chronic hepatitis, liver cirrhosis or control group, indicating that IGF-1R should be an essential signaling for the formation, progression and poor prognosis of HCC^[21].

IGF-1R signaling regulates cell differentiation, organ development, and tissue regeneration during embryonic development. Also, IGF-1R as a critical molecule of the IGF axis was associated with the progression and drug resistance of HCC, and it could promote cell proliferation and activate HCC reprogramming in the liver, especially in patients with chronic liver diseases^[22]. Dysregulation of IGF-1R signaling might activate expression of HCC stemness that leads to hepatocytes malignant transformation, especially in HBV-related HCC^[14] or rat hepatocarcinogenesis with abnormal IGF-1R or CD44 activation^[11,23]. In this study, the relationship between high IGF-1R and increasing P-gp were analyzed, that both were significant obstacle to the successful treatment of HCC with chemotherapy. High IGF-1R mediated mitogenic, differentiating, and anti-

apoptotic features and P-gp was related to MDR of HCC patients, suggesting that high IGF-1R and P-gp expression might affect in the effective treatment of HCC.

Hepatic IGF-1R as a multifunctional regulatory factor plays an importanta vital role in HCC occurrence and progression. Abnormal activation of IGF-1R signaling promotes the proliferation, dissemination, and aggressive behaviors of HCC cells^[24,25]. In this study, higher IGF-1R expression is consistent with the level in HCC tissues, circulating blood, and HepG2 cells. Down-regulating IGF-1R has an anti-cancer effect through increasing apoptosis, inhibiting growth of HCC cell proliferation, decreasing migration or invasion of HepG2 cells, and cell cycle arrest in the G1 phase, because of IGF-1R as an essential transmembrane protein for the related-pathway activation *via* its tyrosine kinase^[26], suggesting that oncogenic IGF-IR could be a molecule-targeted for HCC therapy.

Acquiring resistance to chemotherapy remains a significant hurdle to effective HCC treatment^[27,28]. Biomarker P-gp as an MDR-associated protein 1 (MRP1/ABCC1) is considered a prime factor for MDR induction^[29]. The critical role of nuclear factor-kappa B (NF-κB) is to induce MDR in tumor models characterized precisely by innate or acquired MDR, particularly HCC^[4,30]. Different pharmacological approaches have been employed to reduce the expression/activation of this transcriptional factor and thus to restore chemosensitivity. Scientific evidence was found by the most significant clinical trials regarding NF-κB and new perspectives on the possibility to consider this transcriptional factor a valid drug target in neoplastic diseases^[31]. Both high P-gp and IGF-1R levels are common in HCC. In this study, the specific sgRNA plus anti-tumor drugs had higher inhibitory effects on HCC cells, and the drug IC₅₀ values were significantly decreased with inhibiting cell proliferation of HCC in the sgRNA group, indicating that HCC cells transfected with sgRNA could be more sensitive to anti-cancer drugs *via* inhibiting the NF-κB pathway^[4,32].

CONCLUSION

In conclusion, the high IGF-1R or P-gp expression has been confirmed related to the progression or therapeutic effect of HCC. Although the accurate mechanism for IGF-1R reactivation in HCC remains to be explored, specific edited oncogenic IGF-1R gene is promising for inhibiting proliferation, altering biological behaviors, or as potential modulators for reversal MDR of HCC cells. Further studies should clarify the exact relationship between up-regulating IGF-1R and MDR formation. However, IGF-1R as a critical signaling molecule of the IGF axis might be a novel effective target for inhibiting HCC growth or reversal MDR of HCC.

ARTICLE HIGHLIGHTS

Research background

⁵ Hepatocellular carcinoma (HCC) is one of the leading causes of cancer-related death, with a high incidence and mortality rate in China. ² Insulin-like growth factor 1 receptor (IGF-1R) signaling triggers cell proliferation, liver growth, and tissue regeneration during embryonic development. Unbalanced IGF-1R signaling can promote HCC cell proliferation, and regulating IGF-1R gene transcription should be useful as a potential therapy targeting HCC.

Research motivation

IGF-1R as key signaling of the IGF axis in HCC progression was investigated in sera or tissues from HBV-related HCC patients, analyzed the relationship between IGF-1R and multi-drug resistance (MDR), and edited the IGF-1R gene for down-regulating expression to confirm effects on proliferation and a potential therapeutic role for HCC cells.

Research objectives

The expressing status and clinicopathological characteristics of IGF-1R or P-glycoprotein (P-gp) were investigated in the circulating blood and tissues of HCC patients, and editing IGF-1R gene at the mRNA transcription level to observe effects on

biological behaviors HepG2 cells, and synergistic role with anti-cancer drugs on reversal MDR of HCC.

Research methods

Comparative analysis of IGF-1R and P-gp expression in tissues or sera of HCC patients were analyzed by immunohistochemistry and confirmed by Western blotting. Specific sgRNA was screened among editing IGF-1R gene with Crispr/Cas9 system and then transfected into HepG2 cells. CCK-8, scratch wound test detected HCC cell proliferation, migration, invasion, and transwell assay, respectively.

Research results

Abnormal over-expression of IGF-1R and P-gp were confirmed in tissues or sera of HCC patients, with a positive closely correlation between IGF-1R and P-gp and related to HBV infection or vascular invasion during HCC progression. HepG2 cell biological features were altered by specific IGF-1R-sgRNA with down-regulation, cell proliferation inhibition, cell invasion or migration potential decreasing and enhancing cell susceptibility to anti-cancer drugs.

Research conclusions

Based on this these studies, high IGF-1R or P-gp expression has been confirmed related to the progression or therapeutic effect of HCC. Although the accurate mechanism for IGF-1R reactivation in HCC remains be explored, specific edited oncogenic IGF-1R gene is promising for inhibiting proliferation, altering biological features, or as potential modulators for reversal MDR of HCC cells.

Research perspectives

Abnormal expression of hepatic IGF-1R level was associated with HCC progression. Inhibiting IGF-1R expression could markedly effect on biological behaviors of HCC cell

proliferation, migration or invasion, cell apoptosis, and drug susceptibility, suggesting that the IGF-1R gene should be a promising targeted molecule for HCC therapy.

ACKNOWLEDGEMENTS

All authors thank Prof. Zhizhen Dong for comments on earlier manuscript drafts.

REFERENCES

- 1 **McGlynn KA**, Petrick JL, El-Serag HB. Epidemiology of Hepatocellular Carcinoma. *Hepatology* 2021; **73 Suppl 1**: 4-13 [PMID: 32319693 DOI: 10.1002/hep.31288]
- 2 **Singal AG**, Lampertico P, Nahon P. Epidemiology and surveillance for hepatocellular carcinoma: New trends. *J Hepatol* 2020; **72**: 250-261 [PMID: 31954490 DOI: 10.1016/j.jhep.2019.08.025]
- 3 **Huang DQ**, El-Serag HB, Loomba R. Global epidemiology of NAFLD-related HCC: trends, predictions, risk factors and prevention. *Nat Rev Gastroenterol Hepatol* 2021; **18**: 223-238 [PMID: 33349658 DOI: 10.1038/s41575-020-00381-6]
- 4 **Shi Y**, Wang SY, Yao M, Sai WL, Wu W, Yang JL, Cai Y, Zheng WJ, Yao DF. Chemosensitization of HepG2 cells by suppression of NF- κ B/p65 gene transcription with specific-siRNA. *World J Gastroenterol* 2015; **21**: 12814-12821 [PMID: 26668505 DOI: 10.3748/wjg.v21.i45.12814]
- 5 **Forbes BE**, Blyth AJ, Wit JM. Disorders of IGFs and IGF-1R signaling pathways. *Mol Cell Endocrinol* 2020; **518**: 111035 [PMID: 32941924 DOI: 10.1016/j.mce.2020.111035]
- 6 **Blyth AJ**, Kirk NS, Forbes BE. Understanding IGF-II Action through Insights into Receptor Binding and Activation. *Cells* 2020; **9** [PMID: 33053840 DOI: 10.3390/cells9102276]
- 7 **Chen RS**, Zhang XB, Zhu XT, Wang CS. The crosstalk between IGF-1R and ER- α in the proliferation and anti-inflammation of nucleus pulposus cells. *Eur Rev Med Pharmacol Sci* 2020; **24**: 5886-5894 [PMID: 32572901 DOI: 10.26355/eurrev_202006_21481]
- 8 **Refolo MG**, D'Alessandro R, Lippolis C, Carella N, Cavallini A, Messa C, Carr BI. IGF-1R tyrosine kinase inhibitors and Vitamin K1 enhance the antitumor effects of

Regorafenib in HCC cell lines. *Oncotarget* 2017; **8**: 103465-103476 [PMID: 29262576 DOI: 10.18632/oncotarget.21403]

9 **Han X**, Wang X, Zhao B, Chen G, Sheng Y, Wang W, Teng M. MicroRNA-187 inhibits tumor growth and metastasis via targeting of IGF-1R in hepatocellular carcinoma. *Mol Med Rep* 2017; **16**: 2241-2246 [PMID: 28627639 DOI: 10.3892/mmr.2017.6788]

10 **Wang L**, Yao M, Zheng W, Fang M, Wu M, Sun J, Dong Z, Yao D. Insulin-like Growth Factor I Receptor: A Novel Target for Hepatocellular Carcinoma Gene Therapy. *Mini Rev Med Chem* 2019; **19**: 272-280 [PMID: 30360707 DOI: 10.2174/1389557518666181025151608]

11 **Yan XD**, Yao M, Wang L, Zhang HJ, Yan MJ, Gu X, Shi Y, Chen J, Dong ZZ, Yao DF. Overexpression of insulin-like growth factor-I receptor as a pertinent biomarker for hepatocytes malignant transformation. *World J Gastroenterol* 2013; **19**: 6084-6092 [PMID: 24106410 DOI: 10.3748/wjg.v19.i36.6084]

12 **Kumar A**, Jaitak V. Natural products as multidrug resistance modulators in cancer. *Eur J Med Chem* 2019; **176**: 268-291 [PMID: 31103904 DOI: 10.1016/j.ejmech.2019.05.027]

13 **Aleem E**, Nehrbass D, Klimek F, Mayer D, Bannasch P. Upregulation of the insulin receptor and type I insulin-like growth factor receptor are early events in hepatocarcinogenesis. *Toxicol Pathol* 2011; **39**: 524-543 [PMID: 21411721 DOI: 10.1177/0192623310396905]

14 **Dong Z**, Yao M, Wang L, Yan X, Gu X, Shi Y, Yao N, Qiu L, Wu W, Yao D. Abnormal expression of insulin-like growth factor-I receptor in hepatoma tissue and its inhibition to promote apoptosis of tumor cells. *Tumour Biol* 2013; **34**: 3397-3405 [PMID: 23797814 DOI: 10.1007/s13277-013-0912-y]

15 **Ceballos MP**, Rigalli JP, Céré LI, Semeniuk M, Catania VA, Ruiz ML. ABC Transporters: Regulation and Association with Multidrug Resistance in Hepatocellular Carcinoma and Colorectal Carcinoma. *Curr Med Chem* 2019; **26**: 1224-1250 [PMID: 29303075 DOI: 10.2174/0929867325666180105103637]

16 **Bie CQ**, Liu XY, Cao MR, Huang QY, Tang HJ, Wang M, Cao GL, Yi TZ, Wu SL, Xu WJ, Tang SH. Lentivirus-mediated RNAi knockdown of insulin-like growth factor-1

receptor inhibits the growth and invasion of hepatocellular carcinoma via down-regulating midkine expression. *Oncotarget* 2016; **7**: 79305-79318 [PMID: 27813495 DOI: 10.18632/oncotarget.13027]

17 **Xie DY**, Ren ZG, Zhou J, Fan J, Gao Q. Critical appraisal of Chinese 2017 guideline on the management of hepatocellular carcinoma. *Hepatobiliary Surg Nutr* 2017; **6**: 387-396 [PMID: 29312973 DOI: 10.21037/hbsn.2017.11.01]

18 **Zhang HJ**, Yao DF, Yao M, Huang H, Wu W, Yan MJ, Yan XD, Chen J. Expression characteristics and diagnostic value of annexin A2 in hepatocellular carcinoma. *World J Gastroenterol* 2012; **18**: 5897-5904 [PMID: 23139605 DOI: 10.3748/wjg.v18.i41.5897]

19 **Hua Y**, Wang C, Huang J, Wang K. A simple and efficient method for CRISPR/Cas9-induced mutant screening. *J Genet Genomics* 2017; **44**: 207-213 [PMID: 28416245 DOI: 10.1016/j.jgg.2017.03.005]

20 **Qin Y**, Liu HJ, Li M, Zhai DH, Tang YH, Yang L, Qiao KL, Yang JH, Zhong WL, Zhang Q, Liu YR, Yang G, Sun T, Yang C. Salidroside improves the hypoxic tumor microenvironment and reverses the drug resistance of platinum drugs via HIF-1 α signaling pathway. *EBioMedicine* 2018; **38**: 25-36 [PMID: 30396856 DOI: 10.1016/j.ebiom.2018.10.069]

21 **Xu J**, Bie F, Wang Y, Chen X, Yan T, Du J. Prognostic value of IGF-1R in lung cancer: A PRISMA-compliant meta-analysis. *Medicine (Baltimore)* 2019; **98**: e15467 [PMID: 31083179 DOI: 10.1097/MD.00000000000015467]

22 **Ngo MT**, Jeng HY, Kuo YC, Diony Nanda J, Brahmadhi A, Ling TY, Chang TS, Huang YH. The Role of IGF/IGF-1R Signaling in Hepatocellular Carcinomas: Stemness-Related Properties and Drug Resistance. *Int J Mol Sci* 2021; **22** [PMID: 33669204 DOI: 10.3390/ijms22041931]

23 **Fang M**, Yao M, Yang J, Zheng WJ, Wang L, Yao DF. Abnormal CD44 activation of hepatocytes with nonalcoholic fatty accumulation in rat hepatocarcinogenesis. *World J Gastrointest Oncol* 2020; **12**: 66-76 [PMID: 31966914 DOI: 10.4251/wjgo.v12.i1.66]

24 **Ji Y**, Wang Z, Chen H, Zhang L, Zhuo F, Yang Q. Serum from Chronic Hepatitis B Patients Promotes Growth and Proliferation via the IGF-II/IGF-IR/MEK/ERK

Signaling Pathway in Hepatocellular Carcinoma Cells. *Cell Physiol Biochem* 2018; **47**: 39-53 [PMID: 29763915 DOI: 10.1159/000489744]

25 **Chung W**, Kim M, de la Monte S, Longato L, Carlson R, Slagle BL, Dong X, Wands JR. Activation of signal transduction pathways during hepatic oncogenesis. *Cancer Lett* 2016; **370**: 1-9 [PMID: 26433160 DOI: 10.1016/j.canlet.2015.09.016]

26 **Cheng W**, Huang PC, Chao HM, Jeng YM, Hsu HC, Pan HW, Hwu WL, Lee YM. Glypican-3 induces oncogenicity by preventing IGF-1R degradation, a process that can be blocked by Grb10. *Oncotarget* 2017; **8**: 80429-80442 [PMID: 29113314 DOI: 10.18632/oncotarget.19035]

27 **Yao WF**, Liu JW, Sheng GL, Huang DS. Blockade of IGF-IR exerts anticancer effects in hepatocellular carcinoma. *Mol Med Rep* 2011; **4**: 719-722 [PMID: 21567089 DOI: 10.3892/mmr.2011.486]

28 **Chen S**, Cao Q, Wen W, Wang H. Targeted therapy for hepatocellular carcinoma: Challenges and opportunities. *Cancer Lett* 2019; **460**: 1-9 [PMID: 31207320 DOI: 10.1016/j.canlet.2019.114428]

29 **Wang Y**, Jiang F, Jiao K, Ju L, Liu Q, Li Y, Miao L, Li Z. De-methylation of miR-148a by arsenic trioxide enhances sensitivity to chemotherapy via inhibiting the NF- κ B pathway and CSC like properties. *Exp Cell Res* 2020; **386**: 111739 [PMID: 31759055 DOI: 10.1016/j.yexcr.2019.111739]

30 **Qiu Y**, Dai Y, Zhang C, Yang Y, Jin M, Shan W, Shen J, Lu M, Tang Z, Ju L, Wang Y, Jiao R, Xia Y, Huang G, Yang L, Li Y, Zhang J, Wong VKW, Jiang Z. Arsenic trioxide reverses the chemoresistance in hepatocellular carcinoma: a targeted intervention of 14-3-3 η /NF- κ B feedback loop. *J Exp Clin Cancer Res* 2018; **37**: 321 [PMID: 30572915 DOI: 10.1186/s13046-018-1005-y]

31 **Hwang JW**, Cho H, Lee JY, Jeon Y, Kim SN, Lee SJ, Bae GU, Yoon S, Jeon R, Kim YK. The synthetic ajoene analog SPA3015 induces apoptotic cell death through crosstalk between NF- κ B and PPAR γ in multidrug-resistant cancer cells. *Food Chem Toxicol* 2016; **96**: 35-42 [PMID: 27449564 DOI: 10.1016/j.fct.2016.07.020]

32 **Tovar V**, Cornella H, Moeini A, Vidal S, Hoshida Y, Sia D, Peix J, Cabellos L, Alsinet C, Torrecilla S, Martinez-Quetglas I, Lozano JJ, Desbois-Mouthon C, Solé M, Domingo-Domenech J, Villanueva A, Llovet JM. Tumour initiating cells and IGF/FGF signalling contribute to sorafenib resistance in hepatocellular carcinoma. *Gut* 2017; **66**: 530-540 [PMID: 26658144 DOI: 10.1136/gutjnl-2015-309501]

4%

SIMILARITY INDEX

PRIMARY SOURCES

1	pubmed.ncbi.nlm.nih.gov Internet	65 words — 1%
2	www.researchgate.net Internet	27 words — 1%
3	www.jcancer.org Internet	26 words — 1%
4	bsdwebstorage.blob.core.windows.net Internet	25 words — 1%
5	encyclopedia.pub Internet	25 words — 1%
6	link.springer.com Internet	14 words — < 1%
7	Min Yao, Wenli Sai, Wenjie Zheng, Li Wang, Zhizhen Dong, Dengfu Yao. "Secretory Clusterin as a Novel Molecular-targeted Therapy for Inhibiting Hepatocellular Carcinoma Growth", Current Medicinal Chemistry, 2020 Crossref	13 words — < 1%
8	www.tandfonline.com Internet	13 words — < 1%

EXCLUDE QUOTES	ON	EXCLUDE SOURCES	< 12 WORDS
EXCLUDE BIBLIOGRAPHY	ON	EXCLUDE MATCHES	< 12 WORDS