

78590_Auto_Edited check.doc

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 78590

Manuscript Type: CASE REPORT

Giant bilateral scrotal lipoma with abnormal somatic fat distribution: A case report

Chen Y *et al.* Giant bilateral scrotal lipoma

Yu Chen, Xiu-Ning Li, Xian-Lin Yi, Tang Yong

Abstract

BACKGROUND

Scrotal lipoma is exceedingly rare, so its origin is still unknown. Injury is suggested as a potential factor, but the cause remains unclear. It is difficult to determine the origin of these tumours. Previous studies have suggested that these tumours may be congenital or that they originate from small fatty particles around the cord. Other studies have suggested that these tumours originate from testicles or the tunica.

CASE SUMMARY

A 66-years elderly male with giant scrotal mass. B-ultrasound, computed tomography and magnetic resonance imaging examinations are typical lipoma-like changes, which were confirmed by postoperative pathological analysis. Abnormal somatic fat distribution was confirmed by images. no recurrence after follow-up for 2 years. Previously published English-language literature was reviewed, and a history of inguinal or pelvic surgery was reported in 6 studies (total 21 cases). To our knowledge, this is the heaviest bilateral scrotal lipoma in the English-language literature, and it has the longest postoperative follow-up time. More importantly, the origin of these tumours is reviewed and discussed.

CONCLUSION

Giant bilateral scrotal lipoma in elderly may as part of the symptoms of abnormal somatic fat distribution.

Key Words: Lipoma; Crotum; Bilateral; Extratesticular tumour; Fat distribution; Case report

Yu C, L XN, Y XL, Yong T. Giant bilateral scrotal lipoma with abnormal somatic fat distribution: A case report. *World J Clin Cases* 2022; In press

Core Tip: A case of giant bilateral scrotal lipoma was reported, and the literature was reviewed. A 66-year-old male with a giant scrotal mass was evaluated. Surgical treatment was performed, and scrotal lipoma were confirmed by postoperative pathological analysis. This is the heaviest bilateral scrotal lipoma in the English-language literature, and it has the longest postoperative follow-up time. More importantly, the origin of these tumours is reviewed and discussed.

INTRODUCTION

In 1780, Pelletan described a fatty tumour of the cord^[1]. In 1883, Roswell Park reported a case of intrascrotal lipoma and concurrently reviewed three reported cases^[1,2]. Cecil^[1] reported a case of intrascrotal lipoma in 1927.

Scrotal lipoma is exceedingly rare, so its origin is still unknown^[2,3]. Injury is suggested as a potential factor, but the cause remains unclear. It is difficult to determine the origin of these tumours. Previous studies have suggested that these tumours may be congenital^[4] or that they originate from small fatty particles around the cord. Other studies have suggested that these tumours originate from testicles or the tunica^[2].

These tumours are very large. A 75-cm lipoma from the scrotum to the knee was reported by Kocher^[1]. Regarding weight, Brossard reported a tumour that weighed 9 pounds, and Bonney reported a 20-pound fibrolipoma in 1930^[2]. However, no pure

lipoma was observed^[1]. A study published in 1930 reported a giant scrotal lipoma weighing 480-1100 g^[2,5]. Seidu *et al*^[5] reported a large unilateral scrotal mass that weighed 1100 g in 2020.

Herein, a large bilateral scrotal lipoma with abnormal accumulation of body surface fat was reported. To our knowledge, this is the heaviest bilateral scrotal lipoma in the English-language literature, and it has the longest postoperative follow-up time. More importantly, the origin of these tumours is reviewed and discussed.

CASE PRESENTATION

Chief complaints

Progressive enlargement of bilateral scrotum for more than one year.

History of present illness

progressive enlargement of bilateral scrotum for more than one year. At first, the bilateral scrotal tumours were small (approximately the size of peanuts); then, they became larger. The patients reported slight itching, but no pain or other discomfort. Both sides of his scrotum were obviously enlarged, approximately 13 cm in diameter, and the skin of the scrotum was tight, with no redness or varicose veins.

History of past illness

The patient had undergone electroresection of the prostate and bladder lithotripsy for benign prostatic hyperplasia three years ago in our clinic.

Physical examination

It was failure to touch spermatic cord and testes with negative light transmission test (Figure 1A).

Laboratory examinations

Serum ² beta-human chorionic gonadotropin, alpha-fetoprotein and lactate dehydrogenase levels were normal. Abnormally decreased platelets were shown from the first operation (Table 1). A bone marrow test was performed, and immune thrombocytopenia was diagnosed.

Imaging examinations

Scrotal ultrasonography showed a slightly higher echogenic mass. The computed tomography (CT) scan showed diffuse fat accumulation in the scrotum, and its density was not uniform. The CT value range was approximately -66 to -94 Hu. The testicles on both sides were normal in size and compressed by the scrotal wall (Figure 1B-D). A magnetic resonance imaging (MRI) scan showed bilateral patchy abnormal signals on both scrotum, high signals on T1WI and T2WI scans, a low signal on a T2WI lipid-suppressing sequence, no obvious diffusion-restricted signal changes on DWI and ADC, and no obvious abnormal enhancement on an enhanced scan (Figure 1E and F).

FINAL DIAGNOSIS

Giant bilateral scrotal lipoma with abnormal somatic fat distribution.

TREATMENT

Surgical treatment was performed. The tumour squeezed the bilateral testis, epididymis and spermatic cord tightly against the inner wall of the scrotum, and the boundary between the tumour and the tumour was obvious. The capsules of both tumours were intact, lobulated, light yellow, and soft (Figure 1A-C and 1G,). The cut surface was off-white and slightly tough. The total weight of bilateral scrotal tumours reached 995 g (Figure 1H). Bilateral scrotal lipomas were confirmed by postoperative pathology (Figure 1F). follow-up only postoperative, no radiotherapy and chemotherapy.

OUTCOME AND FOLLOW-UP

The patient was followed up for two years and no recurrence was found.

DISCUSSION

Scrotal lipoma is very rare, and only approximately 20 cases have been reported^[3,6]. In most cases, the aetiology and origination are still unknown. Some patients are young^[7], and the cause is suggested to be congenital^[4]. However, some patients were older than 60 years, as shown in Table 2. An elderly patient with abnormal fat somatic distribution was reported herein, and its origin was discussed in this study.

Florante *et al*^[8] used the term "extratesticular" to describe scrotal tumours that do not directly affect the testes themselves. While 95% of testicular tumours are malignant, most paratesticular tumours, including scrotal lipoma, are benign^[9].

Large scrotal lipomas often present with scrotal enlargement and discomfort^[10-12]. A large right scrotal mass that weighed 1100 g was reported^[5]. Yamamichi *et al*^[13] reported a case of scrotal lipoma with a maximum diameter of 14 cm combined with liposarcoma (weight 250 g)^[13]. The lesion described in the current report is the heaviest bilateral scrotal lipoma in the English-language literature^[10-12], which was 995 g in weight (Figure 1H).

Scrotal lipomas can be divided into three types^[14]: (1) Originating from the posterior spermatic cord and spreading into the scrotum; (2) Originating from the inside or outside of the spermatic cord, and (3) originating from fat lobules of the scrotal dartos tunica. The latter lipomas are called primary scrotal lipomas, and they usually occur in young individuals^[4].

However, abnormal somatic fat distribution may be an important influencing factor in the current case. The CT and MRI results of this patient showed liposis of the lower abdomen, perineum and thigh but no liposis in the chest wall or pelvic cavity (Figure 1 B-F). Excess fat around the penis was shown on MRI (Figure 1 D). A fat test was performed, and only apolipoprotein-A1 was slightly increased (Table 1). Multiple metabolic enzyme abnormalities were also shown, which lasted for three years in this case (Table 1). These results showed that bilateral scrotal lipoma in elderly individuals may be a symptom of abnormal somatic fat distribution caused by metabolic diseases.

Because the scrotal tissue is loose and extensible, which is more suitable for fat accumulation, bilateral scrotal lipoma developed. Recently, a giant bilateral scrotal lipoma along with multiple lipomas was also reported^[3]. Indirect evidence can also be provided by many patients with obesity^[15].

The aetiology of scrotal lipoma is still unclear. In addition to congenital factors^[4], local pluripotent cells develop into adipocyte lines^[15]. Moreover, injury may be a factor when the lesion originates from the spermatic cord or inguinal canal in elderly individuals^[2]. We reviewed the English-language literature, and a history of inguinal or pelvic surgery was reported in 6 patients, including the current patient (21 patients total, Table 2). However, a small mass similar to a peanut was observed before this patient's first operation three years ago. There was one case of contralateral inguinal surgery in the literature^[8]. However, these results suggest that inguinal or pelvic surgery is an accelerating factor, if not an initiating factor.

Blood diseases may also play a role in the tumorigenesis of scrotal lipoma. This current case was diagnosed as primary thrombocytopenia due to decreased platelets for more than 3 years. Szmigielski *et al*^[16] reported a case of a large scrotal lipoma that arose after venous thrombosis and pulmonary embolism. The scrotal mass in the 67-year-old patient was approximately 9 cm × 11 cm after taking oral anticoagulants for 3 mo.

Ultrasonography and MRI play an important role in the evaluation of scrotal lipoma^[11]. Ultrasound is the first choice for an investigation tool. MRI is the most sensitive method to distinguish a benign lesion from a malignant lesion. High T1 signal strength is a characteristic of fat-containing tumours. Nevertheless, it is difficult to distinguish low-grade liposarcoma from benign lipoma based on MRI findings alone^[17]. Complete removal of a tumour by surgery and pathological examination can confirm lipoma^[11]. Lipoma is homogenous, and its CT values are between -50 and -150 Hounsfield units; this information can help in the identification of liposarcoma^[16].

If lipomas are excluded, more than 50% of spermatic cord tumours are malignant^[18]. If the tumour grows rapidly beyond 10 cm, liposarcoma should be considered in the differential diagnosis^[19]. To our knowledge, this is the longest

reported postoperative follow-up time. Yamamichi *et al*^[13] reported a case and follow-up for six months^[13]. Kaplanoglu *et al*^[12] reported a case of intrascrotal lipoma (weight 550 g) with a diameter of 10 cm^[12]. No tumour recurrence was observed after six months of follow-up. The main limitation of this study is that it is a case report, and there are no genomic data.

CONCLUSION

To our knowledge, this is the heaviest bilateral scrotal lipoma and the longest postoperative follow-up time in English-language literature. Bilateral scrotal lipoma in elderly maybe as part of the symptoms of abnormal somatic fat distribution, which caused by metabolic diseases. Inguinal or pelvic surgery is an accelerating factor, if not an initiating factor. Blood system diseases may also play a role in the tumorigenesis.

2%

SIMILARITY INDEX

PRIMARY SOURCES

1	Benjamin J Vowles, Bruce Rodda, David Ringrose. "Evaluation of intubating adjuncts to videolaryngoscopy in a simulated difficult airway: A manikin cross-over study", Anaesthesia and Intensive Care, 2021 <small>Crossref</small>	12 words — 1%
2	cms.galenos.com.tr <small>Internet</small>	12 words — 1%
3	editorialrestauro.com.mx <small>Internet</small>	10 words — 1%

EXCLUDE QUOTES ON
EXCLUDE BIBLIOGRAPHY ON

EXCLUDE SOURCES < 10 WORDS
EXCLUDE MATCHES < 10 WORDS