

78891_Auto_Edited.docx

**Sun-burn induced upper limb lymphedema 11 years following breast cancer surgery:
a case report**

a case report

Min Li, Jun Guo, Rong Zhao, Jinnan Gao, Ming Li, Linying Wang

Abstract

BACKGROUND

Upper arm lymphedema is a common complication one year after breast cancer surgery, which profoundly impacts patients' quality of life.

CASE SUMMARY

We reported a case of lymphedema induced by prolonged sun exposure 11 years after breast cancer surgery.

CONCLUSION

Breast screening, patient education and follow-up after hospital discharge could help to prevent upper-arm lymphedema.

Key Words: Sun-burn; Breast cancer lymphedema; Breast neoplasms; Case report

Li M, Guo J, Zhao R, Gao J, Li M, Wang L. Sun-burn induced upper limb lymphedema 11 years following breast cancer surgery: a case report. *World J Clin Cases* 2022; In press

Core Tip: Upper arm lymphedema is a common complication one year after breast cancer surgery, which profoundly impacts patients' quality of life. We reported a case of lymphedema induced by prolonged sun exposure 11 years after breast cancer surgery.

INTRODUCTION

⁴ Breast cancer is one of the most common malignancies, accounting for 15% of female tumors ^[1]. Since 2000, the incidence in China has been increasing with a 1% annual increment ^[2]. Although survival has been significantly improved due to advances in treatment, the quality of life was affected, sometimes due to lymphedema on the affected side ^[3]. ¹ Lymphedema is the main complication after cancer treatment, affecting 12–28% of the patients after axillary dissection and 3.5–7.5% after sentinel node biopsy ^[4]. Patients have upper limb lymph accumulation due to lymphatic vessel rupture and deformation. Lymphedema causes shoulder joint dysfunction and affects patients' mental well-being ^[5–6]. Research shows that approximately 1/5th of patients in China develop secondary upper limb lymphedema within one year of having breast cancer surgery ^[7]. Although various approaches have been used, lymphedema remains the most common chronic complication of breast cancer treatment, with only palliative therapies available ^[8]. Therefore, prevention against modifiable risk factors has become critical. Risk factors for lymphedema include overweight/obesity, high blood pressure,

axillary lymph node dissection, chemotherapy, local radiotherapy [9], limb overload, trauma and infection [10].

Postoperative infection, delayed healing and improper nursing can hinder lymphatic vessel reconstruction and the establishment of collateral circulation [11]. When using heat therapy, individuals should use common sense and proceed with caution. If there is swelling in the at-risk limb or increased swelling in the lymphedematous limb, stop using heat, such as a hot tub or sauna. Recently, we cured a patient with severe limb lymphedema following solar dermatitis with massive infection, 11 years after right breast cancer surgery. We hereby report this case with the patient's informed consent.

CASE PRESENTATION

Chief complaints

She complained of progressive swelling and heaviness of the right upper limb after a 4-days (Approximately 15 h) of outdoor fieldwork under the direct sun. At that time, the patient wore a short-sleeved shirt with no covering on both upper limbs. She experienced chest wall burning pain on the seventh day and came to seek help on the ninth day. On 9th September 2021, she was admitted with forearm erythema, desquamation, and a body temperature of 39.4 °C. Prior to this, the patient's upper limbs were normal, and his right upper limb was always protected, with no mosquito bites, infections or injuries. When the patient was exposed to the hot sun nine days ago, she mistakenly believed that she had been in surgery for more than 10 years and did not need to protect his upper limb, which resulted in swelling and pain in his right upper limb and sunburn on his left upper limb.

History of present illness

Figure 1 depicts the affected limb on admission. According to the group standard of the Chinese Nursing Society for lymphedema evaluation (T/CNAS05-2020), combined with the symptoms reported voluntarily by patients. Following admission, we performed a Breast Cancer & Lymphedema Symptom Experience Index (BCLE-SEI) evaluation and measured the circumference. The difference in the circumference between the affected and healthy sides was > 3 cm and diagnosed as severely sunburned lymphedema of the right upper limb ^[12]. Supplement Table 1 shows the circumference measurements of both arms.

History of past illness

The right-handed female patient, aged 61, underwent modified radical mastectomy for the right breast cancer (with chemotherapy and radiotherapy of the axilla and chest) in our department 11 years ago. She continued a regular follow-up (The patient was followed up every three months for two years after the operation, every six months for the next 3-5 years and once a year after five years of operation). After breast surgery, there was no complaint of right upper limb discomfort; therefore, arm circumference measurement was not performed regularly.

Personal and family history

Without

Physical examination

Comparison of circumferences of left and right upper limbs

Laboratory examinations

She was also diagnosed with a chest wall infection on the affected side after blood tests

Imaging examinations

without

FINAL DIAGNOSIS

Breast cancer lymphedema

TREATMENT

Based on bacterial culture results, an eight days 400 U penicillin (b.i.d) therapy was initiated along with a standard of daily dressing and close monitoring of the blood circulation of the infected skin. The patient was advised to wear loose dresses to avoid friction. On the ninth day after the infection was cured, a physiotherapist started a lymphatic drainage massage daily for 40 min. The centripetal massage was also performed twice daily for 30 min with a sleeve pressure of < 40 mmHg. After the physiotherapy, the patient was instructed to wear elastic sleeves. The circumference measurements were taken daily at 5:00 pm to monitor the recovery (Supplement Table 2). On the 15th day of admission, the patient was discharged after a comprehensive examination of fully recovered limb and function. Figure 2 shows the limbs on the discharge day. We followed the patient telephonically at two weeks, one month, three months, and six months after discharge, and the patient reported no change in any function or circumference of the right limb. Throughout her hospital stay and follow-up calls, she was encouraged to discuss her negative feelings with the care team and to be actively involved in making decisions about treatment and monitoring and prevention of lymphedema.

OUTCOME AND FOLLOW-UP

The patient got a comprehensive recovery, and there was no complaint of discomfort in the follow-up visit.

DISCUSSION

One of the most common post-surgical complications among breast cancer survivors is lymphedema, which can evolve into irreversible interstitial fibrosis [13]. Therefore, the screening and monitoring of lymphedema are important. The most common clinical manifestations of lymphedema include infection, swelling, pain, discomfort, and reduced joint dexterity^[14]. American National Lymphedema Network declaration indicates that patients should be advised to avoid air traveling, maceration, skin puncture (blood collection), compression (blood pressure measurement), skin infection and extreme temperatures of the affected limb^[15]. The patient, in this case, had lymphedema due to high temperature. The evaluation is primarily based on the symptoms of patients. In the early stages, patients typically present with a subtle onset of arm or hand swelling, accompanied by a sensation of heaviness in the limb. Swelling in the middle and late stages cannot be relieved by raising or applying external pressure to the affected limb. The scope of involvement is broad and has an orange-peel appearance^[16]. The Chinese Nursing Society's group standard for prevention and care of lymphedema after breast cancer surgery (T/CNAS 05-2020) states that all breast cancer patients should be screened for lymphedema risk before being discharged from the hospital after surgery.

Lymphedema risk scales should be used following a breast cancer operation to identify high-risk patients. Furthermore, for high-risk patients, compliance with preventive behaviors should be monitored at least every six months for the first two years after the operation, and whether lymphedema developed or not should be evaluated. All patients did not have typical presentations. People who are at high risk

should accept the objective physical examination, which may include limb measurements, magnetic resonance imaging (MRI) [17], lymphoscintigraphy [18], indocyanine green (ICG) [19], and bioimpedance spectroscopy (BIS). In a retrospective study of over 165,000 hospital admissions for lymphedema, the majority of cases (92%) were associated with cellulitis in the United States between 2012 and 2017 [20, 21]. Research proves that prolonged exposure of affected limbs to ultraviolet radiation stimulates inflammatory mediators to dilate and infiltrate the blood vessel and damages the skin barrier function, leading to local oozing and even infection. This might be due to high protein deposition in the interstitial tissues that accelerates the growth and reproduction of bacteria [22, 23].

The case indicates that patients should be informed to avoid sun exposure shortly or long after breast cancer surgery, especially in regions where people culturally like beaches and outdoor activities. Patients' knowledge, attitude and behavior play a key role in preventing lymphedema in breast cancer patients [24]. The medical care team should disseminate evidence-based knowledge of lymphedema, guide and train patients to avoid risks, and follow up and closely monitor their practice following hospital discharge.

Baseline upper limb circumference is an objective measurement used to monitor the prevention, occurrence and outcome of lymphedema in breast cancer patients. In this case, overall circumference changes before and after the occurrence of lymphedema and after the treatment could not be evaluated, suggesting the need to record complete upper limb measurements for the rehabilitation of breast cancer patients.

Currently, there are no internationally recognized effective treatments for lymphedema [24]. ¹ The conservative treatment of established lymphedema consists of a multimodality regimen that includes general self-care measures (exercise, skin care), compression therapy (compression bandaging, compression garments, intermittent pneumatic compression), and physiotherapy (e.g., simple lymphatic drainage, manual lymphatic drainage, complete decongestive therapy) [25, 26]. In China, upper arm lymphedema

could be significantly prevented if breast cancer screening is promoted in the community, based on the success of western countries for early detection and less radical surgical treatment [27].

CONCLUSION

We reported a case of lymphedema caused by prolonged sun exposure 11 years after breast cancer surgery and radiotherapy. Breast screening, patient education and follow-up after hospital discharge could help to prevent upper-arm lymphedema.

Footnotes

Informed consent statement: The patient has signed the informed consent form and uploaded it before treatment, and the study has passed the review of our hospital's ethics committee.

Conflict-of-interest statement: None

CARE Checklist (2016) statement: Uploaded

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Provenance and peer review: Unsolicited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review started: July 20, 2022

First decision: August 22, 2022

Article in press:

Specialty type: Nursing

Country/Territory of origin: China

Peer-review report's scientific quality classification

Grade A (Excellent): 0

Grade B (Very good): 0

Grade C (Good): C

Grade D (Fair): D

Grade E (Poor): 0

P-Reviewer: Vinh-Hung V, Martinique; Yap RVC, Philippines **S-Editor:** **L-Editor:** **P-Editor:**

Figure Legends




Figure 1. Patients affected right limb on admission (on 9th September 2021)

Supplement Table 1. Comparison of circumferences of left and right upper limbs on admission (cm)

Measurement points

Thumb web

The wrist

10 cm below the elbow

10 cm above the elbow

Left (unaffected)

20.4

17.9

21.3

26.3

Right (affected)

23.2

20.0

27.1

31.2

Difference

2.8

2.1

5.8

4.9

=====

Figure 2. Comparison of the affected limb on discharge

Supplement Table 2. Circumferences of the affected limb during hospital stay (cm)

Hospital stay

Thumb web

The wrist

10 cm under the elbow

10 cm above the elbow

Day 1

23.2

20.0

27.1

31.2

Day 3

22.5

19.2

26.0

29.8

Day 5

22.0

18.9

25.4

29.3

Day 7

21.8

18.5

24.7

29.0

Day 9

21.5

18.4

24.0

28.8

Day 11

21.3

18.4

22.5

28.2

Day 13

21.1

18.2

22.3

28

Day 15

21.1

18.2

22.0

27.1

6%

SIMILARITY INDEX

PRIMARY SOURCES

1	link.springer.com Internet	69 words — 4%
2	Katherine A. Jackson, Joseph L. Feldman, Jane M. Armer. "Lymphedema Risk Reduction and Management", Elsevier BV, 2022 Crossref	16 words — 1%
3	f6publishing.blob.core.windows.net Internet	14 words — 1%
4	www.ncbi.nlm.nih.gov Internet	13 words — 1%